

Posterior & Lateral Compartments of the Leg

Dr. Nisreen Rajeh

Associate Professor of Anatomy, KAU

2026



The leg الساق

- Is the **middle part of the lower limb**, that is the part between the knee and the ankle.



Fascial compartments of the Leg

- Leg deep (crural) fascia is continuous above with the deep fascia of the thigh.
- Attaches to anterior and medial borders of the **tibia**, continuous with its periosteum.
- **Anterior & posterior intermuscular septa** pass from deep surface of deep fascia & attaches to corresponding margins of the fibula.
- Interosseous membrane & fibrous septa divide the leg into **3 compartments**; anterior, posterior (largest) and lateral.

Leg

Cross Section - Fascial Compartments

Anterior

Extensor muscles
 Tibialis anterior
 Extensor digitorum
 Extensor hallucis
 Fibularis (peroneus) tertius
 Anterior tibial artery and
 Deep fibular (peroneal)

Anterior
 intermuscular septum

Lateral compartment

Fibularis
 (peroneus) longus
 Fibularis (peroneus)
 brevis muscle
 Superficial fibular
 (peroneal) nerve

Posterior
 intermuscular septum

Fibula

Crural

Crural fascia

Interosseous

Tibia

Deep posterior compartment

Deep posterior
 compartment
 Flexor digitorum
 Tibialis
 Flexor hallucis
 Popliteus
 Posterior tibial artery and
 Tibial nerve
 Fibular
 (peroneal) artery

Transverse
 intermuscular septum

Superficial posterior


Superficial posterior
 compartment
 Soleus
 Gastrocnemius
 Plantaris

Forms of deep fascia in the leg

1-Interosseous membrane:


-Thin but strong membrane connecting **interosseous borders of tibia and fibula together** & provides attachment for neighboring muscles & has openings permit passage of vessels.

-large opening in its upper part for passage of anterior tibial vessels.



2- Retinacula of the ankle:


- thickened band of deep fascia at ankle region, serve to keep long tendons in position.
- Extensor, flexor & peroneal retinaculum.



3-Plantar aponeurosis:

اللفافة الأخمصية


- Is a **thick, dense central part of plantar deep fascia** of the sole of the foot.
- Protect sole from injury, support foot arches.



Posterior compartment of the leg
=Plantar Flexors compartment (calf)




Nerve	Origin	area of skin supply
Posterior cutaneous nerve of the thigh	Sacral plexus; S1,S2,S3	Popliteal fossa& upper part of back of the leg.
Lateral cutaneous nerve of the calf.	Branch of common peroneal nerve	Upper part of posterolateral surface of the leg.
Sural nerve	Branch of tibial nerve	Lower part of posterolateral surface of the leg.
Saphenous nerve	Branch of Femoral nerve	Posteromedial surface of the leg



Muscles in post.compartment

- ❑ **Deep transverse septum** divides posterior compartment of the **leg** into **large superficial group** and **deep small group** of muscles.
- ❑ Superficial group of muscles is forming the prominence of calf.



Content

❖ Muscles (7): A-superficial group(3).


1-gastrocnemius 2-plantaris
3-Soleus (plantar flexors)

B- deep group(4).

4-Popliteus 5-Flexor
digitorum longus 6- Flexor hallucis
longus 7- Tibialis posterior.

❖ Blood supply: **Posterior Tibial artery.**

❖ Nerve supply: **Tibial nerve**



Muscles of the back of the leg(flexors):

A-Superficial group (larger)

1-Gastrocnemius: is the most superficial.

O : Lateral head: from lateral aspect of lateral Condyle of femur.

Medial head(Larger): from popliteal surface of femur above medial condyle.


I : Large, fleshy bellies, join common tendon tendo-calcaneus (Achilles tendon), attached to post aspect of calcaneum.

N : Tibial nerve.


A : planter **flexion** of foot at ankle joint when knee is extended + **flexion** of knee joint. (acts on knee & ankle).
Raises heel during walking

GASTRONECMIUS

Lateral Head
Medial Head








Posterior Cadaver Leg Muscles

1. Gastrocnemius
(medial head)
2. Gastrocnemius
(lateral head)




2-Soleus


- ❑ Broad, flat muscle, lies **anterior** to gastrocnemius.
- O:** inverted V-shaped origin from **soleal line** on the post. Surface of **tibia**.& from upper $\frac{1}{4}$ of the post.surface of shaft of **fibula**.
- I:** join **common tendon** ;tendo-calcaneus , attached to post aspect of calcaneus.
- N:** **Tibial nerve**
- A:** Planter flexor independent of position of knee, needed in walking & running, (raising the heel off the ground.)




B




C



D



Soleal line



**Tendo-calcaneus is a
common tendon for 2
muscles only;**
-Gastrocnemius
-Soleus




Tendoachilles

Tendon for
plantaris

KEN
HUB

© www.kenhub.com



Plantar flexors of the ankle joint, provide the main powerful propulsive force in walking, and running by using the foot as a lever and raising the heel off the ground.

Basic Anatomy 509





FIGURE 10.66 The foot as a simple lever (A) and as a segmented lever (B). Floor prints of a normal foot and a flat foot are also shown.



Side view of lower leg



Rupture tendo-calcaneus



- common in middle-age men, frequently occurs in **tennis players**.
- rupture occurs at narrowest part
- Sever pain is felt
- Active plantar flexion is impossible.
- tendon should be sutured and the leg immobilized.



3-Plantaris (feeble muscle)

❖ Has a **small fusiform belly** with **long-narrow tendon**. Sometimes double or may be absent.



O: from **lateral supracondylar ridge of femur**

I: the long tendon descends **obliquely** in the interval between gastrocnemius & soleus & then attached to **post. surface of calcaneum** on medial side of tendo-calcaneus.

N: **Tibial nerve**

A: It assists in planter **flexion** of foot at ankle joint + **flexion** of knee joint. (ankle / knee).








The long narrow tendon of plantaris is commonly used in reconstructive surgery of the tendons of the hand. Weak muscle!



- ❖ All the muscles of the superficial calf are producing plantar flexion of ankle joint and flexion of knee joint **EXCEPT soleus** act only on ankle joint.
- ❖ All the muscles of the superficial calf are supplied by **tibial nerve**.
- ❖ All the superficial group of post.calf are **plantar flexors** of ankle joint.

B-Deep group of muscles



❑ 1-Popliteus muscle (flat, triangular muscle)

O : from lateral surface of lateral condyle of femur & by few fibers from *lateral meniscus* (within the capsule of knee joint). **It arises inside capsule of knee joint.**

I : fibers pass downward & medially&


Attached to **posterior surface of tibia above soleal line.**

N : **Tibial nerve**

Action: weak Flexor of the leg at the knee joint (by lateral rotation of the femur on tibia).

(Unlocking the knee joint: initial stage of leg flexion)

تتفرد هذه العضلة بهذه الوظيفة




Locking of the knee joint

- Produced by **quadriceps femoris** muscle & **biceps femoris**.
- It occurs in **erect position** to allow the knee to remain in full extension.

Unlocking of the knee joint


- Produced by **Popliteus** muscle
- It occurs during **initial stage of knee flexion**.
- Occurs as a result of **lateral rotation of the femur on the tibia (if the foot is on the ground)** during the **initial stage of flexion**.



POPLITEUS



2-Tibialis posterior (small, centrally located)



- O** : Posterior surface of shaft of tibia & fibula & interosseous membrane.
- I** : its tendon passes **behind medial malleolus**, deep to (behind) flexor retinaculum, Into the sole of foot, inserted mainly in **tuberosity of navicular bone**. + neighboring bones.
- N** : **Tibial nerve**
- A** : **plantar flexion of the foot** + **Inverts the foot** at subtalar & transverse tarsal joints + **supports medial longitudinal arch** of the foot.



Tibialis Posterior

REHAB
My Patient.com









FIGURE 10.52 Anterior view of the ankles and feet of a 29-year-old woman showing inversion (A) and eversion (B) of the right foot.


Transverse
tarsal joint



Tarso-
metatarsal
joint

Subtalar
joint



Hindfoot Anatomy






3-Flexor digitorum Longus (toes)

- O** : From the medial part of posterior surface of the shaft of **tibia**, below soleal line.(so it is **medial** muscle in the post.leg)
- I** : its tendon passes **behind** **medial malleolus**,  **deep(behind)** flexor retinaculum, and enters **sole** of foot. The main tendon divides into 4 tendons & attached in bases of the distal phalanges of the lateral 4 toes. (Each tendon passes through an opening in the corresponding tendon of flexor digitorum brevis).
- N** : **Tibial nerve**
- A** : 1-Flexion of **distal phalanges of lateral 4 toes**,
2-assists in **plantar flexion of foot**, 3-supports the **medial & lat. longitudinal arches of the foot.**









Drake: Gray's Anatomy for Students, 2nd Edition.
Copyright © 2009 by Churchill Livingstone, an imprint of Elsevier, Inc. All rights reserved.
Figure 6.110 Fibrous digital sheaths.

The *quadratus plantae* is inserted into the main tendon of FDL before its division. *The 4 lumbricals* arise from the 4 tendons of FDL.



Attachments of Muscles of Leg: Posterior View



- **Above soleal line:**

popliteus insertion

- **From soleal line:**

soleus origin

- **Below soleal line:**

flexor digitorum longus origin


3- Flexor hallucis longus (hallucis=big toe)


O: From the lower two-thirds of the posterior surface of the **shaft of fibula**. (so it is a lateral muscle in the post leg) 💡

I: Its tendon passes behind the medial malleolus deep to **(behind)** flexor retinaculum, grooving post. surface of talus, passes forward on the sole of foot. It gives strong slip to the tendon of FDL. It is inserted into the base of the distal phalanx of the big toe (1st toe)

N: Tibial nerve

A: Flexion of distal phalanx of big toe & assists in plantar flexion+ supports **medial longitudinal arch** in the foot.






- 1) **Flexors of the thigh:** Pectineus, iliacus, psoas, sartorius, rectus femoris.
- 2) **Extensors of the thigh:** hamstring portion of adductor magnus. & **gluteus maximus**, long head of biceps femoris, Semimembranosus, Semitendinosus.
- 3) **Lateral rotators of the thigh:** Obturator externus, sartorius, adductor longus, brevis & magnus. , Piriformis, Gemellus Superior , Gemellus inferior, Obturator internus, Quadratus Femoris.
- 4) **Medial rotators of the thigh:** iliacus, psoas. + anterior fibers of gluteus medius.
- 5) **Adductors of the thigh:** Pectineus, adductor longus, adductor brevis, adductor magnus & gracilis.
- 6) **Abductor of the thigh:** Sartorius, Gluteus medius and gluteus minimus
- 7) **Flexors of the knee:** Sartorius , Gracilis, biceps femoris, Semitendinosus , Semimembranosus, . And popliteus (deep flexor) , Gastrocnemius , Plantaris
- 8) **Extensors of the knee:** Quadriceps femoris.
- 9) **Plantar flexors:** Gastrocnemius , Plantaris , Soleus , Tibialis posterior , Flexor digitorum Longus , Flexor hallucis longus . Peroneus longus, Peroneus brevis
- 10) **Invertors of the foot:** Tibialis posterior.
- 11) **Evertors of the foot:** Peroneus longus , Peroneus brevis ,


❖ Arteries of the post. compartment of the leg.

❑ Posterior Tibial Artery:


- ❖ Is one of terminal branches of Popliteal artery.
- ❖ Larger and more direct branch.
- ❖ Accompanying **Tibial nerve** in the back of the leg.
- ❖ It begins : at lower border of popliteus muscle, then passes deep to gastrocnemius, soleus & deep transverse fascia. It lies above on the posterior surface of tibialis posterior muscle and on posterior surface of tibia below.
- ❖ End: in the lower part of leg is covered by skin & fascia, it passes behind medial malleolus deep to flexor retinaculum & ends by dividing into medial and lateral plantar arteries (arteries of the sole).



(i) Posterior Leg



(ii) Anterior Leg



❖ Branches of Posterior tibial artery: مهم

1) **Peroneal (fibular) artery**

- Large artery
- Arises close to the origin of the posterior tibial artery.
- It descends behind the fibula.
- It gives *muscular branches* to lateral compartment of the leg and a *nutrient artery* to *Fibula*.
- It ends by taking part in anastomosis around ankle joint.
- A *perforating branch* pierces the interosseous membrane to reach lower part of front of the leg.


2) Muscular branches to muscles of **posterior compartment** of the leg.

3) Anastomotic branches; share in anastomosis around **ankle joint**.

4) Medial and lateral plantar arteries; terminal branches to **sole of foot**.

5) Nutrient artery to the **tibia**.



Muscles of Leg (Deep Dissection)
Posterior View



Veins:

Venae comitantes of the post.tibial artery join those of ant.tibial artery in the popliteal fossa to form the **popliteal vein**.

III-B







FIGURE 10.43 Deep structures in the posterior aspect of the right leg.

Posterior tibial artery & tibial nerve together on the posterior leg.

□ Venae comitantes of posterior tibial artery:


□ It joins those of the anterior tibial artery in the **popliteal fossa** to form the **popliteal vein**.



Nerves of the back of the leg

□ Tibial nerve

- ❖ **Origin:** the **largest terminal Medial** branch of sciatic nerve in the lower 1/3 of the back of the thigh.
- ❖ **Course:**
- ❖ Arises in the lower third of thigh.
- ❖ It descends downwards in **the popliteal fossa** & descends deep to soleus muscle to enter the posterior aspect of the leg.
- ❖ It lies on the posterior **surface of the tibialis posterior** & lower down the leg, on the **posterior surface of tibia**.




❖ It accompanies **posterior tibial artery** in the back of the leg. They pass together behind the medial malleolus and covered here by **flexor retinaculum** & divides into its 2 terminal branches.


❖ **End:** divides into **medial & lateral plantar nerves** in the medial foot.

Tibial
nerve

Posterior
tibial artery



Medial plantar & lateral plantar arteries are the terminal two branches of posterior tibial artery. Together with **medial and lateral plantar nerves**.



❖ Branches of tibial nerve:

1-Cutaneous: a. Sural nerve:


- It descends between the two heads of gastrocnemius.
- The **sural communication branch** of *common peroneal nerve* usually unites with the nerve.
- It supplies the skin of **the back of the leg and the calf**.
- It runs with the **small saphenous vein** behind the lateral malleolus.
- It supplies the skin **on the lateral border of the foot and little toe**.

b. Medial calcaneal branch: supplies skin over **medial surface of the heel**.

2-Muscular : **Leg:** to all muscles on the back of the leg: gastrocnemius (both heads), plantaris, soleus & popliteus, Tibialis posterior, FHL and FDL.

+**Sole of foot**

3-Articular : to knee and ankle joints.




Region of the ankle:

Clinically, ankle is a common site for fractures, sprains and dislocations.




posterior aspect: Medial side

- **Ankle:** transition zone between leg & foot.
- **Retinacula** : are thickening of the deep fascia that keep long tendons around the ankle in position & act as pulleys.



Attachment of Flexor retinaculum



- Extends from the posterior border of **medial malleolus** to medial border of **calcaneus**.
- It holds& protect tendons which enter the sole **behind the medial malleolus**.




**Structures that pass behind the Medial Malleolus, deep to flexor retinaculum;
from Medial to Lateral;**

- 1- **T**ibialis posterior tendon (T)
- 2- Flexor **d**igitorum longus tendon (d)
- 3- Posterior tibial **a**rtery with vinae- comitants. (v)
- 4- Tibial **n**erve (n)
- 5- Flexor **h**allucis longus. (h)

Tom does very nice hat **تحفظ**
بالترتيب (v=vessels)





Summary

Superficial group:

- 1.gastrocnemeus (knee/ankle)
- 2.Plantaris (knee/ ankle)
- 3.Soleus (ankle)

Deep group:

1. Popliteus (knee):**unlocking**
2. Tibialis posterior (ankle)
3. Flexor digitorum longus (ankle & lat 4 toes)
4. Flexor hallucis longus (ankle & 1st toe)



Tibial nerve

Lateral compartment of the leg = Evertors

Content:

Muscles:

- 1) Peroneus (fibularis) longus.
- 2) Peroneus (fibularis) brevis.

Blood supply: branches from **peroneal artery**.

Nerve supply: *Superficial Peroneal nerve*.



1) Peroneus(fibularis) longus muscle

O : from the upper two-thirds of the lateral surface of the fibula.

I : the tendon runs behind **lateral malleolus** & held in position by *superior peroneal retinaculum*. Then runs on lateral surface of calcaneum, where it held in position by *inferior peroneal retinaculum*, then tendon deviates medially to reach sole of the foot & It is inserted into the medial cuneiform & the base of the first metatarsal.

N : **Superficial Peroneal nerve.**

A : Plantar flexion of the foot(weak) + **E**version of foot at subtalar & transverse tarsal joints+ supports lateral longitudinal & transverse arch of the foot.

A

Common fibular nerve

Interosseous
membrane

Fibularis
longus

Anterior
border of fibula

Fibularis
brevis


B

Groove on
inferior aspect
of cuboid


Medial
cuneiform
Metatarsal I

Fibularis
brevis
tendon

Fibularis
longus
tendon



**Peroneus longus =
fibularis longus**




Drake: Gray's Anatomy for Students, 2nd Edition.
Copyright © 2009 by Churchill Livingstone, an imprint of Elsevier Inc. All rights reserved.

Figure 6.110 Fibrous digital sheaths.

2) Peroneus (fibularis)brevis

- O** : from the lower two-thirds of the lateral surface of fibula.
- I** : The tendon passes **behind & directly** in contact with the **lateral malleolus**, & held in position by *sup.peroneal retinaculum* & *inf.peroneal retinaculum* & inserted into **the tubercle on the base of 5th metatarsal**.
- N** : **Superficial peroneal nerve**.
- A** : **Plantar flexion** of the foot(weak) + **E**version of foot + supports **lateral longitudinal arch**.



**Peroneus brevis tendon,
base of 5th metatarsal**



Tibialis posterior	tuberosity of navicular bone. + neighboring bones.
Flexor digitorum longus	bases of the <u>distal phalanges</u> of the lateral 4 toes.
Flexor hallucis longus	base of the <u>distal phalanx</u> of the big toe (1 st toe)
Peroneus longus	medial cuneiform & the base of the 1 st <u>metatarsal</u> .
Peroneus brevis	base of 5 th <u>metatarsal</u> .

Nerve of the front and lateral compartment of the leg & dorsum of foot.

□ Common peroneal nerve

Origin: -It is the smaller lateral branch of the two terminal branches of sciatic nerve.


-It begins in the **lower third** of the thigh.

Course:

- It descends downwards in the popliteal fossa
- It then passes behind the **head of fibula** and curves laterally around the **neck of fibula**.

End: at the lateral aspect of **neck of fibula** it divides into:

- 1- **Superficial peroneal nerve**; to lat. leg
- 2- **Deep peroneal nerve**; to ant. leg





❑ Branches:

- **Cutaneous:** 1) *Sural communicating branch*
2) The lateral cutaneous nerve of the calf.
- **Muscular** : To short head of *biceps femoris*.
- **Articular** : to the knee joint.

❑ Superficial Peroneal nerve

- ❖ Is one of terminal branches of common peroneal nerve arises in the substance of **peroneus longus** muscle at the level of **neck of fibula**.
- ❖ It descends between **peroneus longus & brevis** and in the lower leg *becomes cutaneous*.
- ❖ **Branches:** 1. **Muscular branches** to *peroneus longus & brevis*. 2. **Cutaneous branches** :medial & lateral branches to skin of lower part of the front of the leg & dorsum of the foot, dorsal surfaces of skin of all toes Except the adjacent sides of the 1st & 2nd toes and the lateral side of little toe.






❑ Arterial supply of lateral compartment of the leg

❑ Peroneal artery

- ❑ Large artery *arises from post tibial artery*, close to its origin.
- ❑ It descends within the substance of FHL.
- ❑ Gives many muscular branches and ends by taking part in anastomosis around ankle joint.



Region of the ankle:

Clinically, ankle is a common site for fractures, sprains and dislocations.





posterior aspect: Lateral side

❑ Structures pass behind the lateral malleolus *superficial* to superior peroneal retinaculum

Sural nerve and small saphenous vein.

❑ Structures pass behind the lateral malleolus *deep* to superior peroneal retinaculum

Peroneus longus & brevis tendons; both tendons share a common synovial sheath: lower down, beneath inferior peroneal retinaculum, they have separate sheaths.




Observe the difference on medial and lateral side?

Tendon Sheaths of Ankle

Lateral View

Sup. peroneal
retinaculum

Inf. peroneal
retinaculum



❖ Remember :

- Tibialis posterior causes **inversion** of foot
- Peroneus longus and brevis causes **Eversion** of foot.

	Blood supply	Nerve supply
1)Anterior compartment of thigh.	Femoral artery	Femoral nerve
2)Medial compartment of the thigh	Profunda femoris artery & obturator artery.	Obturator nerve
3)Gluteal region	Superior gluteal & inferior gluteal arteries	Superior gluteal & inferior gluteal nerves. Nerve to qadratus femoris &nerve to obturator internus
4)Posterior thigh	Profunda femoris artery	Sciatic nerve components
5)posterior leg	Posterior Tibial artery	Tibial nerve
6)Lateral leg	Peroneal artery	Superficial peroneal nerve

Navicular bone	Medial cuneiform & base of 1st metatarsal	Base of the 5th metatarsal
Tibialis posterior	Tibialis anterior	Peroneus tertius
	P. longus	Peroneus brevis



(Formative assessment)

2) Regarding the beginning of popliteal vein, it is normally begins at :

- A. lower part of adductor hiatus.
- B. Lower border of popliteus muscle.
- C. Upper border of popliteus muscle.
- D. Subsartorial (adductor) canal.

2) One of the following muscles **is not** supplied by tibial nerve:

- A. Semitendinosus
- B. Popliteus
- C. Short head of biceps femoris
- D. Biceps femoris.

3) Peroneal artery, is a large artery which is a branch of:

- A. Popliteal artery.
- B. Anterior tibial artery.
- C. Posterior tibial artery.
- D. Femoral artery.

5)Tendocalcaneus at the back of the leg receives the insertion of:

A)Soleus and popliteus muscles.

B)Soleus and plantaris muscles.

C)Soleus and gastrocnemius muscles.

D)Gastrocnemius and plantaris muscles.

6) One of the following muscles everts the foot at the subtalar joint:

A) Soleus

B) Plantaris.

C) Tibialis posterior.

D) Peroneus brevis.

7-Popliteus muscle is supplied by which of the following nerves?

- A. Popliteal
- B. Peroneal
- C. Tibial
- D. Common peroneal

Unlocking of the knee joint is the action of •
which muscle?

A. Plantaris

B. Popliteus

C. Gastrocnemius

D. Biceps femoris



اليقين هو ان تدعو الله بشيء وكل الأسباب حولك
توحي بعدم تحقيقه. ولكن بداخلك يقين تام بأن الله
سيستجيب....