

TEMPORAL TURNOVER OF BEE-FLOWER INTERACTIONS

Imperial College London

Jia Le, Lim

jialelim@hotmail.com

Department of Life Sciences, Silwood Park Campus, Imperial College London



Dynamics of bee-flower interactions

Bee-flower interactions differ drastically between subsequent seasons or even weeks. However, this high interaction turnover has been largely overlooked by previous studies, which assume a static picture of pollination networks. Understanding how and why bee-flower interactions vary over time is crucial in the conservation of bees, and in the continuation of their pollination services. Moreover, climate change can result in fluctuating flowering times, thereafter indirectly sabotaging bee-flower interactions.

We ask...

- Does temperature, precipitation or humidity affect bee-flower interaction turnover?
- Does bee-flower interaction turnover differ between seasons?
- Does bee-flower interaction turnover of the tropics differ from those of the temperate regions?

Data

- *Species level:* More than 200 different species of plants and bees
- *Interactions level:* More than 600 unique interactions and more than 3500 recorded interactions across 39 months

Aim: To investigate the effect of climate on bee-flower interaction turnover.

Calculating turnover

Whittaker's dissimilarity index, β_{int} , reflects the difference between two successive monthly networks.



$$\beta_{int} = \frac{a + b + c}{(2a + b + c)/2} - 1$$

where:

- a is the number of interactions shared between the two networks
- b and c are the number of unique interactions in the two networks

β_{int} ranges from 0 to 1. A higher β_{int} reflects a higher difference between monthly networks.

Preliminary Results



Conclusions and Ongoing work

- Monthly bee-flower interaction turnover is high.
- There are no significant differences between turnover rates within seasons and between seasons.
- There is a cyclic pattern in turnover of bee-flower interactions.

Possible challenges

- Insufficient data; However, β_{int} is rarely affected by small sample sizes.
- If climate does not affect bee-flower interaction turnover, other factors to be considered include bee body size and lifespan.

References:

- CaraDonna, P. J., Petry, W. K., Brennan, R. M., et. al. (2017). Interaction rewiring and the rapid turnover of plant?pollinator networks. *Ecol. Letters*, 20: 385?394. doi:10.1111/ele.12740
- Simone, C.R., Simpson, B., Tidon, R., Neff, J., Pawar, S. (2015). Aseasonal pollinators connect seasonal modules in pollination networks from the Cerrado, a highly diverse seasonally dry neotropical savanna. [Not published]
- Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012). The dissimilarity of species interaction networks. *Ecol. Letters*, 11, 564-575

Acknowledgements: Special thanks to M.C. BoaVentura and S. C. Cappellari for making their data set available for the analyses, as well as Samraat Pawar for his supervision and advice throughout this project.