

Analytical Gradients for the Range-Separated Random Phase Approximation Correlation Energies Using a Lagrangian Framework

Bastien Mussard^a, János G. Ángyán^a, Péter G. Szalay^b

^a CRM², Université de Lorraine, Nancy, France

^b Laboratory of Theoretical Chemistry, Eötvös Loránd University (ELTE), Budapest, Hungary

`bastien.mussard@univ-lorraine.fr`

9 juin 2013

Analytical Gradients for the Range-Separated Random Phase Approximation Correlation Energies Using a Lagrangian Framework

Bastien Mussard^a, János G. Ángyán^a, Péter G. Szalay^b

^a CRM², Université de Lorraine, Nancy, France

^b Laboratory of Theoretical Chemistry, Eötvös Loránd University (ELTE), Budapest, Hungary

bastien.mussard@univ-lorraine.fr

9 juin 2013

Observation


- ▶ DFT : good at short range
- ▶ WF methods : suitable at long range

Observation

- ▶ DFT : good at short range
- ▶ WF methods : suitable at long range

The idea **Mix the two :**

$$E^{\text{RSH}} = \min_{\phi} \left\{ \langle \phi | \hat{T} + \hat{V}_{ne} + \hat{W}_{ee}^{\text{lr}, \mu} | \phi \rangle + E_{\text{Hxc}}^{\text{sr}, \mu}[n_{\phi}] \right\}$$



Observation


- ▶ DFT : good at short range
- ▶ WF methods : suitable at long range

The idea Mix the two :

$$E^{\text{RSH}} = \min_{\phi} \left\{ \langle \phi | \hat{T} + \hat{V}_{ne} + \hat{W}_{ee}^{\text{lr}, \mu} | \phi \rangle + E_{\text{Hxc}}^{\text{sr}, \mu}[n_{\phi}] \right\}$$

where ϕ_0 via a Euler-Lagrange equation

with : $\hat{H}_0 = \hat{T} + \hat{V}_{ne} + \hat{V}_{\text{Hx, HF}}^{\text{lr}} + \hat{V}_{\text{Hxc}}^{\text{sr}}$



Observation


- DFT : good at short range
- WF methods : suitable at long range

The idea Mix the two :

$$E^{\text{RSH}} = \min_{\phi} \left\{ \langle \phi | \hat{T} + \hat{V}_{ne} + \hat{W}_{ee}^{\text{lr}, \mu} | \phi \rangle + E_{\text{Hxc}}^{\text{sr}, \mu}[n_{\phi}] \right\}$$

where ϕ_0 via a Euler-Lagrange equation

with : $\hat{H}_0 = \hat{T} + \hat{V}_{ne} + \hat{V}_{\text{Hx, HF}}^{\text{lr}} + \hat{V}_{\text{Hxc}}^{\text{sr}}$



Long-range correlation

$$\begin{aligned} E &= E^{\text{RSH}} + E_c^{\text{lr}} = \langle \mathbf{d}^{(0)} \mathbf{f}^{\text{lr}} \rangle - \langle \mathbf{d}^{(0)} \mathbf{D} \mathbf{C}^{\text{lr}} \rangle + E_{\text{Hxc}}^{\text{sr}, \mu}[n_{\phi_0}] + E_c^{\text{lr}} \\ &= \langle \mathbf{d}^{(0)} (\mathbf{f}^{\text{lr}} + \mathbf{f}^{\text{sr}}) \rangle - \langle \mathbf{d}^{(0)} \mathbf{D} \mathbf{C} \rangle + E_c^{\text{lr}} \end{aligned}$$

Random Phase Approximation

RPA

- ▶ especially useful for dispersion energies (vdW forces)
- ▶ many different *flavors*, the simplest being :

$$\begin{cases} E_c^{\text{dRPA-I}} &= \langle \mathbf{K} \mathbf{T} \rangle \\ 0 &= 2(\mathbf{K} + \mathbf{K} \mathbf{T} + \mathbf{T} \mathbf{K} + \mathbf{T} \mathbf{K} \mathbf{T}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon) \end{cases}$$


Random Phase Approximation

RPA

- ▶ especially useful for dispersion energies (vdW forces)
- ▶ many different *flavors*, the simplest being :

$$\begin{cases} E_c^{\text{dRPA-I}} = \langle \mathbf{KT} \rangle \\ 0 = 2(\mathbf{K} + \mathbf{KT} + \mathbf{TK} + \mathbf{TKT}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon) \end{cases}$$

A non-variational method




Random Phase Approximation

RPA

- ▶ especially useful for dispersion energies (vdW forces)
- ▶ many different *flavors*, the simplest being :

$$\begin{cases} E_c^{\text{dRPA-I}} = \langle \mathbf{KT} \rangle \\ 0 = 2(\mathbf{K} + \mathbf{KT} + \mathbf{TK} + \mathbf{TKT}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon) \end{cases} = \mathbf{R}(\mathbf{T})$$

A non-variational method



Random Phase Approximation

RPA


- ▶ especially useful for dispersion energies (vdW forces)
- ▶ many different *flavors*, the simplest being :

$$\begin{cases} E_c^{\text{dRPA-I}} = \langle \mathbf{KT} \rangle \\ 0 = 2(\mathbf{K} + \mathbf{KT} + \mathbf{TK} + \mathbf{TKT}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon) \end{cases}$$

$$= \langle \mathbf{IT} \rangle$$

$$= \mathbf{R}(\mathbf{T})$$

A non-variational method



A Simple Gradient : HF

Hellmann-Feynman theorem

- ▶ valid for **variational** wavefunctions

$$\frac{\partial E}{\partial \kappa} = \langle \Psi | \frac{\partial \hat{H}}{\partial \kappa} | \Psi \rangle$$

A Simple Gradient : HF

Hellmann-Feynman theorem

- ▶ valid for **variational** wavefunctions

$$\frac{\partial E}{\partial \kappa} = \langle \Psi | \frac{\partial \hat{H}}{\partial \kappa} | \Psi \rangle$$

HF gradient

$$\frac{\partial E_{HF}}{\partial \kappa} = \langle HF | \frac{\partial \hat{H}}{\partial \kappa} | HF \rangle$$

A Simple Gradient : HF

Hellmann-Feynman theorem

- ▶ valid for **variational** wavefunctions

$$\frac{\partial E}{\partial \kappa} = \langle \Psi | \frac{\partial \hat{H}}{\partial \kappa} | \Psi \rangle$$

HF gradient

$$\hat{H} = \underbrace{S_{\mu\alpha}^{-1} h_{\alpha\beta} S_{\nu\beta}^{-1}}_{\text{blue}} \chi_{\mu}^{\dagger} \chi_{\nu} + \frac{1}{2} \underbrace{S_{\mu\alpha}^{-1} S_{\nu\beta}^{-1}}_{\text{purple}} (\alpha\beta|\gamma\delta) \underbrace{S_{\gamma\sigma}^{-1} S_{\delta\lambda}^{-1}}_{\text{green}} \chi_{\mu}^{\dagger} \chi_{\nu}^{\dagger} \chi_{\sigma} \chi_{\lambda}$$

$$\frac{\partial E_{HF}}{\partial \kappa} = \langle HF | \frac{\partial \hat{H}}{\partial \kappa} | HF \rangle$$

A Simple Gradient : HF

Hellmann-Feynman theorem

- ▶ valid for **variational** wavefunctions

$$\frac{\partial E}{\partial \kappa} = \langle \Psi | \frac{\partial \hat{H}}{\partial \kappa} | \Psi \rangle$$

HF gradient

$$\hat{H} = \underbrace{S_{\mu\alpha}^{-1} h_{\alpha\beta} S_{\nu\beta}^{-1}}_{\text{blue}} \chi_{\mu}^{\dagger} \chi_{\nu} + \frac{1}{2} \underbrace{S_{\mu\alpha}^{-1} S_{\nu\beta}^{-1}}_{\text{green}} (\alpha\beta|\gamma\delta) \underbrace{S_{\gamma\sigma}^{-1} S_{\delta\lambda}^{-1}}_{\text{green}} \chi_{\mu}^{\dagger} \chi_{\nu}^{\dagger} \chi_{\sigma} \chi_{\lambda}$$

$$\begin{aligned} \frac{\partial E_{\text{HF}}}{\partial \kappa} &= \langle \text{HF} | \frac{\partial \hat{H}}{\partial \kappa} | \text{HF} \rangle \\ &= \underbrace{\delta h_{\alpha\beta}}_{\text{blue}} P_{\alpha\beta} + \frac{1}{2} \underbrace{\delta(\mu\lambda|\nu\sigma)}_{\text{green}} (P_{\mu\lambda} P_{\nu\sigma} - P_{\mu\sigma} P_{\nu\lambda}) + \underbrace{\delta S_{\mu\nu} S_{\nu\lambda}^{-1}}_{\text{green}} F_{\lambda\sigma} P_{\sigma\mu} \end{aligned}$$

- ▶ HF gradient *straightforwardly* found by the Hellman-Feynman theorem

A Simple Gradient : HF

Hellmann-Feynman theorem

- ▶ valid for **variational** wavefunctions

$$\frac{\partial E}{\partial \kappa} = \langle \Psi | \frac{\partial \hat{H}}{\partial \kappa} | \Psi \rangle$$


HF gradient

$$\hat{H} = \underbrace{S_{\mu\alpha}^{-1} h_{\alpha\beta} S_{\nu\beta}^{-1}}_{\text{blue}} \chi_{\mu}^{\dagger} \chi_{\nu} + \frac{1}{2} \underbrace{S_{\mu\alpha}^{-1} S_{\nu\beta}^{-1}}_{\text{green}} (\alpha\beta|\gamma\delta) \underbrace{S_{\gamma\sigma}^{-1} S_{\delta\lambda}^{-1}}_{\text{grey}} \chi_{\mu}^{\dagger} \chi_{\nu}^{\dagger} \chi_{\sigma} \chi_{\lambda}$$

$$\begin{aligned} \frac{\partial E_{HF}}{\partial \kappa} &= \langle HF | \frac{\partial \hat{H}}{\partial \kappa} | HF \rangle \\ &= \underbrace{\delta h_{\alpha\beta}}_{\text{blue}} P_{\alpha\beta} + \frac{1}{2} \underbrace{\delta(\mu\lambda|\nu\sigma)}_{\text{green}} (P_{\mu\lambda} P_{\nu\sigma} - P_{\mu\sigma} P_{\nu\lambda}) + \underbrace{\delta S_{\mu\nu} S_{\nu\lambda}^{-1}}_{\text{grey}} F_{\lambda\sigma} P_{\sigma\mu} \end{aligned}$$

- ▶ HF gradient *straightforwardly* found by the Hellman-Feynman theorem

For non-variational wavefunctions... (RPA)



A Simple Gradient : HF

Hellmann-Feynman theorem

- valid for **variational** wavefunctions

$$\frac{\partial E}{\partial \kappa} = \langle \Psi | \frac{\partial \hat{H}}{\partial \kappa} | \Psi \rangle$$


HF gradient

$$\hat{H} = \sum_{\mu\alpha}^{-1} h_{\alpha\beta} S_{\nu\beta}^{-1} \chi_{\mu}^{\dagger} \chi_{\nu} + \frac{1}{2} \sum_{\mu\alpha}^{-1} S_{\nu\beta}^{-1} (\alpha\beta|\gamma\delta) S_{\gamma\sigma}^{-1} S_{\delta\lambda}^{-1} \chi_{\mu}^{\dagger} \chi_{\nu}^{\dagger} \chi_{\sigma} \chi_{\lambda}$$

$$\begin{aligned} \frac{\partial E_{HF}}{\partial \kappa} &= \langle HF | \frac{\partial \hat{H}}{\partial \kappa} | HF \rangle \\ &= \delta h_{\alpha\beta} P_{\alpha\beta} + \frac{1}{2} \delta (\mu\lambda|\nu\sigma) (P_{\mu\lambda} P_{\nu\sigma} - P_{\mu\sigma} P_{\nu\lambda}) + \delta S_{\mu\nu} S_{\nu\lambda}^{-1} F_{\lambda\sigma} P_{\sigma\mu} \end{aligned}$$

- HF gradient *straightforwardly* found by the Hellman-Feynman theorem

For non-variational wavefunctions... (RPA)




Work with an **alternative** object that **is** variational


Remember : for a non-variational method :

Remember : for a non-variational method : energy
 $E(\mathbf{P})$

Lagrangian Framework




Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$



Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$


- ▶ introduce the Lagrangian $\mathcal{L}(\mathbf{P}, \lambda) = E(\mathbf{P}) + \langle \lambda \mathbf{R}(\mathbf{P}) \rangle$ ($= E(\mathbf{P})$)



Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$

- ▶ introduce the Lagrangian $\mathcal{L}(\mathbf{P}, \lambda) = E(\mathbf{P}) + \langle \lambda \mathbf{R}(\mathbf{P}) \rangle$ ($= E(\mathbf{P})$)
- ▶ stationary conditions for \mathcal{L}


Lagrangian Framework



Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$

- ▶ introduce the Lagrangian $\mathcal{L}(\mathbf{P}, \lambda) = E(\mathbf{P}) + \langle \lambda \mathbf{R}(\mathbf{P}) \rangle$ ($= E(\mathbf{P})$)
- ▶ stationary conditions for \mathcal{L} $\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \lambda} = \mathbf{R}(\mathbf{P}) = 0$

Lagrangian Framework




Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$

- ▶ introduce the Lagrangian $\mathcal{L}(\mathbf{P}, \lambda) = E(\mathbf{P}) + \langle \lambda \mathbf{R}(\mathbf{P}) \rangle$ ($= E(\mathbf{P})$)
- ▶ stationary conditions for \mathcal{L} $\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \lambda} = \mathbf{R}(\mathbf{P}) = 0$

$$\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \mathbf{P}} = \frac{\partial E(\mathbf{P})}{\partial \mathbf{P}} + \langle \lambda \frac{\partial \mathbf{R}(\mathbf{P})}{\partial \mathbf{P}} \rangle = 0$$

Lagrangian Framework




Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$

- ▶ introduce the Lagrangian $\mathcal{L}(\mathbf{P}, \lambda) = E(\mathbf{P}) + \langle \lambda \mathbf{R}(\mathbf{P}) \rangle$ ($= E(\mathbf{P})$)
- ▶ stationary conditions for \mathcal{L} $\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \lambda} = \mathbf{R}(\mathbf{P}) = 0$

$$\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \mathbf{P}} = \frac{\partial E(\mathbf{P})}{\partial \mathbf{P}} + \langle \lambda \frac{\partial \mathbf{R}(\mathbf{P})}{\partial \mathbf{P}} \rangle = 0$$

non zero !



Remember : for a non-variational method : energy rules for \mathbf{P}
 $E(\mathbf{P})$ $\mathbf{R}(\mathbf{P}) = 0$

- ▶ introduce the Lagrangian $\mathcal{L}(\mathbf{P}, \lambda) = E(\mathbf{P}) + \langle \lambda \mathbf{R}(\mathbf{P}) \rangle$ ($= E(\mathbf{P})$)
- ▶ stationary conditions for \mathcal{L} $\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \lambda} = \mathbf{R}(\mathbf{P}) = 0$

$$\frac{\partial \mathcal{L}(\mathbf{P}, \lambda)}{\partial \mathbf{P}} = \frac{\partial E(\mathbf{P})}{\partial \mathbf{P}} + \langle \lambda \frac{\partial \mathbf{R}(\mathbf{P})}{\partial \mathbf{P}} \rangle = 0$$

non zero !

RPA case

- ▶ non-variational parameters : $E(\mathbf{T}, \mathbf{C})$
- ▶ three constraints : $\mathbf{R}(\mathbf{T}, \mathbf{C}) = 0$, $(\mathbf{f})_{ai} = 0$, $(\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) = 0$
- ▶ three Lagrangian multipliers : λ , \mathbf{z} and \mathbf{x}

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

Computing λ

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

Computing λ

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$\frac{\partial \mathcal{L}}{\partial \mathbf{T}} = \mathbf{I} + \lambda \frac{\partial \mathbf{R}}{\partial \mathbf{T}} = \mathbf{0}$$

RPA Gradient

Computing λ

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$\frac{\partial \mathcal{L}}{\partial \mathbf{T}} = \mathbf{I} + \lambda \frac{\partial \mathbf{R}}{\partial \mathbf{T}} = \mathbf{0}$$

Computing \mathbf{x} and \mathbf{z}

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{K} \mathbf{M} \rangle + \langle \mathbf{K}' \mathbf{N} \rangle + \langle \mathbf{J} \mathbf{O} \rangle + \underbrace{\langle \epsilon \mathbf{T} \rangle + \langle \mathbf{T} \epsilon \rangle}_{\langle \mathbf{d}^{(2)} \mathbf{f} \rangle} + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

RPA Gradient

Computing λ

$$\frac{\partial \mathcal{L}}{\partial \mathbf{T}} = \mathbf{I} + \lambda \frac{\partial \mathbf{R}}{\partial \mathbf{T}} = \mathbf{0}$$

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$0 = 2(\mathbf{K} + \mathbf{K} \mathbf{T} + \mathbf{T} \mathbf{K} + \mathbf{T} \mathbf{K} \mathbf{T}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon)$$

Computing \mathbf{x} and \mathbf{z}

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{K} \mathbf{M} \rangle + \langle \mathbf{K}' \mathbf{N} \rangle + \langle \mathbf{J} \mathbf{O} \rangle + \underbrace{\langle \epsilon \mathbf{T} \rangle + \langle \mathbf{T} \epsilon \rangle}_{\langle \mathbf{d}^{(2)} \mathbf{f} \rangle} + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

RPA Gradient

Computing λ

$$\frac{\partial \mathcal{L}}{\partial \mathbf{T}} = \mathbf{I} + \lambda \frac{\partial \mathbf{R}}{\partial \mathbf{T}} = \mathbf{0}$$

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$0 = 2(\mathbf{K} + \mathbf{K} \mathbf{T} + \mathbf{T} \mathbf{K} + \mathbf{T} \mathbf{K} \mathbf{T}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon)$$

Computing \mathbf{x} and \mathbf{z}

$$\frac{\partial \mathcal{L}}{\partial \mathbf{C}} = \dots = \mathbf{\Theta} + \tilde{\mathbf{\Theta}}(\mathbf{z}) + 2\mathbf{x} = \mathbf{0}$$

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{K} \mathbf{M} \rangle + \langle \mathbf{K}' \mathbf{N} \rangle + \langle \mathbf{J} \mathbf{O} \rangle + \underbrace{\langle \epsilon \mathbf{T} \rangle + \langle \mathbf{T} \epsilon \rangle}_{\langle \mathbf{d}^{(2)} \mathbf{f} \rangle} + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

RPA Gradient

Computing λ

$$\frac{\partial \mathcal{L}}{\partial \mathbf{T}} = \mathbf{I} + \lambda \frac{\partial \mathbf{R}}{\partial \mathbf{T}} = \mathbf{0}$$

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$0 = 2(\mathbf{K} + \mathbf{K} \mathbf{T} + \mathbf{T} \mathbf{K} + \mathbf{T} \mathbf{K} \mathbf{T}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon)$$

Computing \mathbf{x} and \mathbf{z}

$$\frac{\partial \mathcal{L}}{\partial \mathbf{C}} = \dots = \mathbf{\Theta} + \tilde{\mathbf{\Theta}}(\mathbf{z}) + 2\mathbf{x} = \mathbf{0}$$

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{K} \mathbf{M} \rangle + \langle \mathbf{K}' \mathbf{N} \rangle + \langle \mathbf{J} \mathbf{O} \rangle + \underbrace{\langle \epsilon \mathbf{T} \rangle + \langle \mathbf{T} \epsilon \rangle}_{\langle \mathbf{d}^{(2)} \mathbf{f} \rangle} + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

► yields a set of two equations, solved for \mathbf{z} and \mathbf{x} :

$$\begin{cases} \left(\mathbf{\Theta} - \mathbf{\Theta}^\dagger + \mathbf{f} \mathbf{z} - \mathbf{z} \mathbf{f} + 4\mathbf{g}(\mathbf{z}) + 4\mathbf{W}[\mathbf{z}] \right)_{ai} = 0 \\ (1 + \tau_{pq}) \left(\mathbf{\Theta} + \tilde{\mathbf{\Theta}}(\mathbf{z}) \right)_{pq} = -4(\mathbf{x})_{pq} \end{cases}$$

RPA Gradient

Computing λ

$$\frac{\partial \mathcal{L}}{\partial \mathbf{T}} = \mathbf{I} + \lambda \frac{\partial \mathbf{R}}{\partial \mathbf{T}} = \mathbf{0}$$

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{I} \mathbf{T} \rangle + \langle \lambda \mathbf{R} \rangle + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$0 = 2(\mathbf{K} + \mathbf{K} \mathbf{T} + \mathbf{T} \mathbf{K} + \mathbf{T} \mathbf{K} \mathbf{T}) + (\epsilon \mathbf{T} + \mathbf{T} \epsilon)$$


Computing \mathbf{x} and \mathbf{z}

$$\mathcal{L}(\mathbf{T}, \lambda, \mathbf{C}, \mathbf{z}, \mathbf{x}) = \langle \mathbf{K} \mathbf{M} \rangle + \langle \mathbf{K}' \mathbf{N} \rangle + \underbrace{\langle \mathbf{J} \mathbf{O} \rangle + \langle \epsilon \mathbf{T} \rangle + \langle \mathbf{T} \epsilon \rangle}_{\langle \mathbf{d}^{(2)} \mathbf{f} \rangle} + \langle \mathbf{z} \mathbf{f} \rangle + \langle \mathbf{x} (\mathbf{C}^\dagger \mathbf{S} \mathbf{C} - \mathbf{1}) \rangle$$

$$\frac{\partial \mathcal{L}}{\partial \mathbf{C}} = \dots = \mathbf{\Theta} + \tilde{\mathbf{\Theta}}(\mathbf{z}) + 2\mathbf{x} = \mathbf{0}$$

► yields a set of two equations, solved for \mathbf{z} and \mathbf{x} :

$$\begin{cases} \left(\mathbf{\Theta} - \mathbf{\Theta}^\dagger + \mathbf{f} \mathbf{z} - \mathbf{z} \mathbf{f} + 4\mathbf{g}(\mathbf{z}) + 4\mathbf{W}[\mathbf{z}] \right)_{ai} = 0 \\ (1 + \tau_{pq}) \left(\mathbf{\Theta} + \tilde{\mathbf{\Theta}}(\mathbf{z}) \right)_{pq} = -4(\mathbf{x})_{pq} \end{cases}$$



RPA Gradient

Once multipliers are known gradients obtained from **derivatives of \mathcal{L}**

RPA Gradient

Once multipliers are known gradients obtained from **derivatives of \mathcal{L}**

$$(E_c^{\text{RPA}})^x = \mathcal{L}^x = \langle (\mathbf{z} + \mathbf{d}^{(2)}) \mathbf{f}^x \rangle + \langle \mathbf{M} \mathbf{K}^x \rangle + \langle \mathbf{N} \mathbf{K}'^x \rangle + \langle \mathbf{O} \mathbf{J}^x \rangle + \langle \mathbf{x} \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

RPA Gradient

Once multipliers are known gradients obtained from **derivatives of \mathcal{L}**

$$(E_c^{\text{RPA}})^x = \mathcal{L}^x = \langle (\mathbf{z} + \mathbf{d}^{(2)}) \mathbf{f}^x \rangle + \langle \mathbf{M} \mathbf{K}^x \rangle + \langle \mathbf{N} \mathbf{K}'^x \rangle + \langle \mathbf{O} \mathbf{J}^x \rangle + \langle \mathbf{x} \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

$$(E^{\text{RSH}})^x = \langle \mathbf{d}^{(0)} \mathbf{f}^x \rangle - \langle \mathbf{d}^{(0)} \mathbf{D} \mathbf{C}^x \rangle - 2 \langle \epsilon \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

RPA Gradient

Once multipliers are known gradients obtained from **derivatives of \mathcal{L}**

$$(E_c^{\text{RPA}})^x = \mathcal{L}^x = \langle (\mathbf{z} + \mathbf{d}^{(2)}) \mathbf{f}^x \rangle + \langle \mathbf{M} \mathbf{K}^x \rangle + \langle \mathbf{N} \mathbf{K}'^x \rangle + \langle \mathbf{O} \mathbf{J}^x \rangle + \langle \mathbf{x} \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

$$(E^{\text{RSH}})^x = \langle \mathbf{d}^{(0)} \mathbf{f}^x \rangle - \langle \mathbf{d}^{(0)} \mathbf{D} \mathbf{C}^x \rangle - 2 \langle \epsilon \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

Careful derivation of $(\mathbf{f}^x - \mathbf{D} \mathbf{C}^x)$

Once multipliers are known gradients obtained from **derivatives of \mathcal{L}**

$$(E_c^{\text{RPA}})^x = \mathcal{L}^x = \langle (\mathbf{z} + \mathbf{d}^{(2)}) \mathbf{f}^x \rangle + \langle \mathbf{M} \mathbf{K}^x \rangle + \langle \mathbf{N} \mathbf{K}'^x \rangle + \langle \mathbf{O} \mathbf{J}^x \rangle + \langle \mathbf{x} \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

$$(E^{\text{RSH}})^x = \langle \mathbf{d}^{(0)} \mathbf{f}^x \rangle - \langle \mathbf{d}^{(0)} \mathbf{D} \mathbf{C}^x \rangle - 2 \langle \epsilon \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

Careful derivation of $(\mathbf{f}^x - \mathbf{D} \mathbf{C}^x)$

- ▶ long-range part

$$(\mathbf{d}^{(0)} + \mathbf{d}^{(2)} + \mathbf{z}) \mathbf{h}^x + \left(\frac{1}{2} \mathbf{d}^{(0)} + \mathbf{d}^{(2)} + \mathbf{z} \right) \mathbf{d}^{(0)}_{\text{int}} \text{LR}^x$$

Once multipliers are known gradients obtained from **derivatives of \mathcal{L}**

$$(E_c^{\text{RPA}})^x = \mathcal{L}^x = \langle (\mathbf{z} + \mathbf{d}^{(2)}) \mathbf{f}^x \rangle + \langle \mathbf{M} \mathbf{K}^x \rangle + \langle \mathbf{N} \mathbf{K}'^x \rangle + \langle \mathbf{O} \mathbf{J}^x \rangle + \langle \mathbf{x} \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

$$(E^{\text{RSH}})^x = \langle \mathbf{d}^{(0)} \mathbf{f}^x \rangle - \langle \mathbf{d}^{(0)} \mathbf{D} \mathbf{C}^x \rangle - 2 \langle \epsilon \mathbf{C}^\dagger \mathbf{S}^x \mathbf{C} \rangle$$

Careful derivation of $(\mathbf{f}^x - \mathbf{D} \mathbf{C}^x)$

- ▶ long-range part

$$(\mathbf{d}^{(0)} + \mathbf{d}^{(2)} + \mathbf{z}) \mathbf{h}^x + \left(\frac{1}{2} \mathbf{d}^{(0)} + \mathbf{d}^{(2)} + \mathbf{z} \right) \mathbf{d}^{(0)} \text{int}^{\text{LR}x}$$

- ▶ short-range part is actually $E_{\text{Hxc}}^{\text{SR}}[\tilde{\rho}]^x$:


$$E_{\text{Hxc}}^{\text{SR}}[\tilde{\rho}]^x = E_{\text{Hxc}}^{\text{SR}}[\rho^{(0)}]^x + \int \frac{\partial E_{\text{Hxc}}^{\text{SR}}[\rho^{(0)}]}{\partial \rho^{(0)}} (\tilde{\rho} - \rho^{(0)})$$

Preliminary Results

Test the implementation

- ▶ perfect agreement with numerical calculations
- ▶ timings are as expected (same behavior as MP2)

Geometry Optimization



Conclusion & Outlook

The **Lagrangian framework** has successfully been applied to derive **RPA gradients**

Outlook

- ▶ Implementation has been done in Molpro
(useful parallel with **MP2 gradients**)
- ▶ Gradients of *mixed* RPA energies need further coding
(e.g. Szabo-Ostlund variant)
- ▶ Geometry optimization seem to work
- ▶ Extension to **density fitting** seems straightforward