	. The probability of a leap year selected at random contain 53					
Su	nday is:					
70501	` '	, ,	(c) 2/7	` '		
	_			marble is drawn at		
rar		•	wing a black ba			
200	` '	` '	(c) $0/5$	` '		
		1. ¹⁷		0.85. What is the		
pro	•	t will not rain t				
	` '	` '	, ,	(d) none of these		
				cted from the numbers		
(1,		5) is a multiple		(1) 1 (0		
_			(c) 2/15			
5.				w three coins?		
_			(c) 8			
	•	•	e number selec	ted at random from the		
nu	mbers (1,2,3,		() 1010	- (1)		
_	` '	, ,	• • •	5 (d) none of these		
1.			of an event and			
•	` ') 0 (d) no			
		• •	are given; choo	ose the correct answer		
tot	that which is	-	() 7 (5	(1)		
_				(d) none of these.		
			nultaneously, tl	han the probability of		
_	tting at least t	•	1 > 1	(1) 4 (0)		
	` '	` '	(c) ½	` '		
				ters of the word		
•				e letter chosen has:		
	(a) 6/13	(b) //13	(c) 1	(d) none of these.		
	A 11	E' 1.1				
				etting an even number.		
(A)	2/3	(B) 1	(C) 5/6	(D) 1/2		
12. Two coins are thrown at the same time. Find the probability of						
	tting both hea			ш. р. олишту от		
_	3/4 (B) 1/4		(D) 0			
()	(-) .,	(-) ., _	(-)			
13	. Two dice are	thrown simult	aneously. The	probability of getting a		

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	1/9
14-17 H100 000 1100 000 000 000	rds are numbe rime number.	red from 1 to	100. Find the	e probability of
(A) 3/4	(B) 27/50	(C) 1/4	(D) 29/100
of drawing				If the probability en the number of
(A) 5	(B) 10	(C) 15	(D) 20	
taken out a	of 600 bulbs co at random from tive bulb is:			
(A) 143/15	60 (<mark>B) 147</mark>	<mark>7/150</mark> (C) 1/25	(D) 1/50
mixed thou the probab (A) 9/100	ility that the nu (B) 1/10	ard is drawn fr umber on card (C) 3/10	om this box is a perfect (D) 19/1	randomly, then square. 00
18. What i (A) 1/7	s the probability (B) 53/366	The second secon		n a leap year? 7/366
probability	l is drawn from of getting a ki (B) 3/26 (ng of red suit.		2 cards. Find the
equally like 1,2,312	ely to come to then the prob (B) 1/12	rest pointing t ability that it v	o one of the	number an odd number is:
its outcom result i.e. t probability	e each time. A	ryan wins if al three tails and I lose the gam	I the tosses I loses other e.	times and noting give the same wise. Then the

2

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:						
He fine a section of the same in the second	(B) 31/365	The second secon	(D) 1/133225			
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that x^2 < 2 is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$						
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is $2/3$, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
Then the proba		multiple of 3 and) natural numbers. 4 is:			
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$			
			es are 50, 70, 82,			
93, and 20. The a) 25.79		c) 25.29	d) 25.69			
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15			d) 13, 16			
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $\frac{1}{2}$	b) $\frac{1}{3}$	c) $^1\!/_4$ d 3. The value of I	d) $\frac{1}{6}$			
) 27 d)	9			
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?						

3

a) 3	b) 4	c) 5	d) 7
•			

32.Out of the following values, which one is not possible in probability?

a)
$$P(x) = 1$$

b)
$$\sum x P(x) = 3$$

c)
$$P(x) = 0.5$$

a)
$$P(x) = 1$$

c) $P(x) = 0.5$
b) $\sum x P(x) = 3$
d) $P(x) = -0.5$

33.If E(x) = 2 and E(z) = 4, then E(z - x) = ?

d) Insufficient data

34. The covariance of two independent random variable is

d) Undefined

35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is?

d) Insufficient data

36.If P(x) = 0.5 and x = 4, then E(x) = ?

37.In a discrete probability distribution, the sum of all probabilities is always?

d) Undefined

38.If the probability of hitting the target is 0.4, find mean and variance.

39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance?

a) 0.6, 0.24

b) 6, 2.4 c) 0.4, 0.16

d) 4, 1.6

40. Find the mean of tossing 8 coins.

a) 2

b) 4

c) 8

d) 1

41. What is the mean and variance for standard normal distribution?

The second secon		s 1 b) Mean is 1 and varian s ∞ d) Mean is ∞ and varia	
42.Variance a) E(X)		ariable X is given by c) E(X2) - (E(X))2	· d) (E(X))2
43.Mean of a) E(X)		ble X is given by c) E(X2) - (E(X))2	d) (E(X))2
44.Mean of	a constant 'a' is	S	

a) 0 b) a c) a/2 d) 1


45. Variance of a constant 'a' is _____ .
a) 0 b) a c) a/2 d) 1

46. Find the mean and variance of X?

Х	0	1	2	3	4
f(x)	1/9	2/9	3/9	2/9	1/9

a) 2, 4/3 b) 3, 4/3 c) 2, 2/3 d) 3, 2/3

47. Find the expectation of a random variable X?



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

CS CamScanner

5

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard Deviation?

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}