




Seep Pahuja
Biology Educator
(NEET UG)

 seep.pahuja

 Seep Pahuja Biology - Official Unacademy



LAST 10 YEARS

NEET PYQ

Chapter 6 : Anatomy of Flowering Plants

1. In the seeds of cereals, the outer covering of endosperm separates the embryo by a protein-rich layer called
(1) Coleoptile [NEET-2025]
(2) Coleorhiza
(3) Integument
(4) Aleurone layer
2. Find the statement that is NOT correct with regard to the structure of monocot stem. [NEET-2025]
(1) Hypodermis is parenchymatous.
(2) Vascular bundles are scattered.
(3) Vascular bundles are conjoint and closed.
(4) Phloem parenchyma is absent.
3. Which of the following simple tissues are commonly found in the fruit walls of nuts and pulp of pear? [Re-NEET-2024]
(1) Sclereids
(2) Fibres
(3) Parenchyma
(4) Collenchyma
4. Given Given below are two statements: [Re-NEET-2024]
Statement I: In a dicotyledonous leaf, the adaxial epidermis generally bears more stomata than the abaxial epidermis.
Statement II: In a dicotyledonous leaf, the adaxially placed palisade parenchyma is made up of elongated cells, which are arranged vertically and parallel to each other.
In the light of the above statements, choose the correct answer from the options given below:
(1) Statement I is true but Statement II is false
(2) Statement I is false but Statement II is true
(3) Both Statement I and Statement II are true
(4) Both Statement I and Statement II are false

5. Formation of interfascicular cambium from fully developed parenchyma cells is an example for **[NEET-2024]**

- (1) Differentiation
- (2) Redifferentiation
- (3) Dedifferentiation
- (4) Maturation

6. Given below are two statements: **[NEET-2024]**

Statement I : Parenchyma is living but collenchyma is dead tissue.

Statement II : Gymnosperms lack xylem vessels but presence of xylem vessels is the characteristic of angiosperms.

In the light of the above statements, choose the correct answer from the options given below:

- (1) Both Statement I and Statement II are true
- (2) Both Statement I and Statement II are false
- (3) Statement I is true but Statement II is false
- (4) Statement I is false but Statement II is true

7. Bulliform cells are responsible for **[NEET-2024]**

- (1) Inward curling of leaves in monocots.
- (2) Protecting the plant from salt stress.
- (3) Increased photosynthesis in monocots.
- (4) Providing large spaces for storage of sugars.

8. The transverse section of a plant part showed polyarch, radial and exarch xylem, with endodermis and pericycle. The plant part is identified as : **[NEET-2023-MANIPUR]**

- (1) Dicot stem
- (2) Monocot stem
- (3) Monocot root
- (4) Dicot root

9. Consider the following plant tissues: **[NEET-2023-MANIPUR]**

- (A) Axillary buds
- (B) Fascicular vascular cambium
- (C) Interfascicular cambium
- (D) Cork cambium
- (E) Intercalary meristem

Identify the lateral meristems among the above.

- | | |
|--------------------------------|--------------------------------|
| (1) (A), (B), (C) and (E) only | (2) (A), (B), (D) and (E) only |
| (3) (A), (C) and (D) only | (4) (B), (C) and (D) only |

10. Given below are two statements :

[NEET-2023]

Statement I : Endarch and exarch are the terms often used for describing the position of secondary xylem in the plant body.

Statement II : Exarch condition is the most common feature of the root system.

In the light of the above statements, choose the correct answer from the options given below:

- (1) Both Statement I and Statement II are true
- (2) Both Statement I and Statement II are false
- (3) Statement I is correct but Statement II is false
- (4) Statement I is incorrect but Statement II is true

11. The type of tissue commonly found in the fruit wall of nuts is :

[NEET-2022]

- (1) Parenchyma
- (2) Collenchyma
- (3) Sclerenchyma
- (4) Sclereid

12. Match List-I with List-II.

[NEET-2021]

List -I

List - II

- | | |
|---|--------------------------|
| (a) Cells with active cell division capacity | (i) Vascular tissues |
| (b) Tissue having all cells similar in structure and function | (ii) Meristematic tissue |
| (c) Tissue having different types of cells | (iii) Sclereids |
| (d) Dead cells with highly thickened walls and narrow lumen | (iv) Simple tissue |

(a) (b) (c) (d)

- (1) (ii) (iv) (i) (iii)
- (2) (iv) (iii) (ii) (i)
- (3) (i) (ii) (iii) (iv)
- (4) (iii) (ii) (iv) (i)

13. Match List -I with List - II.

[NEET-2021]

List -I

List - II

- | | |
|----------------------|-------------------------|
| (a) Lenticels | (i) Phellogen |
| (b) Cork cambium | (ii) Suberin deposition |
| (c) Secondary cortex | (iii) Exchange of gases |
| (d) Cork | (iv) Phelloderm |

(a) (b) (c) (d)

- (1) (iv) (i) (iii) (ii)
- (2) (iii) (i) (iv) (ii)
- (3) (ii) (iii) (iv) (i)
- (4) (iv) (ii) (i) (iii)

14. Select the correct pair. [NEET-2021]
- (1) Large colorless empty cells in the epidermis of grass leaves – Subsidiary cells
 - (2) In dicot leaves, vascular bundles are surrounded by large thick-walled cells – Conjunctive tissue
 - (3) Cells of medullary rays that form part of cambial ring – Interfascicular cambium
 - (4) Loose parenchyma cells rupturing the epidermis and forming a lens- shaped opening in bark – Spongy parenchyma
15. Identify the incorrect statement. [NEET-2020]
- (1) Due to deposition of tannins, resins, oils etc., heart wood is dark in colour
 - (2) Heart wood does not conduct water but gives mechanical support
 - (3) Sapwood is involved in conduction of water and minerals from root to leaf
 - (4) Sapwood is the innermost secondary xylem and is lighter in colour
16. Large, empty colourless cells of the adaxial epidermis along the veins of grass leaves are [NEET-2020(COVID-19)]
- (1) Lenticels
 - (2) Guard cells
 - (3) Bundle sheath cells
 - (4) Bulliform cells
17. Grass leaves curl inwards during very dry weather. Select the most appropriate reason from the following : [NEET-2019]
- (1) Closure of stomata
 - (2) Flaccidity of bulliform cells
 - (3) Shrinkage of air spaces in spongy mesophyll
 - (4) Tyloses in vessels
18. Phloem in gymnosperms lacks : [NEET-2019]
- (1) Albuminous cells and sieve cells
 - (2) Sieve tubes only
 - (3) Companion cells only
 - (4) Both sieve tubes and companion cells
19. Regeneration of damaged growing grass following grazing is largely due to :- [NEET-2019(Odisha)]
- (1) Lateral meristem
 - (2) Apical meristem
 - (3) Intercalary meristem
 - (4) Secondary meristem
20. Casparian strips occur in [NEET-2018]
- (1) Epidermis
 - (2) Pericycle
 - (3) Cortex
 - (4) Endodermis

21. Secondary xylem and phloem in dicot stem are produced by [NEET-2018]
(1) Apical meristems (2) Vascular cambium
(3) Phellogen (4) Axillary meristems
22. Stomata in grass leaf are [NEET-2018]
(1) Dumb-bell shaped (2) Kidney shaped
(3) Rectangular (4) Barrel shaped
23. The vascular cambium normally gives rise to [NEET-2017]
(1) Primary phloem (2) Secondary xylem
(3) Periderm (4) Phelloderm
24. Which of the following is made up of dead cells? [NEET-2017]
(1) Collenchyma (2) Phellem
(3) Phloem (4) Xylem parenchyma
25. Specialised epidermal cells surrounding the guard cells are called :- [NEET-I 2016]
(1) Complementary cells (2) Subsidiary cells
(3) Bulliform cells (4) Lenticels
26. Cortex is the region found between :- [NEET-II 2016]
(1) Endodermis and pith
(2) Endodermis and vascular bundle
(3) Epidermis and stele
(4) Pericycle and endodermis
27. The balloon-shaped structures called tyloses :- [NEET-II 2016]
(1) Are extensions of xylem parenchyma cells into vessels
(2) Are linked to the ascent of sap through xylem vessels
(3) Originate in the lumen of vessels
(4) Characterize the sapwood

ANSWER KEY

1. (4)
2. (1)
3. (1)
4. (2)
5. (3)
6. (4)
7. (1)
8. (3)
9. (4)
- 10.(4)
- 11.(4)
- 12.(1)
- 13.(2)
- 14.(3)
- 15.(4)
- 16.(4)
- 17.(2)
- 18.(4)
- 19.(3)
- 20.(4)
- 21.(2)
- 22.(1)
- 23.(2)
- 24.(2)
- 25.(2)
- 26.(3)
- 27.(1)