

Quiz 1

[Back to Week 1](#)


8/10 points
earned (80%)

Quiz passed!



1 / 1
points

1.

Consider the data set given below

```
1 x <- c(0.18, -1.54, 0.42, 0.95)
```

And weights given by

```
1 w <- c(2, 1, 3, 1)
```

Give the value of μ that minimizes the least squares equation

$$\sum_{i=1}^n w_i (x_i - \mu)^2$$



0.0025



0.300



0.1471



Correct Response

```
1 sum(x * w)/sum(w)
```

```
1 ## [1] 0.1471
```

☐ 1.077



0 / 1
points

2.

Consider the following data set

```
1 x <- c(0.8, 0.47, 0.51, 0.73, 0.36, 0.58, 0.57, 0.85, 0.44, 0.42)
2 y <- c(1.39, 0.72, 1.55, 0.48, 1.19, -1.59, 1.23, -0.65, 1.49, 0.05)
```

Fit the regression through the origin and get the slope treating y

as the outcome and x as the regressor. (Hint, do not center the data since we want regression through the origin, not through the means of the data.)



-1.713



Incorrect Response



0.8263



0.59915



-0.04462



1 / 1
points

3.

Do `data(mtcars)` from the `datasets` package and fit the regression

model with `mpg` as the outcome and `weight` as the predictor. Give

the slope coefficient.

☐ 30.2851

☐ -9.559

☒ -5.344

Correct Response

```
1 data(mtcars)
2 summary(lm(mpg ~ wt, data = mtcars))
```

```
1 ##
2 ## Call:
3 ## lm(formula = mpg ~ wt, data = mtcars)
4 ##
5 ## Residuals:
6 ##      Min       1Q   Median       3Q      Max
7 ## -4.543 -2.365 -0.125  1.410  6.873
8 ##
9 ## Coefficients:
10 ##              Estimate Std. Error t value Pr(>|t|)
11 ## (Intercept)   37.285      1.878   19.86  < 2e-16 ***
12 ## wt           -5.344      0.559   -9.56  1.3e-10 ***
13 ## ---
14 ## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
15 ##
16 ## Residual standard error: 3.05 on 30 degrees of freedom
17 ## Multiple R-squared:  0.753, Adjusted R-squared:  0.745
18 ## F-statistic: 91.4 on 1 and 30 DF, p-value: 1.29e-10
```

```
1 attach(mtcars)
2 cor(mpg, wt) * sd(mpg)/sd(wt)
```

```
1 ## [1] -5.344
```

```
1 detach(mtcars)
```

☐ 0.5591



0 / 1
points

Consider data with an outcome (Y) and a predictor (X). The standard deviation of the predictor is one half that of the outcome. The correlation between the two variables is .5. What value would the slope coefficient for the regression model with Y as the outcome and X as the predictor?

☒ 0.25

Incorrect Response

☐ 3

☐ 4

☐ 1



1 / 1
points

5.

Students were given two hard tests and scores were normalized to have empirical mean 0 and variance 1. The correlation between the scores on the two tests was 0.4. What would be the expected score on Quiz 2 for a student who had a normalized score of 1.5 on Quiz 1?

☐ 0.16

☐ 1.0

☒ 0.6

Correct Response

This is the classic regression to the mean problem. We are expecting the score to get multiplied by 0.4. So

```
1 1.5 * 0.4
```

```
1 ## [1] 0.6
```



0.4

1 / 1
points

6.

Consider the data given by the following

```
1 x <- c(8.58, 10.46, 9.01, 9.64, 8.86)
```

What is the value of the first measurement if x were normalized (to have mean 0 and variance 1)?



9.31



-0.9719

**Correct Response**

```
1 ((x - mean(x))/sd(x))[1]
```

```
1 ## [1] -0.9719
```



8.58



8.86

1 / 1
points

7.

Consider the following data set (used above as well). What is the intercept for fitting the model with x as the predictor and y as the outcome?

```
1 x <- c(0.8, 0.47, 0.51, 0.73, 0.36, 0.58, 0.57, 0.85, 0.44, 0.42)
2 y <- c(1.39, 0.72, 1.55, 0.48, 1.19, -1.59, 1.23, -0.65, 1.49, 0.05)
```

☐ 2.105

☒ 1.567



Correct Response

```
1 coef(lm(y ~ x))[1]
```

```
1 ## (Intercept)
2 ##          1.567
```

☐ -1.713

☐ 1.252



1 / 1
points

8.

You know that both the predictor and response have mean 0. What can be said about the intercept when you fit a linear regression?

☒ It must be identically 0.



Correct Response

The intercept estimate is $\bar{Y} - \beta_1 \bar{X}$ and so will be zero.

☐ It must be exactly one.

☐ Nothing about the intercept can be said from the information given.

☐ It is undefined as you have to divide by zero.



1 / 1
points

9.

Consider the data given by

```
1 x <- c(0.8, 0.47, 0.51, 0.73, 0.36, 0.58, 0.57, 0.85, 0.44, 0.42)
```

What value minimizes the sum of the squared distances between these points and itself?

- ☐ 0.8
- ☐ 0.44
- ☒ 0.573

Correct Response

This is the least squares estimate, which works out to be the mean in this case.

```
1 mean(x)
```

```
1 ## [1] 0.573
```

- ☐ 0.36



1 / 1
points

10.

Let the slope having fit Y as the outcome and X as the predictor be denoted as β_1 . Let the slope from fitting X as the outcome and Y as the predictor be denoted as γ_1 . Suppose that you divide β_1 by γ_1 ; in other words consider β_1/γ_1 . What is this ratio always equal to?

- ☐ 1
- ☒ $Var(Y)/Var(X)$

Correct Response

The $\beta_1 = Cor(Y, X)SD(Y)/SD(X)$ and $\gamma_1 = Cor(Y, X)SD(X)/SD(Y)$.

Thus the ratio is then $Var(Y)/Var(X)$.

☐ $2SD(Y)/SD(X)$

☐ $Cor(Y, X)$

