

# Company Overview

Yulu is India's leading micro-mobility service provider, which offers unique vehicles for the daily commute. Starting off as a mission to eliminate traffic congestion in India, Yulu provides the safest commute solution through a user-friendly mobile app to enable shared, solo and sustainable commuting. Yulu zones are located at all the appropriate locations (including metro stations, bus stands, office spaces, residential areas, corporate offices, etc) to make those first and last miles smooth, affordable, and convenient!

## Problem Statement

Yulu has recently suffered considerable dips in its revenues. They have contracted a consulting company to understand the factors on which the demand for these shared electric cycles depends. Specifically, they want to understand the factors affecting the demand for these shared electric cycles in the Indian market.

## Solution Approach

- Data Exploration
  - UVA, BVA, MVA
- Hypothesis Test
  - Z Test / T Test
  - Chi square Test
  - Anova

## Detailed Breakdown

- Establishing a relation between the dependent and independent variable (Dependent "Count" & Independent: Workingday, Weather, Season etc)
- Select an appropriate test to check whether:
  - Working Day has effect on number of electric cycles rented?
  - No. of cycles rented similar or different in different seasons?
  - No. of cycles rented similar or different in different weather?
  - Weather is dependent on season?
  - Holiday has effect on number of electric cycles rented?
  - No. of cycles rented similar or different in different temperature, windspeed, humidity and actual temperature levels?
- Hypothesis Test Framework
  - Set up Null Hypothesis ( $H_0$ )
  - State the alternate hypothesis ( $H_1$ )

- Check assumptions of the test (Normality, Equal Variance).
- Check using Histogram, Q-Q plot, statistical methods like levene's test, Shapiro-wilk test
- Set a significance level (alpha)
- Calculate test Statistics and P value
- Decision to accept or reject null hypothesis.
- Inference from the analysis

```
In [197... import pandas as pd
import numpy as np
import matplotlib.pyplot as plt
import seaborn as sns

from scipy.stats import norm, ttest_rel, ttest_ind, kstest, chi2, chi2_contingen
from itertools import combinations
```

```
In [10]: # data = pd.read_csv(r'F:\Muthu_2023\Personal\NextStep\DS Course\Scaler\Business-
data = pd.read_csv(r'E:\Nextstep\Scaler\Business-Case-Study\Yulu\Dataset\bike_sh
```

## EDA

```
In [4]: data.head()
```

```
Out[4]:
```

	datetime	season	holiday	workingday	weather	temp	atemp	humidity	windspee
--	----------	--------	---------	------------	---------	------	-------	----------	----------

0	2011-01-01 00:00:00	1	0	0	1	9.84	14.395	81	0
---	---------------------	---	---	---	---	------	--------	----	---

1	2011-01-01 01:00:00	1	0	0	1	9.02	13.635	80	0
---	---------------------	---	---	---	---	------	--------	----	---

2	2011-01-01 02:00:00	1	0	0	1	9.02	13.635	80	0
---	---------------------	---	---	---	---	------	--------	----	---

3	2011-01-01 03:00:00	1	0	0	1	9.84	14.395	75	0
---	---------------------	---	---	---	---	------	--------	----	---

4	2011-01-01 04:00:00	1	0	0	1	9.84	14.395	75	0
---	---------------------	---	---	---	---	------	--------	----	---



### Inference:

- Total: 12 Columns
- Target Variables: 'casual', 'registered', 'count'

```
In [5]: data.info()
```

```
<class 'pandas.core.frame.DataFrame'>
RangeIndex: 10886 entries, 0 to 10885
Data columns (total 12 columns):
#   Column      Non-Null Count  Dtype
---  -
0   datetime    10886 non-null  object
1   season      10886 non-null  int64
2   holiday     10886 non-null  int64
3   workingday  10886 non-null  int64
4   weather     10886 non-null  int64
5   temp        10886 non-null  float64
6   atemp       10886 non-null  float64
7   humidity    10886 non-null  int64
8   windspeed   10886 non-null  float64
9   casual      10886 non-null  int64
10  registered  10886 non-null  int64
11  count       10886 non-null  int64
dtypes: float64(3), int64(8), object(1)
memory usage: 1020.7+ KB
```

### Inference:

- No null values in all the columns
- datetime column is not in datetime64 format, hence conversion required
- All are numerical columns, some may be binary (holiday, working day, weather, etc.,)

```
In [11]: for i in ['season', 'holiday', 'workingday', 'weather']:
        print(i, ': ', data[i].unique())
```

```
season : [1 2 3 4]
holiday : [0 1]
workingday : [0 1]
weather : [1 2 3 4]
```

### \*Inference:\*

- Holiday and Working day are binary columns
- Season and Weather are categorical with 4 categories

```
In [12]: data.describe()
```

```
Out[12]:
```

	season	holiday	workingday	weather	temp	ater
count	10886.000000	10886.000000	10886.000000	10886.000000	10886.000000	10886.000000
mean	2.506614	0.028569	0.680875	1.418427	20.23086	23.6550
std	1.116174	0.166599	0.466159	0.633839	7.79159	8.4746
min	1.000000	0.000000	0.000000	1.000000	0.82000	0.7600
25%	2.000000	0.000000	0.000000	1.000000	13.94000	16.6650
50%	3.000000	0.000000	1.000000	1.000000	20.50000	24.2400
75%	4.000000	0.000000	1.000000	2.000000	26.24000	31.0600
max	4.000000	1.000000	1.000000	4.000000	41.00000	45.4550

```
In [15]: data['datetime'].min(), data['datetime'].max()
```

```
Out[15]: ('2011-01-01 00:00:00', '2012-12-19 23:00:00')
```

**\*Inference:** Dataset contains 2 years of data

## Preprocessing

```
In [12]: data['date'] = pd.to_datetime(data['datetime']).dt.date
data['time'] = pd.to_datetime(data['datetime']).dt.time
data['day'] = pd.to_datetime(data['datetime']).dt.day_name()
data['year'] = pd.to_datetime(data['date']).dt.year
data['hour'] = pd.to_datetime(data['datetime']).dt.hour
```


## UVA

```
In [9]: print('Total no. of days: ', data['date'].nunique())
```

Total no. of days: 456

```
In [17]: (pd.to_datetime(data['date']).dt.day_name().value_counts()).plot(kind='bar')
```


```
Out[17]: <Axes: >
```



**\*Inference:**

- Above plot doesn't give any insight as it is a time series data from '2011-01-01 00:00:00', '2012-12-19 23:00:00' and it is recorded every 1 hour
- The univariate analysis on the given dataset gives the details about the conditions of the environment doesn't provides much insights to increase revenue


```
In [70]: plt.pie(data['weather'].value_counts(), labels = list(data['weather'].unique()),
```



#### **\*Inference:\***

- Weather 1 and 2 are predominant throughout the years
- Considering the strategies involving weather 1 and 2 will potentially reflect on revenue

```
In [119... plt.figure(figsize=(12,6))
plt.subplot(1,4,1)
sns.boxplot(data=data, y = 'temp')
plt.subplot(1,4,2)
sns.boxplot(data=data, y = 'atemp')
plt.subplot(1,4,3)
sns.boxplot(data=data, y = 'humidity')
plt.subplot(1,4,4)
sns.boxplot(data=data, y = 'windspeed')
plt.tight_layout()
```



**\*Inference:\***


- No IQR outliers in temperature, absolute temperature and humidity columns
- IQR outlier is detected for windspeed indicating the values > 30 rarely occurs
- Range: (Excluding outliers)
  - $0 < \text{Temp} < 40$
  - $0 < \text{aTemp} < 45$
  - $10 < \text{Humidity} < 100$
  - $0 < \text{windspeed} < 30$

In [172...

```
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.histplot(data['casual'], kde=True)
plt.subplot(1,3,2)
sns.histplot(data['registered'], kde=True)
plt.subplot(1,3,3)
sns.histplot(data['count'], kde=True)
```

Out[172...

<Axes: xlabel='count', ylabel='Count'>




**\*Inference:\***

- Distribution of usage by casual users, registered users, overall users are all right skewed

## BVA

```
In [29]: #Date Vs Count
data.groupby('day')[['casual', 'registered', 'count']].mean().plot(kind='bar')
sns.barplot(data=data, x = 'day', y='casual', estimator='mean', hue='registered')
```

```
Out[29]: <Axes: xlabel='day'>
```




### \*Inference:\*

- Casual users are comparatively very much higher on weekends and lesser on weekdays
- Registered users are less during weekends
- Registered users are predominantly office goers or students
- Due to this behavior, the total count approximately remains constant throughout the days except Sunday
- It confirms the outside activity of the users are less on Sunday
- Prediction of Casual users during weekends is very much required to optimize the demand and supply

```
In [79]: data.groupby('year')[['casual', 'registered']].mean().plot(kind='bar', stacked=True)
```

```
Out[79]: <Axes: xlabel='year'>
```



```
In [101... # YoY percentage increase
df_grp = data.groupby('year')[['casual', 'registered', 'count']].sum().pct_change()
print('YoY increase of casual users: ', round(df_grp['casual'].iloc[1]), '%')
print('YoY increase of registered users: ', round(df_grp['registered'].iloc[1]), '%')
print('YoY increase of overall users: ', round(df_grp['count'].iloc[1]), '%')
```

```
YoY increase of casual users:  52 %
YoY increase of registered users:  70 %
YoY increase of overall users:  67 %
```


Inference:

- Average number of casual and registered users are increased in the year 2012 compared to 2011
- The percentage increase of users is measured to be 52, 70 and 67% for casual, registered and overall users respectively

```
In [163... plt.figure(figsize=(12,6))
data.groupby('hour')[['casual', 'registered', 'count']].mean().plot(kind = 'bar')
plt.show()
```

```
<Figure size 2000x1600 with 0 Axes>
```






**\*Inference:\***

- Bell curve pattern is observed for casual users usage with time
- Average number of Casual users peaks between 13 - 17 hrs
- Double bell curve pattern is observed for registered users with time
- Registered users peak at 7-8 hrs at morning and 17-18 hrs at evening

```
In [62]: # Season Vs Count
ax = data.groupby(['year', 'season'])[['casual', 'registered', 'count']].sum().p
data.groupby(['year', 'season'])[['registered']].sum().plot(kind='line', ax=ax,
```

```
Out[62]: <Axes: xlabel='year,season'>
```



**\*Inference:\***


- Analysing the 2 years of data separately, both years show
  - Much higher trend in Fall followed by Summer and Winter
  - Spring shows very much lesser trend
  - YoY increasing trend for Registered users is noticed which is responsible for the YoY increase in the count
  - Concentrating on increasing the registered users would help to increase the revenue

```
In [35]: data.groupby('season')['date'].nunique()
```

```
Out[35]: season
1      114
2      114
3      114
4      114
Name: date, dtype: int64
```

```
In [60]: # Holiday vs Count
data.groupby(['holiday'])[['casual', 'registered']].mean().plot(kind='barh', sta
```

```
Out[60]: <Axes: ylabel='holiday'>
```




**\*Inference:\***

- On Holidays, average number of casual users > average number of registered users

```
In [59]: # Working day vs Count
data.groupby(['workingday'])[['casual', 'registered']].mean().plot(kind='barh',
```

```
Out[59]: <Axes: ylabel='workingday'>
```




### \*Inference:\*

- On Working days, average number of registered users > average number of casual users

```
In [76]: # Weather vs Count
# Weather corresponds to time not day
plt.figure(figsize=(12,6))
ax = plt.subplot(1,2,1)
data.groupby(['year', 'weather'])[['casual', 'registered', 'count']].mean().plot
plt.title('Usage based on weather (Year wise)')
ax = plt.subplot(1,2,2)
data.groupby(['weather'])[['casual', 'registered', 'count']].mean().plot(kind='b
plt.title('Usage based on weather')
```

Out[76]: Text(0.5, 1.0, 'Usage based on weather')



### \*Inference:\*

- The usage of bikes based on weather is in the order 1 > 2 > 4 > 3
- Interestingly the usage of bikes by registered users is high during Heavy rain climate than light rain, further analysis required
- The average number of usage increases between years 2011 and 2012

```
In [63]: data.groupby(['year', 'weather'])['date'].nunique()
```


```
Out[63]: year  weather
2011    1         212
        2         166
        3          90
2012    1         222
        2         180
        3          97
        4           1
Name: date, dtype: int64
```

**\*Inference:\***

- Strategies including weather 1 and 2 makes significant impact in the revenue than weather 3
- In the dataset, Weather 4 is used in only one record, the insight due to weather 4 doesn't make significant impact
  - The insight "the average usage of bikes by registered users is high during Heavy rain climate than light rain" cannot be concluded

```
In [107... # Temp vs casual and registered
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.scatterplot(data=data, x = 'temp', y='casual')
plt.subplot(1,3,2)
sns.scatterplot(data=data, x = 'temp', y='registered')
plt.subplot(1,3,3)
sns.scatterplot(data=data, x = 'temp', y='count')
```

```
Out[107... <Axes: xlabel='temp', ylabel='count'>
```




**\*Inference:\***

- There is an increasing trend for the rise in temperature for casual users
- No significant pattern is observed when there is rise in temperature for registered users
- Categorizing the temperature data to different level might so show some correlation (further analysis is required)

```
In [114... # aTemp vs casual and registered
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.scatterplot(data=data, x= 'atemp', y='casual')
plt.subplot(1,3,2)
sns.scatterplot(data=data, x = 'atemp', y='registered')
plt.subplot(1,3,3)
sns.scatterplot(data=data, x = 'atemp', y='count')
```

Out[114... <Axes: xlabel='atemp', ylabel='count'>




**\*Inference:\***

- There is an increasing trend for the rise in actual temperature for casual users
- No significant pattern is observed when there is rise in actual temperature for registered users
- Categorizing the actual temperature data to different level might so show some correlation (further analysis is required)

```
In [115... # Temp vs casual and registered
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.scatterplot(data=data, x = 'humidity', y='casual')
plt.subplot(1,3,2)
sns.scatterplot(data=data, x = 'humidity', y='registered')
plt.subplot(1,3,3)
sns.scatterplot(data=data, x = 'humidity', y='count')
```

Out[115... <Axes: xlabel='humidity', ylabel='count'>




**\*Inference:\***

- There is a slight decreasing trend for the rise in humidity level for casual users
- No significant pattern is observed when there is a rise in humidity level for registered users
- Categorizing the humidity level data to different levels might show some correlation (further analysis is required)

```
In [116... # Temp vs casual and registered
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.scatterplot(data=data, x = 'windspeed', y='casual')
plt.subplot(1,3,2)
sns.scatterplot(data=data, x = 'windspeed', y='registered')
plt.subplot(1,3,3)
sns.scatterplot(data=data, x = 'windspeed', y='count')
```

```
Out[116... <Axes: xlabel='windspeed', ylabel='count'>
```




**\*Inference:\***

- No significant pattern is observed when there is a change in wind speed for casual and registered users
- Categorizing the wind speed data to different levels might show some correlation (further analysis is required)

```
In [124... # Weather vs Temp
sns.kdeplot(data = data, x = 'season', y = 'temp')
```

```
Out[124... <Axes: xlabel='season', ylabel='temp'>
```






## MVA

```
In [126... # Usage, Weather,
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.scatterplot(data=data, x = 'temp', y='casual', hue = 'weather')
plt.subplot(1,3,2)
sns.scatterplot(data=data, x = 'temp', y='registered', hue = 'weather')
plt.subplot(1,3,3)
sns.scatterplot(data=data, x = 'temp', y='count', hue = 'weather')
```


```
Out[126... <Axes: xlabel='temp', ylabel='count'>
```






```
In [125... # Usage, Weather,
plt.figure(figsize=(12,6))
plt.subplot(1,3,1)
sns.scatterplot(data=data, x = 'temp', y='casual', hue = 'season')
plt.subplot(1,3,2)
sns.scatterplot(data=data, x = 'temp', y='registered', hue = 'season')
plt.subplot(1,3,3)
sns.scatterplot(data=data, x = 'temp', y='count', hue = 'season')
```

Out[125... <Axes: xlabel='temp', ylabel='count'>



```
In [133... plt.figure(figsize=(12,6))
plt.subplot(2,2,1)
plt.hist(data['temp'], bins = 10)
plt.subplot(2,2,2)
plt.hist(data['atemp'], bins = 10)
plt.subplot(2,2,3)
plt.hist(data['humidity'], bins = 10)
plt.subplot(2,2,4)
plt.hist(data['windspeed'], bins = 10)
```

Out[133... (array([1.313e+03, 4.083e+03, 2.827e+03, 1.540e+03, 6.960e+02, 2.800e+02,  
1.070e+02, 3.100e+01, 6.000e+00, 3.000e+00]),  
array([ 0. , 5.69969, 11.39938, 17.09907, 22.79876, 28.49845,  
34.19814, 39.89783, 45.59752, 51.29721, 56.9969 ]),  
<BarContainer object of 10 artists>)



## Numerical to Categorical Features


- Temp:
  - Low: <15
  - Med: 15 - 30
  - High: >30
- aTemp:
  - Low: <15
  - Med: 15 - 30
  - High: >30
- Humidity:
  - Low: < 40
  - Med: 40 - 85
  - High: >85
- windspeed:
  - Low: 0 - 10
  - Med: 11 - 25
  - High: >25

```
In [13]: bins = [0, 15, 30, 50]
label = ['Low', 'Med', 'High']
data['temp_cat'] = pd.cut(data['temp'], bins=bins, labels=label)
data['temp_cat'].value_counts()
```

```
Out[13]: Med      6249
Low       3393
High      1244
Name: temp_cat, dtype: int64
```

```
In [136... data.groupby('temp_cat')[['casual', 'registered', 'count']].mean().plot(kind='ba
```

```
Out[136... <Axes: xlabel='temp_cat'>
```



**\*Inference:\***


- Average number of casual and registered users increases with an increase in temperature
- It is evident that there is a linear relationship between the temperature and average number of users

```
In [14]: bins = [0, 15, 30, 50]
label = ['Low', 'Med', 'High']
data['atemp_cat'] = pd.cut(data['atemp'], bins=bins, labels=label)
data['atemp_cat'].value_counts()
```

```
Out[14]: Med      5674
High     3250
Low      1962
Name: atemp_cat, dtype: int64
```

```
In [140... data.groupby('atemp_cat')[['casual', 'registered', 'count']].mean().plot(kind='b
```

```
Out[140... <Axes: xlabel='atemp_cat'>
```



**\*Inference:\***


- Average number of both casual and registered users increases with an increase in absolute temperature
- It is evident that there is a linear relationship between the absolute temperature and average number of users

```
In [15]: bins = [0, 40, 85, 100]
label = ['Low', 'Med', 'High']
data['humidity_cat'] = pd.cut(data['humidity'], bins=bins, labels=label)
data['humidity_cat'].value_counts()
```

```
Out[15]: Med      7715
Low       1616
High      1533
Name: humidity_cat, dtype: int64
```

```
In [141...] data.groupby('humidity_cat')[['casual', 'registered', 'count']].mean().plot(kind
```

```
Out[141...] <Axes: xlabel='humidity_cat'>
```



**\*Inference:\***


- Average number of both casual and registered users increases with the decrease in humidity level
- It is evident that the average number of users is negatively correlated with humidity

```
In [234...] bins = [0, 10, 25, 100]
label = ['Low', 'Med', 'High']
data['windspeed_cat'] = pd.cut(data['windspeed'], bins=bins, labels=label)
data['windspeed_cat'].value_counts()
```

```
Out[234...] Med      5698
Low       3026
High        849
Name: windspeed_cat, dtype: int64
```

```
In [142...] data.groupby('windspeed_cat')[['casual', 'registered', 'count']].mean().plot(kin
```

```
Out[142...] <Axes: xlabel='windspeed_cat'>
```




**\*Inference:\***

- Average number of both casual and registered users slightly increases from low to medium windspeed
- No significant change in the average number of users between medium and high windspeed conditions

```
In [261... plt.figure(figsize=(12,6))
temp_data = data[['season', 'holiday', 'workingday', 'weather', 'temp', 'atemp',
sns.heatmap(temp_data.corr(), annot=True)
```

```
Out[261... <Axes: >
```



Inference:

- Average number of casual and registered users are highly positively correlated with temp, atemp
- Humidity and weather are positively correlated
- Humidity and windspeed are negatively correlated
- Average number of causal users is negatively correlated with working day

# Hypothesis Tests

## 1. Problem Statement

Working Day has effect on number of electric cycles rented

**\*Solution Approach:\***

- Null Hypothesis:  $\mu_1 = \mu_2$
- Alternate Hypothesis:  $\mu_1 > \mu_2$ 
  - $\mu_1$  - Average no. of cycles rented during working day
  - $\mu_2$  - Average no. of cycles rented during non working day
- Significance level: 5%
- Comparison between Average no. of cycles rented (*Numerical*) and working day (*Category with 2 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
- Hence, **2 Sample T Test**

In [194...

```
def check_normality(samples):
    for i in range(len(samples)):
        stat, p_value = shapiro(samples[i])
```

```

if p_value < 0.05:
    print("Reject Null Hypothesis. Pval is ", p_value, ". Hence, Sample"
else:
    print("Fail to Reject Null Hypothesis. Pval is ", round(p_value,2) ,

```

```

In [195... def plot_dist(samples):
    plt.figure(figsize=(12,6))
    for i in range(len(samples)):
        plt.subplot(1, len(samples), i+1)
        sns.kdeplot(samples[i])
        plt.title('Sample ' + str(i+1))

```

```

In [ ]: df = data.groupby('date').agg({'workingday': 'min', 'count': 'mean'})
sample1 = df[df['workingday'] == 1]['count']
sample2 = df[df['workingday'] != 1]['count']

```

## 1.1 Check For Assumptions

```

In [26]: check_normality([sample1, sample2])


```

Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution  
 Fail to Reject Null Hypothesis. Pval is 0.08 . Hence, Sample 2 follows normal distribution

```

In [60]: plot_dist([sample1, sample2])

```



### \*Inference:\*

- Average no. of cycles rented during non-working day follows normal distribution
- Average no. of cycles rented during working day doesn't follow normal distribution
- Distribution plot confirms the above point

## 1.2. Perform T-Test

```

In [181... alpha = 0.05
print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
tstat, p_value = ttest_ind(sample1, sample2, alternative='greater')

```



```
print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
if p_value < 0.05:
    print('Reject Null Hypothesis. Hence, Average number of cycles rented during
else:
    print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented
```

Sample-1 Mean: 192.28 Sample-2 Mean: 188.33

T-Stat: 0.51 P-Val: 0.31

Fail to Reject Null Hypothesis. Hence, Average number of cycles rented during working day is equal to non working day

Inference:

- Since the test is between a numerical and categorical variable (with 2 categories), **2 sample t-test is selected**
- Alternate hypothesis is chosen as  $u_1 > u_2$  instead of  $u_1 < u_2$
- Fail to Reject Null Hypothesis. Hence, **Average number of cycles rented during working day is equal to non working day**

## 2. Problem Statement

No. of cycles rented similar or different in different seasons

**\*Solution Approach:\***

- Null Hypothesis: **Average no. of cycles rented are equal for all seasons**
- Alternate Hypothesis: **Average no. of cycles rented is different for atleast one season**
- Comparison between Average no. of cycles rented (*Numerical*) and Seasons (*Category with 4 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
  - **Shapiro Test**
- **Variance Test**
  - Check for Homogeneity of variances
  - **Levene Test**
- **One Way ANOVA Test**
- Significance level: 5%


```
In [40]: count_season = []
for i in range(1, data['season'].nunique()+1):
    count_season.append(list(data[data['season'] == i]['count']))
```

### 2.1. Check for Assumptions

```
In [44]: check_normality(count_season)
```

Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution  
 Reject Null Hypothesis. Pval is 6.039093315091269e-39 . Hence, Sample 2 doesn't follow normal distribution  
 Reject Null Hypothesis. Pval is 1.043458045587339e-36 . Hence, Sample 3 doesn't follow normal distribution  
 Reject Null Hypothesis. Pval is 1.1301682309549298e-39 . Hence, Sample 4 doesn't follow normal distribution

In [58]: `plot_dist(count_season)`



```
In [67]: def Check_Variiances(samples):
          stat, p_value = levene(samples[0], samples[1], samples[2], samples[3])
          if p_value < 0.05:
              print("Reject Null Hypothesis. Pval is ", p_value ,". Hence, the variance is significantly different")
          else:
              print("Fail to Reject Null Hypothesis. Pval is ", round(p_value,2) ,". Hence, the variance is not significantly different")
```

In [68]: `Check_Variiances(count_season)`

Reject Null Hypothesis. Pval is 1.0147116860043298e-118 . Hence, the variance of at least one sample is significantly different

#### **\*Inference:**

- For Anova, the assumptions are failed. All the samples doesn't follow normal distribution and homogeneity of variances
- Perform Anova and also Kruskal wallis test

## 2.2. Perform One Way Anova Test

```
In [86]: # One Way Anova
          alpha = 0.05
          stat, p_value = f_oneway(count_season[0], count_season[1], count_season[2], count_season[3])
          #print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2), 'Sample-3 Mean: ', round(sample3.mean(),2), 'Sample-4 Mean: ', round(sample4.mean(),2))
          print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
          if p_value < alpha:
              print('Reject Null Hypothesis. Hence, Average no. of cycles rented is different for at least one sample')
```

```

else:
    print('Fail to Reject Null Hypothesis. Hence, Average no. of cycles rented is

```

Test Statistic: 236.95 P-Val: 6.164843386499654e-149

Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast one season

**\*Inference:\***

- One way Anova test concluded that **Average number of cycles rented is significantly different for atleast one season**
- In order to find the seasons where the Average number of cycles rented is significantly different we need to perform 2 sample t test

## 2.3. Perform Kruskal Wallis Test

```

In [73]: # Kruskal Wallis Test
alpha = 0.05
stat, p_value = kruskal(count_season[0], count_season[1], count_season[2], count
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(samp
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Median no. of cycles rented is differe
else:
    print('Fail to Reject Null Hypothesis. Hence, Median no. of cycles rented is

```

Test Statistic: 699.67 P-Val: 2.479008372608633e-151

Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one season

**\*Inference:\***

- Since the assumptions of one way anova test is not met, the results can not concluded directly from that test
- Kruskal wallis test, concludes that the **Median no. of cycles rented is different for atleast one season**

## 2.4. Perform 2 Sample T Test between dependent and each independent variable

```

In [97]: alpha = 0.05
for sample in count_season:
    print('Average no. of cycles rented in season ', str(i), ': ', np.mean(samp

for idx1, idx2 in list(combinations(np.arange(len(count_season)),2)):
    tstat, p_value = ttest_ind(count_season[idx1], count_season[idx2], alternati
    print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, Average number of cycles rented is
    else:
        print('Fail to Reject Null Hypothesis. Hence, Average number of cycles r

```

Average no. of cycles rented in season 4 : 116.34326135517499  
 Average no. of cycles rented in season 4 : 215.25137211855105  
 Average no. of cycles rented in season 4 : 234.417124039517  
 Average no. of cycles rented in season 4 : 198.98829553767374  
 T-Stat: -22.42 P-Val: 1.6578587340400098e-106  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 0 and 1  
 T-Stat: -26.26 P-Val: 3.4038504355310974e-143  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 0 and 2  
 T-Stat: -19.76 P-Val: 5.236417429066781e-84  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 0 and 3  
 T-Stat: -3.64 P-Val: 0.00027431561172498644  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 1 and 2  
 T-Stat: 3.25 P-Val: 0.001157968169413171  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 1 and 3  
 T-Stat: 6.98 P-Val: 3.294359667247495e-12  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 2 and 3

**\*Inference:\***

- Average number of cycles rented is significantly different between each seasons

### 3. Problem Statement

No. of cycles rented similar or different in different weather

**\*Solution Approach:\***

- Null Hypothesis: **Average no. of cycles rented are equal for all weather conditions**
- Alternate Hypothesis: **Average no. of cycles rented is different for atleast one weather**
- Comparison between Average no. of cycles rented (*Numerical*) and Weather (*Category with 4 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
  - **Shapiro Test**
- **Variance Test**
  - Check for Homogeneity of variances
  - **Levene Test**
- **One Way ANOVA Test**
- Significance level: 5%

In [101...

```

count_weather = []
for i in range(1, data['weather'].nunique()): # Ignoring weather 4 as it is present
    count_weather.append(list(data[data['weather'] == i]['count']))
  
```

### 3.1. Check for Assumptions


```
In [102... check_normality(count_weather)
```

Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 9.781063280987223e-43 . Hence, Sample 2 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 3.876090133422781e-33 . Hence, Sample 3 doesn't follow normal distribution

```
In [103... plot_dist(count_weather)
```



```
In [105... stat, p_value = levene(count_weather[0], count_weather[1], count_weather[2])
if p_value < 0.05:
    print("Reject Null Hypothesis. Pval is ", p_value, ". Hence, the variance of
else:
    print("Fail to Reject Null Hypothesis. Pval is ", round(p_value,2), ". Hence
```

Reject Null Hypothesis. Pval is 6.198278710731511e-36 . Hence, the variance of at least one sample is significantly different

**\*Inference:\***

- For Anova, the assumptions are failed. All the samples don't follow normal distribution and homogeneity of variances
- Perform Anova and also Kruskal wallis test

### 3.2. Perform One Way Anova Test

```
In [107... # One Way Anova
alpha = 0.05
stat, p_value = f_oneway(count_weather[0], count_weather[1], count_weather[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(samp
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Average no. of cycles rented is differ
else:
    print('Fail to Reject Null Hypothesis. Hence, Average no. of cycles rented i
```

Test Statistic: 98.28 P-Val: 4.976448509904196e-43

Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast one weather

**\*Inference:\***

- One way Anova test concluded that **Average number of cycles rented is significantly different for atleast one weather**
- In order to find the weather condition at which the Average number of cycles rented is significantly different we need to perform 2 sample t test

### 3.3. Perform Kruskal Wallis Test

```
In [108... # Kruskal Wallis Test
alpha = 0.05
stat, p_value = kruskal(count_weather[0], count_weather[1], count_weather[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one weather')
else:
    print('Fail to Reject Null Hypothesis. Hence, Median no. of cycles rented is not different for atleast one weather')
```

Test Statistic: 204.96 P-Val: 3.122066178659941e-45

Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one weather

Inference:

- Since the assumptions of one way anova test is not met, the results can not be concluded directly from that test
- Kruskal wallis test, concludes that the Median no. of cycles rented is different for atleast one weather conditions

### 3.4. Perform 2 Sample T Test between dependent and each independent variable

```
In [109... alpha = 0.05
for sample in count_weather:
    print('Average no. of cycles rented in season ', str(i), ': ', np.mean(sample))

for idx1, idx2 in list(combinations(np.arange(len(count_weather)),2)):
    tstat, p_value = ttest_ind(count_weather[idx1], count_weather[idx2], alternative='two-sided')
    print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, Average number of cycles rented is different for atleast one weather')
    else:
        print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented is not different for atleast one weather')
```

Average no. of cycles rented in season 3 : 205.23679087875416

Average no. of cycles rented in season 3 : 178.95553987297106

Average no. of cycles rented in season 3 : 118.84633294528521

T-Stat: 6.49 P-Val: 9.098916216508542e-11

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 0 and 1

T-Stat: 13.05 P-Val: 1.4918709771846279e-38

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 0 and 2

T-Stat: 9.53 P-Val: 2.7459673190273646e-21

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between seasons 1 and 2

Inference:

- Average number of cycles rented is significantly different between each weather conditions

## 4. Problem Statement

Is Weather dependent on Season ?

**\*Solution Approach:\***

- Null Hypothesis: **Weather and Season are independent**
- Alternate Hypothesis: **Weather and Season are not independent**
- Comparison between Season (*Category with 4 categories*) and Weather (*Category with 4 categories*)
- **Chi Square Contingency Test**
- Significance level: 5%

```
In [ ]: def chi_test(ds1, ds2, alpha):
        df_conti = pd.crosstab(ds1, ds2)
        stat, p_value, dof, exp = chi2_contingency(df_conti)
        if p_value < alpha:
            print('Reject Null Hypothesis. Hence, Weather and Season are dependent')
        else:
            print('Fail to Reject Null Hypothesis. Weather and Season are independent')
```

```
In [190]: alpha = 0.05
chi_test(data[data['weather']<4]['weather'], data['season'], alpha):
```

Reject Null Hypothesis. Hence, Weather and Season are dependent

## 5. Problem Statement

Check the dependency of Average no. of cycles rented and all the categorical features?

### 5.1. Holiday Vs Average no. of cycles rented

**\*Solution Approach:\***

- Null Hypothesis:  **$u_1 = u_2$**
- Alternate Hypothesis:  **$u_1 > u_2$**

- u1 - Average no. of cycles rented during Holiday
- u2 - Average no. of cycles rented during non Holiday
- Significance level: 5%
- Comparison between Average no. of cycles rented (*Numerical*) and Holiday (*Category with 2 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
- Hence, **2 Sample T Test**


### 5.1.1 Check for Assumptions

```
In [ ]: df = data.groupby('date').agg({'holiday': 'min', 'count': 'mean'})
sample1 = df[df['holiday'] == 1]['count']
sample2 = df[df['holiday'] != 1]['count']
```

```
In [198... check_normality([sample1, sample2])
```

Reject Null Hypothesis. Pval is 1.3982287782710046e-05 . Hence, Sample 1 does n't follow normal distribution  
Fail to Reject Null Hypothesis. Pval is 0.08 . Hence, Sample 2 follows normal distribution

```
In [199... plot_dist([sample1, sample2])
```



#### \*Inference:\*

- Average no. of cycles rented during non holiday follows normal distribution
- Average no. of cycles rented during holiday doesn't follow normal distribution
- Distribution plot confirms the above point

### 5.1.2 Perform T-Test

```
In [200... alpha = 0.05
print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
tstat, p_value = ttest_ind(sample1, sample2, alternative='greater')
print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
```



```

if p_value < 0.05:
    print('Reject Null Hypothesis. Hence, Average number of cycles rented during
else:
    print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented

```

Sample-1 Mean: 192.28 Sample-2 Mean: 188.33

T-Stat: 0.51 P-Val: 0.30658555817068833

Fail to Reject Null Hypothesis. Hence, Average number of cycles rented during working day is equal to non working day

**\*Inference:\***

- Since the test is between a numerical and categorical variable (with 2 categories), 2 sample t-test is selected
- Alternate hypothesis is chosen as  $u_1 > u_2$  instead of  $u_1 < u_2$
- Fail to Reject Null Hypothesis. Hence, Average number of cycles rented during holiday day is equal to non holiday

## 6. Problem Statement

No. of cycles rented similar or different in different temperature bins

**\*Solution Approach:\***

- Null Hypothesis: **Average no. of cycles rented are equal for all temperature bins**
- Alternate Hypothesis: **Average no. of cycles rented is different for atleast one temperature bin**
- Comparison between Average no. of cycles rented (*Numerical*) and temperature bins (*Category with 4 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
  - **Shapiro Test**
- **Variance Test**
  - Check for Homogeneity of variances
  - **Levene Test**
- **One Way ANOVA Test**
- Significance level: 5%

In [209...

```

count_temp = []
for item in list(data['temp_cat'].unique()):
    count_temp.append(list(data[data['temp_cat'] == item]['count']))

```

### 6.1. Check for Assumptions

In [211...

```

check_normality(count_temp)

```


Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 2 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 2.3293470155990732e-18 . Hence, Sample 3 doesn't follow normal distribution

```
C:\Users\Muthukumar\AppData\Roaming\Python\Python311\site-packages\scipy\stats\morestats.py:1816: UserWarning: p-value may not be accurate for N > 5000.
warnings.warn("p-value may not be accurate for N > 5000.")
```

In [212... `plot_dist(count_temp)`



```
In [213... def Check_Variances_mod(samples):
    stat, p_value = levene(samples[0], samples[1], samples[2])
    if p_value < 0.05:
        print("Reject Null Hypothesis. Pval is ", p_value ,". Hence, the variance is significantly different")
    else:
        print("Fail to Reject Null Hypothesis. Pval is ", round(p_value,2) ,". Hence, the variance is not significantly different")
```

In [214... `Check_Variances_mod(count_temp)`

Reject Null Hypothesis. Pval is 1.3487721880363727e-133 . Hence, the variance of at least one sample is significantly different

#### **\*Inference:\***

- For Anova, the assumptions are failed. All the samples doesn't follow normal distribution and homogeneity of variances
- Perform Anova and also Kruskal wallis test

## 6.2. Perform One Way Anova Test

```
In [215... # One Way Anova
alpha = 0.05
stat, p_value = f_oneway(count_temp[0], count_temp[1], count_temp[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Average no. of cycles rented is different for at least one season')
else:
    print('Fail to Reject Null Hypothesis. Hence, Average no. of cycles rented is not significantly different for at least one season')
```

Test Statistic: 874.52 P-Val: 0.0

Reject Null Hypothesis. Hence, Average no. of cycles rented is different for at least one season

#### \*Inference:\*

- One way Anova test concluded that **\*Average number of cycles rented is significantly different for atleast one temperature bin\***
- In order to find the temperature where the Average number of cycles rented is significantly different we need to perform 2 sample t test

### 6.3. Perform Kruskal Wallis Test

```
In [217... # Kruskal Wallis Test
alpha = 0.05
stat, p_value = kruskal(count_temp[0], count_temp[1], count_temp[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one temp bin')
else:
    print('Fail to Reject Null Hypothesis. Hence, Median no. of cycles rented is same for all temp bins')
```

Test Statistic: 1650.27 P-Val: 0.0

Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one temp bin

Inference:

- Since the assumptions of one way anova test is not met, the results can not be concluded directly from that test
- Kruskal wallis test, concludes that the Median no. of cycles rented is different for atleast one temp bin

### 6.4. Perform 2 Sample T Test between dependent and each independent variable

```
In [219... alpha = 0.05
for sample in count_temp:
    print('Average no. of cycles rented in season ', str(i), ': ', np.mean(count_temp[sample]))

for idx1, idx2 in list(combinations(np.arange(len(count_temp)),2)):
    tstat, p_value = ttest_ind(count_temp[idx1], count_temp[idx2], alternative='two-sided')
    print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, Average number of cycles rented is different for atleast one temp bin')
    else:
        print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented is same for all temp bins')
```

Average no. of cycles rented in season 2 : 110.04892425582081  
 Average no. of cycles rented in season 2 : 207.43206913106096  
 Average no. of cycles rented in season 2 : 334.274115755627  
 T-Stat: -27.46 P-Val: 6.882380020634517e-160  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between temperature bin 0 and 1  
 T-Stat: -48.55 P-Val: 0.0  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between temperature bin 0 and 2  
 T-Stat: -21.98 P-Val: 8.19521197158562e-104  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between temperature bin 1 and 2

Inference:

- Average number of cycles rented is significantly different between each temperature bins

## 7. Problem Statement

No. of cycles rented similar or different in different actual temperature bins

**\*Solution Approach:\***

- Null Hypothesis: **Average no. of cycles rented are equal for all actual temperature bins**
- Alternate Hypothesis: **Average no. of cycles rented is different for atleast one actual temperature bin**
- Comparison between Average no. of cycles rented (*Numerical*) and actual temperature bins (*Category with 3 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
  - **Shapiro Test**
- **Variance Test**
  - Check for Homogeneity of variances
  - **Levene Test**
- **One Way ANOVA Test**
- Significance level: 5%

```
In [222... count_atemp = []
for item in list(data['atemp_cat'].unique()):
    count_atemp.append(list(data[data['atemp_cat'] == item]['count']))
```


### 7.1. Check for Assumptions

```
In [223... check_normality(count_atemp)
```

Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution  
 Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 2 doesn't follow normal distribution  
 Reject Null Hypothesis. Pval is 6.537652891420251e-31 . Hence, Sample 3 doesn't follow normal distribution

```
C:\Users\Muthukumar\AppData\Roaming\Python\Python311\site-packages\scipy\stats\_morestats.py:1816: UserWarning: p-value may not be accurate for N > 5000.
warnings.warn("p-value may not be accurate for N > 5000.")
```

In [224... `plot_dist(count_temp)`



```
In [213... def Check_Variances_mod(samples):
    stat, p_value = levene(samples[0], samples[1], samples[2])
    if p_value < 0.05:
        print("Reject Null Hypothesis. Pval is ", p_value ,". Hence, the variance is significantly different")
    else:
        print("Fail to Reject Null Hypothesis. Pval is ", round(p_value,2) ,". Hence, the variance is not significantly different")
```

In [225... `Check_Variances_mod(count_atemp)`

Reject Null Hypothesis. Pval is 2.2555487020783844e-150 . Hence, the variance of atleast one sample is significantly different

#### **\*Inference:\***

- For Anova, the assumptions are failed. All the samples doesn't follow normal distribution and homogeneity of variances
- Perform Anova and also Kruskal wallis test

## 7.2. Perform One Way Anova Test

```
In [226... # One Way Anova
alpha = 0.05
stat, p_value = f_oneway(count_atemp[0], count_atemp[1], count_atemp[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast one temp bin')
else:
    print('Fail to Reject Null Hypothesis. Hence, Average no. of cycles rented is not significantly different for atleast one temp bin')
```

Test Statistic: 1008.35 P-Val: 0.0

Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast one temp bin

### \*Inference:\*

- One way Anova test concluded that Average number of cycles rented is significantly different for atleast one actual temperature bin
- In order to find the temperature where the Average number of cycles rented is significantly different we need to perform 2 sample t test

## 7.3. Perform Kruskal Wallis Test

```
In [227... # Kruskal Wallis Test
alpha = 0.05
stat, p_value = kruskal(count_atemp[0], count_atemp[1], count_atemp[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one temp bin')
else:
    print('Fail to Reject Null Hypothesis. Hence, Median no. of cycles rented is same for all temp bins')
```

Test Statistic: 1846.6 P-Val: 0.0

Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one temp bin

Inference:

- Since the assumptions of one way anova test is not met, the results can not be concluded directly from that test
- Kruskal wallis test, concludes that the Median no. of cycles rented is different for atleast one actual temperature bin

## 7.4. Perform 2 Sample T Test between dependent and each independent variable

```
In [228... alpha = 0.05
for sample in count_temp:
    print('Average no. of cycles rented in season ', str(i), ': ', np.mean(count_temp[sample]))

for idx1, idx2 in list(combinations(np.arange(len(count_atemp)),2)):
    tstat, p_value = ttest_ind(count_atemp[idx1], count_atemp[idx2], alternative='two-sided')
    print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, Average number of cycles rented is different for atleast one temp bin')
    else:
        print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented is same for all temp bins')
```

Average no. of cycles rented in season 2 : 89.24617737003058  
 Average no. of cycles rented in season 2 : 169.62495593937257  
 Average no. of cycles rented in season 2 : 291.6686153846154  
 T-Stat: -20.48 P-Val: 9.160970366148517e-91  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between temperature bin 0 and 1  
 T-Stat: -41.34 P-Val: 0.0  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between temperature bin 0 and 2  
 T-Stat: -31.41 P-Val: 1.8596348077188179e-205  
 Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between temperature bin 1 and 2

**\*Inference:\***

- Average number of cycles rented is significantly different between each actual temperature bins

## 8. Problem Statement

No. of cycles rented similar or different in different range of windspeed

**\*Solution Approach:\***

- Null Hypothesis: **Average no. of cycles rented are equal for all range of windspeed**
- Alternate Hypothesis: **Average no. of cycles rented is different for atleast one range of windspeed**
- Comparison between Average no. of cycles rented (*Numerical*) and range of windspeed (*Category with 3 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
  - **Shapiro Test**
- **Variance Test**
  - Check for Homogeneity of variances
  - **Levene Test**
- **One Way ANOVA Test**
- Significance level: 5%

```
In [237... count_ws = []
for item in list(data['windspeed_cat'].dropna().unique()):
    count_ws.append(list(data[data['windspeed_cat'] == item]['count']))
```

```
In [238... for i in range(len(count_ws)):
    print(len(count_ws[i]))
```

3026  
 5698  
 849

```
In [236... data['windspeed_cat'].dropna().unique()
```

```
Out[236...] ['Low', 'Med', 'High']
Categories (3, object): ['Low' < 'Med' < 'High']
```

## 8.1. Check for Assumptions

```
In [239...] check_normality(count_ws)
```


Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 2 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 6.79376919669305e-23 . Hence, Sample 3 doesn't follow normal distribution

C:\Users\Muthukumar\AppData\Roaming\Python\Python311\site-packages\scipy\stats\morestats.py:1816: UserWarning: p-value may not be accurate for N > 5000.  
warnings.warn("p-value may not be accurate for N > 5000.")

```
In [240...] plot_dist(count_ws)
```



```
In [241...] Check_Variiances_mod(count_ws)
```

Reject Null Hypothesis. Pval is 1.1009171815057205e-08 . Hence, the variance of atleast one sample is significantly different

**\*Inference:\***

- For Anova, the assumptions are failed. All the samples doesn't follow normal distribution and homogeneity of variances
- Perform Anova and also Kruskal wallis test

## 8.2. Perform One Way Anova Test

```
In [243...] # One Way Anova
alpha = 0.05
stat, p_value = f_oneway(count_ws[0], count_ws[1], count_ws[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(samp
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Average no. of cycles rented is differ
```



```
else:
    print('Fail to Reject Null Hypothesis. Hence, Average no. of cycles rented i
```

Test Statistic: 61.49 P-Val: 2.9212488566229153e-27

Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast windspeed range

**\*Inference:**

- One way Anova test concluded that Average number of cycles rented is significantly different for atleast one windspeed range
- In order to find the windspeed range where the Average number of cycles rented is significantly different we need to perform 2 sample t test

### 8.3. Perform Kruskal Wallis Test

In [244...

```
# Kruskal Wallis Test
alpha = 0.05
stat, p_value = kruskal(count_ws[0], count_ws[1], count_ws[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one windspeed range')
else:
    print('Fail to Reject Null Hypothesis. Hence, Median no. of cycles rented is
```

Test Statistic: 146.64 P-Val: 1.4350374917697305e-32

Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one windspeed range

**\*Inference:**

- Since the assumptions of one way anova test is not met, the results can not be concluded directly from that test
- Kruskal wallis test, concludes that the Median no. of cycles rented is different for atleast one windspeed range

### 8.4. Perform 2 Sample T Test between dependent and each independent variable

In [245...

```
alpha = 0.05
for sample in count_ws:
    print('Average no. of cycles rented in season ', str(i), ': ', np.mean(sample))

for idx1, idx2 in list(combinations(np.arange(len(count_ws)),2)):
    tstat, p_value = ttest_ind(count_ws[idx1], count_ws[idx2], alternative='two-sided')
    print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, Average number of cycles rented is different for atleast one windspeed range')
    else:
        print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented is
```

Average no. of cycles rented in season 2 : 165.66027759418375

Average no. of cycles rented in season 2 : 209.1123201123201

Average no. of cycles rented in season 2 : 213.35689045936397

T-Stat: -10.68 P-Val: 1.9179099323170704e-26

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between windspeed range 0 and 1

T-Stat: -7.17 P-Val: 8.740907653897123e-13

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between windspeed range 0 and 2

T-Stat: -0.62 P-Val: 0.5352388697851858

Fail to Reject Null Hypothesis. Hence, Average number of cycles rented is equal between windspeed range 1 and 2

**\*Inference:\***

- Average number of cycles rented is significantly different between windspeed range 0 and 1 & windspeed range 0 and 2
- Average number of cycles rented is significantly equal between windspeed range 1 and 2

## 9. Problem Statement

No. of cycles rented similar or different in different humidity levels

**\*Solution Approach:\***

- Null Hypothesis: **Average no. of cycles rented are equal for all humidity levels**
- Alternate Hypothesis: **Average no. of cycles rented is different for atleast one humidity level**
- Comparison between Average no. of cycles rented (*Numerical*) and humidity levels (*Category with 3 categories*)
- **Normality Test**
  - Check for Average no. of cycles rented follow Normal distribution
  - **Shapiro Test**
- **Variance Test**
  - Check for Homogeneity of variances
  - **Levene Test**
- **One Way ANOVA Test**
- Significance level: 5%

In [246...

```
count_humidity = []
for item in list(data['humidity_cat'].dropna().unique()):
    count_humidity.append(list(data[data['humidity_cat'] == item]['count']))
```

### 9.1. Check for Assumptions

In [249...

```
check_normality(count_humidity)
```


Reject Null Hypothesis. Pval is 0.0 . Hence, Sample 1 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 4.90454462513686e-44 . Hence, Sample 2 doesn't follow normal distribution

Reject Null Hypothesis. Pval is 1.0363253878974854e-24 . Hence, Sample 3 doesn't follow normal distribution

```
C:\Users\Muthukumar\AppData\Roaming\Python\Python311\site-packages\scipy\stats\_morestats.py:1816: UserWarning: p-value may not be accurate for N > 5000.  
warnings.warn("p-value may not be accurate for N > 5000.")
```

In [250... `plot_dist(count_humidity)`



In [241... `Check_Variances_mod(count_ws)`

Reject Null Hypothesis. Pval is 1.1009171815057205e-08 . Hence, the variance of atleast one sample is significantly different

**\*Inference:**

- For Anova, the assumptions are failed. All the samples doesn't follow normal distribution and homogeneity of variances
- Perform Anova and also Kruskal wallis test

## 9.2. Perform One Way Anova Test

```
In [251... # One Way Anova  
alpha = 0.05  
stat, p_value = f_oneway(count_humidity[0], count_humidity[1], count_humidity[2])  
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))  
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)  
if p_value < alpha:  
    print('Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast one humidity level')  
else:  
    print('Fail to Reject Null Hypothesis. Hence, Average no. of cycles rented is same for all humidity levels')
```

Test Statistic: 482.02 P-Val: 2.741782831381464e-201

Reject Null Hypothesis. Hence, Average no. of cycles rented is different for atleast one humidity level

**\*Inference:**

- One way Anova test concluded that **Average number of cycles rented is significantly different for atleast one humidity level**
- In order to find the humidity level where the Average number of cycles rented is significantly different we need to perform 2 sample t test

### 9.3. Perform Kruskal Wallis Test

In [252...

```
# Kruskal Wallis Test
alpha = 0.05
stat, p_value = kruskal(count_humidity[0], count_humidity[1], count_humidity[2])
#print('Sample-1 Mean: ', round(sample1.mean(),2), 'Sample-2 Mean: ', round(sample2.mean(),2))
print('Test Statistic: ', round(stat,2), 'P-Val: ', p_value)
if p_value < alpha:
    print('Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one humidity level')
else:
    print('Fail to Reject Null Hypothesis. Hence, Median no. of cycles rented is not significantly different for atleast one humidity level')
```

Test Statistic: 1061.6 P-Val: 2.996871369661641e-231

Reject Null Hypothesis. Hence, Median no. of cycles rented is different for atleast one humidity level

Inference:

- Since the assumptions of one way anova test is not met, the results can not be concluded directly from that test
- Kruskal wallis test, concludes that the Median no. of cycles rented is different for atleast one humidity level

### 9.4. Perform 2 Sample T Test between dependent and each independent variable

In [253...

```
alpha = 0.05
for sample in count_humidity:
    print('Average no. of cycles rented in season ', str(i), ': ', np.mean(sample))

for idx1, idx2 in list(combinations(np.arange(len(count_humidity)),2)):
    tstat, p_value = ttest_ind(count_humidity[idx1], count_humidity[idx2], alternative='two-sided')
    print('T-Stat: ', round(tstat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between humidity level 0 and 1')
    else:
        print('Fail to Reject Null Hypothesis. Hence, Average number of cycles rented is not significantly different between humidity level 0 and 1')
```

Average no. of cycles rented in season 2 : 190.4756966947505

Average no. of cycles rented in season 2 : 96.85192433137638

Average no. of cycles rented in season 2 : 288.89789603960395

T-Stat: 19.78 P-Val: 2.358356748074402e-85

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between humidity level 0 and 1

T-Stat: -19.93 P-Val: 1.3306807309895657e-86

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between humidity level 0 and 2

T-Stat: -32.53 P-Val: 2.174595591792677e-200

Reject Null Hypothesis. Hence, Average number of cycles rented is significantly different between humidity level 1 and 2

**\*Inference:\***

- Average number of cycles rented is significantly different between each humidity levels

## 10. Problem Statement

Check for dependency between each categorical features

**\*Solution Approach:\***

- Null Hypothesis: **Categorical variables are independent**
- Alternate Hypothesis: **Categorical variables are dependent**
- **Chi-square Independency Test**
- Significance level: 5%


In [270...

```
cat_features = ['season', 'workingday', 'weather', 'temp_cat', 'atemp_cat', 'hum
df = pd.DataFrame(data = np.ones((len(cat_features),len(cat_features))) * -1, co
for idx1, idx2 in list(combinations(cat_features,2)):
    contingency_table = pd.crosstab(data[idx1], data[idx2])
    stat, p_value, a, exp = chi2_contingency(contingency_table)
    print('Stat: ', round(stat,2), 'P-Val: ', p_value)
    if p_value < 0.05:
        print('Reject Null Hypothesis. Hence, ', str(idx1), ' and ', str(idx2),
    else:
        print('Fail to Reject Null Hypothesis. Hence, ', str(idx1), ' and ', str
df.loc[idx1, idx2] = round(p_value,2)
df.loc[idx2, idx1] = round(p_value,2)
```

Stat: 2.57 P-Val: 0.4626148207703564  
 Fail to Reject Null Hypothesis. Hence, season and workingday are independent  
 Stat: 49.16 P-Val: 1.549925073686492e-07  
 Reject Null Hypothesis. Hence, season and weather are dependent  
 Stat: 5897.24 P-Val: 0.0  
 Reject Null Hypothesis. Hence, season and temp\_cat are dependent  
 Stat: 6630.76 P-Val: 0.0  
 Reject Null Hypothesis. Hence, season and atemp\_cat are dependent  
 Stat: 441.6 P-Val: 3.154697752402428e-92  
 Reject Null Hypothesis. Hence, season and humidity\_cat are dependent  
 Stat: 191.15 P-Val: 1.4477127726644647e-38  
 Reject Null Hypothesis. Hence, season and windspeed\_cat are dependent  
 Stat: 16.16 P-Val: 0.0010502165960627732  
 Reject Null Hypothesis. Hence, workingday and weather are dependent  
 Stat: 9.57 P-Val: 0.008342423420746568  
 Reject Null Hypothesis. Hence, workingday and temp\_cat are dependent  
 Stat: 9.73 P-Val: 0.0077190014112448494  
 Reject Null Hypothesis. Hence, workingday and atemp\_cat are dependent  
 Stat: 1.74 P-Val: 0.4193926395523214  
 Fail to Reject Null Hypothesis. Hence, workingday and humidity\_cat are independent  
 Stat: 3.36 P-Val: 0.1865186782185527  
 Fail to Reject Null Hypothesis. Hence, workingday and windspeed\_cat are independent  
 Stat: 152.29 P-Val: 2.5393088306954665e-30  
 Reject Null Hypothesis. Hence, weather and temp\_cat are dependent  
 Stat: 277.81 P-Val: 4.6243537419201974e-57  
 Reject Null Hypothesis. Hence, weather and atemp\_cat are dependent  
 Stat: 2048.43 P-Val: 0.0  
 Reject Null Hypothesis. Hence, weather and humidity\_cat are dependent  
 Stat: 44.31 P-Val: 6.422935286585923e-08  
 Reject Null Hypothesis. Hence, weather and windspeed\_cat are dependent  
 Stat: 8197.02 P-Val: 0.0  
 Reject Null Hypothesis. Hence, temp\_cat and atemp\_cat are dependent  
 Stat: 330.26 P-Val: 3.2088658507065833e-70  
 Reject Null Hypothesis. Hence, temp\_cat and humidity\_cat are dependent  
 Stat: 55.9 P-Val: 2.103733252175866e-11  
 Reject Null Hypothesis. Hence, temp\_cat and windspeed\_cat are dependent  
 Stat: 808.21 P-Val: 1.2813030257313233e-173  
 Reject Null Hypothesis. Hence, atemp\_cat and humidity\_cat are dependent  
 Stat: 82.52 P-Val: 5.1018430943325584e-17  
 Reject Null Hypothesis. Hence, atemp\_cat and windspeed\_cat are dependent  
 Stat: 660.66 P-Val: 1.1468186720611514e-141  
 Reject Null Hypothesis. Hence, humidity\_cat and windspeed\_cat are dependent

In [273... `sns.heatmap(df, vmin = 0, annot=True)`

Out[273... `<Axes: >`



**\*Inference:\***

- In the above heatmap, dark indicates dependency (pval~0) and light color indicates independency (pval>0)
- It is evident that almost all the categorical features are dependent except working day

## Business Insights and Recommendations

### Weather-Based Insights:

- Observation:
  - Light weather conditions (Weather 1 and 2) see high usage of the cycles.
- Recommendation:
  - Focus marketing efforts and offer discounts or reduced costs during Weather 1 and 2 conditions to capitalize on high demand
  - Hypothesis test also concludes that Average number of cycles rented is significantly different for weather 1 and 2. Hence the strategies involving weather 1 and 2 will make a significant impact whereas the weather 2 doesn't.
  - Utilize dynamic pricing models that adjust costs based on real-time weather conditions, prioritizing Weather 1 for promotions to maximize revenue.

### Weekly usage-Based Insights:

- Observation:
  - Registered users are predominant on weekdays indicates that they rent cycles for commuting to work/students
  - Casual users are predominant on weekends indicating that they commute for leisure
  - Users count during holiday and working day also confirms the above point
- Recommendation:
  - Increase efforts to grow the registered user base to boost weekday revenue, as they tend to have higher usage rates
  - Introduce weekday subscription plans or loyalty programs for registered users to encourage frequent use
  - On weekends, tailor promotions and offers to attract casual users with leisure-oriented marketing
  - Design weekend packages that appeal to casual users, such as group discounts or event partnerships

### **YoY Growth Insights:**

- Observation:
  - Year-over-year (YoY) user growth indicates that current promotion strategies are effective.
- Recommendation:
  - Continue and refine existing promotional strategies that have proven successful.
  - Analyze which promotional channels and messages have driven the most growth and amplify those efforts.

### **Time-Based Insights:**

- Observation:
  - The average number of casual users peaks between 13:00 and 17:00, indicating high demand during these hours.
- Recommendation:
  - Implement dynamic pricing during peak hours to manage demand and maximize revenue. Adjust pricing to reflect the higher value of cycles during peak usage times
  - Set up a pricing model that increases rates during peak hours (13:00-17:00) while offering discounts during off-peak times to balance demand throughout the day

### **Temperature-Based Insights:**

- Observation:
  - Cycle usage is positively correlated with temperature
- Recommendation:
  - Hypothesis testing confirms that the average number of cycles rented varies significantly with temperature. Hence the strategies involving temperature bin will make a significant impact



- Incorporate temperature data into dynamic pricing models. Increase prices during favorable temperature conditions (higher usage) and consider lowering prices or offering incentives when conditions are less ideal.
- Utilize accurate weather prediction APIs to anticipate demand and adjust pricing accordingly.

#### **Humidity-Based Insights:**

- Observation:
  - Cycle usage is negatively correlated with humidity
- Recommendation:
  - Hypothesis testing confirms that the average number of cycles rented varies significantly with humidity. Hence the strategies involving humidity will make a significant impact
  - Incorporate humidity data into dynamic pricing models. Adjust pricing based on humidity levels to encourage usage during less favorable conditions.
  - Utilize accurate weather prediction APIs to anticipate demand and adjust pricing accordingly.

#### **Day-Based Insights:**

- Observation:
  - The average number of cycles rented is not significantly different between working days and non-working days
- Recommendation:
  - The strategies involving working day doesn't significantly impact
  - Maintain consistent pricing across working and non-working days

#### **Season-Based Insights:**

- Observation:
  - The average number of cycles rented varies significantly between seasons
- Recommendation:
  - The strategies involving each seasons will make significantly impact
  - Increase prices during peak seasons with high demand and offer discounts or promotions during off-peak seasons to encourage usage
  - Create seasonal marketing campaigns that highlight the benefits of cycling in each season

**Prepared by Muthukumar G**