

Matrix Multiplication

Madhavan Mukund

<https://www.cmi.ac.in/~madhavan>

Mathematics for Data Science 1
Week 11

Matrices

- A matrix is a two dimensional table
 - $r \times c$ matrix — r rows, c columns

Matrices

- A matrix is a two dimensional table
 - $r \times c$ matrix — r rows, c columns
 - Assume rows are numbered $\{0, 1, \dots, r-1\}$, columns are numbered $\{0, 1, \dots, c-1\}$

Matrices

- A matrix is a two dimensional table
 - $r \times c$ matrix — r rows, c columns
 - Assume rows are numbered $\{0, 1, \dots, r-1\}$, columns are numbered $\{0, 1, \dots, c-1\}$
 - Graph with n nodes, $n \times n$ adjacency matrix, entries are from $\{0, 1\}$

Matrices

- A matrix is a two dimensional table
 - $r \times c$ matrix — r rows, c columns
 - Assume rows are numbered $\{0, 1, \dots, r-1\}$, columns are numbered $\{0, 1, \dots, c-1\}$
 - Graph with n nodes, $n \times n$ adjacency matrix, entries are from $\{0, 1\}$
- Example: Freight traffic by rail between major cities: Bangalore, Chennai, Delhi, Hyderabad, Kolkata, Mumbai

Matrices

- A matrix is a two dimensional table
 - $r \times c$ matrix — r rows, c columns
 - Assume rows are numbered $\{0, 1, \dots, r-1\}$, columns are numbered $\{0, 1, \dots, c-1\}$
 - Graph with n nodes, $n \times n$ adjacency matrix, entries are from $\{0, 1\}$
- Example: Freight traffic by rail between major cities: Bangalore, Chennai, Delhi, Hyderabad, Kolkata, Mumbai
 - Number the cities:
0-Bangalore, 1-Chennai, 2-Delhi,
3-Hyderabad, 4-Kolkata,
5-Mumbai

Matrices

- A matrix is a two dimensional table
 - $r \times c$ matrix — r rows, c columns
 - Assume rows are numbered $\{0, 1, \dots, r-1\}$, columns are numbered $\{0, 1, \dots, c-1\}$
 - Graph with n nodes, $n \times n$ adjacency matrix, entries are from $\{0, 1\}$
- Example: Freight traffic by rail between major cities: Bangalore, Chennai, Delhi, Hyderabad, Kolkata, Mumbai

- Number the cities:
0-Bangalore, 1-Chennai, 2-Delhi,
3-Hyderabad, 4-Kolkata,
5-Mumbai

- Represent freight volume for a month as a 6×6 matrix

	0	1	2	3	4	5
0	0	694	828	384	247	479
1	642	0	919	575	402	673
2	768	734	0	231	595	540
3	731	606	156	0	351	804
4	825	607	316	490	0	998
5	196	580	339	588	394	0

Adding matrices

- Suppose we have freight volumes for first and second half of financial year.

April–September							October–March						
	0	1	2	3	4	5		0	1	2	3	4	5
0	0	694	828	384	247	479	0	0	851	626	280	399	365
1	642	0	919	575	402	673	1	544	0	479	269	432	933
2	768	734	0	231	595	540	2	867	804	0	681	326	398
3	731	606	156	0	351	804	3	727	976	418	0	667	294
4	825	607	316	490	0	998	4	894	390	247	547	0	314
5	196	580	339	588	394	0	5	914	147	574	859	524	0

Adding matrices

- Suppose we have freight volumes for first and second half of financial year.

April–September							October–March						
	0	1	2	3	4	5		0	1	2	3	4	5
0	0	694	828	384	247	479	0	0	851	626	280	399	365
1	642	0	919	575	402	673	1	544	0	479	269	432	933
2	768	734	0	231	595	540	2	867	804	0	681	326	398
3	731	606	156	0	351	804	3	727	976	418	0	667	294
4	825	607	316	490	0	998	4	894	390	247	547	0	314
5	196	580	339	588	394	0	5	914	147	574	859	524	0

- How do we compute the freight volumes for the entire financial year?

Adding matrices

- Suppose we have freight volumes for first and second half of financial year.

April–September							October–March						
	0	1	2	3	4	5		0	1	2	3	4	5
0	0	694	828	384	247	479	0	0	851	626	280	399	365
1	642	0	919	575	402	673	1	544	0	479	269	432	933
2	768	734	0	231	595	540	2	867	804	0	681	326	398
3	731	606	156	0	351	804	3	727	976	418	0	667	294
4	825	607	316	490	0	998	4	894	390	247	547	0	314
5	196	580	339	588	394	0	5	914	147	574	859	524	0

- How do we compute the freight volumes for the entire financial year?
- Add the corresponding entries in the two tables
 - Total freight volume from 2 (Delhi) to 4 (Kolkata) is $595 + 326 = 921$

Adding matrices

- For a matrix M , $M[i,j]$ is the entry in row i , column j

Adding matrices

- For a matrix M , $M[i,j]$ is the entry in row i , column j
- Let A and B represent the volumes in the two half-years

	0	1	2	3	4	5
0	0	694	828	384	247	479
1	642	0	919	575	402	673
2	768	734	0	231	595	540
3	731	606	156	0	351	804
4	825	607	316	490	0	998
5	196	580	339	588	394	0

	0	1	2	3	4	5
0	0	851	626	280	399	365
1	544	0	479	269	432	933
2	867	804	0	681	326	398
3	727	976	418	0	667	294
4	894	390	247	547	0	314
5	914	147	574	859	524	0

Adding matrices

- For a matrix M , $M[i,j]$ is the entry in row i , column j
- Let A and B represent the volumes in the two half-years
- Let C represent the annual volume

	0	1	2	3	4	5
0	0	694	828	384	247	479
1	642	0	919	575	402	673
2	768	734	0	231	595	540
3	731	606	156	0	351	804
4	825	607	316	490	0	998
5	196	580	339	588	394	0

	0	1	2	3	4	5
0	0	851	626	280	399	365
1	544	0	479	269	432	933
2	867	804	0	681	326	398
3	727	976	418	0	667	294
4	894	390	247	547	0	314
5	914	147	574	859	524	0

Adding matrices

- For a matrix M , $M[i,j]$ is the entry in row i , column j
- Let A and B represent the volumes in the two half-years
- Let C represent the annual volume
- For each i,j , $C[i,j] = A[i,j] + B[i,j]$

	0	1	2	3	4	5
0	0	694	828	384	247	479
1	642	0	919	575	402	673
2	768	734	0	231	595	540
3	731	606	156	0	351	804
4	825	607	316	490	0	998
5	196	580	339	588	394	0

	0	1	2	3	4	5
0	0	851	626	280	399	365
1	544	0	479	269	432	933
2	867	804	0	681	326	398
3	727	976	418	0	667	294
4	894	390	247	547	0	314
5	914	147	574	859	524	0

	0	1	2	3	4	5
0	0	1545	1454	664	646	844
1	1186	0	1398	844	834	1606
2	1635	1538	0	912	921	938
3	1458	1582	574	0	1018	1098
4	1719	997	563	1037	0	1312
5	1110	727	913	1447	918	0

Adding matrices

- For a matrix M , $M[i,j]$ is the entry in row i , column j
- Let A and B represent the volumes in the two half-years
- Let C represent the annual volume
- For each i,j , $C[i,j] = A[i,j] + B[i,j]$
- More concisely, $C = A + B$ — matrix addition

	0	1	2	3	4	5
0	0	694	828	384	247	479
1	642	0	919	575	402	673
2	768	734	0	231	595	540
3	731	606	156	0	351	804
4	825	607	316	490	0	998
5	196	580	339	588	394	0

	0	1	2	3	4	5
0	0	851	626	280	399	365
1	544	0	479	269	432	933
2	867	804	0	681	326	398
3	727	976	418	0	667	294
4	894	390	247	547	0	314
5	914	147	574	859	524	0

	0	1	2	3	4	5
0	0	1545	1454	664	646	844
1	1186	0	1398	844	834	1606
2	1635	1538	0	912	921	938
3	1458	1582	574	0	1018	1098
4	1719	997	563	1037	0	1312
5	1110	727	913	1447	918	0

Multiplying matrices

- Can we multiply matrices the same way?

Multiplying matrices

- Can we multiply matrices the same way?
- For each i,j , $C[i,j] = A[i,j] \times B[i,j]$

		0	1	2	3
	0	0	6	8	3
A	1	4	0	9	7
	2	6	3	0	1
	3	1	0	5	0

		0	1	2	3
	0	0	1	2	8
B	1	5	0	4	2
	2	6	0	0	1
	3	2	6	4	0

		0	1	2	3
	0	0	6	16	24
C	1	20	0	36	14
	2	36	0	0	1
	3	2	0	20	0

Multiplying matrices

- Can we multiply matrices the same way?
- For each i,j , $C[i,j] = A[i,j] \times B[i,j]$
- This turns out to be not very useful

		0	1	2	3
	0	0	6	8	3
A	1	4	0	9	7
	2	6	3	0	1
	3	1	0	5	0

		0	1	2	3
	0	0	1	2	8
B	1	5	0	4	2
	2	6	0	0	1
	3	2	6	4	0

		0	1	2	3
	0	0	6	16	24
C	1	20	0	36	14
	2	36	0	0	1
	3	2	0	20	0

Multiplying matrices

- Can we multiply matrices the same way?
- For each i,j , $C[i,j] = A[i,j] \times B[i,j]$
- This turns out to be not very useful
- Instead, we compute $C[i,j]$ in a more complicated way

	0	1	2	3
0	0	6	8	3
1	4	0	9	7
2	6	3	0	1
3	1	0	5	0

	0	1	2	3
0	0	1	2	8
1	5	0	4	2
2	6	0	0	1
3	2	6	4	0

Multiplying matrices

- Can we multiply matrices the same way?
- For each i, j , $C[i, j] = A[i, j] \times B[i, j]$
- This turns out to be not very useful
- Instead, we compute $C[i, j]$ in a more complicated way
- Assume $r = c = n$. Let

$$\begin{aligned} C[i, j] = & A[i, 0] \cdot B[0, j] + \\ & A[i, 1] \cdot B[1, j] + \\ & \dots + \\ & A[i, n-1] \cdot B[n-1, j] \end{aligned}$$

	0	1	2	3
0	0	6	8	3
1	4	0	9	7
2	6	3	0	1
3	1	0	5	0

	0	1	2	3
0	0	1	2	8
1	5	0	4	2
2	6	0	0	1
3	2	6	4	0

Multiplying matrices

- Can we multiply matrices the same way?
- For each i, j , $C[i, j] = A[i, j] \times B[i, j]$
- This turns out to be not very useful
- Instead, we compute $C[i, j]$ in a more complicated way
- Assume $r = c = n$. Let

$$\begin{aligned} C[i, j] = & A[i, 0] \cdot B[0, j] + \\ & A[i, 1] \cdot B[1, j] + \\ & \dots + \\ & A[i, n-1] \cdot B[n-1, j] \end{aligned}$$

- For instance,
 $C[1, 3] = 4 \cdot 8 + 0 \cdot 2 + 9 \cdot 1 + 7 \cdot 0 = 41$

	0	1	2	3
0	0	6	8	3
1	4	0	9	7
2	6	3	0	1
3	1	0	5	0

	0	1	2	3
0	0	1	2	8
1	5	0	4	2
2	6	0	0	1
3	2	6	4	0

Multiplying matrices

- Can we multiply matrices the same way?
- For each i, j , $C[i, j] = A[i, j] \times B[i, j]$
- This turns out to be not very useful
- Instead, we compute $C[i, j]$ in a more complicated way
- Assume $r = c = n$. Let
$$C[i, j] = A[i, 0] \cdot B[0, j] + A[i, 1] \cdot B[1, j] + \dots + A[i, n-1] \cdot B[n-1, j]$$
- For instance,
$$C[1, 3] = 4 \cdot 8 + 0 \cdot 2 + 9 \cdot 1 + 7 \cdot 0 = 41$$

		0	1	2	3
	0	0	6	8	3
A	1	4	0	9	7
	2	6	3	0	1
	3	1	0	5	0

		0	1	2	3
	0	0	1	2	8
B	1	5	0	4	2
	2	6	0	0	1
	3	2	6	4	0

		0	1	2	3
	0	84	18	36	20
C	1	68	46	36	41
	2	17	12	28	54
	3	30	1	2	13

Multiplying matrices

- Matrix product: $C = A \times B$

		0	1	2	3
	0	0	6	8	3
A	1	4	0	9	7
	2	6	3	0	1
	3	1	0	5	0

		0	1	2	3
	0	0	1	2	8
B	1	5	0	4	2
	2	6	0	0	1
	3	2	6	4	0

		0	1	2	3
	0	84	18	36	20
C	1	68	46	36	41
	2	17	12	28	54
	3	30	1	2	13

Multiplying matrices

■ Matrix product: $C = A \times B$

■ For $r = c = n$,

$$\begin{aligned} C[i, j] = & A[i, 0] \cdot B[0, j] + \\ & A[i, 1] \cdot B[1, j] + \\ & \dots + \\ & A[i, n-1] \cdot B[n-1, j] \end{aligned}$$

	0	1	2	3
A 0	0	6	8	3
1	4	0	9	7
2	6	3	0	1
3	1	0	5	0

	0	1	2	3
B 0	0	1	2	8
1	5	0	4	2
2	6	0	0	1
3	2	6	4	0

	0	1	2	3
C 0	84	18	36	20
1	68	46	36	41
2	17	12	28	54
3	30	1	2	13

Multiplying matrices

- Matrix product: $C = A \times B$

- For $r = c = n$,

$$\begin{aligned} C[i, j] = & A[i, 0] \cdot B[0, j] + \\ & A[i, 1] \cdot B[1, j] + \\ & \dots + \\ & A[i, n-1] \cdot B[n-1, j] \end{aligned}$$

- More concisely, $C[i, j] = \sum_{k=0}^{n-1} A[i, k] \cdot B[k, j]$

		0	1	2	3
A	0	0	6	8	3
	1	4	0	9	7
	2	6	3	0	1
	3	1	0	5	0

		0	1	2	3
B	0	0	1	2	8
	1	5	0	4	2
	2	6	0	0	1
	3	2	6	4	0

		0	1	2	3
C	0	84	18	36	20
	1	68	46	36	41
	2	17	12	28	54
	3	30	1	2	13

Multiplying matrices

- Matrix product: $C = A \times B$

- For $r = c = n$,

$$\begin{aligned} C[i, j] = & A[i, 0] \cdot B[0, j] + \\ & A[i, 1] \cdot B[1, j] + \\ & \dots + \\ & A[i, n-1] \cdot B[n-1, j] \end{aligned}$$

- More concisely, $C[i, j] = \sum_{k=0}^{n-1} A[i, k] \cdot B[k, j]$

- Don't require both A and B to be $n \times n$

- Each row entry of A must have a matching column entry in B

- If A is $m \times n$ and B is $n \times p$, $A \times B$ is $m \times p$

		0	1	2	3
A	0	0	6	8	3
	1	4	0	9	7
	2	6	3	0	1
	3	1	0	5	0

		0	1	2	3
B	0	0	1	2	8
	1	5	0	4	2
	2	6	0	0	1
	3	2	6	4	0

		0	1	2	3
C	0	84	18	36	20
	1	68	46	36	41
	2	17	12	28	54
	3	30	1	2	13

Multiplying matrices

- Matrix product: $C = A \times B$

- For $r = c = n$,

$$\begin{aligned} C[i, j] = & A[i, 0] \cdot B[0, j] + \\ & A[i, 1] \cdot B[1, j] + \\ & \dots + \\ & A[i, n-1] \cdot B[n-1, j] \end{aligned}$$

- More concisely, $C[i, j] = \sum_{k=0}^{n-1} A[i, k] \cdot B[k, j]$

- Don't require both A and B to be $n \times n$
 - Each row entry of A must have a matching column entry in B
 - If A is $m \times n$ and B is $n \times p$, $A \times B$ is $m \times p$

		0	1	2	3
	0	0	6	8	3
A	1	4	0	9	7
	2	6	3	0	1

		0	1	2
	0	0	1	2
B	1	5	0	4
	2	6	0	0
	3	2	6	4

		0	1	2
	0	84	18	36
C	1	68	46	36
	2	17	12	28

Transitive closure

- A is an adjacency matrix
 - $A[i,j] = 1$ if and only if there is a direct edge (path of length 1) from i to j

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$
- Algebra of boolean values

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$
- Algebra of boolean values
 - **True** is 1, **False** is 0

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$
- Algebra of boolean values
 - **True** is 1, **False** is 0
 - Logical **or** is represented by $+$:
 $0 + 0 = 0$, $0 + 1 = 1 + 0 = 1 + 1 = 1$

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$
- Algebra of boolean values
 - **True** is 1, **False** is 0
 - Logical **or** is represented by $+$:
 $0 + 0 = 0$, $0 + 1 = 1 + 0 = 1 + 1 = 1$
 - Logical **and** is represented by \times :
 $1 \times 1 = 1$, $0 \times 0 = 0 \times 1 = 1 \times 0 = 0$

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$
- Algebra of boolean values
 - True is 1, False is 0
 - Logical or is represented by $+$:
 $0 + 0 = 0$, $0 + 1 = 1 + 0 = 1 + 1 = 1$
 - Logical and is represented by \times :
 $1 \times 1 = 1$, $0 \times 0 = 0 \times 1 = 1 \times 0 = 0$

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or } (A[i, 1] \text{ and } A[1, j]) \text{ or } \dots (A[i, n-1] \text{ and } A[n-1, j])$

Transitive closure

- A is an adjacency matrix
 - $A[i, j] = 1$ if and only if there is a direct edge (path of length 1) from i to j
- A^2 represents paths of length 2
 - $A^2[i, j] = 1$ if, for some k , $A[i, k] = 1$ and $A[k, j] = 1$
- Algebra of boolean values
 - True is 1, False is 0
 - Logical or is represented by $+$:
 $0 + 0 = 0$, $0 + 1 = 1 + 0 = 1 + 1 = 1$
 - Logical and is represented by \times :
 $1 \times 1 = 1$, $0 \times 0 = 0 \times 1 = 1 \times 0 = 0$

	0	1	2	3	4	5
0	0	1	0	0	1	0
1	0	0	1	0	0	0
2	1	0	0	0	0	0
3	0	0	0	0	1	0
4	1	0	0	1	0	0
5	0	0	0	1	0	0

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or } (A[i, 1] \text{ and } A[1, j]) \text{ or } \dots (A[i, n-1] \text{ and } A[n-1, j])$
- $A^2[i, j] = A[i, 0] \times A[0, j] + A[i, 1] \times A[1, j] + \dots + A[i, n-1] \times A[n-1, j]$

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or}$
 $(A[i, 1] \text{ and } A[1, j]) \text{ or}$
 \dots
 $(A[i, n-1] \text{ and } A[n-1, j])$
- $A^2[i, j] = A[i, 0] \times A[0, j] +$
 $A[i, 1] \times A[1, j] +$
 $\dots +$
 $A[i, n-1] \times A[n-1, j]$

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or}$
 $(A[i, 1] \text{ and } A[1, j]) \text{ or}$
 \dots
 $(A[i, n-1] \text{ and } A[n-1, j])$
- $A^2[i, j] = A[i, 0] \times A[0, j] +$
 $A[i, 1] \times A[1, j] +$
 $\dots +$
 $A[i, n-1] \times A[n-1, j]$
- So $A^2[i, j] = \sum_{k=0}^{n-1} A[i, k] \times A[k, j]$

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or}$
 $(A[i, 1] \text{ and } A[1, j]) \text{ or}$
 \dots
 $(A[i, n-1] \text{ and } A[n-1, j])$
- $A^2[i, j] = A[i, 0] \times A[0, j] +$
 $A[i, 1] \times A[1, j] +$
 $\dots +$
 $A[i, n-1] \times A[n-1, j]$
- So $A^2[i, j] = \sum_{k=0}^{n-1} A[i, k] \times A[k, j]$
- In other words, $A^2 = A \times A$

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or}$
 $(A[i, 1] \text{ and } A[1, j]) \text{ or}$
 \dots
 $(A[i, n-1] \text{ and } A[n-1, j])$

- $A^2[i, j] = A[i, 0] \times A[0, j] +$
 $A[i, 1] \times A[1, j] +$
 $\dots +$
 $A[i, n-1] \times A[n-1, j]$

- So $A^2[i, j] = \sum_{k=0}^{n-1} A[i, k] \times A[k, j]$

- In other words, $A^2 = A \times A$

- Likewise, $A^3[i, j] = 1$ if, for some k ,
 $A^2[i, k] = 1$ and $A[k, j] = 1$

- Paths of length 3

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or}$
 $(A[i, 1] \text{ and } A[1, j]) \text{ or}$
 \dots
 $(A[i, n-1] \text{ and } A[n-1, j])$

- $A^2[i, j] = A[i, 0] \times A[0, j] +$
 $A[i, 1] \times A[1, j] +$
 $\dots +$
 $A[i, n-1] \times A[n-1, j]$

- So $A^2[i, j] = \sum_{k=0}^{n-1} A[i, k] \times A[k, j]$

- In other words, $A^2 = A \times A$

- Likewise, $A^3[i, j] = 1$ if, for some k ,
 $A^2[i, k] = 1$ and $A[k, j] = 1$

- Paths of length 3

- $A^3 = A^2 \times A$

Transitive closure

- $A^2[i,j] = (A[i,0] \text{ and } A[0,j]) \text{ or}$
 $(A[i,1] \text{ and } A[1,j]) \text{ or}$
 \dots
 $(A[i,n-1] \text{ and } A[n-1,j])$

- $A^2[i,j] = A[i,0] \times A[0,j] +$
 $A[i,1] \times A[1,j] +$
 $\dots +$
 $A[i,n-1] \times A[n-1,j]$

- So $A^2[i,j] = \sum_{k=0}^{n-1} A[i,k] \times A[k,j]$

- In other words, $A^2 = A \times A$

- Likewise, $A^3[i,j] = 1$ if, for some k ,
 $A^2[i,k] = 1$ and $A[k,j] = 1$

- Paths of length 3

- $A^3 = A^2 \times A = A \times A^2$

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or}$
 $(A[i, 1] \text{ and } A[1, j]) \text{ or}$
 \dots
 $(A[i, n-1] \text{ and } A[n-1, j])$

- $A^2[i, j] = A[i, 0] \times A[0, j] +$
 $A[i, 1] \times A[1, j] +$
 $\dots +$
 $A[i, n-1] \times A[n-1, j]$

- So $A^2[i, j] = \sum_{k=0}^{n-1} A[i, k] \times A[k, j]$

- In other words, $A^2 = A \times A$

- Likewise, $A^3[i, j] = 1$ if, for some k ,
 $A^2[i, k] = 1$ and $A[k, j] = 1$

- Paths of length 3

- $A^3 = A^2 \times A = A \times A^2$

- In general, $A^{\ell+1} = A^\ell \times A = A \times A^\ell$

- Paths of length $\ell+1$

Transitive closure

- $A^2[i, j] = (A[i, 0] \text{ and } A[0, j]) \text{ or } (A[i, 1] \text{ and } A[1, j]) \text{ or } \dots (A[i, n-1] \text{ and } A[n-1, j])$

- $A^2[i, j] = A[i, 0] \times A[0, j] + A[i, 1] \times A[1, j] + \dots + A[i, n-1] \times A[n-1, j]$

- So $A^2[i, j] = \sum_{k=0}^{n-1} A[i, k] \times A[k, j]$

- In other words, $A^2 = A \times A$

- Likewise, $A^3[i, j] = 1$ if, for some k , $A^2[i, k] = 1$ and $A[k, j] = 1$

- Paths of length 3

- $A^3 = A^2 \times A = A \times A^2$

- In general, $A^{\ell+1} = A^\ell \times A = A \times A^\ell$

- Paths of length $\ell+1$

- Finally, $A^+ = A + A^2 + \dots + A^{n-1}$

- $A^+[i, j] = 1$ if i, j connected by path of length 1 or path of length 2 or \dots or

- path of length $n-1$

Summary

- Matrix addition is defined elementwise
 - $C[i,j] = A[i,j] + B[i,j]$
 - A , B , C have same dimensions $r \times c$

Summary

- Matrix addition is defined elementwise
 - $C[i,j] = A[i,j] + B[i,j]$
 - A, B, C have same dimensions $r \times c$
- Matrix multiplication is a more complicated operation
 - Let A be $m \times n$ and B be $n \times p$
 - $$C[i,j] = \sum_{k=0}^{n-1} A[i,k] \times B[k,j]$$
 - C has dimensions $m \times p$

Summary

- Matrix addition is defined elementwise
 - $C[i,j] = A[i,j] + B[i,j]$
 - A, B, C have same dimensions $r \times c$
- Matrix multiplication is a more complicated operation
 - Let A be $m \times n$ and B be $n \times p$
 - $$C[i,j] = \sum_{k=0}^{n-1} A[i,k] \times B[k,j]$$
 - C has dimensions $m \times p$
- Using Boolean algebra, describe transitive closure using matrix multiplication
 - A , adjacency matrix, paths of length 1
 - $A^{\ell+1} = A^{\ell} \times A$, paths of length ℓ
 - Transitive closure, $A^+ = A + A^2 + \dots A^{n-1}$