
Integrated Diabetes Report (IDR) – insulin pen

Date: 14/02/2023 - 27/02/2023	CGM (% active): NightScout/xDrip (95%)	Insulin pen (avg. injections/day): 4.1
Patient: ClientId 1650	Sex: Male	DOB: *manual input*
Type: *manual input*	Duration: *manual input*	Physician: *manual input*

Time in ranges



Time above range 40% (goal <25%)
Time below range 3.4% (goal <4%)


Glucose statistics
Average glucose: 9.48 mmol/L
Glucose management indicator (GMI, goal <7%): 7.4%
Glucose variability (CV, goal ≤36%): 42.2%
Minimum glucose: 2.2 mmol/L (24/02/2023 08:36)
Maximum glucose: 22.3 mmol/L (17/02/2023 17:11)

Carbohydrate statistics
Average carb intake (daily): 233 g (46 entries)


Physical activity statistics
Average exercise (daily): 34 minutes
Average steps (daily): 25 573

Diabetes complications
Type: *manual input*
Duration: *manual input*


24-hour glucose percentile profile



Insulin by time of day



Insulin profile



Parameter	Calculated	Real	Unit
TAID - Total average insulin dose	50.0	65.6	IU/day
ABaD - Average basal dose	25.0	29.7	IU/day
ABoD - Average bolus dose	29.1	35.9	IU/day
ICR - Insulin-to-carb ratio	10.0	6.6	g/IU
CD - Correction dose	5.8	6.3	IU/day
ICF - Insulin correction factor	2.0	0.5 (n = 5)	mmol/L per IU
IDF - Insulin dose for food	23.3	29.6	IU/day

Data notes

- CGM readings: 3839 of 4032 expected (95.2%).
- Insulin: 13 basal doses (usual dose 32 IU; 8 logged with a positive sign and reclassified as basal), 44 boluses (36 meal, 8 correction).
- Rule: negative amount = basal; positive amount equal to the usual basal dose (±15%) = basal; other positive = bolus. A bolus within ±45 min of a carb entry = meal bolus.
- 10 carb entries have no bolus within ±45 min.
- All glucose values are in mmol/L (1 mmol/L = 18 mg/dL). GMI = 3.31 + 0.43056 × mean glucose (mmol/L).
- Minimum or maximum glucose is at the CGM measuring limit (about 2.2 or 22.2 mmol/L), so the true value may have been lower or higher.
- Real values (daily averages): ABaD = basal insulin, IDF = meal-bolus insulin, CD = correction-bolus insulin, ABoD = IDF + CD, TAID = ABaD + ABoD. Real ICR = meal carbs / meal insulin.
- Real ICF = median glucose drop (mmol/L) per IU, 3 h after an isolated correction bolus: no carb entry (from 30 min before), no other bolus and fewer than 10 exercise minutes (≥100 steps/min), and a CGM reading at both ends. Of 8 correction boluses, 5 were usable (excluded: carbs 2, exercise 1); at least 5 are needed, otherwise the report shows «insufficient data».
- Calculated values (paper formulas, glucose in mmol/L): TDD = factor (0.5 IU/kg) × weight; basal = 50% of TDD; ICR = 500/TDD; ICF = 100/TDD (the mmol/L form of the 1800 rule); IDF = daily carbs / ICR; CD = (BG at each bolus - 5.6 mmol/L) / ICF, summed per day; ABoD = IDF + CD.
- Calculated TAID is the weight-based TDD; calculated ABoD is based on the actual carbohydrate intake and glucose, so calculated ABaD + ABoD may differ from TAID.
- TDD from body weight 100 kg (entered manually).
- Physical activity from minute steps (Xiaomi Mi Band): averages over the 12 days with at least 90% of minutes recorded (tracker coverage 96%). Exercise = minutes with at least 100 steps per minute.