

Math HW6

Matthew Vanaman

04-08-2019


6.2, 6.3a, 6.4a, 6.5a, 6.6

6.1


(a)

$$f(x) = \int_{-1.96}^{1.96} \frac{1}{\sqrt{2\pi}} e^{-0.5^2} dx$$


Left




Right



Midpoint



Trapezoidal



(b)

Left

Answer:

$$A \approx \frac{0.392}{\sqrt{2\pi}} \left(\sum_{i=0}^9 e^{-0.5^2 \left(0.392i - 1.96 \right)^2} \right).$$

Work:

The left Riemann Sum formula is:

$$A \approx \sum_{i=0}^{n-1} f\left(a + \frac{(b-a)i}{n}\right).$$

We want to find the sums for 10 partitions between -1.96 and 1.96. Which means that $n = 10$, $a = -1.96$, and $b = 1.96$. We also plug in the pdf for f . The Riemann Sum formula gets substituted in for x in $f(x)$:

$$\begin{aligned} A &\approx \sum_{i=0}^{n-1} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(a + \frac{(b-a)i}{n} \right)^2}, \\ &\approx \sum_{i=0}^{10-1} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-1.96 + \frac{(1.96 - (-1.96))i}{10} \right)^2}, \\ &\approx \sum_{i=0}^9 \frac{1}{\sqrt{2\pi}} e^{-0.5 \left((0.392i - 1.96)0.392 \right)^2}, \end{aligned}$$

0.392 does not depend on index, and neither does the constant $\frac{1}{\sqrt{(2\pi)}}$:

$$\approx \frac{1}{\sqrt{2\pi}} (0.392) \left(\sum_{i=0}^9 e^{-0.5 \left((0.392i - 1.96)0.392 \right)^2} \right),$$

$$\approx \frac{0.392}{\sqrt{2\pi}} \left(\sum_{i=0}^9 e^{-0.5^2 \left(0.392i - 1.96 \right)^2} \right).$$

```
# calculation:
(1.96 + 1.96) / 10
```

```
[1] 0.392
```

Right:

Answer:

$$A \approx \frac{0.392}{\sqrt{2\pi}} \left(\sum_{i=1}^{10} e^{-0.5^2 \left(0.392i - 1.96 \right)^2} \right).$$

Work:

Do the same thing here except the index is from 1 to n instead of from 0 to $n - 1$.

$$\begin{aligned} A &\approx \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-1.96 + \frac{1.96 - (-1.96)}{10} i \left(\frac{1.96 - (-1.96)}{10} \right) \right)^2} \right), \\ &\approx \sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left((0.392i - 1.96) 0.392 \right)^2}, \\ &\approx \frac{0.392}{\sqrt{2\pi}} \left(\sum_{i=1}^{10} e^{-0.5^2 \left(0.392i - 1.96 \right)^2} \right). \end{aligned}$$

Midpoint:

Answer:

$$A \approx 0.392 \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-2.16 + 0.392i \right)^2} \right).$$

Work:

$$A \approx \sum_{i=0}^{n-1} f \left(a + \frac{(b-a)(i+0.5)}{n} \right).$$

We want to find the sums for 10 partitions between -1.96 and 1.96. Which means that $n = 10$, $a = -1.96$, and $b = 1.96$. We also plug in the pdf for f . The Riemann Sum formula gets substituted in for x in $f(x)$:

$$\begin{aligned} A &\approx \left(\sum_{i=0}^{n-1} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(a + \frac{(b-a)(i+0.5)}{n} \left(\frac{b-a}{n} \right) \right)^2} \right), \\ &\approx \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-1.96 + \frac{1.96 - (-1.96)}{10} \left(\frac{(i+0.5)}{1} \right) \left(\frac{1.96 - (-1.96)}{10} \right) \right)^2} \right), \end{aligned}$$

$$\begin{aligned}
&\approx \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-1.96 + 0.392(i-0.5)(0.392) \right)^2} \right), \\
&\approx 0.392 \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-1.96 + 0.392i - 0.196 \right)^2} \right), \\
&\approx 0.392 \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-2.16 + 0.392i \right)^2} \right), \\
&\approx 0.392 \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-2.16 + 0.392i \right)^2} \right).
\end{aligned}$$

Calculations

0.392 * 0.5

[1] 0.196

-1.96 - 0.196

[1] -2.16

Trapezoidal

Answer:

$$A \approx \left(\frac{0.196}{\sqrt{2\pi}} \right) \sum_{i=0}^9 \frac{e^{-0.5(0.392i-1.96)^2} + e^{-0.5(0.392i-0.564)^2}}{2}.$$

Work:

$$\begin{aligned}
A &\approx \sum_{i=0}^{n-1} \frac{b-a}{n} \frac{\frac{1}{\sqrt{2\pi}} e^{\left(a + \frac{(b-a)i}{n} \right)^2}}{2} + \frac{1}{\sqrt{2\pi}} e^{\left(a + \frac{(b-a)i+1}{n} \right)^2} \frac{b-a}{n}, \\
&\approx \sum_{i=0}^{10-1} \frac{1.96 - (-1.96)}{10} \frac{\frac{1}{\sqrt{2\pi}} e^{\left((-1.96) + \frac{(1.96 - (-1.96))i}{10} \right)^2}}{2} + \frac{1}{\sqrt{2\pi}} e^{\left((-1.96) + \frac{(1.96 - (-1.96))i+1}{10} \right)^2} \frac{1}{2}, \\
&\approx \sum_{i=0}^9 (0.392) \frac{\frac{1}{\sqrt{2\pi}} e^{-0.5(0.392i-1.96)^2} + \frac{1}{\sqrt{2\pi}} e^{-0.5(0.392i-0.564)^2}}{2}, \\
&\approx \frac{0.392}{2} \left(\frac{1}{\sqrt{2\pi}} \right) \sum_{i=0}^9 \frac{e^{-0.5(0.392i-1.96)^2} + e^{-0.5(0.392i-0.564)^2}}{2}, \\
&\approx \left(\frac{0.196}{\sqrt{2\pi}} \right) \sum_{i=0}^9 \frac{e^{-0.5(0.392i-1.96)^2} + e^{-0.5(0.392i-0.564)^2}}{2}.
\end{aligned}$$

0.392 / 2

[1] 0.196

(c)

Left

Answer:

0.947

Work:

Just plug in to x each level of n from 0 - 9:

$$A \approx \frac{0.392}{\sqrt{2\pi}} \left(\sum_{i=0}^9 e^{-0.5^2 \left(\frac{0.392i - 1.96}{0.392} \right)^2} \right).$$

```
e <- exp(1) # get euler's number
(0.392 / (sqrt(2 * pi))) * (e^(-0.5 * ((0.392 * 0) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 2) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 3) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 4) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 5) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 6) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 7) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 8) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 9) - 1.96)^2)
)

[1] 0.947
```

Right

Answer:

0.947

** Work:**

Same thing, except runs from 1-10 instead of 0-9. Not surprising that it gives you the same answer:

```
e <- exp(1) # get euler's number
(0.392 / (sqrt(2 * pi))) * (e^(-0.5 * ((0.392 * 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 2) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 3) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 4) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 5) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 6) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 7) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 8) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 9) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 10) - 1.96)^2)
)

[1] 0.947
```

Midpoint

Answer:

0.951

Work:

$$A \approx 0.392 \left(\sum_{i=1}^{10} \frac{1}{\sqrt{2\pi}} e^{-0.5 \left(-2.16 + 0.392i \right)^2} \right).$$

```
e <- exp(1) # get euler's number
(0.392 / (sqrt(2 * pi))) * (e^(-0.5 * ((0.392 * 1) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 2) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 3) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 4) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 5) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 6) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 7) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 8) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 9) - 2.16)^2) +
  e^(-0.5 * ((0.392 * 10) - 2.16)^2)
)
```

[1] 0.951

Trapezoidal

Answer:

0.947

Work:

$$A \approx \left(\frac{0.196}{\sqrt{2\pi}} \right) \sum_{i=0}^9 \frac{e^{-0.5(0.392i-1.96)^2} + e^{-0.5(0.392i-0.564)^2}}{2}.$$

```
e <- exp(1) # get euler's number
(0.196 / (sqrt(2 * pi))) * (e^(-0.5 * ((0.392 * 0) - 1.96)^2) +
  e^(-0.5 * (0.392 * (0 + 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 1) - 1.96)^2) +
  e^(-0.5 * (0.392 * (1 + 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 2) - 1.96)^2) +
  e^(-0.5 * (0.392 * (2 + 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 3) - 1.96)^2) +
  e^(-0.5 * (0.392 * (3 + 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 4) - 1.96)^2) +
  e^(-0.5 * (0.392 * (4 + 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 5) - 1.96)^2) +
  e^(-0.5 * (0.392 * (5 + 1) - 1.96)^2) +
  e^(-0.5 * ((0.392 * 6) - 1.96)^2) +
  e^(-0.5 * (0.392 * (6 + 1) - 1.96)^2)
)
```

```

e^(-0.5 * (0.392 * (6 + 1) - 1.96)^2) +
e^(-0.5 * ((0.392 * 7) - 1.96)^2) +
e^(-0.5 * (0.392 * (7 + 1) - 1.96)^2) +
e^(-0.5 * ((0.392 * 8) - 1.96)^2) +
e^(-0.5 * (0.392 * (8 + 1) - 1.96)^2) +
e^(-0.5 * ((0.392 * 9) - 1.96)^2) +
e^(-0.5 * (0.392 * (9 + 1) - 1.96)^2)
)

```

```
[1] 0.947
```

6.2

(a)

Answer:

Work:

$$\int x^{100} + 3e^x - 7(4^x)dx,$$