

Solutions to Sheet 7

Exercise 1

Let $A \rightarrow B$ be a homomorphism of rings, let M be an A -module and let N be a B -module.

1. Show that the map

$$\mathrm{Hom}_A(M, N) \rightarrow \mathrm{Hom}_B(B \otimes_A M, N), \quad \varphi \mapsto (b \otimes m \mapsto b\varphi(m))$$

is a well-defined isomorphism.

2. Show that the map

$$M \otimes_A N \rightarrow (M \otimes_A B) \otimes_B N, \quad m \otimes n \mapsto (m \otimes 1) \otimes n$$

is a well-defined isomorphism.

3. Deduce that $S^{-1}M_1 \otimes_A S^{-1}M_2 \cong S^{-1}M_1 \otimes_{S^{-1}A} S^{-1}M_2$ for two A -modules M_1, M_2 and a multiplicative subset $S \subset A$.

Solution.

- 1.

Exercise 2

Let A be a ring. We define the *support* of an A -module M as $\mathrm{Supp}(M) := \{\mathfrak{p} \in \mathrm{Spec}(A) \mid M_{\mathfrak{p}} \neq 0\}$.

1. Assume M is finitely generated. Show that $\mathrm{Supp}(M) = \{\mathfrak{p} \in \mathrm{Spec}(A) \mid M \otimes_A k(\mathfrak{p}) \neq 0\}$, where $k(\mathfrak{p}) = \mathrm{Quot}(A/\mathfrak{p})$.
2. Assume M, N are finitely generated A -modules. Show $\mathrm{Supp}(M \otimes_A N) = \mathrm{Supp}(M) \cap \mathrm{Supp}(N)$.

Solution.

1. We will show that $M_{\mathfrak{p}} \neq 0$ if and only if $M \otimes_A k(\mathfrak{p}) \neq 0$. The map $A \rightarrow k(\mathfrak{p})$ factors through the map $A_{\mathfrak{p}} \rightarrow k(\mathfrak{p})$, and we find $M \otimes_A k(\mathfrak{p}) = M_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$, this directly gives the implication $M \otimes_A k(\mathfrak{p}) \neq 0 \implies M_{\mathfrak{p}} \neq 0$.

For the other direction, we use Nakayama's Lemma. It (or at least one version of it) states that if $N \neq 0$ is a finitely generated module over a local ring B with maximal ideal I , we have $IN \neq N$. In our situation, if we assume $M_{\mathfrak{p}} \neq 0$, Nakayama says

$$M_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}) \cong M_{\mathfrak{p}}/\mathfrak{p}M_{\mathfrak{p}} \neq 0.$$

Done.

2. We'll show that $(M \otimes_A N) \otimes k(\mathfrak{p}) \neq 0$ if and only if $M \otimes_A k(\mathfrak{p}) \neq 0$ and $N \otimes_A k(\mathfrak{p}) \neq 0$. Exercise 1.2 gives the isomorphism

$$(M \otimes_A k(\mathfrak{p})) \otimes_{k(\mathfrak{p})} (N \otimes_A k(\mathfrak{p})) \cong M \otimes_A (N \otimes_A k(\mathfrak{p})) \cong (M \otimes_A N) \otimes_A k(\mathfrak{p}).$$

From here we can directly check the desired equivalence.

Exercise 3

Let A be a ring, let $S \subset A$ be a multiplicative subset and let M, N be A -modules.

1. Assume that M is finitely presented A -module. Show that the map

$$S^{-1} \operatorname{Hom}_A(M, N) \rightarrow \operatorname{Hom}_{S^{-1}A}(S^{-1}M, S^{-1}N), \quad \varphi/s \mapsto (m/t \mapsto \varphi(m)/st)$$

is a well-defined isomorphism.

2. Construct a counterexample to the above if M is only assumed to be finitely generated.

Solution.

1. First, note that we always (without hypothesis on M) obtain such a map. This directly follows (for example) from exercise 1.1 with $B = S^{-1}A$.

Now we have to show that this is an isomorphism if M is finitely presented. As usual, we write M as part of a short exact sequence

$$0 \rightarrow A^m \rightarrow A^n \rightarrow M \rightarrow 0.$$

Now we use that $\operatorname{Hom}_A(-, N)$ is right-exact. Hence applying $\operatorname{Hom}_A(-, N)$ yields an exact sequence

$$0 \rightarrow 0 \rightarrow \operatorname{Hom}_A(M, N) \rightarrow \operatorname{Hom}_A(A^n, N) \cong N^n \rightarrow \operatorname{Hom}_A(A^m, N) \cong A^m.$$

Localizing at S is exact, so we obtain

$$0 \rightarrow 0 \rightarrow S^{-1} \operatorname{Hom}_A(M, N) \rightarrow (S^{-1}N)^n \rightarrow (S^{-1}N)^m.$$

Similarly, we can localize at S first and then apply $\operatorname{Hom}_{S^{-1}A}(S^{-1}(-), S^{-1}N)$, which yields the exact sequence

$$0 \rightarrow 0 \rightarrow \operatorname{Hom}_{S^{-1}A}(S^{-1}M, S^{-1}N) \rightarrow (S^{-1}N)^n \rightarrow (S^{-1}N)^m.$$

Now we can use the 5-lemma again!

$$\begin{array}{ccccccc} 0 & \longrightarrow & 0 & \longrightarrow & S^{-1} \operatorname{Hom}_A(M, N) & \longrightarrow & (S^{-1}A)^n \longrightarrow (S^{-1}A)^m \\ \parallel & & \parallel & & \downarrow \cdot \text{iso} & & \parallel & \parallel \\ 0 & \longrightarrow & 0 & \longrightarrow & \operatorname{Hom}_{S^{-1}A}(S^{-1}M, S^{-1}N) & \longrightarrow & (S^{-1}A)^n \longrightarrow (S^{-1}A)^m \end{array}$$

2. A standard example seems to be the following. Let $A = k[x, y_1, y_2, \dots]$ be the polynomial ring in variables indexed by \mathbb{N} . Let $M = A/(y_1, y_2, \dots)$, $N = A/(xy_1, x^2y_2, \dots)$ and $S = \{1, x, x^2, \dots\}$. Now let's compare both sides of the morphism. Note that M is generated by 1, so that any A -linear morphism $\varphi : M \rightarrow N$ is uniquely determined by the value of $\varphi(1) \in N$. Now we have $0 = y_1\varphi(1) = y_2\varphi(1) = \dots$, which shows that any lift $\tilde{\varphi}(1) \in R$ is infinitely divisible by x , hence $\varphi(1) = 0$. On the left hand side, we find that $S^{-1}M \cong S^{-1}N \cong k[x^{\pm 1}]$, so there are many $S^{-1}A$ -linear morphisms $S^{-1}M \rightarrow S^{-1}N$.

Exercise 4

Let A be a principal domain and let $f \in A \setminus \{0\}$ be a non-unit. Show that the $A[T]$ -module $(f, T) \subset A[T]$ is not flat.

Solution. Consider the map given by multiplication with f , which we will denote as $\varphi : A \rightarrow A$. It is injective. Note that $A \cong A[T]/(T)$. We want to show that $(f, T) \otimes_{A[T]} A$ is not injective, showing that (f, T) is not flat. We have an isomorphism (of $A[T]$ -modules)

$$(f, T) \otimes_{A[T]} A \cong (f, T)/T(f, T),$$

and $(f, T) \otimes \varphi$ corresponds to the endomorphism given by multiplication with f under this identification. Now, $T \neq 0$ in $(f, T)/T(f, T)$, but $fT = \varphi(T) = 0$.