

Solutions to Sheet 12

Exercise 1

Let I be an ideal of a number field K . Show that there is a finite field extension L of K such that $I\mathcal{O}_L$ is a principal ideal of \mathcal{O}_L .

Solution. By finiteness of $\text{Cl}(K)$ there is some integer m such that $[I]^m = [I^m] = [(1)] \in \text{Cl}(K)$, i.e., $I^m = (\alpha)$ is a principal ideal. We put $L = K(\alpha^{1/m})$. Now $\alpha^{1/m} \in \mathcal{O}_L$, and we have

$$(I\mathcal{O}_L)^m = I^m\mathcal{O}_L = \alpha\mathcal{O}_L = (\alpha^{1/m})^m\mathcal{O}_L.$$

After decomposing $I\mathcal{O}_L$ and $\alpha^{1/m}\mathcal{O}_L$ into prime factors, we see that this equation implies $I\mathcal{O}_L = \alpha^{1/m}\mathcal{O}_L$.

Exercise 2

Let $K = \mathbb{Q}(\sqrt{2}, \sqrt{3})$ and set

$$\Gamma = \{(1 + \sqrt{2})^i(2 + \sqrt{3})^j(\sqrt{2} + \sqrt{3})^k \mid i, j, k \in \mathbb{Z}\}.$$

Show that Γ is a subgroup of \mathcal{O}_K^\times and that $[\mathcal{O}_K^\times : \Gamma] < \infty$.

Solution. Write u, v, w for the respective factors, so that $\Gamma = u^{\mathbb{Z}}v^{\mathbb{Z}}w^{\mathbb{Z}}$. Note that $N_{K/\mathbb{Q}}(u) = N_{K/\mathbb{Q}}(v) = 1$ and $N_{K/\mathbb{Q}}(w) = -1$, so that indeed, u, v, w are units and Γ is a subgroup of \mathcal{O}_K^\times . One quickly verifies that K is totally real. Indeed, it is Galois and there is an embedding $K \hookrightarrow \mathbb{R}$ (now all other embeddings are obtained by shifting with elements in the Galois group). Hence, by Dirichlet's unit theorem,

$$\mathcal{O}_K^\times \cong \mu(K) \times \mathbb{Z}^{r+s-1} = \pm 1 \times \mathbb{Z}^3.$$

On the other hand, Γ is free of rank 3. Indeed, $u \in \mathbb{Q}(\sqrt{2})^\times$, $v \in \mathbb{Q}(\sqrt{3})^\times$ and $w \in \mathbb{Q}(\sqrt{2}, \sqrt{3})^\times \setminus (\mathbb{Q}(\sqrt{2})^\times \cup \mathbb{Q}(\sqrt{3})^\times)$. Let $\mathcal{O}_{K, > 0}^\times \cong \mathbb{Z}^3$ be the (free) group of positive units $K \hookrightarrow \mathbb{R}$. Now Γ is a free subgroup of full rank in this group, and in particular, it has finite index. Inclusion $\mathcal{O}_{K, > 0}^\times \hookrightarrow \mathcal{O}_K^\times$ also has finite index, hence $\Gamma \hookrightarrow \mathcal{O}_K^\times$ has finite index.

Exercise 3

Let K be a totally real number field, i.e., one that has **only** real embeddings. Let

$$T \subset \text{Hom}(K, \mathbb{R}) = \{\tau : K \rightarrow \mathbb{R} \mid \tau \text{ is a field homomorphism}\}$$

be a proper non-mepty subset. Show that there exists $u \in \mathcal{O}_K^\times$ such that $0 < \tau(u) < 1$ for $\tau \in T$ and $\tau(u) > 1$ for $\tau \in \text{Hom}(K, \mathbb{R}) \setminus T$.

Solution. Let $\sigma_1, \dots, \sigma_r : K \rightarrow \mathbb{R}$ be the real embeddings of K (in our case $r = n = [K : \mathbb{Q}]$). From the proof of Dirichlet's unit theorem, we know that the map

$$\mathcal{L} : \mathcal{O}_K^\times \rightarrow \mathbb{R}^r, \quad u \mapsto (\log |\sigma_1(u)|, \dots, \log |\sigma_r(u)|)$$

is a group homomorphism from \mathcal{O}_K to \mathbb{R}^{r-1} . It's image lies in the sub vector space V given by

$$V = \left\{ (x_1, \dots, x_r)^t \in \mathbb{R}^r \mid \sum_{i=1}^r x_i = 0 \right\},$$

and its kernel is given by $\mu(K)$, the group of roots of unity in K (in our case this is $\cong \{\pm 1\}$). Also, the image $\mathcal{L}(\mathcal{O}_K^\times)$ has full rank in V , i.e., $\mathcal{L}(\mathcal{O}_K^\times) \otimes_{\mathbb{Z}} \mathbb{R} \cong V$ (it is a lattice in V).

Without loss of generality we can assume that $T = \{\sigma_1, \dots, \sigma_q\}$ for some $1 \leq q < r$. Let $Q \subset \mathbb{R}^r$ be the quadrant given by

$$Q = \{(x_1, \dots, x_r)^t \in \mathbb{R}^r \mid x_i < 0 \text{ for } i = 1, \dots, q \text{ and } x_i > 0 \text{ for } i = q+1, \dots, r\}.$$

The intersection $Q \cap V$ is non-empty by construction, and one readily verifies that there is a point $x \in Q \cap \mathcal{L}(\mathcal{O}_K^\times)$. Now choose some preimage $u \in \mathcal{O}_K^\times$ of x . As u satisfies $|\sigma_i(u)| < 1$ for $1 \leq i \leq q$ and $|\sigma_i(u)| > 1$ for $q < i \leq r$, the element u^2 satisfies all constraints.

Exercise 4

Let K be a number field, let I be a non-zero ideal of \mathcal{O}_K and let $C \in \text{Cl}(K)$. Use theorem 5.3 to show that there exists a non-zero ideal J of \mathcal{O}_K such that $I + J = \mathcal{O}_K$ and $C = [J]$.