

RADI.jl

Olivier Sulpis and Matthew P. Humphreys

March 26, 2020

Abstract

RADI.jl is a Julia implementation of RADI: the 1-D Reaction-Advection-Diffusion-Irrigation Diagenetic Sediment Module. Here, we define the variables and the equations used in this implementation. So far, only one solute (dissolved oxygen) and one solid (particulate organic carbon) are included and documented here.

1 Parameters

1.1 Time

Time units are always in years.

- T (`stoptime` in a) is the total time that the model runs for.
- dt (`interval` in a) is the time resolution (i.e. the interval between each timestep).
- t (`timesteps` in a) refers to the array of modelled timepoints.

The model therefore runs from time 0 to T in intervals of dt .

1.2 Sediment column

1.2.1 Structure

Depth units are always in metres.

- Z (`z_max` in m) is the total height of the sediment column being modelled.
- dz (`z_res` in m) is the depth resolution (i.e. the height of each model layer).
- z (`depths` in m) refers to the array of modelled depths within the sediment.

The model layers are therefore at depths within the sediment from 0 to Z in increments of dz , where 0 represents the interface between the surface sediment and overlying seawater.

1.2.2 Overlying water

Properties of the overlying water can be set by the user:

- $[O_2]_w$ (`oxy_w` in $\text{mol} \cdot \text{m}^{-3}$) is the seawater dissolved oxygen concentration.
- F_c (`Foc` in $\text{mol} \cdot \text{m}^{-1} \cdot \text{a}^{-1}$) is the flux of particulate organic carbon arriving at the seafloor.
- δ (`dbl` in m) is the diffusive boundary layer thickness.
- T_w (`T` in $^{\circ}\text{C}$) is the water temperature.

Note that T is also used for the total model runtime...

1.2.3 Sediment properties

In the current set-up of RADI.jl, the porosity parameters ϕ_{∞} and ϕ_0 are set to 0.74 and 0.85 respectively (`phiInf` and `phi0`, both dimensionless), and $\beta = 33 \text{ m}^{-1}$ (`beta`). These values were obtained by fitting real data from station 7, mooring 3 of cruise NBP98-2 [Sayles et al., 2001]. **Should these (ϕ_0 , ϕ_{∞} and β) be inputs that the user can easily adjust?**

The porosity profile (ϕ , `phi`, dimensionless) is parameterised following Boudreau [1996b]:

$$\phi = \phi_{\infty} + (\phi_0 - \phi_{\infty}) \exp(-\beta z) \quad (1)$$

The corresponding “solid porosity” (ϕ_s , `phiS`, dimensionless) is:

$$\phi_s = 1 - \phi \quad (2)$$

RADI.jl also uses the related convenience variable `phiS_phi` = ϕ_s/ϕ .

Following Boudreau [1996a], the sediment tortuosity (θ , `tort`, dimensionless) is:

$$\theta = \sqrt{1 - 2 \log \phi} \quad (3)$$

RADI.jl also uses the related convenience variable `tort2` = θ^2 .

Following Archer et al. [2002], the surface sediment bioturbation coefficient B_0 (`D_bio_0` in $\text{m}^2 \cdot \text{a}^{-1}$) is:

$$B_0 = (0.0232 \cdot 10^{-4})(F_c \cdot 10^2)^{0.85} \quad (4)$$

Eq. (4): where do the 10^{-4} and 10^2 multipliers come from? These are in the code but not in the documentation.

The bioturbation coefficient propagates down through the sediment as B_z (`D_bio` in $\text{m}^2 \cdot \text{a}^{-1}$):

$$B_z = B_0 \exp(-z/\lambda_b)[O_2]_w/([O_2]_w + 0.02 \text{ mol} \cdot \text{m}^{-3}) \quad (5)$$

where λ_b (`lambda_b` in m) is the characteristic depth of 0.08 m, following Sayles et al. [2001]. **Where does the magic 0.02 number come from?**

The rate constant for organic matter degradation (k_z , `krefractory` in a^{-1}) is [Archer et al., 2002]:

$$k_z = 80.25 B_0 \exp(-z) \quad (6)$$

2 Variables

2.1 Porewater solutes

Within the sediment porewaters:

- $[O_2]$ (oxy in $\text{mol} \cdot \text{m}^{-3}$) is the dissolved oxygen concentration.

2.2 Solids

Within the sediment itself:

- $[POC]$ (poc in $\text{mol} \cdot \text{m}^{-3}$) is the particulate organic carbon concentration.

3 Master equation

For each modelled variable v at time t and depth z :

$$v_{(t+dt),z} = v_{t,z} + [R(v_{t,z}) + A(v_{t,z}) + D(v_{t,z}) + I(v_{t,z})] \cdot dt \quad (7)$$

where:

- $R(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to **reactions**.
- $A(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to **advection**.
- $D(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to **diffusion**.
- $I(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to **irrigation**.

4 Reaction

Biogeochemical reactions operate on solutes and solids throughout the entire sediment column, including the very top and bottom layers. $R(v_t)$ is the net rate at which v is being consumed (negative R) or created (positive R) by these reactions.

4.1 Organic matter degradation

Organic matter degradation affects dissolved oxygen and particulate organic carbon:

$$R(POC) = -k_z[POC] \quad (8)$$

$$R(O_2) = r(POC) \cdot \phi_s / \phi \quad (9)$$

where the rate constant k_z was defined in Eq. (6), and porosity coefficients ϕ and ϕ_s in Eqs. (1) and (2) respectively. **Eq. (9): should there not be a photosynthetic quotient (C:O₂ ratio) in here?**

5 Advection

By "advection" do we mean things literally moving around within a static sediment (i.e. porewater is advecting relative to the centre of the Earth), or is it that things to move relatively speaking because the sediment surface moves up as new material settles onto it (i.e. porewater is static relative to the centre of the Earth, but advecting relative to the sediment-water interface)?

Advection is modelled differently (1) at the sediment-water interface (i.e. where $z = 0$), (2) within the sediment ($0 < z < Z$), and (3) at the bottom of the sediment ($z = Z$).

5.1 Advection at the sediment-water interface

At sediment depth $z = 0$ only, for solutes (e.g. O_2):

$$A(v_0) = u_0(v_w - v_0)\theta^2/\delta \quad (10)$$

and for solids (e.g. POC):

$$A(v_0) = w_0(F_c - \phi_s w_0 v_0)/(B_0 \phi_s) \quad (11)$$

5.2 Advection within the sediment

At sediment depth z , where $0 < z < Z$, for solutes (e.g. O_2):

$$A(v_z) = -[u_z - (\theta^2 d\phi/\phi - d\{\theta^2\})d(v_z)/\theta^2][v_{(z+dz)} - v_{(z-dz)}]/2dz \quad (12)$$

where $d(v_z)$ is the relevant diffusion coefficient given in section 6.

For solids (e.g. POC):

$$A(v_z) = -[(1 - \sigma_z)v_{(z+dz)} + 2\sigma_z v_z - (1 + \sigma_z)v_{(z-dz)}] \cdot [w_z - dB_z - d\phi_s B_z/\phi_s]/2dz \quad (13)$$

where σ_z (**sigma**) is given by:

$$\sigma_z = 1/\tanh(P_{e_h}) - 1/P_{e_h} \quad (14)$$

in which P_{e_h} (**Peh**) is:

$$P_{e_h} = w_z dz/(2B_z) \quad (15)$$

5.3 Advection at the bottom of the sediment

Solutes (e.g. O_2) are not affected by advection at the bottom of the sediment.

At sediment depth $z = Z$ only, and for solids only (e.g. POC):

$$A(v_Z) = -(w_Z - dB_Z - d\phi_s B_Z/\phi_s)(v_Z - v_{(Z-dz)})\sigma_Z/dz \quad (16)$$

6 Diffusion

Diffusion is modelled differently (1) at the sediment-water interface (i.e. where $z = 0$), (2) within the sediment ($0 < z < Z$), and (3) at the bottom of the sediment ($z = Z$).

Diffusion is controlled by each variable's diffusion coefficient (generically $d(v)$, `D_var` in $\text{m}^2 \cdot \text{s}^{-1}$).

For dissolved oxygen:

$$d(\text{O}_2) = (0.034862 + 0.001409T_w)/\theta^2 \quad (17)$$

where the temperature function is from Li and Gregory [1974], and θ is the tortuosity from Eq. (3).

For particulate organic carbon:

$$d(\text{POC}) = B_z \quad (18)$$

where B_z was defined in Eq. (5).

6.1 Diffusion at the sediment-water interface

At sediment depth $z = 0$ only, for solutes (e.g. O_2):

$$D(v_0) = 2d(v_z) \cdot [v_{dz} - v_0 + \theta_0^2 dz(v_w - v_0)/\delta]/(dz)^2 \quad (19)$$

where θ_0 is the tortuosity at the sediment surface from Eq. (3). `RADI.jl` uses the convenience variable $\text{TR} = 2\theta_0^2 dz/\delta$ to simplify Eq. (19).

Is the factor of 2 in Eq. (19) above incorrect? It doubles the upwards diffusive flux relative to that within the sediment. See also similar note in section 6.3 below.

For solids (e.g. POC):

$$D(v_0) = 2d(v_z) \cdot [v_{dz} - v_0 + (F_c - \phi_s w_0 v_0)/(B_0 \phi_s)]/(dz)^2 \quad (20)$$

6.2 Diffusion within the sediment

At sediment depth z , where $0 < z < Z$, for both solutes and solids:

$$D(v_z) = d(v_z) \cdot (v_{(z+dz)} - 2v_z + v_{(z-dz)})/(dz)^2 \quad (21)$$

where $d(v_z)$ is the relevant diffusion coefficient.

6.3 Diffusion at the bottom of the sediment

At sediment depth $z = Z$ only, for both solutes and solids:

$$D(v_Z) = 2d(v_z) \cdot (-v_Z + v_{(Z-dz)})/(dz)^2 \quad (22)$$

Is the factor of 2 here incorrect? Without it, this equation would be the same as for 'within the sediment', but with no diffusive flux coming up from below. With it, this assumes that whatever diffusive flux comes from above, also comes up from below (i.e. the flux from above is doubled). See also similar note in section 6.1 above.

7 Irrigation

Irrigation only affects the solutes (e.g. O_2), not the solids (e.g. POC). Its effect is consistent throughout the sediment:

$$I(v_z) = \alpha_z (v_w - v_z) \quad (23)$$

The coefficient α_z (**alpha** in a^{-1}) at the sediment-water interface (i.e. α_0 , **alpha_0**) is, following Archer et al. [2002]:

$$\alpha_0 = 11 \left[\text{atan} \left(\frac{5F_c \cdot 10^2 - 400}{400\pi} \right) + 0.5 \right] - 0.9 + \frac{20[O_2]_w}{[O_2]_w + 0.01} \cdot \frac{F_c \cdot 10^2}{F_c \cdot 10^2 + 30} \cdot \exp \left(\frac{-[O_2]_w}{0.01} \right) \quad (24)$$

Where do all these magic numbers come from?

Within the sediment itself:

$$\alpha_z = \alpha_0 \exp[-(z/\lambda_i)^2] \quad (25)$$

where $\lambda_i = 0.05$ m (**lambda_i**) is the **somewhat mysterious** characteristic depth of Archer et al. [2002].

References

- D. E. Archer, J. L. Morford, and S. R. Emerson. A model of suboxic sedimentary diagenesis suitable for automatic tuning and gridded global domains. *Global Biogeochem. Cy.*, 16(1):17–1–17–21, 2002. doi: 10.1029/2000GB001288.
- B. P. Boudreau. The diffusive tortuosity of fine-grained unlithified sediments. *Geochim. Cosmochim. Acta*, 60(16):3139–3142, 1996a. doi: 10.1016/0016-7037(96)00158-5.
- B. P. Boudreau. A method-of-lines code for carbon and nutrient diagenesis in aquatic sediments. *Comput. Geosci.*, 22(5):479–496, June 1996b. doi: 10.1016/0098-3004(95)00115-8.
- Y.-H. Li and S. Gregory. Diffusion of ions in sea water and in deep-sea sediments. *Geochim. Cosmochim. Acta*, 38(5):703–714, 1974. doi: 10.1016/0016-7037(74)90145-8.
- F. L. Sayles, W. R. Martin, Z. Chase, and R. F. Anderson. Benthic remineralization and burial of biogenic SiO_2 , $CaCO_3$, organic carbon, and detrital material in the Southern Ocean along a transect at 170 West. *Deep-Sea Res. Pt II*, 48(19):4323–4383, 2001. doi: 10.1016/S0967-0645(01)00091-1.