

IB Linear Algebra (from lecture 18)

Martin von Hodenberg (mjv43@cam.ac.uk)

December 26, 2021

Linear algebra description etc

This article constitutes my notes for the ‘IB Linear Algebra’ course, held in Michaelmas 2021 at Cambridge. The course was lectured by Prof. Pierre Raphael.

Contents

1	Bilinear forms	1
1.1	Sylvester’s law and sesquilinear forms	4
1.1.1	Sesquilinear forms	8
1.2	Hermitian forms and skew symmetric forms	8

§1 Bilinear forms

Lemma 1.1

We have a bilinear form $\phi : V \times V \rightarrow \mathbb{F}$, where V is a finite vector space and B, B' are bases of V . Let

$$\phi = [\text{Id}]_{B, B'}.$$

Then

$$[\phi]_{B'} = P^T [\phi]_B P.$$

Proof. This is just a special case of the general change of basis formula; see that proof. \square

Definition 1.2 (Congruent matrices)

Two square matrices A, B are said to be **congruent** if there exists an invertible square matrix P such that

$$A = P^T B P.$$

Remark. This defines an equivalence relation.

Definition 1.3 (Symmetric bilinear form)

A bilinear form on V is said to be **symmetric** if

$$\phi(u, v) = \phi(v, u) \quad \forall u, v \in V.$$

Remark. We have encountered this definition before in IA Vectors and Matrices.

1. If A is a square matrix, we say that A is symmetric if $A^T = A$. Equivalently, $A_{ij} = A_{ji}$.
2. ϕ is symmetric iff $[\phi]_B$ is symmetric in *any* basis B .
3. To be able to represent ϕ by a diagonal matrix in some basis B , it is necessary that ϕ is symmetric:

$$P^T A P = D = D^T = P A^T P^T \implies A = A^T \implies \phi \text{ is symmetric.}$$

Definition 1.4 (Quadratic form)

A map $Q : V \rightarrow F$ is said to be a **quadratic form** if there exists a bilinear form $\phi : V \times V \rightarrow F$ such that

$$\forall u \in V, Q(u) = \phi(u, u).$$

Remark (Computation in a basis). Let $B = (e_i)_{1 \leq i \leq n}$ be a basis of V , and let $A = [\phi]_B$. Let $u = \sum_{i=1}^n u_i e_i$, then

$$Q(u) = \phi(u, u) = \phi \left(\sum_{i=1}^n u_i e_i, \sum_{j=1}^n u_j e_j \right) = \sum_{i,j=1}^n u_i u_j \phi(e_i, e_j) = \sum_{i,j=1}^n a_{ij} u_i u_j.$$

(by bilinearity of ϕ) Therefore we essentially have

$$Q(u) = U^T A U, \text{ where } U = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix}.$$

Remark. We can note that

$$Q(u) = U^T A U = \sum_{i,j=1}^n a_{ij} u_i u_j = \sum_{i,j=1}^n \left(\frac{a_{ij} + a_{ji}}{2} \right) u_i u_j = U^T \left(\frac{A + A^T}{2} \right) U.$$

So this representation of A is not necessarily unique.

Proposition 1.5

If $Q : V \rightarrow F$ is a quadratic form, then there exists a unique symmetric bilinear form $\phi : V \times V \rightarrow F$ such that

$$Q(u) = \phi(u, u) \quad \forall u \in V.$$

Proof. (Polarisation identity)

Proof of existence

Let ψ be a bilinear form on V such that

$$\forall u \in V, Q(u) = \psi(u, u).$$

Let $\phi(u, v) = \frac{1}{2}(\psi(u, v) + \psi(v, u))$. Thus we have that:

- ϕ is a bilinear form

- ϕ is symmetric
- $\phi(u, u) = \psi(u, u) = Q(u)$.

This concludes the proof of existence.

Proof of uniqueness

Let ϕ be a symmetric bilinear form such that

$$\forall u \in V, \phi(u, u) = Q(u).$$

Then

$$\begin{aligned} Q(u+v) &= \phi(u+v, u+v) \\ &= \phi(u, u) + \phi(u, v) + \phi(v, u) + \phi(v, v) \text{ by bilinearity} \\ &= Q(u) + 2\phi(u, v) + Q(v) \text{ by symmetry} \end{aligned}$$

From this we get that

$$\phi(u, v) = \frac{1}{2}(Q(u+v) - Q(u) - Q(v)).$$

□

Theorem 1.6 (Diagonalisation of symmetric bilinear forms)

Let $\phi : V \times V \rightarrow F$ be a symmetric bilinear form. ($\dim V = n$). Then there exists a basis B of V such that $[\phi]_B$ is diagonal.

Proof. We proceed by induction on the dimension of V . For $n = 1$ it is trivially true. Suppose the theorem holds for all dimensions $< n$: then

- If $\phi(u, u) = 0 \forall u \in V$, then $\phi = 0$ by the polarisation identity (ϕ is symmetric).
- If $\phi \neq 0$, then there exists a $u \in V \setminus \{0\}$ such that $\phi(u, u) \neq 0$. Let us call $u = e_1$.
- Let U be the 'orthogonal' of e_1 :

$$\begin{aligned} U &= \{v \in V : \phi(e_1, v) = 0\} \\ &= \ker \theta \text{ where } \theta : V \rightarrow F \text{ is given by } \theta(v) = \phi(e_1, v). \end{aligned}$$

Since it is a kernel of a linear map $V \rightarrow F$, U is a vector subspace of V . By the Rank-Nullity theorem, we have

$$\dim V = n = r(\theta) + \text{null } \theta = \dim U + 1.$$

We now claim that $U + \langle e_1 \rangle = U \oplus \langle e_1 \rangle$. Indeed,

$$\begin{aligned} v \in \langle e_1 \rangle \cap U &\implies v = \lambda e_1 \text{ and } \phi(e_1, v) = 0. \\ \implies 0 &= \phi(e_1, v) = \phi(e_1, \lambda e_1) = \lambda \phi(e_1, e_1) \implies \lambda = 0 \implies v = 0. \\ \implies U + \langle e_1 \rangle &= U \oplus \langle e_1 \rangle. \end{aligned}$$

Therefore $V = U \oplus \langle e_1 \rangle$, and pick a basis $B' = (e_2, \dots, e_n)$ such that (e_1, e_2, \dots, e_n) is a basis of V (since the sum is direct). So

$$[\phi]_B = (\phi(e_i, e_j))_{1 \leq i, j \leq n} = \left(\begin{array}{c|c} \phi(e_1, e_1) & 0 \\ \hline 0 & A' \end{array} \right).$$

Therefore $(A')^T = A'$, and $A' = [\phi|_U]_{B'}$ where $\phi|_U$ is the restriction of ϕ onto U . Now we apply the induction hypothesis to find a basis (e'_1, \dots, e'_n) of V such that $[\phi|_U]_B$ is diagonal. So

$$\hat{B} = (e_1, e'_2, \dots, e'_n)$$

is a basis of V , and finally we have that $[\phi]_{\hat{B}}$ is diagonal. □

Example 1.7

Let $V = \mathbb{R}^3$, and

$$\begin{aligned} Q(x_1, x_2, x_3) &= x_1^2 + x_2^2 + 2x_3^2 + 2x_1x_2 + 2x_1x_3 - 2x_2x_3 \\ &= x^T Ax \text{ where } A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 1 & -1 & 2 \end{pmatrix} \end{aligned}$$

There are two ways to diagonalise our quadratic form:

1. Diagonalise using the algorithm we developed in the previous proof.
2. Complete the square.

Let's complete the square:

$$\begin{aligned} Q(x_1, x_2, x_3) &= (x_1 + x_2 + x_3)^2 + x_3^2 - 4x_2x_3 \\ &= (x_1 + x_2 + x_3)^2 + (x_3 - 2x_2)^2 - (2x_2)^2 \\ &= (x'_1)^2 + (x'_2)^2 + (x'_3)^2. \end{aligned}$$

Therefore

$$P^T AP = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

To find P , note that

$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & -2 & 1 \\ 0 & -2 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = P^{-1} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

§1.1 Sylvester's law and sesquilinear forms

Recall our theorem that a symmetric bilinear form has a diagonal basis.

Corollary 1.8

Let $F = \mathbb{C}$, and ϕ be a symmetric bilinear form of rank r on $V \times V$ where $\dim V = n$. Then there exists a basis B of V such that the matrix

$$[\phi]_B = \left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array} \right).$$

Proof. Pick $E = (e_1, \dots, e_n)$ such that

$$[\phi]_E = \begin{pmatrix} a_1 & & 0 \\ & \ddots & \\ 0 & & a_n \end{pmatrix}.$$

Order the a_i such that

$$\begin{cases} a_i \neq 0 & 1 \leq i \leq r \\ a_i = 0 & i > r \end{cases}$$

For $i \leq r$, let $\sqrt{a_i}$ be a choice of complex root for a_i . Let

$$\begin{cases} v_i = \frac{e_i}{\sqrt{a_i}} & 1 \leq i \leq r \\ v_i = e_i & i > r \end{cases}$$

Then $B = (v_1, \dots, v_r, v_{r+1}, \dots, v_n)$ is a basis of V and we can check that we have diagonalised and normalised ϕ :

$$[\phi]_B = \left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array} \right).$$

□

Corollary 1.9

Every symmetric matrix of $M_n(\mathbb{C})$ is congruent to a *unique* matrix of the form

$$\left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array} \right).$$

Proof. Immediate.

□

Corollary 1.10

Let $F = \mathbb{R}$. Let ϕ be a symmetric bilinear form on $V \times V$ where $\dim V = n$. Then there exists a basis $B = (v_1, \dots, v_n)$ basis of V such that for some $p - q \geq 0$ with $p + q = r(\phi)$,

$$[\phi]_B = \left(\begin{array}{c|c|c} I_p & 0 & 0 \\ \hline 0 & -I_q & 0 \\ \hline 0 & 0 & 0 \end{array} \right).$$

Proof. Pick $E = (e_1, \dots, e_n)$ such that

$$[\phi]_E = \begin{pmatrix} a_1 & & 0 \\ & \ddots & \\ 0 & & a_n \end{pmatrix}.$$

Order the a_i such that

$$\begin{cases} a_i > 0 & 1 \leq i \leq p \\ a_i < 0 & p+1 \leq i \leq p+q \\ a_i = 0 & i > p+q \end{cases}$$

Similarly to before we define

$$\begin{cases} v_i = \frac{e_i}{\sqrt{a_i}} & 1 \leq i \leq p \\ v_i = \frac{e_i}{\sqrt{-a_i}} & p+1 \leq i \leq p+q \\ v_i = e_i & i > p+q \end{cases}$$

Then this basis will do the job. □

Definition 1.11 (Signature of a quadratic form)

For $F = \mathbb{R}$, we define the **signature** of ϕ

$$s(\phi) = p - q.$$

This is also the signature of the associated quadratic form Q . We will soon see that this is well-defined.

Definition 1.12 (Positive definite quadratic/bilinear form)

Let ϕ be a symmetric bilinear form on a real vector space V . We say that

(i) ϕ is positive definite if

$$\phi(u, u) > 0 \quad \forall u \in V \setminus \{0\}.$$

(ii) ϕ is positive semidefinite if

$$\phi(u, u) \geq 0 \quad \forall u \in V \setminus \{0\}.$$

(iii) ϕ is negative definite if

$$\phi(u, u) < 0 \quad \forall u \in V \setminus \{0\}.$$

(iv) ϕ is negative semidefinite if

$$\phi(u, u) \leq 0 \quad \forall u \in V \setminus \{0\}.$$

Example 1.13

The matrix

$$\left(\begin{array}{c|c} I_p & 0 \\ \hline 0 & 0 \end{array} \right)$$

is

- positive definite for $p = n$
- positive semidefinite for $1 \leq p \leq n$.

Theorem 1.14 (Sylvester's law of inertia)

The signature of a quadratic form is well defined.

Formally, if a real symmetric bilinear form is represented in two different ways by two identity blocks of size p, q and p', q' then we have

$$p = p', q = q'.$$

Proof. In order to prove uniqueness of p , it is enough to show that p is the largest dimension of a subspace of V on which ϕ is positive definite. Say $B = (v_1, \dots, v_n)$ and

$$[\phi]_B = I_p - I_q.$$

Let $X = \langle v_1, \dots, v_p \rangle$. Then ϕ is positive definite on X . If we take a $u \in X$ with $u = \sum_{i=1}^p \lambda_i v_i$, then

$$Q(u) = \phi(u, u) = \phi\left(\sum_{i=1}^p \lambda_i v_i, \sum_{j=1}^p \lambda_j v_j\right) = \sum_{i,j=1}^p \lambda_i \lambda_j \phi(v_i, v_j) = \sum_{i=1}^p \lambda_i^2.$$

This is > 0 as long as $u \neq 0$. Suppose that ϕ is positive definite on another subspace X . Let

$$X = \langle v_1, \dots, v_p \rangle, Y = \langle v_{p+1}, \dots, v_n \rangle.$$

Then arguing as above, looking at $[\phi]_B$ we know that ϕ is negative semidefinite on Y . This implies that

$$Y \cap X = \{0\}.$$

Indeed, if $y \in Y \cap X$, then $Q(y) \leq 0$ since $y \in Y$, but this implies that $y = 0$ since $y \in X$. Therefore

$$Y + X = Y \oplus X \implies n = \dim V \geq \dim(Y + X) = \dim Y + \dim X \implies n \geq n - p + \dim X \implies \dim X \leq p.$$

Similarly, we show that q is the largest dimension of a subspace on which ϕ is negative definite. So we have a *geometric characterisation* of p and q , which concludes the proof. \square

Definition 1.15 (Kernel of a bilinear form)

Define the kernel of the bilinear form

$$K = \{v \in V : \forall u \in V, \phi(u, v) = 0\}.$$

Remark. $\dim K + r(\phi) = n$.

One can show using the above notation that there is a subspace T of dimension $m - (p + q) + \min p, q$ such that $\phi_T = 0$. The subspace we want to take consists of all the basis vectors that are killed by the lower corner of $[\phi]_B$, but it also includes the 'cancellations' between I_p and $-I_q$.

Moreover, one can show that the dimension of T is the largest possible dimension of such a subspace.

§1.1.1 Sesquilinear forms

Recall that the standard inner product on \mathbb{C}^n is

$$\langle x, y \rangle = \left\langle \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \right\rangle = \sum_{i=1}^n x_i \bar{y}_i.$$

However note that this map is **not** a bilinear form, since the conjugation of the second coordinate prevents this. However, it is still bilinear 'in a way'. We now explore this further.

Definition 1.16 (Sesquilinear form)

Let V, W be vector spaces over \mathbb{C} . A sesquilinear form on $V \times W$ is a function

$$\phi : V \times W \rightarrow \mathbb{C}.$$

such that

1. $\phi(\lambda_1 v_1 + \lambda_2 v_2, w) = \lambda_1 \phi(v_1, w) + \lambda_2 \phi(v_2, w)$
2. $\phi(\lambda_1 v_1 + \lambda_2 v_2, w) = \bar{\lambda}_1 \phi(v_1, w) + \bar{\lambda}_2 \phi(v_2, w)$

For B a basis of V , C a basis of W , we define the matrix

$$[\phi]_{B,C} = (\phi(v_i, w_j))_{1 \leq i \leq m, 1 \leq j \leq n}.$$

Lemma 1.17

If B basis for V , C a basis for W ,

$$\phi(v, w) = [v]_B^T [\phi]_{B,C} [\bar{w}]_C.$$

Further, if B', C' are other bases for V, W respectively, and $P = [Id]_{B',B}, Q = [Id]_{C',C}$ then

$$[\phi]_{B',C'} = P^T [\phi]_{B,C} \bar{Q}.$$

Proof. Same proofs as for bilinear forms, just slightly modified. □

§1.2 Hermitian forms and skew symmetric forms

We will now generalise symmetric bilinear forms to sesquilinear forms.

Definition 1.18 (Hermitian form)

Let V be a finite dimensional vector space and $\phi : V \times V \rightarrow \mathbb{C}$ be a sesquilinear form. We call ϕ **hermitian** if for all $(u, v) \in V \times V$ we have

$$\phi(u, v) = \overline{\phi(v, u)}.$$

Remark. If ϕ is Hermitian, then $\phi(u, u) = \overline{\phi(u, u)} \implies \phi(u, u) \in \mathbb{R}$. Moreover,

$$\phi(\lambda u, \lambda u) = |\lambda|^2 \phi(u, u).$$

This allows us to talk about negative/ positive definite Hermitian form.

Lemma 1.19

A sesquilinear form $\phi : V \times V \rightarrow \mathbb{C}$ is hermitian if and only if for every basis B of V ,

$$[\phi]_B = [\phi]_B^T.$$

Proof. Let $A = [\phi]_B$ have entries $a_{ij} = \phi(e_i, e_j)$

□