Math 340 • Fall 2026 • Victor Matveev

Polynomial Interpolation Error

A big advantage of Newton's divided difference method of writing down an interpolating polynomial is the ease with which we can analyze its error. In fact, the remainder (error) of interpolation looks exactly like an extra interpolation term obtained by adding one more point:

$$f(x) = \underbrace{f[x_1] + f[x_1,x_2](x-x_1) + \ldots + f[x_1,\ldots,x_n](x-x_1)\ldots(x-x_{n-1})}_{P_{n-1}(x)} + \underbrace{f[x_1,\ldots,x_n,x](x-x_1)\ldots(x-x_n)}_{\text{Remainder / Error, } R_{n-1}(x)}$$

The **proof** of this remainder formula is remarkably simple: consider $\boldsymbol{x}$ in the above expression as another interpolation node, $x \equiv x_{n+1}$; then the above is simply a formula for $P_n(x)$ -- an interpolant matching one extra point of $f(x)$ -- which exists for any value of $x \equiv x_{n+1}$.

It can also be elegantly proven that each of the coefficients above is related to the derivative of $f(x)$, highlighting the analogy between interpolating polynomial and a "distributed" form of Taylor expansion:

$$\exists c \in \text{Interval containing all } \{x_k\}_{k=1}^n, \text{ such that } f[x_1,\ldots,x_n] = \frac{f^{(n-1)}(c)}{(n-1)!}$$

For the remainder, $\boldsymbol{R_{n-1}(x) = f[x_1, \ldots, x_n, x](x-x_1)\ldots(x-x_n)}$, this implies that

$$\exists c \in \text{Interval containing all } \{x_k\}_{k=1}^n \text{ and } x, \text{ such that } R_{n-1}(x) = \frac{f^{(n)}(c)}{n!}(x-x_1)\ldots(x-x_n)$$

Even though we don't know where the argument value $\boldsymbol{c}$ resides, the latter expression is useful when finding the upper bound on the error. Let's consider an example to show how to use the remainder formula to determine error bounds:

EXAMPLE: Construct a **linear** approximation of $\boldsymbol{f(x) = e^x}$ using nodes $\{x_k\} = \{0, 1\}$. Examine its error over the interval [0, 1], in several ways:

1) Compute the error in the middle, at $x = \frac{1}{2}$ (since we expect the error to be largest half-way between $x_{1,2}$)

2) Find exact maximal error on the interval [0, 1].

3) Analyze the error bound using the remainder formula above.

First, let's find the interpolating polynomial (which is a secant line in this case of just two interpolation nodes).

x_k	$f[x_k]$	$f[x_{k-1},x_k]$
0	1	
1	e	$\frac{e-1}{1-0} = e-1$

Thus, our interpolating line is: $f(x) \sim P_1(x) = f[x_1] + f[x_1,x_2](x-x_1) = 1 + (e-1)x$

1) The remainder at $x = 1/2$ then equals: $\left|R_1\left(\frac{1}{2}\right)\right| = \left|f\left(\frac{1}{2}\right) - P_1\left(\frac{1}{2}\right)\right| = \left|e^{1/2} - \left(1+\frac{e-1}{2}\right)\right| \approx \mathbf{0.2104}$

2) Find the exact maximum of remainder, $R_1 \equiv P_1(x) - f(x)$, by setting its derivative to zero:

$$R_1'(x) = f'(x) - P_1'(x) = e^x - (e-1) = 0 \;\rightarrow\; e^x = e-1 \;\rightarrow\; x^* = \ln(e-1) \approx 0.54132$$

$$\max_{[0,1]} R_1(x) = R_1(x^*) = 1 + (e-1)x^* - e^{x^*} \approx \mathbf{0.212} \quad \textbf{Very close to the error in the middle written above!}$$

It is interesting that the actual maximal error is achieved slightly to the right of the mid-point (see figure below)

3) Finally, let's compare this with the upper bound on the error obtained using the relationship between Newton's divided differences and the function derivatives:

$$|R_1(x)| = \left|\frac{f''(c)}{2!}(x-x_1)(x-x_2)\right|$$

$$\leq \max_{[0,1]}\left|\frac{f''(c)}{2!}\right| \cdot \max_{[0,1]}|(x-x_1)(x-x_2)|$$


$$= \max_{[0,1]}\left|\frac{e^c}{2}\right| \cdot \max_{[0,1]}|x(x-1)| = \frac{e}{2}\left|\frac{1}{2}\left(\frac{1}{2}-1\right)\right| = \frac{e}{8} \approx \mathbf{0.34}$$

As expected, the error bound is an over-estimate of the true maximal error.

Note that the first term (the derivative of exponential) is the largest at the right end of the interval, where the error is actually reduced by the proximity to the interpolation point.

Let's compare the maximal true error with the Taylor expansion near $x = 0$, $f(x) = \exp(x) \approx T_1(x) = 1 + x$. The maximal error of this approximation is of course achieved at $x = 1$: $Error = f(1) - T_1(1) = e - 2 \approx 0.72$. This is significantly larger than the error of the polynomial interpolant, but we could have done better by expanding around the

Figure 1: Linear interpolation of $exp(x)$ using nodes {0, +1}. Note that the largest error is slightly to the right of the mid-point, $x = \frac{1}{2}$.



midpoint instead of $x = 0$. In that case, the error would still be higher: $Error = f(1) - T_1(1) \approx 0.2452$.

Conclusion 1: Polynomial interpolation is always more accurate than Taylor expansion, if the goal is to approximate the function with uniform error bound over an entire interval.

Conclusion 2: exact error formula is of course preferable, but the error bound approach gives a simple way to find the "worst-case scenario" error over the **entire** interval, with minimal extra computation.


On the next slide, we will see that our choice of the two interpolation nodes in this figure is a bit silly, given our goals.

Optimal Interpolation Node Choice: Chebyshev Points

From the figure above, it is clear that endpoints (and, more generally, equidistant points) are far from an optimal choice.

In fact, we will prove directly in class that the basis function $|\psi(x)| = |(x - x_1)(x - x_2)|$ appearing in the error formula in the case of two interpolation points is minimized over the interval [0, 1] when $x_{1,2} = \frac{1}{2}\left(1 \pm \cos\frac{\pi}{4}\right) = \frac{1}{2}\left(1 \pm \frac{1}{\sqrt{2}}\right)$

We get a much better fit with this choice, shown in the **Figure 2**.



Parenthetically, we note that the maximal error in this case is achieved at the endpoint $x = 1$, not in the middle of the domain. In principle, we could further reduce the maximal error by moving the nodes slightly to the right, until the error is the same at both endpoints and at the maximal error x-value between the endpoints.