

ELEC3705 Lecture Notes

Oskar Meszaros (z5636164)

September 2026

Contents

1	Why Quantum Mechanics?	2
1.1	Photoelectric Effect	2
1.2	Planck-Einstein Relations	2
1.3	Double-Slit Experiment	2
1.4	Double-Slit with Particles	2
1.5	Wave Function	3
1.6	Schrödinger Equation	4
1.7	Wave Packets	4
1.8	Heisenberg Uncertainty Principle	4
2	The Mathematical Formalism of Quantum Mechanics	5
2.1	Hilbert Space	5
2.2	Operators	5
2.2.1	Linear Operators	5
2.2.2	Adjoint Operators	6
2.2.3	Hermitian Operators	6
2.3	Basis of a Hilbert Space	6
2.4	Observables	6
3	Spin and The Postulates of Quantum Mechanics	7
3.1	Postulate 1	7
3.1.1	Applications	7
3.2	Postulate 2	7
3.2.1	Applications	7
A	Fourier Transform	8

1 Why Quantum Mechanics?

1.1 Photoelectric Effect

The photoelectric effect allows us to extract electrons from a piece of metal using light/high-powered lasers.

1. Below a certain frequency (colour) of light, no electrons are extracted from the metal. This is completely independent of the light intensity.
2. Above a cutoff frequency ω_c , the electrons are extracted from the metal, even at very small light intensities.
3. The energy E of the extracted electrons is proportional to $\omega - \omega_c$.
4. The momentum $p = mv = \sqrt{2mE}$ of the extracted electrons is inversely proportional to the wavelength of light.

1.2 Planck-Einstein Relations

This effect can be explained if we assume that light is quantised in packets of energy, called photons, where

$$E = \hbar\omega = h\nu$$
$$\vec{p} = \hbar\vec{k} = h/\lambda,$$

where $h = 6.626 \times 10^{-34} \text{ J s}$ is the Planck constant, and $\hbar = h/2\pi = 1.054 \times 10^{-34} \text{ J s/rad}$ is the "reduced" Planck constant. These are the Planck-Einstein relations. They describe light as composed of "particles" of energy, determined by the frequency.

1.3 Double-Slit Experiment

The double-slit experiment demonstrated that light is also an electromagnetic-wave, and can be understood by using the linearity of Maxwell's equations.

The light intensity is proportional to $|E|^2$, and if $\vec{E}(\vec{r}) = \vec{E}_1(\vec{r}) + \vec{E}_2(\vec{r})$ from the two sources, then

$$I \propto \left| \vec{E}_1(\vec{r}) + \vec{E}_2(\vec{r}) \right|^2 \neq \left| \vec{E}_1(\vec{r}) \right|^2 + \left| \vec{E}_2(\vec{r}) \right|^2,$$

shows the wave nature of light.

This experiment still holds at extremely low intensities. It is still individual photons that show up on the screen, however when averaged over time it still results in the interference pattern.

1.4 Double-Slit with Particles

This experiment can be repeated using things that are normally thought of as particles, such as electron or neutrons. To find the required wavelength, we can use the de Broglie hypothesis

Definition 1.1. The de Broglie hypothesis states:

"to every particle is associated a wave, which obeys the same Planck-Einstein relations as applicable for photons."

From the relation above, we can derive

$$\lambda = \frac{2\pi}{k} = \frac{h}{p}$$

This is only significant for extremely small objects. The distance between intensity peaks is given by

$$\delta x = \lambda \frac{D}{d},$$

with $\lambda = h/p$, D is the distance between the slits and the screen, and d is the separation of the slits. This means that larger particles have smaller distances between intensity peaks, with all else being the same. To make a meaningful observation, we can

- Increase the wavelength by decreasing the momentum (smaller velocity for constant mass),
- Decrease the distance between the slits,
- Or increase the distance to the screen.

1.5 Wave Function

These two experiments force us to recognise the idea that particles are simultaneously particles and waves, and in fact that waves and particles are not different things. Instead, each wave is made of packets of energy, with is associated with wavelength $\lambda = h/p$ and frequency $\nu = E/h$.

Instead of trying to describe the trajectory of a particle with position $\vec{r}$ and momentum $\vec{p}$, we use a wave function $\psi(\vec{r}, t)$, which contains all the information we can gather on the particle. This wave function is a probability amplitude, such that

$$C \iiint_V |\psi(\vec{r}, t)|^2 d^3\vec{r} = P_V(t),$$

where $P_V(t)$ is the probability of finding the particle in the volume V at time t , and C is a normalisation constant to ensure the value is between 0 and 1, i.e.

$$C \iiint_{-\infty}^{\infty} |\psi(\vec{r}, t)|^2 d^3\vec{r} = 1,$$

i.e. the particle has to be somewhere. Now instead of calculating the trajectory of a particle, we instead calculate the probability of finding the particle in a certain region in space, and how that evolves over time.

We know that Maxwell's equations allow solutions in the form of propagating waves, e.g.

$$\vec{E}(\vec{r}, t) = A e^{i(k\vec{r} - \omega t)}$$

is a plane wave that solve the wave equation of the form

$$\nabla^2 \vec{E} = \frac{1}{v^2} \frac{\partial^2 \vec{E}}{\partial t^2},$$

where $v = \omega/k$. This wave equation contains the partial second derivative of the electric field with respect to both time and space. Taking the second derivative of the plane wave results in the coefficients k^2 and ω^2 which give rise to the $1/v^2$ term in the equation. Note that v is constant.

Consider a free particle. The energy is purely kinetic, so

$$E = \frac{1}{2}mv^2 = \frac{p^2}{2m} = \frac{\hbar^2 k^2}{2m},$$

which we can impose Planck-Einstein to get

$$\hbar\omega = \frac{\hbar^2 k^2}{2m}.$$

An important thing to note is that the energy depends quadratically on k , but linearly on ω . This means that an appropriate wave equation should contain a first derivative with respect to time, and the second derivative with respect to space. Here, v is no longer constant.

1.6 Schrödinger Equation

Schrödinger postulated an equation that describes the time evolution of the wavefunction of quantum object to be

$$i\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi(\vec{r}, t) + V(\vec{r}) \psi(\vec{r}, t)$$

$V(\vec{r})$ is a (position-dependent) potential energy. It is similar to a wave equation, containing the first derivative w.r.t. time, and the second w.r.t. space. The coefficients are chosen to impose the Planck-Einstein relations. Because of the first derivative w.r.t. time, the equation explicitly contains i .

Proof. Consider a free particle not subjected to any potentials or forces ($V(\vec{r}) = 0$):

$$i\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi(\vec{r}, t).$$

This has a solution of the form

$$\psi(\vec{r}, t) = A e^{i(\vec{k} \cdot \vec{r} - \omega t)}$$

with the condition

$$\hbar \omega = \frac{\hbar^2 k^2}{2m}.$$

Inserting the Planck-Einstein relations, the above conditions means

$$E = \frac{p^2}{2m} = \frac{1}{2} m v^2.$$

□

1.7 Wave Packets

These wavefunctions can be combined into a single wave packet by simply summing them together (and hence with 3 wave numbers k). The interference between the different wave components causes the wave components to no longer be spread out equally in space, however it still have recurrences.

These recurrences get more spread out the more wave components are added, up to a limit of infinitely many components, each with a different wave number k and with a coefficient $g(k)$. For simplicity, take the 1D case ($\vec{r} \equiv x$):

$$\psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(k) e^{i(kx - \omega t)} dk.$$

We can find $g(k)$ by setting $t = 0$ (i.e. initial conditions)

$$\psi(x, 0) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(k) e^{ikx} dk,$$

therefore

$$g(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x, 0) e^{-ikx} dx.$$

This means that $g(k)$ is the Fourier transform (in space) of the shape of the wave packet at $t = 0$, i.e. of the probability amplitude of finding the object at point x at $t = 0$.

1.8 Heisenberg Uncertainty Principle

Consider a Gaussian wave packet

$$\psi(x, 0) = \frac{1}{\sqrt{2\pi}\sigma_x} \exp\left(-\frac{x^2}{2\sigma_x^2}\right).$$

It's Fourier transform is also a Gaussian

$$\begin{aligned} g(k) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x, 0) e^{-ikx} dx \\ &= \frac{1}{\sqrt{2\pi}\sigma_x} \exp\left(-\frac{k^2}{2\omega_k^2}\right). \end{aligned}$$

From the properties of Fourier transforms, we know that

$$\sigma_k = 1/\sigma_x \Rightarrow \sigma_k \sigma_x = 1.$$

Imposing the Planck-Einstein relation $p = \hbar k \Rightarrow \sigma_p = \hbar \sigma_k$, we get

$$\boxed{\sigma_p \sigma_x = \hbar}$$

This relation is a simple mathematical property of Fourier transform. This appears as soon as you assume the Planck-Einstein relations, i.e. wave-particle duality. This also holds if you assume the uncertainty principle as a postulate, then wave-particle duality is a consequence of the mathematical properties of Fourier transforms.

2 The Mathematical Formalism of Quantum Mechanics

2.1 Hilbert Space

A function is a vector in an infinite-dimensional vector space. It is sometimes convenient to represent the function using a finite number of "basis functions", so that the vector space necessary to describe the function has finite dimension. This is known as a Hilbert Space.

A vector space requires a suitable basis.

Example 2.1. 3D vector space $\mathbb{R}^3$.

Let $\{\vec{u}_x, \vec{u}_y, \vec{u}_z\}$ be an orthonormal basis in $\mathbb{R}^3$. We can then describe a vector in terms of this basis as

$$\vec{v} = x\vec{u}_x + y\vec{u}_y + z\vec{u}_z,$$

where x , y , and z are coefficients. We then have

$$|\vec{u}_\alpha| = 1, \quad \text{and} \quad \vec{u}_\alpha \cdot \vec{u}_\beta = \delta_{\alpha\beta}.$$

Once we have the basis, we can represent any vector in that basis as a vector of its coefficients in the chosen basis

$$\vec{v} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

We can denote this vector as $|v\rangle$, indicating it is a normal vector. Similarly, we can define the transpose of this vector as $\langle v|$. Note that we also have to take the complex conjugate of the elements in this vector as we are dealing with complex numbers. We then define the dot product of this to be

$$\vec{a} \cdot \vec{b} \rightarrow \vec{a}^T \vec{b} \langle a|b \rangle = \sum_{i=1}^3 a_i^* b_i$$

This is known as Dirac notation.

2.2 Operators

2.2.1 Linear Operators

In a vector space, we can define linear operators (usually matrices) that transform a vector into another

$$|\psi'\rangle = A|\psi\rangle.$$

Note that $\langle\phi|\psi\rangle$ is a number, and $|\phi\rangle\langle\psi|$ is a matrix.

2.2.2 Adjoint Operators

Adjoint operators provide the correspondence between bras and kets

$$|\psi'\rangle = A|\psi\rangle \rightarrow \langle\psi'| = \langle\psi| A^\dagger.$$

Here, $A^\dagger$ is the adjoint of A . It is obtained by transposing and taking the complex conjugate of each element.

2.2.3 Hermitian Operators

If an operator is its own adjoint (i.e. $A = A^\dagger$), then they are called Hermitian operators. They are simple to recognise

- The elements on opposite side of the diagonal are the complex conjugate of each
- The elements on the diagonal are purely real.

2.3 Basis of a Hilbert Space

A set of kets $\{|\nu_i\rangle\}$ is called orthonormal if

$$\langle\nu_i|\nu_j\rangle = \delta_{ij}.$$

The set $\{|\nu_i\rangle\}$ forms a basis of the Hilbert space $\mathcal{H}$ if every $|\psi\rangle$ can be written as

$$|\psi\rangle = \sum_j c_j |\nu_j\rangle.$$

An important property is the closure relation

$$|\psi\rangle = \sum_j c_j |\nu_j\rangle = \sum_j \langle\nu_i|\psi\rangle |\nu_j\rangle = \sum_j |\nu_j\rangle \langle\nu_j|\psi\rangle.$$

For this to be true, it requires

$$\sum_j |\nu_j\rangle \langle\nu_j| = \mathbb{I}.$$

This is called the closure relation. The operators $|\nu_j\rangle \langle\nu_j|$ projects a vector onto the basis state $|\nu_j\rangle$.

2.4 Observables

General linear algebra theorems

- The eigenvalues of a Hermitian operator are real
- The eigenvectors corresponding to different eigenvalues are orthogonal

A Hermitian operator is called observable if its eigenvectors form a basis of the vector space. We call a unitary an operator for which holds

$$U^{-1} = U^\dagger.$$

The eigenvalues of unitary operators are complex number of modulus 1.

3 Spin and The Postulates of Quantum Mechanics

3.1 Postulate 1

The first postulate of quantum mechanics: **At an initial time, the state of a quantum system is specified by a ket that belongs to a suitable vector space.**

This means that any ket

$$|\psi'\rangle = \lambda_1 |\psi_1\rangle + \lambda_2 |\psi_2\rangle,$$

is also a legitimate state of the system. This is known as the superposition principle.

Note that we normally assume that the state vectors are normalised

$$\langle\psi|\psi\rangle = 1.$$

3.1.1 Applications

Assuming a 2D vector space (spin-up and spin-down). We have a basis in this space to be

$$|\uparrow\rangle, |\downarrow\rangle.$$

They must satisfy the following conditions

- Normalisation: $\langle\downarrow|\downarrow\rangle = \langle\uparrow|\uparrow\rangle = 1$
- Orthogonality: $\langle\uparrow|\downarrow\rangle = 0$
- Closure: $|\downarrow\rangle\langle\downarrow| + |\uparrow\rangle\langle\uparrow| = \mathbb{I}$

There are an infinite number of options for this, but the simplest is

$$|\uparrow\rangle = |1\rangle = \begin{pmatrix} 1 & 0 \end{pmatrix}, \quad |\downarrow\rangle = |2\rangle = \begin{pmatrix} 0 & 1 \end{pmatrix}.$$

Then, any ket of the form $\alpha |1\rangle + \beta |2\rangle$, with $\alpha^2 + \beta^2 = 1$ is a legitimate state of the system

$$|\psi\rangle = \begin{pmatrix} \alpha & \beta \end{pmatrix}.$$

3.2 Postulate 2

The second postulate of quantum mechanics: **Every measurable physical quantity is described by an operator, which must be an observable.**

This means you can construct operators in your space ϵ that not observable, i.e. they cannot correspond to measurable physical quantities.

3.2.1 Applications

A Fourier Transform

A Fourier transform decomposes a function into another function that describes the number of occurrences of frequencies within the original function.

$$F(\omega) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt.$$

Fundamentally it is an inner product between $f(t)$ and $e^{-i\omega t}$, with the second vector being characterised by the parameter ω . It tells you how similar the two vectors are to each other, thereby defining the number of occurrences of each frequency.

The Fourier series of a sinusoid is a delta function, indicating that it is infinitely similar at that frequency, and not at all at the others.