ELEC3705 Lecture Notes

Oskar Meszaros (z5636164)

September 2026

Contents

1 Why Quantum Mechanics? 3
- 1.1 Photoelectric Effect . . . 3
- 1.2 Planck-Einstein Relations . . . 3
- 1.3 Double-Slit Experiment . . . 3
- 1.4 Double-Slit with Particles . . . 3
- 1.5 Wave Function . . . 4
- 1.6 Schrödinger Equation . . . 5
- 1.7 Wave Packets . . . 5
- 1.8 Heisenbery Uncertainty Principle . . . 6
- 1.9 Problems . . . 6

2 The Mathematical Formalism of Quantum Mechanics 8
- 2.1 Hilbert Space . . . 8
- 2.2 Operators . . . 8
  - 2.2.1 Linear Operators . . . 8
  - 2.2.2 Adjoint Operators . . . 8
  - 2.2.3 Hermitian Operators . . . 9
- 2.3 Basis of a Hilbert Space . . . 9
- 2.4 Observables . . . 9
- 2.5 Problems . . . 9

3 Spin and The Postulates of Quantum Mechanics 9
- 3.1 Postulate 1 . . . 9
  - 3.1.1 Application . . . 10
- 3.2 Postulate 2 . . . 10
  - 3.2.1 Application . . . 10
- 3.3 Postulate 3 . . . 10
  - 3.3.1 Application . . . 10
- 3.4 Postulate 4 . . . 11
  - 3.4.1 Application . . . 11
- 3.5 Postulate 5 . . . 12
  - 3.5.1 Application . . . 12
- 3.6 Postulate 6 . . . 13
  - 3.6.1 Application . . . 13
- 3.7 Spin Matrices . . . 13
- 3.8 Projection Operator . . . 14
- 3.9 Exponential of a Matrix . . . 14
- 3.10 Hamiltonian Eigenbasis . . . 15
- 3.11 Time Evolution Operator . . . 16
  - 3.11.1 Unitary Operators . . . 17
  - 3.11.2 Time Reversibility . . . 17

3.12 Commutation . . . 17
3.12.1 Ehrenfest Theorem . . . 18
3.13 Pauli Matrices . . . 18
3.14 Non-Hermitian Operators . . . 19
3.15 Problems . . . 19

4 Quantum Mechanics in Real Space **20**
4.1 Schrödinger Equations in Real Space . . . 20
4.2 Stationary States . . . 21
4.3 Low-Energy Truncation . . . 22
4.4 Non-Stationary States . . . 22

5 Quantum Tunnelling **23**
5.1 Particle in a Step-Like Potential . . . 23
5.2 Quantum Tunnelling . . . 24
5.3 Double-Well . . . 25
5.3.1 Approximation 1: Infinite Barrier . . . 25
5.3.2 Approximation 2: Finite Wells . . . 26
5.3.3 Tunnelling Splitting . . . 26
5.3.4 Tunnelling Oscillations . . . 26
5.4 Relation to Spin . . . 27

6 Position and Momentum Operators **27**
6.1 Position Eigenbasis . . . 27
6.2 Momentum Operator . . . 28
6.3 Commutation Relations . . . 28

7 Dynamics and Control of a Quantum Bit **28**
7.1 Time Evolution . . . 29
7.2 Rabi Formula and Oscillation . . . 30
7.3 Time-Dependent Hamiltonian . . . 31
7.4 Rotating Frame . . . 31
7.5 Change of Basis . . . 32
7.6 Magnetic Resonance . . . 33
7.7 Density Operator . . . 33
7.7.1 Properties of the Density Operator . . . 33
7.7.2 Density Operator for Mixed States . . . 34
7.7.3 Populations and Coherences . . . 34
7.7.4 State Purity . . . 35

A Fourier Transform **36**

B Matlab/Python Commands **36**
B.1 Matrix Operations . . . 36
B.1.1 Exponential . . . 36

1 Why Quantum Mechanics?

1.1 Photoelectric Effect

The photoelectric effect allows us to extract electrons from a piece of metal using light/high-powered lasers.

1. Below a certain frequency (colour) of light, no electrons are extracted from the metal. This is completely independent of the light intensity.
2. Above a cutoff frequency ω_c, the electrons are extracted from the metal, even at very small light intensities.
3. The energy E of the extracted electrons is proportional to $\omega - \omega_c$.
4. The momentum $p = mv = \sqrt{2mE}$ of the extracted electrons is inversely proportional to the wavelength of light.

1.2 Planck-Einstein Relations

This effect can be explained is we assume that light in quantised in packets of energy, called photons, where

$$E = \hbar\omega = hv$$
$$\vec{p} = \hbar\vec{k} = h/\lambda,$$

where $h = 6.626 \times 10^{-34}$J s is the Planck constant, and $\hbar = h/2\pi = 1.054 \times 10^{-34}$J s/rad is the "reduced" Planck constant. These are the Planck Einstein relations. They describe light as composed of "particles" of energy, determined by the frequency.

1.3 Double-Slit Experiment

The double-slit experiment demonstrated that light is also an electromagnetic-wave, and can be understood by using the linearity of Maxwell's equations.

The light intensity is proportional to $|E|^2$, and if $\vec{E}(\vec{r}) = \vec{E}_1(\vec{r}) + \vec{E}_2(\vec{r})$ from the two sources, then

$$I \propto \left|\vec{E}_1(\vec{r}) + \vec{E}_2(\vec{r})\right|^2 \neq \left|\vec{E}_1(\vec{r})\right|^2 + \left|\vec{E}_2(\vec{r})\right|^2,$$

shows the wave nature of light.

This experiment still holds at extremely low intensities. It is still individual photons that show up on the screen, however when averaged over time it still results in the interference pattern.

1.4 Double-Slit with Particles

This experiment can be repeated using things that are normally thought of as particles, such as electron or neutrons. To find the required wavelength, we can use the de Broglie hypothesis

Definition 1.1. The de Broglie hypothesis states:
"to every particle is associated a wave, which obeys the same Planck-Einstein relations as applicable for photons."

From the relation above, we can derive

$$\boxed{\lambda = \frac{2\pi}{k} = \frac{h}{p}}$$

This is only significant for extremely small objects. The distance between intensity peaks is given by

$$\delta x = \lambda \frac{D}{d},$$

with $\lambda = h/p$, D is the distance between the slits and the screen, and d is the separation of the slits. This means that larger particles have smaller distances between intensity peaks, with all else being the same. To make a meaningful observation, we can

- Increase the wavelength by decreasing the momentum (smaller velocity for constant mass),
- Decrease the distance between the slits,
- Or increase the distance to the screen.

1.5 Wave Function

These two experiments force us to recognise the idea that particles are simultaneously particles and waves, and in fact that waves and particles are not different things. Instead, each wave is made of packets of energy, with is associated with wavelength $\lambda = h/p$ and frequency $v = E/h$.

Instead of trying to describe the trajectory of a particle with position $\vec{r}$ and momentum $\vec{p}$, we use a wave function $\psi(\vec{r}, t)$, which contains all the information we can gather on the particle. This wave function is a probability amplitude, such that

$$C \iiint_V |\psi(\vec{r}, t)|^2 d^3\vec{r} = P_V(t),$$

where $P_V(t)$ is the probability of finding the particle in the volume V at time t, and C is a normalisation constant to ensure the value is between 0 and 1, i.e.

$$C \iiint_{-\infty}^{\infty} |\psi(\vec{r}, t)|^2 d^3\vec{r} = 1,$$

i.e. the particle has to be somewhere. Now instead of calculating the trajectory of a particle, we instead calculate the probability of finding the particle in a certain region in space, and how that evolves over time.

We know that Maxwell's equations allow solutions in the form of propagating waves, e.g.

$$\vec{E}(\vec{r}, t) = Ae^{i(k\vec{r} - \omega t)}$$

is a plane wave that solve the wave equation of the form

$$\nabla^2 \vec{E} = \frac{1}{v^2} \frac{\partial^2 \vec{E}}{\partial t^2},$$

where $v = \omega/k$. This wave equation contains the partial second derivative of the electric field with respect to both time and space. Taking the second derivative of the plane wave results in the coefficients k^2 and ω^2 which give rise to the $1/v^2$ term in the equation. Note that v is constant.

Consider a free particle. The energy is purely kinetic, so

$$E = \frac{1}{2}mv^2 = \frac{p^2}{2m} = \frac{\hbar^2 k^2}{2m},$$

which we can impose Planck-Einstein to get

$$\hbar\omega = \frac{\hbar^2 k^2}{2m}.$$

An important thing to note is that the energy depends quadratically on k, but linearly on ω. This means that an appropriate wave equation should contain a first derivative with respect to time, and the second derivative with respect to space. Here, v is no longer constant.

1.6 Schrödinger Equation

Schrödinger postulated an equation that describes the time evolution of the wavefunction of quantum object to be

$$\boxed{i\hbar\frac{\partial\psi(\vec{r},t)}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2\psi(\vec{r},t) + V(\vec{r})\psi(\vec{r},t)}$$

$V(\vec{r})$ is a (position-dependent) potential energy. It is similar to a wave equation, containing the first derivative w.r.t. time, and the second w.r.t. space. The coefficients are chosen to impose the Planck-Einstein relations. Because of the first derivative w.r.t. time, the equation explicitly contains i.

Proof. Consider a free particle not subjected to any potentials or forces ($V(\vec{r}) = 0$):

$$i\hbar\frac{\partial\psi(\vec{r},t)}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2\psi(\vec{r},t).$$

This has a solution of the form

$$\psi(\vec{r},t) = Ae^{i(\vec{k}\cdot\vec{r}-\omega t)}$$

with the condition (see Problem 1.4 for a proof of this condition)

$$\hbar\omega = \frac{\hbar^2k^2}{2m}.$$

Inserting the Planck-Einstein relations, the above conditions means

$$E = \frac{p^2}{2m} = \frac{1}{2}mv^2.$$

□

1.7 Wave Packets

These wavefunctions can be combined into a single wave packet by simply summing them together (and hence with 3 wave numbers k). The interference between the different wave components causes the wave components to no longer be spread out equally in space, however it still have recurrences.

These recurrences get more spread out the more wave components are added, up to a limit of infinitely many components, each with a different wave number k and with a coefficient $g(k)$. For simplicity, take the 1D case ($\vec{r} \equiv x$):

$$\psi(x,t) = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\infty} g(k)e^{i(kx-\omega t)}dk.$$

We can find $g(k)$ by setting $t = 0$ (i.e. initial conditions)

$$\psi(x,0) = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\infty} g(k)e^{ikx}dk,$$

therefore

$$g(k) = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\infty} \psi(x,0)e^{-ikx}dx.$$

This means that $g(k)$ is the Fourier transform (in space) of the shape of the wave packet at $t = 0$, i.e. of the probability amplitude of finding the object at point x at $t = 0$.

1.8 Heisenbery Uncertainty Principle

Consider a Gaussian wave packet

$$\psi(x,0) = \frac{1}{\sqrt{2\pi}\sigma_x} \exp\left(-\frac{x^2}{2\sigma_x^2}\right).$$

It's Fourier transform is also a Gaussian

$$\begin{aligned} g(k) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(x,0) e^{-ikx} dx \\ &= \frac{1}{\sqrt{2\pi}\sigma_x} \exp\left(-\frac{k^2}{2\omega_k^2}\right). \end{aligned}$$

From the properties of Fourier transforms, we know that

$$\sigma_k = 1/\sigma_x \quad \Rightarrow \quad \sigma_k \sigma_x = 1.$$

Imposing the Planck-Einstein relation $p = \hbar k \Rightarrow \sigma_p = \hbar \sigma_k$, we get

$$\boxed{\sigma_p \sigma_x = \hbar}$$

This relation is a simple mathematical property of Fourier transform. This appears as soon as you assume the Planck-Einstein relations, i.e. wave-particle duality. This also holds if you assume the uncertainty principle as a postulate, then wave-particle duality is a consequence of the mathematical properties of Fourier transforms.

1.9 Problems

Problem 1.1. A particle of mass $1ng$ moves at $1mm/s$. Estimate its de Broglie wavelength and compare it to atomic dimensions

Answer:

The de Broglie hypothesis states that the wavelength of an object is given by

$$\lambda = \frac{2\pi}{k} = \frac{h}{p},$$

where λ is the wavelength, k is the wave vector, h is the Planck constant, and p is the classical momentum. Plugging in our values we get

$$\begin{aligned} p &= 1 \times 10^{-9} \times 1 \times 10^{-3} \\ &= 1 \times 10^{-12} \\ \Rightarrow \lambda &= \frac{h}{p} \\ &= \frac{6.626 \times 10^{-34}}{1 \times 10^{-12}} \\ &= 6.626 \times 10^{-22} m. \end{aligned}$$

The diameter of a proton is 1.68×10^{-15}, which is several orders of magnitude larger than the wavelength of the particle.

Problem 1.2. Using Heisenberg's uncertainty principle, estimate the minimum spread in velocity of an electron confined to a region of size $0.1nm$.

Answer:

Heisenberg's uncertainty principle states that

$$\sigma_x \sigma_p \geq \frac{\hbar}{2}.$$

Rearranging, we find the deviation of the momentum to be

$$\sigma_p \geq \frac{\sigma_x \hbar}{2}.$$

Plugging in our value for σ_x, we get

$$\begin{aligned}\sigma_p &\geq \frac{\sigma_x \hbar}{2} \\ &= \frac{1 \times 10^{-10} \times 1.054 \times 10^{-34}}{2} \\ &= 5.27 \times 10^{-45}.\end{aligned}$$

We can then solve for the velocity to get

$$\begin{aligned}\sigma_p &= \Delta(mv) \\ \Rightarrow \sigma_v &\geq \frac{5.27 \times 10^{-45}}{1 \times 10^{-10}} \\ &\geq 5.27 \times 10^{-35}.\end{aligned}$$

Problem 1.3. Show that for a Gaussian wave packet, the Fourier transform is also Gaussian, and derive $\sigma_k \sigma_x = 1$.

Problem 1.4. Verify that $\psi(\vec{r}, t) = Ae^{i(\vec{k}\cdot\vec{r} - \omega t)}$ solves the free particle Schrödinger equation only if $\hbar\omega = \frac{\hbar^2 k^2}{2m}$.

Answer:

We have the Schrödinger equation to be

$$i\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2 \psi(\vec{r}, t) + V(\vec{r})\psi(\vec{r}, t).$$

For the free particle case, there are no external interactions with the system, i.e. $V(\vec{r}) = 0$. This gives Schrödinger to be

$$i\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2 \psi(\vec{r}, t).$$

Plugging in our solution (and assuming a one-dimensioned case), we get

$$i\hbar \frac{\partial}{\partial t}\left[Ae^{i(\vec{k}r - \omega t)}\right] = -\frac{\hbar^2}{2m}\nabla^2\left[Ae^{i(\vec{k}r - \omega t)}\right]. \tag{1}$$

Since we are only considering the 1D case, we have the Laplacian to be $\nabla^2 = \frac{\partial^2}{\partial r^2}$, which gives

$$\begin{aligned}\nabla^2\left[Ae^{i(\vec{k}r - \omega t)}\right] &= \frac{\partial^2}{\partial r^2}\left[Ae^{i(\vec{k}r - \omega t)}\right] \\ &= -k^2 Ae^{i(\vec{k}r - \omega t)}.\end{aligned}$$

Now solving the LHS we get

$$\frac{\partial}{\partial t}\left[Ae^{i(\vec{k}r - \omega t)}\right] = -i\omega Ae^{i(\vec{k}r - \omega t)}.$$

Plugging both of these back into 1, we get

$$\begin{aligned} i\hbar\left[-i\omega A e^{i(\vec{k}r-\omega t)}\right] &= \frac{\hbar^2}{2m}\left[-k^2 A e^{i(\vec{k}r-\omega t)}\right] \\ \hbar\omega A e^{i(\vec{k}r-\omega t)} &= \frac{\hbar^2 k^2}{2m} A e^{i(\vec{k}r-\omega t)} \\ \therefore \hbar\omega &= \frac{\hbar^2 k^2}{2m}. \end{aligned}$$

2 The Mathematical Formalism of Quantum Mechanics

2.1 Hilbert Space

A function is a vector in an infinite-dimensioned vector space. It is sometimes convenient to represent the function using a finite number of "basis functions", so that the vector space necessary to describe the function has finite dimension. This is known as a Hilbert Space.

A vector space requires a suitable basis.

Example 2.1. 3D vector space $\mathbb{R}^3$.

Let $\{\vec{u}_x, \vec{u}_y, \vec{u}_z\}$ be an orthonormal basis in $\mathbb{R}^3$. We can then describe a vector in terms of this basis as

$$\vec{v} = x\vec{u}_x + y\vec{u}_y + z\vec{u}_z,$$

where x, y, and z are coefficients. We then have

$$|\vec{u}_\alpha| = 1, \quad \text{and} \quad \vec{u}_\alpha \cdot \vec{u}_\beta = \delta_{\alpha\beta}.$$

Once we have the basis, we can represent any vector in that basis as a vector of its coefficients in the chosen basis

$$\vec{v} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

We can denote this vector as $|v\rangle$, indicating it is a normal vector. Similarly, we can define the transpose of this vector as $\langle v|$. Note that we also have to take the complex conjugate of the elements in this vector as we are dealing with complex numbers. We then define the dot product of this to be

$$\vec{a} \cdot \vec{b} \to \vec{a}^T \vec{b} \, \langle a|b\rangle = \sum_{i=1}^{3} a_i^* b_i$$

This is known as Dirac notation.

2.2 Operators

2.2.1 Linear Operators

In a vector space, we can define linear operators (usually matrices) that transform a vector into another

$$|\psi'\rangle = A\,|\psi\rangle\,.$$

Note that $\langle\phi|\psi\rangle$ is a number, and $|\phi\rangle\,\langle\psi|$ is a matrix.

2.2.2 Adjoint Operators

Adjoint operators provide the correspondence between bras and kets

$$|\psi'\rangle = A\,|\psi\rangle \to \langle\psi'| = \langle\psi|\,A^\dagger.$$

Here, $A^\dagger$ is the adjoint of A. It is obtained by transposing and taking the complex conjugate of each element.

2.2.3 Hermitian Operators

If an operator is it's own adjoint (i.e. $A = A^\dagger$), then they are called Hermitian operators. The are simple to recognise

- The elements on opposite side of the diagonal are the complex conjugate of each
- The elements on the diagonal are purely real.

2.3 Basis of a Hilbert Space

A set of kets $\{|\nu_i\rangle\}$ is called orthonormal if

$$\langle \nu_i | \nu_j \rangle = \delta_{ij}.$$

The set $\{|\nu_i\rangle\}$ forms a basis of the Hilbert space ϵ if every $|\psi\rangle$ can be written as

$$|\psi\rangle = \sum_j c_j \, |\nu_j\rangle \, .$$

An important property is the closure relation

$$|\psi\rangle = \sum_j c_j \, |\nu_j\rangle = \sum_j \langle \nu_i | \psi \rangle \, |\nu_j\rangle = \sum_j |\nu_j\rangle \, \langle \nu_k | \psi \rangle \, .$$

For this to be true, it requires

$$\sum_j |\nu_j\rangle \, \langle \nu_j| = \mathbb{I}.$$

This is called the closure relation. The operators $|\nu_j\rangle \, \langle \nu_j|$ projects a vector onto the basis state $|\nu_j\rangle$.

2.4 Observables

General linear algebra theorems

- The eigenvalues of a Hermitian operator are real
- The eigenvectors corresponding to different eigenvalues are orthogonal

A Hermitian operator is called observable if its eigenvectors form a basis of the vector space. We call a unitary an operator for which holds

$$U^{-1} = U^\dagger.$$

The eigenvalues of unitary operators are complex number of modulus 1.

2.5 Problems

3 Spin and The Postulates of Quantum Mechanics

3.1 Postulate 1

The first postulate of quantum mechanics:

> **First Postulate of Quantum Mechanics 3.1.** At an initial time t_0, the state of a quantum system is specified by a ket $|\psi(t_0)\rangle$ that belongs to a suitable vector space ϵ.

This means that any ket

$$|\psi'\rangle = \lambda_1 \, |\psi_1\rangle + \lambda_2 \, |\psi_2\rangle \, ,$$

is also a legitimate state of the system. This is known as the superposition principle.

Note that we normally assume that the state vectors are normalised

$$\langle \psi | \psi \rangle = 1.$$

3.1.1 Application

Assuming a 2D vector space (spin-up and spin-down). We have a basis in this space to be

$$|\uparrow\rangle, |\downarrow\rangle.$$

They must satisfy the following conditions

- Normalisation: $\langle\downarrow|\downarrow\rangle = \langle\uparrow|\uparrow\rangle = 1$
- Orthogonality: $\langle\uparrow|\downarrow\rangle = 0$
- Closure: $|\downarrow\rangle\langle\downarrow| + |\uparrow\rangle\langle\uparrow| = \mathbb{I}$

There are an infinite number of options for this, but the simplest is

$$|\uparrow\rangle = |1\rangle = \begin{pmatrix}1 & 0\end{pmatrix}, \quad |\downarrow\rangle = |2\rangle = \begin{pmatrix}0 & 1\end{pmatrix}.$$

Then, any ket of the form $\alpha|1\rangle + \beta|2\rangle$, with $\alpha^2 + \beta^2 = 1$ is a legitimate state of the system

$$|\psi\rangle = \begin{pmatrix}\alpha & \beta\end{pmatrix}.$$

3.2 Postulate 2

The second postulate of quantum mechanics:

Second Postulate of Quantum Mechanics 3.1. Every measurable physical quantity $\mathcal{A}$ is described by an operator A, which must be an observable.

This means you can construct operators in your space ϵ that not observable, i.e. they cannot correspond to measurable physical quantities.

3.2.1 Application

The physical quantity measured in the Stern-Gerlach experiment is the z-component of the angular momentum $\mathcal{S}_z$. Since it is found to provide only two discrete results, the operator $\mathcal{S}_z$ must be an observable in a 2-dimensional Hilbert space, i.e. a 2×2 matrix.

3.3 Postulate 3

The third postulate of quantum mechanics:

Third Postulate of Quantum Mechanics. When measuring the value of the quantity $\mathcal{A}$, we can only obtain as a result one of the eigenvalues a_n of the observable A.

For an observable $\mathcal{A}$ with a discrete set of eigenvalues a_n, the corresponding physical quantity must be within that set of eigenvalues, not any arbitrary value. This is called quantization.

3.3.1 Application

Using the first two postulates, and the observed values of $\mathcal{S}_z$ of the Stern-Gerlach experiment, we can deduce that S_z must be a 2×2 observable with eigenvalues $\pm\hbar/2$. We can choose to define S_z as the matrix

$$S_z = \frac{\hbar}{2}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}.$$

Proof. We can prove that this is the correct matrix by multiplying them by the observed values

$$S_z |\uparrow\rangle = \frac{\hbar}{2}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\begin{pmatrix}1\\0\end{pmatrix} = \frac{\hbar}{2}\begin{pmatrix}1\\0\end{pmatrix} = \frac{\hbar}{2}|\uparrow\rangle$$
$$S_z |\downarrow\rangle = \frac{\hbar}{2}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\begin{pmatrix}0\\1\end{pmatrix} = -\frac{\hbar}{2}\begin{pmatrix}0\\1\end{pmatrix} = -\frac{\hbar}{2}|\downarrow\rangle,$$

i.e. $|\uparrow\rangle$ and $|\downarrow\rangle$ are eigenvectors of S_z with eigenvalues $\pm\hbar/2$. □

3.4 Postulate 4

Fourth Postulate of Quantum Mechanics. When measuring the quantity $\mathcal{A}$ on a system described by the state $|\psi\rangle$, the probability of obtaining as a result the eigenvalue a_n is given by

$$\boxed{\mathcal{P}(a_n) = |\langle u_n|\psi\rangle|^2}$$

where $|u_n\rangle$ is the eigenvector of A with eigenvalue a_n.

This means the result of the measurement of a quantum system is (generally) probabilistic.

3.4.1 Application

Consider an atom with spin oriented along $\vec{u}$, entering the Stern-Gerlach apparatus. The state of the atom just before the measurement is

$$\left|\psi(t^-)\right\rangle = \cos\frac{\theta}{2}e^{-\frac{i\phi}{2}}|\downarrow\rangle + \sin\frac{\theta}{2}e^{\frac{i\phi}{2}}|\downarrow\rangle = \begin{pmatrix}\cos\frac{\theta}{2}e^{-\frac{i\phi}{2}}\\ \sin\frac{\theta}{2}e^{\frac{i\phi}{2}}\end{pmatrix}.$$

According to the Born's rule (see Def. 3.1),the probability to find the eigenvalue $+\hbar/2$ corresponding to the eigenvector $|\uparrow\rangle$ is

$$\begin{aligned}\mathcal{P}_\uparrow &= \left|\left\langle\uparrow\middle|\psi(t^-)\right\rangle\right|^2\\ &= \left|(1\quad 0)\begin{pmatrix}\cos\frac{\theta}{2}e^{-\frac{i\phi}{2}}\\ \sin\frac{\theta}{2}e^{\frac{i\phi}{2}}\end{pmatrix}\right|^2\\ &= \cos^2\frac{\theta}{2}.\end{aligned}$$

This result is independent of ϕ, since the apparatus only measures the projection along z.

Definition 3.1. Born's Rule

> The Born rule is a postulate of quantum mechanics that gives the probability that a measurement of a quantum system will yield a given result. It states that the probability density for finding a particle at a given position is proportional to the square of the amplitude of the system's wavefunction at that position. (Wikipedia)

It states that an observable that was measured within a quantum system with normalised wavefunction $|\psi\rangle$ corresponds to a self-adjoint operator A whose spectrum is discrete if the measured result will be one of the eigenvalues λ of A, and the probability of measuring a given eigenvalue λ will equal $\langle\psi| P_i |\psi\rangle$, where P_i is the projection onto the eigenspace of A corresponding to λ_i.

For example, a single structureless particle that can be respresented by a wavefunction ψ that depends on position (x, y, z) and time t has the probability density function p for the result of a measurement of the particle at any particular time t_0 to be

$$p(x, y, z, t_0) = |\psi(x, y, z, t_0)|^2.$$

Example 3.2. Take the spin of a particle $\theta = \frac{\pi}{2}$ (spin in the xy-plane). The chance of finding the particle pointing "up" is

$$\mathcal{P}_\uparrow = \cos^2 \frac{pi}{4} = \frac{1}{2}.$$

3.5 Postulate 5

Fifth Postulate of Quantum Mechanics. If the measurement of $\mathcal{A}$ at time t has given the result a_n, immediately after the measurement the system is found in the normalised projection of the state $|\psi(t^-)\rangle$ before the measurement onto the eigenvector $|u_n\rangle$ corresponding to the measured eigenvalue a_n.

$$|\psi(t^-)\rangle \to |\psi(t^+)\rangle = \frac{P_n |\psi(t^-)\rangle}{\sqrt{\langle\psi(t^-)| P_n \langle\psi(t^-)|}},$$

where $P_n = |u_n\rangle \langle u_n|$ (Section 3.8) is the projection operator onto the eigenstate $|u_n\rangle$. The denominator normalises the wavefunction.

This is known as the collapse of the wavefunction, and is where taking the measurement of a system "collapses" it into one it's possible states, with the probability of collapsing into this state being defined by the previous Postulate. In general, measuring some property of the system changes its state.

3.5.1 Application

The Stern-Gerlach apparatus can be used to prepare a known quantum state. If we pierce a hole in the screen where the "up" atoms would collide, then we know that those atoms are in the state $|\psi(t^+)\rangle = |\uparrow\rangle$. This new "up" only beam can then be fed into another apparatus to verify this.

Now consider the second apparatus being rotated along $\vec{u}$ instead of purely vertical. Even though we are certain that all the particles are in the state $|\uparrow\rangle$, the outcome of the measurement is not certain. This is because the second apparatus is now measuring $\mathcal{S}_u$ instead of $\mathcal{S}_z$.

We know from Born's rule the probability that each of the eigenvalues may appear as the outcome of the measurement, because we know the quantum state $|\psi\rangle$. Therefore, the average value (called the expectation value) of an observable A given the state $|\psi\rangle$ is

$$\langle A \rangle_\psi = \sum_n a_n \mathcal{P}(a_n) = \sum_n a_n |\langle u_n|\psi\rangle|^2 .$$

Note: given two vectors $|\psi\rangle$ and $|\phi\rangle$, $\langle\phi|\psi\rangle = \langle\psi|\phi\rangle^*$. Factorising further we get

$$\begin{aligned}
&= \sum_n a_n |\langle u_n|\psi\rangle|^2 \\
&= \sum_n a_n \langle\psi|u_n\rangle \langle u_n|\psi\rangle \\
&= \sum_n \langle\psi| A |u_n\rangle \langle u_n|\psi\rangle \quad \text{note: } a_n |u_n\rangle = A |u_n\rangle \\
&= \langle\psi| A \left[\sum_n |u_n\rangle \langle u_n|\right] |\psi\rangle \quad \text{closure relation} \\
&= \langle\psi| A |\psi\rangle .
\end{aligned}$$

In the example from before:

$$\begin{aligned}
\langle S_u \rangle_{|\uparrow\rangle} &= \langle\uparrow| S_u |\uparrow\rangle \\
&= (1 \quad 0) \frac{\hbar}{2} \begin{pmatrix} \cos\theta & \sin\theta \cdot e^{-i\phi} \\ \sin\theta & -\cos\theta \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= \frac{\hbar}{2} \cos\theta.
\end{aligned}$$

3.6 Postulate 6

Sixth Postulate of Quantum Mechanics. If a system is in a state $|\psi(t)\rangle$ at time t, its subsequent time evolution is governed by the Schrödinger equation

$$i\hbar\frac{\partial|\psi(t)\rangle}{\partial t} = \mathcal{H}(t)|\psi(t)\rangle$$

where $\mathcal{H}(t)$ is the Hamiltonian of the system, i.e. the operator that describes its total energy.

Quantum mechanics is for the most part deterministic. Given an initial state, the system evolves in a predictable way, according to a linear differential equation called the Schrödinger equation. The probabilistic aspects enter when you try to measure the state.

3.6.1 Application

We have Schrödinger's equation to be (as above)

$$i\hbar\frac{\partial|\psi(t)\rangle}{\partial t} = \mathcal{H}(t)|\psi(t)\rangle\,,$$

where $\mathcal{H}(t)$ is a matrix, $|\psi\rangle$ is a vector, so the Schrödinger equation is a system of linear differential equations. We can write the formal solution as

$$|\psi(t)\rangle = U(t,t_0)|\psi(t_0)\rangle\,,$$

with the condition

$$U(t_0,t_0) = \mathbb{I}.$$

Substituting $|\psi(t)\rangle = U(t,t_0)|\psi(t_0)\rangle$ into the Schrödinger equation gives

$$i\hbar\frac{d}{dt}U(t,t_0) = \mathcal{H}(t)U(t,t_0).$$

Assuming for simplicity that the Hamiltonian is constant, i.e. $\mathcal{H}(t)\to\mathcal{H}$. The differential equation can then be formally integrated to yield

$$\begin{aligned}U(t,t_0) &= \int_{t_0}^{t} -i\frac{\mathcal{H}}{\hbar}U(t',t_0)\\ &= \exp\left[-i\frac{\mathcal{H}}{\hbar}(t-t_0)\right].\end{aligned}$$

Thus, knowing $|\psi(t_0)\rangle$, the quantum state at later times is given by

$$|\psi(t)\rangle = e^{-i\frac{\mathcal{H}}{\hbar}(t-t_0)}|\psi(t_0)\rangle\,,$$

where

$$\boxed{U(t,t_0) = e^{-i\frac{\mathcal{H}}{\hbar}(t-t_0)}}$$

is the time-evolution operator.

3.7 Spin Matrices

In Section 3.3, we had the S_z matrix which described the spin of a particle. We also have the spin projection along x and y to be

$$S_x = \frac{\hbar}{2}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}, \quad S_y = \frac{\hbar}{2}\begin{pmatrix}0 & -i\\ i & 0\end{pmatrix}.$$

(Note that these are the Pauli matrices σ_x, σ_y, σ_z with a prefactor of $\hbar/2$). These matrices have the eigenvectors

$$|X\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle + |\downarrow\rangle\right) \qquad |-X\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle - |\downarrow\rangle\right)$$
$$|Y\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle + i\,|\downarrow\rangle\right) \qquad |-Y\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle - i\,|\downarrow\rangle\right).$$

Consider an arbitrary unit vector $\vec{u}$ in 3-dimensional space. The observable corresponding to the spin projection along $\vec{u}$ is given by

$$\begin{aligned} S_u &= S_x \sin\theta\cos\phi + S_y \sin\theta\sin\phi + S_z\cos\theta \\ &= \frac{\hbar}{2}\begin{pmatrix} \cos\theta & \sin\theta\cdot e^{-i\phi} \\ \sin\theta\cdot e^{i\phi} & -\cos\theta \end{pmatrix}. \end{aligned}$$

The eigenvectors of this spin projection are then

$$\begin{aligned} |U\rangle &= \cos\frac{\theta}{2}e^{-\frac{i\phi}{2}}\,|\uparrow\rangle + \sin\frac{\theta}{2}e^{\frac{i\phi}{2}}\,|\downarrow\rangle \\ |-U\rangle &= -\sin\frac{\theta}{2}e^{-\frac{i\phi}{2}}\,|\uparrow\rangle + \cos\frac{\theta}{2}e^{\frac{i\phi}{2}}\,|\downarrow\rangle. \end{aligned}$$

3.8 Projection Operator

A projection operator extracts the component of a quantum state lying in a chosen direction or subspace. It is used in Born's rule (Def. 3.1) to define the probability of making a specific measurement, or to transform a vector from one basis to another.

In order for a matrix to be a projection matrix, it must be idempotent ($\hat{P}^2 = \hat{P}$), and if it is also Hermitian is called an orthogonal projection matrix (otherwise it is an oblique projection matrix).

Let $|u_n\rangle$ be an eigenstate in some basis. The projection operator is then given by

$$P_n = |u_n\rangle\langle u_n|.$$

To projection a ket $|\chi\rangle$ onto this eigenstate is given by

$$|\chi\rangle_n = P_n|\chi\rangle = |u_n\rangle\langle u_n|\chi\rangle,$$

which gives the probability of finding $|\chi\rangle$ in the eigenstate $|u_n\rangle$. See Problem 3.1 for a proof that $|u_n\rangle\langle u_n|$ acts a projection operator.

3.9 Exponential of a Matrix

We can use Taylor Expansion to calculate some arbitrary function of a matrix, such as e^A.

$$e^x = \sum_{n=0}^{\infty}\frac{x^n}{n!} = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \cdots.$$

By analogy

$$e^A = \sum_{n=0}^{\infty}\frac{A^n}{n!} = \mathbb{I} + A + \frac{A^2}{2} + \frac{A^3}{6} + \cdots.$$

Note: if A is diagonal, A^n is also diagonal, and every function of A is diagonal also

$$A = \begin{pmatrix} a_1 & 0 & 0 \\ 0\cdots & 0 & \\ 0 & 0 & a_N \end{pmatrix} \rightarrow \quad e^A = \begin{pmatrix} e^{a_1} & 0 & 0 \\ 0 & \cdots & 0 \\ 0 & 0 & e^{a_N} \end{pmatrix}.$$

In general, if $|\phi\rangle$ is an eigenvector of A with eigenvalue a, then $|\phi\rangle$ is also an eigenvector of $F(A)$ with eigenvalue $F(a)$.

3.10 Hamiltonian Eigenbasis

The Hamiltonian is an observable physical quantity, so it must be described by a matrix whose eigenvectors form a basis of the vector space. If we choose to use the eigenvectors of the Hamiltonian as our basis, it means the Hamiltonian appears as a diagonal matrix

$$\mathcal{H} = \begin{pmatrix} E_1 & 0 & 0 \\ 0 & \cdots & 0 \\ 0 & 0 & E_N \end{pmatrix},$$

where $E_1, \cdots, E_N$ are the eigenvalues (in this case they are "eigenenergies"). Therefore, in the Hamiltonian eigenbasis, the time evolution operator is simply

$$U(t, t_0) = \begin{pmatrix} e^{-i\frac{E_1}{\hbar}(t-t_0)} & 0 & 0 \\ 0 & \cdots & 0 \\ 0 & 0 & e^{-i\frac{E_N}{\hbar}(t-t_0)} \end{pmatrix}.$$

If you can describe the system in the Hamiltonian eigenbasis, the time evolution operator can calculated by hand.

Example 3.1. Spin Precession

Classical description: Consider a magnetic moment $\vec{M} = \gamma\vec{S}$ placed in a uniform magnetic field $\vec{B}$. If $\vec{S}$ and $\vec{B}$ are not parallel, there is a torque $\vec{T}$ acting on the angular momentum

$$\vec{T} = \gamma\vec{S} \times \vec{B}.$$

The derivative of the angular momentum is equal to the torque

$$\frac{d\vec{S}}{dt} = \vec{T} = \gamma\vec{S} \times \vec{B}.$$

We can then define the quantity of angular momentum

$$\omega = -\gamma B.$$

Quantum Description: The energy of a magnetic moment $\vec{M} = \gamma\vec{S}$ placed in a uniform magnetic field $\vec{B}$ is

$$W = -\gamma\vec{S} \cdot \vec{B}.$$

Assume that $\vec{B} \parallel \hat{z}$. Since there are no other relevant contribution to the energy of the system, the Hamiltonian is

$$\begin{aligned} \mathcal{H} &= -\gamma S_z B \\ &= -\gamma B \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \omega \frac{\hbar}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} \frac{\hbar\omega}{2} & 0 \\ 0 & -\frac{\hbar\omega}{2} \end{pmatrix}. \end{aligned}$$

Note that this Hamiltonian is time-independent if the magnetic field is constant (which we are assuming).

This Hamiltonian then has the two eigenenergies

$$E_1 = \frac{\hbar\omega}{2}, \quad E_2 = -\frac{\hbar\omega}{2}.$$

We can then plug these into the time evolution operator

$$U(t,t_0) = e^{-i\frac{\mathcal{H}}{\hbar}(t-t_0)} = \begin{pmatrix} e^{-i\frac{E_1}{\hbar}(t-t_0)} & 0 \\ 0 & e^{-i\frac{E_2}{\hbar}(t-t_0)} \end{pmatrix} = \begin{pmatrix} e^{-i\frac{\omega}{2}(t-t_0)} & 0 \\ 0 & e^{i\frac{\omega}{2}(t-t_0)} \end{pmatrix}.$$

This was simple because we used the eigenstates of the Hamiltonian as our basis. We can then prepare the spin in the symmetric eigenstate of S_x

$$|\psi(0)\rangle = |X\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle + |\downarrow\rangle\right) = \frac{1}{\sqrt{2}}\begin{pmatrix}1\\1\end{pmatrix}.$$

At a later time t, the state will be

$$|\psi(t)\rangle = U(t,0)\,|\psi(0)\rangle = \begin{pmatrix} e^{-i\frac{\omega}{2}t} & 0 \\ 0 & e^{i\frac{\omega}{2}t} \end{pmatrix}\frac{1}{\sqrt{2}}\begin{pmatrix}1\\1\end{pmatrix} = \frac{1}{\sqrt{2}}\begin{pmatrix} e^{-i\frac{\omega}{2}t} \\ e^{i\frac{\omega}{2}t} \end{pmatrix}.$$

To make sense of the results of a wavefunction, we use the expectation value of the wavefunction, i.e. the wavefunction averaged over time. From the previous example, we want to calculate the state along the three spatial axes to see how it evolves over time.

Example 3.2. Calculate the expectation value of the wavefunction in the previous problem.

Starting with the z axis, we have

$$\begin{aligned}\langle S_z\rangle_{\psi(t)} &= \langle\psi(t)|\,S_z\,|\psi(t)\rangle \\ &= \frac{\hbar}{4}\begin{pmatrix} e^{i\frac{\omega}{2}t} & e^{-i\frac{\omega}{2}t}\end{pmatrix}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\begin{pmatrix} e^{-i\frac{\omega}{2}t} \\ e^{i\frac{\omega}{2}t}\end{pmatrix} \\ &= 0. \end{aligned} \tag{2}$$

Then the x axis,

$$\begin{aligned}\langle S_x\rangle_{\psi(t)} &= \langle\psi(t)|\,S_x\,|\psi(t)\rangle \\ &= \frac{\hbar}{4}\begin{pmatrix} e^{i\frac{\omega}{2}t} & e^{-i\frac{\omega}{2}t}\end{pmatrix}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}\begin{pmatrix} e^{-i\frac{\omega}{2}t} \\ e^{i\frac{\omega}{2}t}\end{pmatrix} \\ &= \frac{\hbar}{4}(e^{i\omega t} + e^{-i\omega t}) = \frac{\hbar}{2}\cos\omega t. \end{aligned} \tag{3}$$

And finally for the y axis

$$\begin{aligned}\langle S_y\rangle_{\psi(t)} &= \langle\psi(t)|\,S_y\,|\psi(t)\rangle \\ &= \frac{\hbar}{4}\begin{pmatrix} e^{i\frac{\omega}{2}t} & e^{-i\frac{\omega}{2}t}\end{pmatrix}\begin{pmatrix}0 & -i\\ i & 0\end{pmatrix}\begin{pmatrix} e^{-i\frac{\omega}{2}t} \\ e^{i\frac{\omega}{2}t}\end{pmatrix} \\ &= \frac{\hbar}{4}(-ie^{i\omega t} + ie^{-i\omega t}) = \frac{\hbar}{2}\sin(\omega t). \end{aligned} \tag{4}$$

From 2, we can see that there is no projection along the vertical (z) axis, meaning it remains within the equatorial (xy) plane. From 3 and 4, we see that the x and y projections oscillate 90 out of phase with frequency ω. This is known as Larmor precession.

3.11 Time Evolution Operator

The time evolution operator $U(t,t_0)$ has some important properties:

- It is a unitary operator

- Its eigenvalues are complex numbers with modulus 1, and it conserves the modulus of any vector it operates on (modulus is the magnitude of the complex number)
- It is time reversible.

3.11.1 Unitary Operators

A unitary operator is a bounded linear operator (a linear transformation between vector spaces that maps bounded sets to bounded sets) on a Hilbert space that satisfies $U^\dagger U = UU^\dagger = \mathbb{I}$, where $U^\dagger$ is the adjoint of U.

Example 3.1. Common Unitary Operators

- The identity function (identity matrix) is trivially a unitary operator
- Rotations in $\mathbb{R}^2$ are the simplest nontrivial example of unitary operators.
- The Fourier operator is a unitary operator (i.e. the operator that performs the Fourier transform with proper normalisation).
- Quantum logic gates are unitary operators, however they are not all Hermitian.

3.11.2 Time Reversibility

For simplicity, take $t_0 = 0$. If we know $U(t,0) = e^{-i\frac{\mathcal{H}}{\hbar}t}$, then

$$U(-t,0) = e^{i\frac{\mathcal{H}}{\hbar}t} = U^{-1}(t,0).$$

This means we can track the system back in time by inverting the time evolution operator. This is especially important for quantum computers as (along with being unitary) this imposes that no information can be lost.

3.12 Commutation

In general, the product of matrices is not commutative. However, we can define the commutator of two operators Λ and Ω as

$$[\Lambda, \Omega] = \Lambda\Omega - \Omega\Lambda.$$

Note that the commutator is itself an operator. When this commutator is 0, we say that the operators commute, so they can be multiplied in any order.

In quantum mechanics, commutation relations have an important meaning. When two observables do not commute, it means the order in which the observables are measured changes the outcomes.

Example 3.1. What is the commutation of S_x and S_z?

Calculating the commutation, we have

$$S_z S_x = \frac{\hbar^2}{4}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix} = \frac{\hbar^2}{4}\begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}$$
$$S_x S_z = \frac{\hbar^2}{4}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix} = \frac{\hbar^2}{4}\begin{pmatrix}0 & -1\\ 1 & 0\end{pmatrix}.$$

The commutator is then

$$[S_z, S_x] = S_z S_x - S_x S_z = \frac{\hbar^2}{4}\begin{pmatrix}0 & 2\\ -2 & 0\end{pmatrix} = i\hbar\frac{\hbar}{2}\begin{pmatrix}0 & -i\\ i & 0\end{pmatrix} = i\hbar S_y.$$

Similarly,

$$[S_x, S_y] = i\hbar S_z$$
$$[S_y, S_z] = i\hbar S_x.$$

This means that in the example where there was two Stern-Gerlach machines, the order in which measurements are made matters.

3.12.1 Ehrenfest Theorem

The expectation value of an observable A at time t is given by

$$\langle A \rangle (t) = \langle \psi(t) | A | \psi(t) \rangle .$$

To see how to evolves with time, we can assume A is time-independent, and differentiate with respect to t to get

$$\frac{d}{dt} \langle \psi(t) | A | \psi(t) \rangle = \left[\frac{d}{dt} \langle \psi(t) | \right] A | \psi(t) \rangle + \langle \psi(t) | A \left[\frac{d}{dt} | \psi(t) \rangle \right] .$$

And from the Schrödinger equation, we have

$$\begin{aligned} \frac{d}{dt} | \psi(t) \rangle = \frac{1}{i\hbar} \mathcal{H} | \psi(t) \rangle \Rightarrow \frac{d}{dt} \langle \psi(t) | &= -\frac{1}{i\hbar} \langle \psi(t) | \mathcal{H} \\ \frac{d}{dt} \langle \psi(t) | A | \psi(t) \rangle = \frac{1}{i\hbar} \langle \psi(t) | A\mathcal{H} - \mathcal{H}A | \psi(t) \rangle \Rightarrow i\hbar \frac{d}{dt} \langle A \rangle &= \langle [A, \mathcal{H}] \rangle . \end{aligned}$$

An observable that commutes with the Hamiltonian is a constant of the motion.

3.13 Pauli Matrices

Let us define the following matrices

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} .$$

These are called the Pauli matrices, and together with the identity matrix $\mathbb{I}$, they form a basis of the vector space of 2×2 matrices. This means that any 2×2 matrix with complex coefficients can be written as

$$A = a_1 \mathbb{I} + a_x \sigma_x + a_y \sigma_y + a_z \sigma_z .$$

Any 2-level system can be "mapped" onto a spin, or equivalently, all quantum mechanics 2-level systems are equivalent. For a "true" spin, the Pauli matrices are proportional to the operators that describe the spin orientation along the 3 axes in real space. There are other 2-level systems for which there is no such meaning.

The three matrices σ_x, σ_y, and σ_z can be grouped into a "vector matrices"

$$\vec{\sigma} = \begin{pmatrix} \sigma_x \\ \sigma_y \\ \sigma_z \end{pmatrix} .$$

This is called vector Pauli matrix. It is useful when calculating the inner product of a real vector in 3 dimensions, with the 3 spin operators.

Example 3.1. The Hamiltonian of a spin in a magnetic field $\vec{B}$ is

$$\mathcal{H} = -\gamma \vec{B} \cdot \vec{S} = -\gamma \frac{\hbar}{2} \vec{B} \cdot \vec{\sigma} = -\gamma \frac{\hbar}{2} \left(B_x \sigma_x + B_y \sigma_y + B_z \sigma_z \right) .$$

3.14 Non-Hermitian Operators

All the operators so far have been Hermitian (except the time evolution operator), and described with physical observables. Now consider the following operators

$$\sigma_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \sigma_- = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

These are clearly non-Hermitian. Therefore, they cannot represent physical observables. But, they are useful tools. For example

$$\sigma_+ |\downarrow\rangle = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} = |\uparrow\rangle .$$

σ_+ is a "flip up" operator. However,

$$\sigma_+ |\uparrow\rangle = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

applying σ_+ to the spin-up results in the zero-vector, with is not a legitimate state vector as it is not normalised. One thing to note is that these matrices can be created by outer product of the spin-up and spin-down eigenstates with themselves, and act as a projection operator.

3.15 Problems

Problem 3.1. Show that $|\psi\rangle \langle\psi|$ acts as a projection operator, and verify the closure relation

$$\sum |\psi\rangle \langle\psi| = 1$$

for an orthonormal basis.

Answer

Let $\hat{P}_\psi \equiv |\psi\rangle \langle\psi|$. When applied to a ket, the bra must bind to the ket, giving

$$\hat{P}_\psi = |\psi\rangle \langle\psi|\chi\rangle = \langle\psi|\chi\rangle |\psi\rangle .$$

Thus it extracts the component of $|\chi\rangle$ along $|\psi\rangle$. If $|\psi\rangle \perp |\chi\rangle$, then $\langle\psi|\chi\rangle = 0$, so $\hat{P}_\psi |\chi\rangle = 0$.

In order for $\hat{P}_\psi$ to be a projection operator, it must satisfy

$$\hat{P}_\psi^2 = \hat{P}_\psi,$$

i.e. applying the operator to the same ket twice is the same as applying only once. If we assume $|\psi\rangle$ is normalised ($\langle\psi|\psi\rangle = 1$), then

$$\begin{aligned} \hat{P}_\psi^2 &= (|\psi\rangle \langle\psi|) (|\psi\rangle \langle\psi|) \\ &= |\psi\rangle \langle\psi|\psi\rangle \langle\psi| \\ &= |\psi\rangle (1) \langle\psi| \\ &= \hat{P}_\psi. \end{aligned}$$

Now checking for Hermiticity, using $(AB)^\dagger = B^\dagger A^\dagger$, $|\psi\rangle^\dagger = \langle\psi|$, and $\langle\psi|^\dagger = |\psi\rangle$ we get

$$\begin{aligned} \hat{P}_\psi^\dagger &= (|\psi\rangle \langle\psi|)^\dagger \\ &= \left(\langle\psi|^\dagger\right) \left(|\psi\rangle^\dagger\right) \\ &= |\psi\rangle \langle\psi| \\ &= \hat{P}_\psi. \end{aligned}$$

Finally, we can check for orthogonality of basis projectors. Let $\{|i\rangle\}$ be an orthonormal basis, $\langle i|j\rangle = \delta_{ij}$. Define $\hat{P}_i = |i\rangle\langle i|$. Then,

$$\begin{aligned}\hat{P}_i\hat{P}_j &= |i\rangle\langle i|j\rangle|j\rangle \\ &= \delta_{ij}|i\rangle\langle j|.\end{aligned}$$

Hence,

$$\hat{P}_i\hat{P}_j = \begin{cases}\hat{P}_i & i = j \\ 0 & i \neq j\end{cases}.$$

This means projections onto different basis directions do not interfere with each other. We can now use these results to verify the closure relation by having the operator act on an arbitrary ket $|\chi\rangle$

$$\begin{aligned}\left(\sum_i |i\rangle\langle j|\right)|\chi\rangle &= \sum_i |i\rangle\langle i|\chi\rangle \\ &= \sum_i c_i|i\rangle, \quad c_i = \langle i|\chi\rangle.\end{aligned}$$

This is the same as the expansion of $|\chi\rangle$ on the complete orthonormal basis $\{|i\rangle\}$

$$|\chi\rangle = \sum_i c_i|i\rangle.$$

Since the two operators act the same on every ket,

$$\sum_i |i\rangle\langle i| = \mathbb{I}.$$

4 Quantum Mechanics in Real Space

4.1 Schrödinger Equations in Real Space

When we are solving the Schrödinger equation, we generally want to solve it in the form

$$\psi(x,t) = \phi(x)\cdot\chi(t),$$

i.e. we want to separate the spatial and temporal components. This gives the S.E. to be

$$i\hbar\phi(x)\frac{\partial\chi(t)}{\partial t} = -\frac{\hbar^2}{2m}\chi(t)\frac{\partial^2}{\partial x^2}\phi(x) + \chi(t)V(x)\phi(x),$$

and dividing both sides by $\phi(x)\cdot\chi(t)$

$$\frac{i\hbar}{\chi(t)}\frac{\partial\chi(t)}{\partial t} = \frac{1}{\phi(x)}\left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2}\phi(x)\right] + V(x).$$

This gives the S.E. to have the temporal-related components on the LHS, and the spatial related components on the RHS. This can only hold if both sides of the equation are a constant, which we set equal to $\hbar\omega$.

We can define the forms of the wavefunctions as below. Starting with the LHS

$$i\hbar\frac{\partial\chi(t)}{\partial t} = \hbar\omega\cdot\chi(t) \Rightarrow \chi(t) = A\cdot\exp(-i\omega t).$$

And with the RHS

$$-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2}\phi(x) + V(x)\phi(x) = \hbar\omega\cdot\phi(x).$$

This is just the "real space" version of the TISE

$$\mathcal{H}\phi(x) = E\phi(x) \quad \rightarrow \quad \left[\frac{-\hbar^2}{2m}\frac{\partial^2}{\partial x^2} + V(x)\right]\phi(x) = E\phi(x),$$

and therefore the function

$$\psi(x,t) = \phi(x)\exp(-i\omega t)$$

is a solution of the Schrödinger equation, if $\phi(x)$ is a solution of the time-independent part. The function $\psi(x,t)$ written above is a stationary state.

4.2 Stationary States

Whilst the function does contain t in the exponent, it isn't time-dependent as this is just a phase-factor which does not matter in isolation.

Proof. Consider an arbitrary ket $|\psi(t)\rangle = |\psi_0\rangle\, e^{-i\omega t}$ and an observable $\mathcal{A}$. The expectation value of the observable is given by

$$\begin{aligned}\langle A\rangle_\psi &= \langle\psi(t)|\,\mathcal{A}\,|\psi(t)\rangle \\ &= \langle\psi_0|\, e^{i\omega t}\mathcal{A}e^{-i\omega t}\,|\psi_0\rangle \\ &= \langle\psi_0|\, A\,|\psi_0\rangle\,,\end{aligned}$$

which is strictly time-independent. □

Quantum systems can be described using density matrices, which are simply the ket and bra multiplied together, $\rho = |\psi\rangle\langle\psi|$. In this case, the density matrix for $|\psi(t)\rangle$ is

$$\rho = |\psi(t)\rangle\langle\psi(t)| = |\psi_0\rangle\, e^{-i\omega t}e^{i\omega t}\langle\psi_0| = |\psi_0\rangle\langle\psi_0|\,,$$

so the phase factor disappears.

As the S.E. is linear, this means a linear equation of its solutions is also a solution

$$\psi(x,t) = \sum_n c_n\psi_n(x)e^{-i\omega_n t}.$$

This means if we know the stationary states $\psi_n(t)$ and their energies $E_n = \hbar\omega_n$, the problem can be easily solved if the initial state is written as a linear combination of stationary states

$$\psi(x,0) = \sum_n c_n\psi_n(x).$$

When considering the wavefunctions as vectors in an infinite-dimensioned space, this is essentially using a diagonal time-evolution operator.

Example 4.1. Consider a system with potential energy being given by

$$V(x) = \begin{cases} 0 & 0 < x < a \\ \infty & \text{otherwise}\end{cases}.$$

The particles cannot access this region outside $(0,a)$ due to the infinite potential required to exceed it. This imposes $\psi = 0$ at and beyond the walls. Note that $\frac{\partial\phi}{\partial x}$ does not need to be continuous. Our general solution (Ansatz) is

$$\psi(x) = Ae^{ikx} + Be^{-ikx}.$$

With our boundary condition at $x = a$, we have

$$\psi(a) = 0 \Rightarrow k_n = \frac{n\pi}{a}, n \in \mathbb{Z}.$$

Therefore, the energy eigenvalues $E_n = \frac{\hbar^2 k_n^2}{2m}$ are

$$E_n = \frac{n^2\pi^2\hbar^2}{2ma^2}.$$

Imposing the normalisation of the wavefunction, $\int |\psi(x)|^2\, dx = 1$, we get

$$\boxed{\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right)}$$

The boundary conditions have the effect of selecting a discrete set of possible momenta, and therefore possible energies, for the particle in the well. The stationary states are standing waves in the well.

4.3 Low-Energy Truncation

Since $\mathcal{H}$ is an observable, its eigenvalues $\psi_n(x)$ form a basis of the Hilbert space. In principle, $n \to \infty$, however in practice this can be truncated to a finite set of N low-energy state, for example becayse the temperature is finite, so the very high-energy states are always unoccupied. Therefore, if the Hilbert space is truncated to a finite dimensioned N, we can write a finite set of basis kets as

$$|\psi_1\rangle = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad |\psi_2\rangle = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix} \quad |\psi_3\rangle = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix},$$

where the Hamiltonian is simply

$$\mathcal{H} = \begin{pmatrix} E_1 & 0 & \cdots & 0 \\ 0 & E_2 & \cdots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & E_N \end{pmatrix}.$$

Each ket $|\psi_n\rangle$ corresponds to the function of space $\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right)$.

4.4 Non-Stationary States

A linear superposition of states is also a legitimate state

$$|\psi(t)\rangle = \sum_n c_n(t)\, |\phi_n\rangle\,.$$

Example 4.1. Evaluate

$$|\psi(0)\rangle = \frac{1}{\sqrt{2}} \left(|\phi_1\rangle + |\phi_2\rangle\right).$$

Since we are using the basis of the eigenstates of the Hamiltonian, the time-evolution operator is diagonal

$$U(t,0) = \begin{pmatrix} e^{-iE_1 t/\hbar} & 0 & \cdots & 0 \\ 0 & e^{iE_2 t/\hbar} & \cdots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & e^{-iE_N t/\hbar} \end{pmatrix}.$$

The time-evolved state is then given by

$$|\psi(t)\rangle = U(t,0)\,|\psi(0)\rangle = \frac{1}{\sqrt{2}}\left(e^{-iE_1t/\hbar}\,|\phi_1\rangle + e^{-iE_2t/\hbar}\,|\phi_2\rangle\right).$$

Multiplying by $e^{iE_1t/\hbar}$, we obtain

$$|\psi(t)\rangle\, e^{iE_1t/\hbar} = \frac{1}{\sqrt{2}}\left(|\phi_1\rangle + e^{-i(E_2-E_1)t/\hbar}\,|\phi_2\rangle\right). \tag{5}$$

Note that the global phase factor on the LHS is completely irrelevant, however the relative phase factor on the RHS is what matters. We can call this ω_{12}, and define it as

$$\omega_{12} = \frac{E_2 - E_1}{\hbar},$$

which we can use in Eq. 5 to get

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}}\left(|\phi_1\rangle + e^{-i\omega_{12}t}\,|\phi_2\rangle\right).$$

The corresponding function of real space is then

$$\psi(x,t) = \frac{1}{\sqrt{2}}\left(\phi_1(x) + e^{-i\omega_{12}t}\phi_2(x)\right),$$

which gives the probability to find the particle at point x at time t to be

$$|\psi(x,t)|^2 = \frac{1}{2}\phi_1^2(x) + \frac{1}{2}\phi_2^2(x) + \phi_1(x)\phi_2(x)\cos(\omega_{12}t).$$

5 Quantum Tunnelling

5.1 Particle in a Step-Like Potential

Consider a piecewise constant potential

$$V(x) = \begin{cases} 0 & x < 0 \\ V_0 & x \geq 0 \end{cases},$$

and a particle with total energy E and momentum $p = \hbar k$, moving from left to right ($x \to \infty$). We have our TISE as

$$\left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2} + (V(x) - E)\right]\phi(x) = 0$$
$$\left[\frac{\partial^2}{\partial x^2} + \frac{2m}{\hbar^2}(E - V(x))\right]\phi(x) = 0$$

In order to properly construct this problem, we have to consider each region separately. Starting with the first region $x < 0$ where $E > V$, we have the particle's total energy greater than the potential energy, meaning there is a kinetic component

$$E - V = \frac{1}{2}mv^2 = \frac{p^2}{2m} = \frac{\hbar^2k^2}{2m}.$$

Taking our Ansatz to be

$$\phi(x) = Ae^{ikx} + Be^{-ikx},$$

we know that the particle is going in the $+x$ direction, which means $B = 0$. Now considering the second region $x \geq 0$, this case is forbidden as the kinetic energy component cannot be negative, i.e. the particle would just bounce and reflect against the step in potential. Quantum mechanically, however, we can write

$$V - E = \frac{\hbar^2 \alpha^2}{2m}, \quad \alpha = \frac{\sqrt{2m(V_0 - E)}}{\hbar},$$

with α taking the place of $-ik$. This then gives our Ansatz to be

$$\phi(x) = Ce^{-\alpha x} + De^{\alpha x}.$$

The first term $Ce^{-\alpha x}$ describes a wavefunction that decays exponentially as the particle tries to go under the step, called an evanescent wave.

In order to construct the full wavefunction, we must have A, B, C, and D chosen such that $\phi(x)$ and $\frac{\partial \phi}{\partial x}$ are continuous (unless the step is infinitely high). The overall problem can then be solved by writing the appropriate solution in each constant potential region, then matching them at the boundary.

Let's consider, as an example, $E > V$ in both regions

$$\phi_1(x) = A_1 e^{ik_1 x} + A_1' e^{-ik_1 x}$$
$$\phi_2(x) = A_2 e^{ik_2 z} + A_2' e^{-ik_2 x}$$

with

$$k_1 = \frac{\sqrt{2mE}}{\hbar}, \quad k_2 = \frac{\sqrt{2m(E - V_0)}}{\hbar},$$

and $A_2' = 0$ since the particle comes from the left. We haven't done the same with A_1' as it may reflect at the step. Matching the conditions at $x = 0$ (the boundary), we get

$$A_1 + A_1' = A_2 \qquad \Rightarrow \phi(x) \text{ is continuous at the boundary}$$
$$ik_1 A_1 - ik_1 A_1' = ik_2 A_2 \qquad \Rightarrow \frac{\partial \phi}{\partial x} \text{ is continuous at the boundary}$$

From this, we can calculate the transmission coefficient T from the amplitude in region II travelling right over the amplitude in region I also travelling right. This gives

$$T \equiv \frac{k_2}{k_1} \left| \frac{A_2}{A_1} \right|^2 = \frac{4k_1 k_2}{(k_1 + k_2)^2}.$$

If we instead take the amplitude travelling left in region I, we get the reflection coefficient R

$$R \equiv \left| \frac{A_1'}{A_1} \right|^2 = 1 - T = 1 - \frac{1k_1 k_2}{(k_1 + k_2)^2}.$$

Note that the condition

$$T + R = 1$$

must always hold, as it enforces the particle existing somewhere in the system. In the classical case, you would expect $R = 0$ because the particle has enough energy to overcome the step, it would just slow down. Instead, it has a non-zero probability of still reflecting back.

5.2 Quantum Tunnelling

Now consider a potential system defined as

$$V(x) = \begin{cases} 0 & x < 0 \\ V_0 & 0 \leq x \leq l \\ 0 & x > l \end{cases}.$$

This gives the system of equations

$$\phi_1(x) = A_1 e^{ik_1 x} + A_1' e^{-ik_1 x}$$
$$\phi_2(x) = A_2 e^{-\alpha x} + A_2' e^{\alpha x}$$
$$\phi_3(x) = A_3 e^{ik_1 x} + A_3' e^{-ik_1 x},$$

with $A_3' = 0$. This can be solved by breaking the problem into two steps, matching the boundary conditions in region I and II at $x = 0$, then regions II and III at $x = l$. This gives

$$T = \frac{16E(V_0 - E)}{V_0^2} e^{-\alpha l}$$
$$= \frac{16E(V_0 - E)}{V_0^2} e^{-l\frac{\sqrt{2m(V_0 - E)}}{\hbar}}.$$


We know that the transmission coefficient $T \neq 0$ because the wavefunction under the barrier is an evanescent wave, so it has an exponentially decaying tail that leaks to the other side of the barrier. This coefficient is proportional to

$$e^{-\alpha l} = e^{-l\frac{\sqrt{2m(V_0 - E)}}{\hbar}},$$

i.e. the probability of tunnelling through the barrier depends exponentially on the width l and on the height of the barrier V_0, measuring from the energy of the incoming particle $V_0 - E$. Also note that the particle can be reflected back with probability $1 - T$, but if it tunnels through the barrier, it comes out with the same energy and momentum as when it arrived. This is called elastic tunnelling.

5.3 Double-Well

An ammonia molecule (NH_3) has an umbrella-like structure



Since the nitrogen is much heavier than the hydrogens, we can think of the nitrogen atom as being fixed at $x = 0$, i.e. the center of mass is effectively where the nitrogen atom is. We will be considering the movement of the three hydrogen atoms. They have two stable positions at $x = \pm b$.

If we want to flip the umbrella upside down, the hydrogen atoms have to overcome the electrostatic repulsion of the nitrogen atom. Thus, there is a potential barrier against flipping the umbrella, but its height is finite. For simplicity, we will approximate the potential as two rectangular wells, with a finite barrier in the middle and infinite walls outside.

5.3.1 Approximation 1: Infinite Barrier

If we approximate the wells as being separated by an infinite potential barrier in the middle, this becomes two independent infinite wells, and each eigenvalue is doubly degenerate. This means that are two orthogonal eigenvectors (one for each well) with the same eigenvalue.

For simplicity we can truncate the Hamiltonian, and assume that $k_B t \ll E_2 - E_1$, so only the 2 lowest energy states $|\phi_{1L}\rangle$, $|\phi_{1R}\rangle$ are populated, which means we have truncated the Hamiltonian to a 2-dimensional

space. We can define our basis as either completely in one well or completely in the other. The two basis states have the same energy E_1, so the Hamiltonian is given by

$$\mathcal{H} = \begin{pmatrix} E_1 & 0 \\ 0 & E_1 \end{pmatrix} = E_1 \mathbb{I}.$$

Since the potential energy is defined with respect to an arbitrary constant, we can set $E_1 = 0$, i.e. measure the energies starting from E_1 upwards. We can also define a different basis, this time with the first ket being positive in both wells, and the second being negative in the first well and positive in the second. We can call these the symmetric and anti-symmetric states, respectively.

$$|\phi_{1S}\rangle = \frac{1}{\sqrt{2}} \left(|\phi_{1R} + |\phi_{1L}\rangle \rangle \right), \quad |\phi_{1A}\rangle = \frac{1}{\sqrt{2}} \left(|\phi_{1R}\rangle - |\phi_{1L}\rangle \right).$$

The states are orthogonal and normalised.

5.3.2 Approximation 2: Finite Wells

Now considering the realistic case, where the barrier between the two wells is not infinite. We know that, in this case, the wavefunction does not go to 0 at the edge, but instead decays exponentially, so it has a finite probability to "leak" to the other side. This is the tunnelling effect. In our symmetric/anti-symmetric basis, both eigenstates now have some leakage under the barrier, but in different amounts, meaning $|\phi_{1S}\rangle$ and $|\phi_{1A}\rangle$ now have different energies. We define $E_A - E_S$ as tunnelling splitting.

5.3.3 Tunnelling Splitting

When the barrier was infinitely high, we had pairs of degenerate states (left/right). With the finite barrier, each pair splits in energy by an amount that depends on how each symmetric/anti-symmetric eigenstate leaks under the barrier, which is tunnelling splitting. This is a measure of the difference in total energy between the two eigenstates for each energy doublet, which increases with higher-energy doublets because they "see" a lower barrier.

5.3.4 Tunnelling Oscillations

We can prepare the system in the initial state

$$|\psi(0)\rangle = \frac{1}{\sqrt{2}} \left(|\phi_{1S}\rangle + |\phi_{1A}\rangle \right) = |\phi_{1R}\rangle .$$

Note that this is the umbrella on the right state, written in the basis of symmetric/anti-symmetric eigenstates of the Hamiltonian. At subsequent times, we'll have

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}} \left(|\phi_{1S}\rangle + e^{-i(E_{1A} - E_{1S})t/\hbar} |\phi_{1A}\rangle \right).$$

This gives the tunnelling oscillations to have the frequency

$$\omega_T = \frac{E_{1A} - E_{1S}}{\hbar}.$$

For ammonia, this is a well defined frequency

$$\omega_T = \frac{E_{1A - E_{1S}}}{\hbar} \approx 2\pi \times^2 4\text{GHz}.$$

5.4 Relation to Spin

In the presence of tunnel coupling, the Hamiltonian of the ammonia molecule (truncated to the lowest-energy doublet) becomes

$$\mathcal{H} = \begin{pmatrix} 0 & -\Delta \\ -\Delta & 0 \end{pmatrix},$$

where

$$2\Delta = E_{1A} - E_{1S}.$$

Tunnel coupling plays the same roles as a magnetic field parallel to x on a spin, which can be seen as the eigenstates $|\phi_{1S}\rangle$ and $|\phi_{1A}\rangle$ look exactly like $|X\rangle$ and $|-X\rangle$ for a spin. The tunnelling oscillations have the same form as the precession in the yz-plane of a spin placed in a magnetic field parallel to x.

Due to the distribution of charge between the atoms, the ammonia molecule as an electric dipole μ, so an electric field applied along the axis of the molecule pulls the umbrella on one side. Therefore, an electric field plays the same role as a magnetic field perpendicular to z, which splits the energy of the natural basis states. Overall, we can write the Hamiltonian as

$$\mathcal{H} = \begin{pmatrix} \epsilon & -\Delta \\ -\Delta & \epsilon \end{pmatrix} = -\Delta\sigma_x + \epsilon\sigma_z,$$

where

$$\begin{cases} 2\Delta = E_{1A} - E_{1S} & \epsilon = 0 \Rightarrow \text{ Tunnel coupling} \\ 2\epsilon = E_{1R} - E_{1L} & \Delta = 0 \Rightarrow \text{ Detuning by electric field} \end{cases}.$$

This means under suitable circumstances (low temperature), a complex system, like an ammonia molecule, is very well described by the same Hamiltonian as that of a spin in a magnetic field that has x and z components.

6 Position and Momentum Operators

6.1 Position Eigenbasis

From Postulate 2 (See Callout 3.1), every physical quantity is described by a Hermitian operator. To do this with position, you would discretise space, for example have a 51-dimensional vector to represent the probability of a particle being in one of 51 positions between $x = 0$ and $x = 1$. We can define a position eigenbasis with orthonormal vectors in this space as

$$|\phi_1\rangle = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad |\phi_2\rangle = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix}, \quad \cdots \quad |\phi_{51}\rangle = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}.$$

Each ket represents a particle that is maximally concentrated on the i-th point along the x-axis (in the limit where the discretization step is infinitesimal, it represents a δ- function at the position x_i).

From Postulate 3 (See Callout 3.3), when measuring the value of the quantity $\mathcal{A}$, we can only obtain as a result one of the eigenvalues a_n of the observable $\mathcal{A}$. For example, if we measure X while the state of the system is $|\phi_2\rangle$, we will get $x = 0.02$. So the X operator should look like

$$X = \begin{pmatrix} x_1 & 0 & \cdots & 0 \\ 0 & x_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & x_N \end{pmatrix} = \begin{pmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0.02 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & 1 \end{pmatrix}.$$

Unlike with spin and particle in a box, the operator that's diagonal in the chosen basis is not the Hamiltonian of the system.

6.2 Momentum Operator

We have the TISE, which is the eigenvalue problem for the Hamiltonian

$$\left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2} + V(x)\right]\phi(x) = E\phi(x).$$

Since the kinetic energy is also given by $\frac{p^2}{2m}$, we can interpret this as meaning that the momentum operator must be

$$P = -i\hbar\frac{\partial}{\partial x}.$$

The derivative is a linear operator, so it must be writable as a matrix. As before, we discretise the space, but now we have to calculate the derivative numerically. We can define Δx to be

$$\Delta x = x_{i+1} - x_i,$$

which gives

$$\frac{\partial\phi(x_i)}{\partial x} \approx \frac{1}{2}\left(\frac{\phi(x_{i+1}) - \phi(x_i)}{\Delta x} + \frac{\phi(x_i) - \phi(x_{i-1})}{\Delta x}\right) = \frac{\phi(x_{i+1}) - \phi(x_{i-1})}{2\Delta x}.$$

This is essentially taking the average of the slope immediately before and immediately after the point x_i. This then gives the matrix representation of the momentum operator to be

$$P = -\frac{i\hbar}{2\Delta x}\begin{pmatrix} 0 & 1 & \cdots & 0 \\ -1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & 1 \\ 0 & \cdots & -1 & 1 \end{pmatrix}.$$

With this, we can write explicitly the Hamiltonian of a particle as

$$\mathcal{H} = \frac{P^2}{2m} + V(x),$$

where P is given by the (non-diagonal) matrix above, and $V(x)$ is easy to write because it's just a diagonal matrix with numerical value of the potential at each point x_i. With this, we can calculate the other operators.

6.3 Commutation Relations

One important thing to note is that X and P do not commute. If we calculate brute-force the commutator using the matrices as above, we get

$$[X, P] = XP - PX = \frac{i\hbar}{2}\begin{pmatrix} 0 & 1 & \cdots & 0 \\ 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & 1 \\ 0 & \cdots & 1 & 1 \end{pmatrix} \approx i\hbar\mathbb{I}.$$

This represents, in matrix form, the Heisenberg uncertainty principle.

7 Dynamics and Control of a Quantum Bit

For systems under suitable conditions, i.e. minimal outside interference (this is usually low temperature), the system can be described by a Hamiltonian of the form

$$\mathcal{H} = \epsilon\sigma_z + \Delta\sigma_x.$$

This is known as a qubit Hamiltonian. This is a two-level quantum system, and can be used to encode quantum digital information

- Unperturbed case: $\mathcal{H}_0 = \epsilon\sigma_z$, with eigenstates $|\uparrow\rangle, |\downarrow\rangle$, and eigenvalues $\pm\epsilon$
- Coupled case: $\mathcal{H} = \epsilon\sigma_z + \Delta\sigma_x$, with eigenvalues $\lambda_{1,2} = E_{h,l} = \pm\sqrt{\epsilon^2 + \Delta^2}$.

Proof. We can find the eigenstates of a coupled Hamiltonian as

$$\begin{aligned} \det(\mathcal{H} - \lambda\mathbb{I}) &= 0 \\ \Rightarrow (\epsilon - \lambda)(-\epsilon - \lambda) - \Delta^2 &= \lambda^2 - \epsilon^2 - \Delta^2 = 0 \\ \Rightarrow \lambda_{1,2} = E_{h,l} &= \pm\sqrt{\epsilon^2 + \Delta^2}. \end{aligned}$$

□

Consider a qubit Hamiltonian where Δ is constant and $\epsilon = -\infty \to +\infty$. The eigenvectors will be

$$|\phi_h\rangle = \sin\frac{\theta}{2}|\downarrow\rangle + \cos\frac{\theta}{2}|\uparrow\rangle, \quad |\phi_l\rangle = \cos\frac{\theta}{2}|\downarrow\rangle - \sin\frac{\theta}{2}|\uparrow\rangle,$$

where $\tan\theta = \frac{\Delta}{\epsilon}$ for $0 \le \theta \le \pi$. This is known as energy anticrossing.

We can map this onto a spin that is being subjected to a time-variable magnetic field along z, and a constant magnetic field along x, so that the total field makes an angle θ with the z axis. The energy anticrossing occurs because, since $\Delta \neq 0$, the eigenvalues never degenerate.

7.1 Time Evolution

From above, we know that we can write the time evolution operator as $U(t, t_0) = e^{-i\frac{\mathcal{H}}{\hbar}(t-t_0)}$ and that it is diagonal if we use the eigenvectors of the Hamiltonian as basis of the Hilbert space, i.e.

$$U(t, t_0) = \begin{pmatrix} e^{-i\frac{E_1}{\hbar}(t-t_0)} & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & e^{-i\frac{E_N}{\hbar}(t-t_0)} \end{pmatrix}.$$

To calculate the time evolution analytically, we must therefore

1. Express the initial state $|\psi(0)\rangle$ in the eigenbasis of the Hamiltonian

$$|\psi(0)\rangle = \sum_n c_n(0)|\phi_n\rangle = \begin{pmatrix} c_1(0) \\ \vdots \\ c_N(0) \end{pmatrix}.$$

2. Apply the time evolution operator and obtain the time-evolved state as

$$|\psi(t)\rangle = \sum_n c_n(0)e^{-i\frac{E_n}{\hbar}t}|\phi_n\rangle = \begin{pmatrix} e^{-i\frac{E_1}{\hbar}t}c_1(0) \\ \vdots \\ e^{-i\frac{E_N}{\hbar}t}c_N(0) \end{pmatrix}$$

Example 7.1. Consider a qubit initially described by the Hamiltonian $\mathcal{H}_0 = \epsilon\sigma_z$ and prepared in its eigenstate $|\uparrow\rangle = |\psi(0^-)\rangle$. At time $t = 0$, switch on the term $\Delta\sigma_x$. Now, $|\psi(0^-)\rangle$ is no longer an eigenstate.

Answer:

1. Express $|\psi(0^-)\rangle = |\uparrow\rangle$ in the new eigenbasis $|\phi_h\rangle, |\phi_l\rangle$:

$$\begin{aligned}\left|\psi(0^-)\right\rangle &= \cos\frac{\theta}{2}|\phi_h\rangle - \sin\frac{\theta}{2}|\phi_l\rangle \\ &= \begin{pmatrix}\cos\frac{\theta}{2} \\ -\sin\frac{\theta}{2}\end{pmatrix}.\end{aligned}$$

2. Time-evolve the coefficients:

$$\begin{aligned}|\psi(t)\rangle &= e^{-i\frac{E_h}{\hbar}t}\cos\frac{\theta}{2}|\phi_h\rangle - e^{-i\frac{E_l}{\hbar}t}\sin\frac{\theta}{2}|\phi_l\rangle \\ &= \begin{pmatrix}e^{-i\frac{E_h}{\hbar}t}\cos\frac{\theta}{2} \\ -e^{-i\frac{E_l}{\hbar}t}\sin\frac{\theta}{2}\end{pmatrix}.\end{aligned}$$

We started in the state $|\uparrow\rangle$. The probability to find the spin in $|\downarrow\rangle$ at time t is $|\langle\downarrow|\psi(t)\rangle|^2$. Express $|\downarrow\rangle$ in the new eigenbasis $|\phi_h\rangle, |\phi_l\rangle$.

$$\begin{aligned}|\downarrow\rangle &= \sin\frac{\theta}{2}|\phi_h\rangle + \cos\frac{\theta}{2}|\phi_l\rangle \\ &= \begin{pmatrix}\sin\frac{\theta}{2} \\ \cos\frac{\theta}{2}\end{pmatrix} \\ \langle\downarrow|\psi(t)\rangle &= \begin{pmatrix}\sin\frac{\theta}{2} & \cos\frac{\theta}{2}\end{pmatrix}\begin{pmatrix}e^{-i\frac{E_h}{\hbar}t}\cos\frac{\theta}{2} \\ -e^{-i\frac{E_l}{\hbar}t}\sin\frac{\theta}{2}\end{pmatrix} \\ &= \sin\frac{\theta}{2}\cos\frac{\theta}{2}e^{-i\frac{E_h}{\hbar}} - \sin\frac{\theta}{2}\cos\frac{\theta}{2}e^{-i\frac{E_l}{\hbar}t} \\ &= \frac{1}{2}\sin\theta\left[e^{-i\frac{E_h}{\hbar}t} - e^{-i\frac{E_l}{\hbar}t}\right].\end{aligned}$$

Note $\sin\theta = \frac{\Delta}{\sqrt{\epsilon^2+\Delta^2}}$.

$$E_h = -E_l = \sqrt{\epsilon^2+\Delta^2} \Rightarrow e^{-i\frac{E_h}{\hbar}t} - e^{-i\frac{E_l}{\hbar}t} = 2i\sin\left(\frac{\sqrt{\epsilon^2+\delta^2}}{\hbar}t\right).$$

$$\boxed{|\langle\downarrow|\psi(t)\rangle|^2 = i\frac{\Delta^2}{\sqrt{\epsilon^2+\Delta^2}}\sin\left(\frac{\sqrt{\epsilon^2+\Delta^2}}{\hbar}t\right).}$$

7.2 Rabi Formula and Oscillation

In Example 7.1, we found the probability of finding a particle in a certain state after applying an external field. Simplifying a bit further, we found

$$\mathcal{P}_\downarrow = |\langle\downarrow|\psi(t)\rangle|^2 = \frac{\Delta^2}{\epsilon^2+\Delta^2}\sin^2\left(\frac{\sqrt{\epsilon^2+\Delta^2}}{\hbar}t\right).$$

This is known as Rabi Formula. The probability oscillates at frequency

$$\omega_R = 2\frac{\sqrt{\epsilon^2+\Delta^2}}{\hbar},$$

which is equal to the energy difference $E_h - E_l$ divided by $\hbar$. This oscillation is known as Rabi Oscillation.

It can be interpreted as a spin starting along z, then precesses around the new total field when B_x is switched on. This also means that we can never get $\mathcal{P}_\downarrow = 1$, as this would require $B_x \to \infty$.

7.3 Time-Dependent Hamiltonian

In certain systems, the Hamiltonian can be explicitly time-dependent (e.g. a spin in a time-varying magnetic field). The time-evolution operator is given by

$$U(t,t_0) = \int_{t_0}^{t} e^{-i\frac{\mathcal{H}(t')}{\hbar}} dt',$$

but the integral is impossible to analytically solve. Instead, we discretise the time in small intervals δt, during which the Hamiltonian is approximated as a constant. Calling $t_n = t_0 + n\delta t$, we get

$$\begin{aligned} U(t_{n+1},t_n) &= e^{-i\frac{\mathcal{H}(t_n)}{\hbar}\Delta t} \\ |\psi(t_{n+1})\rangle &= U(t_{n+1},t_n)\,|\psi(t_n)\rangle \\ |\psi(t_N)\rangle &= \left(\prod_{n=0}^{N-1} U(t_{n+1},t_n)\right)|\psi(t_0)\rangle\,. \end{aligned}$$

This is easy to program, however with small time steps it is very computationally expensive.

Consider a Hamiltonian with a periodic time dependence

$$\mathcal{H}(t) = \mathcal{H}_0 + \mathcal{H}_1\cos(\omega t),$$

with period $T = \frac{2\pi}{\omega}$. Since $\mathcal{H}(t+T) = \mathcal{H}(t)$, we just need to calculate the time evolution operator that takes us from the start to the end of the period, splitting the period T into N small intervals $T = N\Delta t$

$$U(t+T,t) = \prod_{n=0}^{N-1} U(t_{n+1},t_n).$$

$U(t+T,t)$ is called the Floquet operator. We can use this to calculate the state after each period as

$$|\psi(t+kT)\rangle = U^k(t+T,t)\,|\psi(t)\rangle\,.$$

This means we can calculate $U(t+T,t)$ once, then keep multiplying by the same time evolution operator.

7.4 Rotating Frame

As with classical mechanics, a time-dependent Hamiltonian can be transformed into a time-independent one (but not always) by performing a change of basis.

Consider a magnetic dipole $\vec{M}$ placed in a static vertical magnetic field $\vec{B}_0 = B_0\hat{z}$, plus a rotating magnetic field in the xy-plane, $\vec{B}_1 = B_1(\cos\omega t\hat{x} + \sin\omega t\hat{y})$, giving a Hamiltonian of

$$\mathcal{H}(t) = -\vec{M}\cdot\vec{B}(t) = -M_zB_z - M_xB_1\cos\omega t - M_yB_1\sin\omega t.$$

The trick is to define a rotating reference frame $OXYZ$ that rotates at frequency ω around the z-axis. The vector $\vec{B}_1$ appears static in this rotating frame. This is similar to frame of reference in SR.

We can define

- $\omega_0 = \gamma B_0$: Larmor precession frequency of the spin in the static field $\vec{B}_0$.
- $\omega_1 = \gamma B_1$: Larmor frequency the spin would have if it were placed in a static field $\vec{B}_1$.
- $\Delta\omega = \omega - \omega_0$: The "detuning" between precession frequency and rotating field.

This gives our Hamiltonian to be

$$\begin{aligned} \mathcal{H}(t) &= \gamma B_0S_z + \gamma B_1(S_x\cos\omega t - S_y\sin\omega t) \\ &= \omega_0S_z + \omega_1\left(\frac{e^{i\omega t}+e^{-i\omega t}}{2}S_x + \frac{e^{i\omega t}-e^{-i\omega t}}{2i}S_y\right) \\ &= \frac{\hbar}{2}\begin{pmatrix}\omega_0 & \omega_1e^{-i\omega t}\\ \omega_1e^{i\omega t} & -\omega_0\end{pmatrix}. \end{aligned}$$

The generic state at time t in the σ_z basis ($|\uparrow\rangle, |\downarrow\rangle$),

$$|\psi(t)\rangle = c_\uparrow(t)\,|\uparrow\rangle + c_\downarrow(t)\,|\downarrow\rangle = \begin{pmatrix} c_\uparrow(t) \\ c_\downarrow(t) \end{pmatrix}.$$

Our time evolution is then given by

$$\begin{aligned} i\hbar\frac{d\,|\psi(t)\rangle}{dt} &= \mathcal{H}(t)\,|\psi(t)\rangle \\ \Rightarrow i\hbar\frac{d}{dt}\begin{pmatrix} c_\uparrow(t) \\ c_\downarrow(t) \end{pmatrix} &= \frac{\hbar}{2}\begin{pmatrix} \omega_0 & \omega_1 e^{-i\omega t} \\ \omega_1 e^{i\omega t} & -\omega_0 \end{pmatrix}\begin{pmatrix} c_\uparrow(t) \\ c_\downarrow(t) \end{pmatrix} \\ &\begin{cases} i\frac{d}{dt}c_\uparrow(t) = \frac{\omega_0}{2}c_\uparrow(t) + \frac{\omega_1}{2}e^{-i\omega t}c_\downarrow(t) \\ i\frac{d}{dt}c_\downarrow(t) = \frac{\omega_1}{2}e^{i\omega t}c_\uparrow(t) - \frac{\omega_0}{2}c_\downarrow(t) \end{cases} \end{aligned}$$

7.5 Change of Basis

A rotation by an angle ϕ around an axis $\vec{u} = u_x\hat{x} + u_y\hat{y} + u_z\hat{z}$, with $|\vec{u}|^2 = 1$ is expressed in the 2D Hilbert space by the operator

$$R_{\vec{u},\phi} = e^{i\frac{\phi}{2}\sigma_{\vec{u}}},$$

where $\sigma_{\vec{u}} = \vec{u}\cdot\vec{\sigma}$ is the Pauli matrix along the unit vector $\vec{u}$. Note that $R_{\vec{u},\phi}$ is unitary. We change our basis by rotating by ωt around $\hat{z}$, so we need

$$R_{\hat{z},\omega t} = e^{i\frac{\omega t}{2}\sigma_z} = \begin{pmatrix} e^{i\omega t/2} & 0 \\ 0 & e^{-i\omega t/2} \end{pmatrix}.$$

We can define $\left|\tilde{\psi}(t)\right\rangle = R_{\vec{z},\omega t}\,|\psi(t)\rangle = \begin{pmatrix} \tilde{c}_\uparrow(t) \\ \tilde{c}_\downarrow(t) \end{pmatrix}$, which gives

$$\Rightarrow |\psi(t)\rangle = R^{-1}_{\vec{z},\omega t}\left|\tilde{\psi}(t)\right\rangle = \begin{pmatrix} e^{-i\omega t/2} & 0 \\ 0 & e^{i\omega t/2} \end{pmatrix}\begin{pmatrix} \tilde{c}_\uparrow(t) \\ \tilde{c}_\downarrow(t) \end{pmatrix}.$$

Therefore,

$$\begin{aligned} c_\uparrow(t) = e^{-i\omega t/2}\tilde{c}_\uparrow(t) \Rightarrow \frac{d}{dt}c_\uparrow(t) &= -\frac{i\omega}{2}\tilde{c}(t)e^{-i\frac{\omega t}{2}} + e^{-i\frac{\omega t}{2}}\tilde{c}_\uparrow(t) \\ c_\downarrow(t) = e^{i\omega t/2}\tilde{c}_\downarrow(t) \Rightarrow \frac{d}{dt}c_\downarrow(t) &= \frac{i\omega}{2}\tilde{c}_\downarrow(t)e^{i\frac{\omega t}{2}} + e^{i\frac{\omega t}{2}}\frac{d}{dt}\tilde{c}_\downarrow(t). \end{aligned}$$

We can substitute these expressions for $c_{\uparrow\downarrow}(t)$ and $\frac{dc_{\uparrow\downarrow}(t)}{dt}$ into the Schrödinger equation to get

$$\begin{aligned} i\frac{d}{dt}\tilde{c}_\uparrow(t) &= -\frac{\Delta\omega}{2}\tilde{c}_\uparrow(t) + \frac{\omega_1}{2}\tilde{c}_\downarrow(t) \\ i\frac{d}{dt}\tilde{c}_\downarrow(t) &= \frac{\omega_1}{2}\tilde{c}_\uparrow(t) + \frac{\Delta\omega}{2}\tilde{c}_\downarrow(t). \end{aligned}$$

This is the same as writing

$$i\hbar\frac{d\left|\tilde{\psi}(t)\right\rangle}{dt} = \tilde{\mathcal{H}}\left|\tilde{\psi}(t)\right\rangle,$$

where

$$\tilde{\mathcal{H}} = \frac{\hbar}{2}\begin{pmatrix} -\Delta\omega & \omega_1 \\ \omega_1 & \Delta\omega \end{pmatrix} = -\frac{\hbar}{2}\Delta\omega\sigma_z + \frac{\hbar}{2}\omega_1\sigma_x.$$

In this frame, the Hamiltonian appears time-independent, and all the time-dependence has been moved to the reference frame.

7.6 Magnetic Resonance

The Hamiltonian in the rotating frame looks very similar to a qubit Hamiltonian with

$$\epsilon = -\frac{\hbar}{2}\Delta\omega, \quad \Delta = \frac{\hbar}{2}\omega_1.$$

Assuming the initial state $|\psi(0)\rangle \equiv \left|\tilde{\psi}(t)\right\rangle = |\uparrow\rangle$. The probability to flip the state to $|\downarrow\rangle$ is obtained from the Rabi formula as

$$\mathcal{P}_\downarrow = |\langle\downarrow|\psi(t)\rangle|^2 = \frac{\omega_1^2}{(\Delta\omega)^2 + \omega_1^2}\sin^2\left(\frac{\sqrt{(\Delta\omega)^2+\omega_1^2}}{2}t\right).$$

In the case $\Delta\omega = 0$, i.e. B_1 rotates at the same frequency as the Larmor precession of the spin in the field B_0, then $\mathcal{P}_\downarrow$ can reach 1, which is called magnetic resonance. In the case $\Delta\omega \gg \omega_1, \mathcal{P}_\downarrow \approx 0$, which is called off-resonance, where the spin state is left untouched. At resonances, the Rabi frequency is equal to ω_1, which means the B_0 field is cancelled out in the rotating frame, so the spin precesses at frequency ω_1 around the field B_1.

7.7 Density Operator

A well-defined quantum state, such as the output of a Stern-Gerlach experimentr, is known as pure state as we know with certainty which state it is in. However, in the case that we haven't prepare the input of a Stern-Gerlach machine as a pure state, the output is completely random. Since we can't assign the input state a well-defined ket, the best we can do is describe it as some mixture of $|\uparrow\rangle$ and $|\downarrow\rangle$ with probabilities $\mathcal{P}_\uparrow$ and $\mathcal{P}_\downarrow$, which is known as a mixed state.

For pure states, we can define a density operator, or density matrix as

$$\rho = |\psi\rangle\langle\psi|.$$

Expanding, we have ρ to be

$$\begin{aligned}
|\psi\rangle &= \begin{pmatrix} c_1 \\ \vdots \\ c_N \end{pmatrix} \\
\Rightarrow \rho &= \begin{pmatrix} c_1 \\ \vdots \\ c_N \end{pmatrix}\begin{pmatrix} c_1^* & \cdots & c_N^* \end{pmatrix} \\
&= \begin{pmatrix} c_1c_1^* & \cdots & c_1c_N^* \\ \vdots & \ddots & \vdots \\ c_Nc_1^* & \cdots & c_Nc_N^* \end{pmatrix},
\end{aligned}$$

or we can call $|u_n\rangle$ the basis vectors of the Hilbert space, we have

$$\rho_{pn} = \langle u_p|\rho|u_n\rangle = c_pc_n^*.$$

The density matrix is Hermitian, and a projector onto the normalised state $|\psi\rangle$ and therefore $\rho = \rho^2$, because

$$\rho^2 = |\psi\rangle\langle\psi|\psi\rangle\langle\psi| = |\psi\rangle\langle\psi| = \rho.$$

See Problem 3.1 for a proof.

7.7.1 Properties of the Density Operator

- Since $|\psi\rangle$ is normalised, then $\mathrm{Tr}(\rho) = 1$
- Since $\rho = \rho^2$, $\mathrm{Tr}(\rho^2) = 1$

- The expectation value $\langle A\rangle_\psi$ is given by $\langle A\rangle_\psi = \mathrm{Tr}(\rho A)$, since

$$\begin{aligned}\langle A\rangle_\psi &= \langle\psi| A |\psi\rangle \\ &= \sum_{n,p} c_n^* c_p A_{np} \\ &= \sum_{n,p} \langle u_p| \rho |u_n\rangle \langle u_n| A |u_p\rangle \\ &= \sum_p \langle u_p| \rho A |u_p\rangle .\end{aligned}$$

- The time evolution of the density matrix

$$\begin{aligned}\frac{d}{dt}\rho(t) &= \left[\frac{d}{dt}|\psi(t)\rangle\right]\langle\psi(t)| + |\psi(t)\rangle\left[\frac{d}{dt}\langle\psi(t)|\right] \\ &= \frac{1}{i\hbar}\mathcal{H}(t)|\psi(t)\rangle\langle\psi(t)| - \frac{1}{i\hbar}|\psi(t)\rangle\langle\psi(t)|\mathcal{H}(t) \\ &= \frac{1}{i\hbar}[\mathcal{H}(t),\rho(t)] .\end{aligned}$$

- For pure states, the density operator contains the same information as the state vector.

7.7.2 Density Operator for Mixed States

Consider a mixed state. The information we know about the system is that if we measure it in the z-basis, we get $\mathcal{P}_\uparrow = \frac{1}{2} = \mathcal{P}_\downarrow$. We can still write $\langle S_z\rangle = \frac{\hbar}{2}\mathcal{P}_\uparrow - \frac{\hbar}{2}\mathcal{P}_\downarrow = 0$, which would be satisfied if we define

$$\rho = \begin{pmatrix}\mathcal{P}_\uparrow & 0\\ 0 & \mathcal{P}_\downarrow\end{pmatrix},$$

since, using this, we are able to retain the property

$$\langle S_z\rangle = \mathrm{Tr}(\rho S_z).$$

In general, we define the density operator for mixed states as

$$\phi = \begin{pmatrix}\mathcal{P}_1 & \cdots & 0\\ \vdots & \ddots & \vdots\\ 0 & \cdots & \mathcal{P}_N\end{pmatrix},$$

where $\mathcal{P}_n$ is the probability of obtaining the eigenvalue a_n when measuring A. This does not require knowing the precise quantum state at the start of the experiment. Note that it now no longer holds true that $\rho = \rho^2$.

7.7.3 Populations and Coherences

The diagonal elements of ρ are called populations. They describe the probability $\mathcal{P}_n$ of finding the system in the state $|u_n\rangle$ when performing a measurement. This holds for both pure and mixed states.

The off-diagonal elements are called coherences. They describe the phase relationship between different components of a quantum superposition of states.

Example 7.1. We have the two states in the σ_z basis given by

$$|X\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle) = \frac{1}{\sqrt{2}}\begin{pmatrix}1\\1\end{pmatrix}, \quad |Y\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle + i|\downarrow\rangle).$$

These correspond to the density matrices

$$|X\rangle \Rightarrow \rho_X = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$
$$|Y\rangle \Rightarrow \rho_Y \frac{1}{2}\begin{pmatrix} 1 & -i \\ i & 1 \end{pmatrix}.$$

The populations are the same, however the coherences are different.

Consider an NMR (nuclear magnetic resonance) experiment, where a large ensemble of identical spins are initially all prepared in the pure state $|X\rangle$. We can describe this ensemble with

$$\rho_X = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

After preparation, each spin precesses at the Larmor frequency $\omega_0 = \gamma B_0$. In reality, each spin sees a slightly different and unknown $B_0(\vec{r})$ field at its location, so after some time we lose track of what exactly is the phase of each spin, and they end up pointing randomly in the xy-plane. We can describe this new ensemble with

$$\rho = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

since we know that each spin is somewhere in the xy-plane, and the coherences have decayed to zero. This phenomenon by which randomness and lack of knowledge of the precise details of the environment of quantum systems results in a loss of knowledge of which pure states the system is in is called decoherence, however it can still be described by a density matrix.

7.7.4 State Purity

Consider the following density matrix

$$\rho_\psi = \frac{1}{2}\begin{pmatrix} 1 & 0.5 \\ 0.5 & 1 \end{pmatrix} = \begin{pmatrix} 0.5 & 0.25 \\ 0.25 & 0.5 \end{pmatrix},$$

from which we get

$$\rho_\psi^2 = \begin{pmatrix} 0.3125 & 0.25 \\ 0.25 & 0.3125 \end{pmatrix}, \quad \mathrm{Tr}\left(\rho_\psi^2\right) = 0.625 < 1,$$

which shows that ρ_ψ does not describe a pure state. However, it still has nonzero coherences. It means you know something about the phase, however there is some uncertainty. The fully mixed state $\rho = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ would have

$$\mathrm{Tr}(\rho^2) = 0.5,$$

which is less than $\mathrm{Tr}\left(\rho_\psi^2\right)$. $\mathrm{Tr}(\rho^2)$ is called state purity.

A Fourier Transform

A Fourier transform decomposes a function into another function that describes the number of occurrences of frequencies within the original function.

$$F(\omega) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt.$$

Fundamentally it is an inner product between $f(t)$ and $e^{-i\omega t}$, with the second vector being characterised by the parameter ω. It tells you how similar the two vectors are to each other, thereby defining the number of occurrences of each frequency.

The Fourier series of a sinusoid is a delta function, indicating that it is infinitely similar at that frequency, and not at all at the others.

B Matlab/Python Commands

B.1 Matrix Operations

B.1.1 Exponential

Command: The exponential of a matrix can be calculated using

```
expm(A)
```

Python has a similar command

```
scipy.linalg.expm(A)
```

Example: Given an initial state `psi_0` at time $t = 0$, the quantum state `psi_t` at a later time t can be found with

(MATLAB):

```
psi_t = expm(-1i*H*t/hbar) * psi_0
```

(Python):

```
psi_t=scipy.linalg.expm(-1j * H * t / hbar) @ psi_0
```