Banner appropriate to article type will appear here in typeset article

Capillary effects on the self-similar growth of immiscible Rayleigh–Taylor mixing

Michael D. Walker[1]†, and Stefan S. Nixon[2]

[1]Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, NJ 08544, USA

[2]Department of Aeronautics, Imperial College London, London SW7 2AZ, UK

(Received xx; revised xx; accepted xx)

In immiscible Rayleigh–Taylor turbulence, the growth parameter α increases as the initial Weber number We_0 decreases, approximately as $\mathrm{We}_0^{-1/2}$ in recent simulations. This paper derives the form that such a dependence can take in a self-similar mixing layer. Complete self-similarity leaves no dimensionless group containing the surface tension, so We_0 can enter only through corrections in the ratio of the capillary length ℓ_σ to the layer width h. When the self-similar moment analysis of Ristorcelli & Clark (*J. Fluid Mech.*, vol. 507, 2004, p. 213) is extended to immiscible fluids, the direct capillary term in the mid-plane scalar-flux budget is the inverse instantaneous Weber number, $(\ell_\sigma/h)^2/2\alpha$; it reduces α in proportion to We_0^{-1}. With fixed profile constants, the same budget relaxes from any departure approximately as h^{-40}, so an increase that decays slowly with h requires an indirect effect of surface tension on the flow structure. A correction linear in ℓ_σ/h gives $\alpha - \alpha_\infty \propto \mathrm{We}_0^{-1/2}$ at fixed h/λ_0, and a spectral closure in which surface tension sets the inner cut-off of a smooth interface produces this form. High-resolution simulations compared at equal mixing width support the form $\alpha - \alpha_{\mathrm{We}_0=\infty} \approx 8.9\,\ell_\sigma/h$, whereas exponents extracted at fixed time drift with the time of comparison.

Key words: buoyancy-driven instability, turbulent mixing, multiphase flow

1. Introduction

The late-time growth of a Rayleigh–Taylor (RT) mixing layer is usually written in the form of the constant-acceleration ansatz $h \approx \alpha \mathrm{A_t} g t^2$, where h is the layer width, g the acceleration, $\mathrm{A_t}$ the Atwood number and α a growth parameter that depends on the initial conditions (Youngs 1984; Dimonte *et al.* 2004). Ristorcelli & Clark (2004) (hereafter RC04) showed that self-similarity of the second-moment equations gives $\dot{h}^2 = \mathrm{A_t} g C_\mathrm{o} h$, so that $\alpha = C_\mathrm{o}/4$ is a functional of the unclosed self-similar profiles rather than a universal constant. Immiscible fluids grow faster. Experiments, many of them with immiscible fluids, give larger α than simulations of miscible mixing (Dimonte & Schneider 2000; Dimonte *et al.* 2004), and at $\mathrm{A_t} \approx 0.5$ miscibility reduced α by about a factor of two in direct comparisons (Roberts &

† Email address for correspondence: michael.walker@princeton.edu

Abstract must not spill onto p.2




Figure 1: Growth parameter $\alpha(t) = \dot{h}^2/4\mathrm{A_t}gh$ in the immiscible simulations of Walker *et al.* (2025, 2026, 2027*a*) at $\mathrm{We_0} = 50$, 100, 200 and ∞, with 1120^3 points, $\mathrm{A_t} = 0.5$ and $\mathrm{Re_0} = \infty$: (*a*) against t/τ_0, with a miscible simulation for reference; (*b*) against h/λ_0 after the peak of α, on logarithmic axes.

Jacobs 2016). Cook *et al.* (2004) (hereafter CCM04) argued from the mid-plane mass flux, where density fluctuations take their largest possible value, that immiscible fluids should grow faster, and recent incompressible, low-order volume-of-fluid simulations find α increasing with surface tension (Zhao & Li 2026; Zhao *et al.* 2026).

Compressible, high-order diffuse-interface simulations at $\mathrm{A_t} = 0.5$ (Brill *et al.* 2025; Walker *et al.* 2025, 2026, 2027*a*) give $\alpha \approx 0.039$ for immiscible fluids without surface tension and $\alpha \approx 0.028$ for miscible fluids, and a further increase as the initial Weber number $\mathrm{We_0}$ decreases (figure 1*a*). The trend was reported as approximately $\alpha \propto \mathrm{We_0^{-1/2}}$ on dimensional grounds (Walker *et al.* 2026). Linear theory predicts the opposite tendency, since surface tension stabilises short wavelengths (Chandrasekhar 1981; Sohn 2009). This paper derives the form that a $\mathrm{We_0}$ dependence can take in a self-similar layer, by extending the RC04 analysis to immiscible fluids with surface tension, and compares it with the simulations at equal mixing width.

2. Constraints from dimensional analysis

Two fluids of densities $\rho_h > \rho_\ell$, separated by an interface with surface tension coefficient σ, are accelerated by gravity g in the $-z$ direction. With $\bar\rho = (\rho_h + \rho_\ell)/2$, $\Delta\rho = \rho_h - \rho_\ell$ and $\mathrm{A_t} = \Delta\rho/2\bar\rho$, an initial perturbation of shortest wavelength λ_0 defines $\tau_0 = (\lambda_0/\mathrm{A_t}g)^{1/2}$ and

$$\mathrm{We_0} = \frac{\bar\rho\, g\, \lambda_0^2}{\sigma}\,, \qquad \ell_\sigma = \left(\frac{\sigma}{\Delta\rho\, g}\right)^{1/2}, \qquad \frac{\ell_\sigma}{\lambda_0} = (2\mathrm{A_t We_0})^{-1/2}\,, \tag{2.1}$$

where ℓ_σ is the cut-off length of linear theory, $\gamma^2 = \mathrm{A_t}gk\,(1 - k^2\ell_\sigma^2)$ (Chandrasekhar 1981; Sohn 2009). The width is the full width $h = 6\int \bar V(1-\bar V)\,\mathrm{d}z$, which equals the total width of a linear profile, with $\bar V$ the plane-averaged volume fraction of heavy fluid, and the growth parameter is measured instantaneously as $\alpha(t) = \dot h^2/4\mathrm{A_t}gh$, which is constant for the RC04 solution with its virtual origin. Dimensional analysis gives $\alpha = \Psi(t/\tau_0; \mathrm{A_t}, \mathrm{We_0}, \mathrm{Re_0}, \dots)$ and admits any dependence on $\mathrm{We_0}$. The quantities $\sigma/\bar\rho$ and $\mathrm{A_t}g$ define the length ℓ_σ but no dimensionless group, so a second length is needed. Under complete self-similarity, the only one available is h, the initial wavelength is forgotten, and the relevant group is the

instantaneous Weber number

$$\mathrm{We}(t) \equiv \frac{\bar{\rho}\,\dot{h}^2 h}{\sigma} = 4\mathrm{A_t}\,\alpha\,\mathrm{We}_0\Big(\frac{h}{\lambda_0}\Big)^2, \qquad \Big(\frac{\ell_\sigma}{h}\Big)^2 = \frac{2\alpha}{\mathrm{We}(t)}\,, \tag{2.2}$$

which grows as t^4. If a self-similar state exists with h as its only outer length, α tends to a limit α_∞ independent of σ, and a We_0 dependence survives only as a correction in ℓ_σ/h that decays as the layer grows, an incomplete similarity in the sense of Barenblatt (1996). A persistent power law in We_0 is excluded. One such correction follows from a velocity scale set by capillarity and buoyancy alone, $U_\sigma \propto (\mathrm{A_t} g \ell_\sigma)^{1/2}$. Adding U_σ^2 to the self-similar $\dot{h}^2 = 4\alpha_\infty \mathrm{A_t} g h$ gives

$$\alpha(t) = \alpha_\infty + K\,\frac{\ell_\sigma}{h} = \alpha_\infty + K\,(2\mathrm{A_t We}_0)^{-1/2}\,\frac{\lambda_0}{h}\,, \tag{2.3}$$

with K a constant. This is the form in which a $\mathrm{We}_0^{-1/2}$ dependence is consistent with dimensional analysis: at fixed h/λ_0, as a correction that decays as h^{-1}. A power law $\alpha \propto \mathrm{We}_0^{-1/2}$ would give $\alpha \to 0$ as $\sigma \to 0$, contrary to $\alpha \approx 0.039$ at $\mathrm{We}_0 = \infty$.

3. Mid-plane scalar-flux budget

Following RC04, this approach begins with the Boussinesq form. Let $c = +1$ in the heavy fluid and $c = -1$ in the light fluid, so that $\rho = \bar{\rho}(1 + \mathrm{A_t} c)$. Let $\mathbf{u} = (u, v, w)$ be the velocity, μ the dynamic viscosity, p the kinematic pressure (pressure divided by $\bar{\rho}$), and $\mathbf{f} = (\sigma/\bar{\rho})\,\kappa\,\hat{\mathbf{n}}\,\delta_s$ the capillary force per unit mass (Brackbill *et al.* 1992), with $\hat{\mathbf{n}} = \nabla c/|\nabla c|$, curvature $\kappa = -\nabla\cdot\hat{\mathbf{n}}$ and δ_s the surface delta function. Angle brackets denote averages over horizontal planes, $c = C + c'$ with $C = \langle c\rangle$, and the subscript '0' denotes the mid-plane $z = 0$. Because $c^2 = 1$ everywhere,

$$\langle c'c'\rangle = 1 - C^2\,, \tag{3.1}$$

so the molecular mixing parameter of RC04 vanishes, the scalar-variance equation is satisfied, and $\langle c'c'\rangle_0 = 1$; for miscible fluids, $\langle c'c'\rangle_0 = 1 - \theta_0$, with θ_0 the mixedness. Integrating the mean indicator equation from $-\infty$ to the mid-plane with the self-similar profile $C = f_C(z/h)$ gives $\langle wc'\rangle_0 = -2\dot{h}/R_f$ (§ S1 of the supplementary material), where $R_f = \Delta\rho\,\dot{h}/|\langle\rho' w'\rangle_0|$ depends, for a self-similar layer, only on the shape of $\bar{V}$; it is 8.0–8.7 for linear, error-function and hyperbolic-tangent profiles. The scalar-flux equation (RC04, their equation 2.8) with surface tension is

$$\frac{\partial}{\partial t}\langle wc'\rangle + \frac{\partial}{\partial z}\langle wwc'\rangle = -\langle ww\rangle\frac{\partial C}{\partial z} - \mathrm{A_t} g\langle c'c'\rangle - \Big\langle c'\frac{\partial p}{\partial z}\Big\rangle + \langle c' f_z\rangle + \frac{\mu}{\bar{\rho}}\langle c'\nabla^2 w\rangle\,. \tag{3.2}$$

Taking the divergence of the momentum equation gives a Poisson equation for the pressure fluctuation p',

$$\nabla^2 p' = -\mathrm{A_t} g\,\frac{\partial c'}{\partial z} - \frac{\partial^2(u_i u_j - \langle u_i u_j\rangle)}{\partial x_i\,\partial x_j} + \nabla\cdot(\mathbf{f} - \langle\mathbf{f}\rangle)\,, \tag{3.3}$$

where u_i and x_i are the velocity and position components. Because (3.3) is linear, p' splits into buoyancy, nonlinear and capillary parts, $p' = p_B + p_N + p_\sigma$, each driven by one of the three source terms. The buoyancy part is set by the indicator field alone, $p_B = -\mathrm{A_t} g\,\nabla^{-2}(\partial c'/\partial z)$, where ∇^{-2} is the inverse Laplacian, and it opposes the force that drives it: a heavy parcel raises the pressure in the fluid below it, which pushes back against its descent. Its contribution to the pressure–scalar correlation in (3.2) is therefore proportional to the buoyancy production

itself,

$$-\mathrm{A_t}g\langle c'c'\rangle - \left\langle c'\frac{\partial p_B}{\partial z}\right\rangle = -\mathrm{A_t}g\,(1-\chi)\langle c'c'\rangle\,, \qquad \chi \equiv \frac{1}{\langle c'c'\rangle}\left\langle c'\,\frac{\partial^2}{\partial z^2}\nabla^{-2}c'\right\rangle, \tag{3.4}$$

so the buoyancy pressure cancels a fraction χ of the buoyancy production. For a statistically homogeneous field with Fourier coefficients $\hat{c}_{\mathbf{k}}$, $\chi = \sum_{\mathbf{k}}(k_z^2/k^2)|\hat{c}_{\mathbf{k}}|^2/\sum_{\mathbf{k}}|\hat{c}_{\mathbf{k}}|^2$ is the fraction of the indicator variance at vertical wavenumbers. It depends only on the shape of the structures: $\chi = 0$ for vertical columns, which slide past one another without a pressure response; $\chi = 1/3$ for isotropic structures; and $\chi \to 1$ for horizontal layers, whose weight is carried entirely by the pressure. In the same way, the capillary pressure removes the gradient part of the capillary force. The two capillary terms in (3.2) combine into $\langle c'(\mathcal{P}\mathbf{f})_z\rangle$, where $\mathcal{P}\mathbf{f} = \mathbf{f} - \langle\mathbf{f}\rangle - \nabla p_\sigma$ and $\mathcal{P}$ is the projection onto divergence-free fields (the Leray–Helmholtz projection). The Laplace pressure of a drop in equilibrium is a pure gradient, which $\mathcal{P}$ removes, so only the part of the capillary force that can drive a flow enters the budget. Let d_T, d_N, d_μ and d_σ denote the mid-plane turbulent-transport, nonlinear-pressure, viscous and capillary terms, normalised by $\dot{h}^2/h$ and signed so that a positive value is a sink of $-\langle wc'\rangle_0$. On the mid-plane, (3.2) becomes an ordinary differential equation for h,

$$\frac{2}{R_f}\,\ddot{h} = \mathrm{A_t}g\,(1-\chi_0)\langle c'c'\rangle_0 - \left(\hat{D} - \frac{1}{R_f} + d_\sigma\right)\frac{\dot{h}^2}{h}\,,$$

$$\hat{D} = \frac{1}{R_f} + d_T + d_N + d_\mu - \hat{w}\,f_C'(0)\,, \tag{3.5}$$

with $\hat{w} = \langle ww\rangle_0/\dot{h}^2$ and $f_C'(0) = h\,\partial C/\partial z|_0$. For $\dot{h}^2 \propto h$, $\ddot{h} = \dot{h}^2/2h$ and

$$\dot{h}^2 = \mathrm{A_t}g\,h\,\frac{\langle c'c'\rangle_0\,(1-\chi_0)}{\hat{D}+d_\sigma}\,, \qquad C_\mathrm{o} \equiv \frac{\langle c'c'\rangle_0\,(1-\chi_0)}{\hat{D}} = 4\alpha_\infty\,. \tag{3.6}$$

Without surface tension, this is the RC04 result $\dot{h}^2 = \mathrm{A_t}gC_\mathrm{o}h$, with the constant written as mid-plane statistics: a purity $\langle c'c'\rangle_0$, which is unity for immiscible fluids, a buoyancy efficiency $1-\chi_0$, and a normalised inertia and drag $\hat{D}$.

4. Direct effect of surface tension

In RC04, every energy-containing quantity scales with h. The curvature and the interface area per unit volume that contribute to $\langle c'(\mathcal{P}\mathbf{f})_z\rangle$ then scale as $1/h$, and the capillary group is the inverse instantaneous Weber number,

$$d_\sigma \sim \frac{\sigma}{\bar{\rho}\,h\,\dot{h}^2} = \frac{1}{\mathrm{We}(t)} = \frac{1}{2\alpha}\left(\frac{\ell_\sigma}{h}\right)^2. \tag{4.1}$$

Writing $d_\sigma = \hat{d}_\sigma(\ell_\sigma/h)^2$ and expanding (3.6) for $d_\sigma \ll \hat{D}$ (§ S2 of the supplementary material),

$$\alpha(t) = \alpha_\infty\left[1 - a_\sigma\left(\frac{\ell_\sigma}{h}\right)^2\right] = \alpha_\infty\left[1 - \frac{a_\sigma}{2\mathrm{A_t}\mathrm{We}_0}\left(\frac{\lambda_0}{h}\right)^2\right], \qquad a_\sigma = \frac{\hat{d}_\sigma}{\hat{D}} > 0\,. \tag{4.2}$$

The sign follows from the restoring character of the capillary force: at the tip of a falling spike, where $c' > 0$, it points upward, and at the cap of a rising bubble, where $c' < 0$, it points downward, so $c'f_z > 0$ and $d_\sigma > 0$. The direct effect of surface tension is therefore a correction proportional to We_0^{-1} that decays as h^{-2} and reduces α. At the end of the simulations of Walker *et al.* (2025, 2026, 2027*a*) (see § 7), $\mathrm{We}(t) \approx (3\text{–}7)\times 10^3$ (figure 2*a*), so $d_\sigma \sim 10^{-4}$ while $\hat{D} = O(1)$.

Focus on Fluids articles must not exceed this page length




Figure 2: (*a*) Capillary group $d_\sigma \sim 1/\mathrm{We}(t)$, (4.1), in the simulations of Walker *et al.* (2025, 2026, 2027*a*) at $\mathrm{We}_0 = 50$, 100 and 200. (*b*) Exponent ν of the correction $(\ell_\sigma/h)^\nu$ against the interface fractal dimension D_s: complete self-similarity, (4.2); the Hinze-scale estimate, (4.4); and the spectral closure, (6.3), with $\ell_{\mathrm{in}} = \ell_\sigma$ and $\ell_{\mathrm{in}} = \ell_H$. The shaded band marks $\nu \approx 1$ from the simulations at equal mixing width.

Small-scale interfacial structure does not change this conclusion. Turbulence should be unable to deform the interface at scales below the Hinze scale ℓ_H (Kolmogorov 1949; Hinze 1955). With the inviscid dissipation scaling of RC04, $\varepsilon_0 = C_\varepsilon \dot{h}^3/h$, and $\dot{h}^2 = 4\alpha \mathrm{A_t} g h$,

$$\ell_H = C_H \left(\frac{\sigma}{\bar{\rho}}\right)^{3/5} \varepsilon^{-2/5}, \qquad \frac{\ell_H}{h} = K_H \left(\frac{\ell_\sigma}{h}\right)^{6/5}, \qquad K_H = C_H\, C_\varepsilon^{-2/5} (2\alpha)^{-3/5}, \tag{4.3}$$

with $C_H \approx 0.725$, so that $K_H \approx 3.3$ for $\alpha = 0.04$ and $C_\varepsilon = 1$. The Hinze scale therefore also separates from h as the layer grows (Chertkov *et al.* 2005), and the capillary force on structures near ℓ_H is balanced mainly by Laplace pressure. The Hinze scale evaluated with the dissipation rate of the simulations (Walker *et al.* 2027*b*) follows this scaling (figure S1 of the supplementary material). For $6 \leqslant t/\tau_0 \leqslant 9.4$, it decreases as $h^{-0.11}$ to $h^{-0.35}$ for the three Weber numbers, around the $h^{-1/5}$ of (4.3), and a fit with that slope gives $K_H \approx 2.3$; with the late-time values of α in the simulations, 0.08–0.13 rather than 0.04, (4.3) gives $K_H \approx 1.6$–2.2. At $t = 9.4\tau_0$, $\ell_H/\ell_\sigma \approx 0.8$–$0.9$ and $\ell_H \approx 0.6$–1.1 times the grid spacing, so the Hinze scale, the capillary length and the grid spacing are comparable. Across Weber numbers, ℓ_H varies as $\mathrm{We}_0^{-0.45}$ rather than $\mathrm{We}_0^{-3/5}$, so the dissipation rate also depends on We_0. The capillary term can instead be estimated at ℓ_H, as the ratio of capillary to buoyancy pressure at that scale weighted by the relative amplitude of the indicator fluctuation there. If the second-order structure function of the indicator scales as r^{2q} (§ 6), (4.3) gives

$$\frac{\sigma/\ell_H}{\Delta\rho\, g\, \ell_H} \left(\frac{\ell_H}{h}\right)^q = \left(\frac{\ell_\sigma}{\ell_H}\right)^2 \left(\frac{\ell_H}{h}\right)^q \propto \left(\frac{\ell_\sigma}{h}\right)^\nu, \qquad \nu = \frac{6q-2}{5} = \frac{7-3D_s}{5}, \tag{4.4}$$

where D_s is the fractal dimension of the interface. For $D_s \geqslant 2$, the exponent is at most $1/5$; for the values $D_s \approx 2.6$–2.7 reported for scalar iso-surfaces in turbulence (Sreenivasan 1991), it is negative, and the correction would grow with h. The sign is again restoring.

5. Relaxation towards self-similarity

Equation (3.5) also describes how a departure from $\dot{h}^2 \propto h$ decays when the profile constants are fixed. Writing $y = \dot{h}^2$, so $\ddot{h} = \frac{1}{2}\,\mathrm{d}y/\mathrm{d}h$, and setting $d_\sigma = 0$,

$$\frac{\mathrm{d}y}{\mathrm{d}h} + (R_f\hat{D} - 1)\,\frac{y}{h} = R_f \mathrm{A_t} g\,(1-\chi_0)\langle c'c'\rangle_0\,, \qquad \text{so that} \qquad \alpha(t) - \alpha_\infty \propto h^{-R_f\hat{D}}\,. \tag{5.1}$$

With $\alpha_\infty \approx 0.035$, $\chi_0 \approx 1/3$ and $R_f \approx 8.0$, the values without surface tension, (3.6) gives $\hat{D} \approx 4.8$ and $R_f\hat{D} \approx 40$. Departures from self-similar growth therefore decay as a high power of h, and a correction that decays as h^{-1}, such as (2.3), cannot be a transient of the moment balance with fixed constants. Together with the sign of (4.2), this shows that an increase of α with σ requires an indirect effect of surface tension on the flow structure: either a slow evolution of χ_0 and $\hat{D}$ with h, or an additional term in the balance that scales as ℓ_σ/h. If surface tension instead fixed the profile constants during the transition, when ℓ_σ/λ_0 is not small, α_∞ itself would depend on We_0 and the excess would not decay with h.

6. A spectral closure

The inner cut-off of the indicator spectrum provides an indirect effect of the required form (§ S4 of the supplementary material). For a two-phase indicator, the second-order structure function $\langle[c(\mathbf{x}+\mathbf{r}) - c(\mathbf{x})]^2\rangle$ is proportional to the probability that a segment of length r crosses the interface. That probability scales with the interface area per unit volume measured at resolution r, which is proportional to $h^{-1}(h/r)^{D_s-2}$, so the structure function scales as r^{3-D_s} (Sreenivasan 1991). Defining q by a structure function proportional to r^{2q} and p by a spectrum $E_c(k) \propto k^{-p}$, with $p = 2q + 1$,

$$D_s = 3 - 2q = 4 - p\,. \tag{6.1}$$

A smooth interface has $D_s = 2$ and the Porod spectrum k^{-2} (Porod 1951); an indicator with the Obukhov–Corrsin spectrum $k^{-5/3}$ (Obukhov 1949; Corrsin 1951) has $D_s = 7/3$. Suppose that only the indicator variance at the energy-containing scales, $1/h \leqslant k \leqslant 2/h$, drives the energy-containing flux, the flux produced at small scales being destroyed locally as in the $k^{-7/3}$ range of the scalar-flux co-spectrum (Lumley 1967). Replacing $\langle c'c'\rangle_0$ in (3.6) by that fraction of the variance, for a spectrum that extends from $1/h$ to an inner cut-off $1/\ell_{\mathrm{in}}$, gives

$$\alpha(h) = \frac{\alpha_\infty}{1 - (\ell_{\mathrm{in}}/h)^{3-D_s}} \approx \alpha_\infty\left[1 + \left(\frac{\ell_{\mathrm{in}}}{h}\right)^{3-D_s}\right]. \tag{6.2}$$

Without surface tension, an immiscible indicator has no inner cut-off, $\ell_{\mathrm{in}} \to 0$. Surface tension sets ℓ_{in} to the smaller of ℓ_σ, below which buoyancy cannot deform the interface, and ℓ_H, below which turbulence cannot. By (4.3), $\ell_{\mathrm{in}} = \ell_\sigma$ for $h/\ell_\sigma \lesssim K_H^5$ and $\ell_{\mathrm{in}} = \ell_H$ beyond. In the simulations, $K_H^5 \approx 60$ and h/ℓ_σ reaches 100–215, so ℓ_H is slightly smaller than ℓ_σ at late times; the two cut-offs differ by less than 25 % (figure S1 of the supplementary material), and the simulations cannot distinguish the two branches of (6.3), which for $D_s = 2$ give $\nu = 1$ and $\nu = 6/5$. The correction (6.2) increases α and, with (2.1) and (4.3),

$$\frac{\alpha - \alpha_\infty}{\alpha_\infty} \approx \begin{cases} (2\mathrm{A_t We}_0)^{-(3-D_s)/2}\,(\lambda_0/h)^{3-D_s}\,, & h/\ell_\sigma \lesssim K_H^5\,, \\ K_H^{3-D_s}\,(2\mathrm{A_t We}_0)^{-3(3-D_s)/5}\,(\lambda_0/h)^{6(3-D_s)/5}\,, & h/\ell_\sigma \gtrsim K_H^5\,. \end{cases} \tag{6.3}$$

For $D_s = 7/3$, this gives $\alpha - \alpha_\infty \propto \mathrm{We}_0^{-1/3}$ at fixed h/λ_0 in the intermediate regime and $\propto \mathrm{We}_0^{-2/5}$ asymptotically. A $\mathrm{We}_0^{-1/2}$ dependence requires $D_s = 2$ with $\ell_{\mathrm{in}} = \ell_\sigma$, or $D_s = 13/6$

Origin	Basis	ν	Coefficient	We_0 dependence	Effect σ on α
Direct capillary term	Moment equations, Eq. (4.2)	2	$-\alpha_\infty a_\sigma$	$\propto -\mathrm{We}_0^{-1}$	decrease
Capillary term at ℓ_H	Hinze-scale estimate, Eq. (4.4)	$(7-3D_s)/5$	—	$\propto -\mathrm{We}_0^{-\nu/2}$	decrease
Inner cut-off $\ell_\mathrm{in}=\ell_\sigma$	Spectral closure, Eq. (6.3)	$3-D_s$	α_∞	$\propto \mathrm{We}_0^{-(3-D_s)/2}$	increase
Inner cut-off $\ell_\mathrm{in}=\ell_H$	Spectral closure, Eq. (6.3)	$\frac{6}{5}(3-D_s)$	$\alpha_\infty K_H^{3-D_s}$	$\propto \mathrm{We}_0^{-3(3-D_s)/5}$	increase
Capillary velocity	Dimensional analysis, Eq. (2.3)	1	K	$\propto \mathrm{We}_0^{-1/2}$	increase
Simulations, equal h	Fit, $7 \leqslant h/\lambda_0 \leqslant 12.5$, Eq. (7.1)	0.97	8.9 (7.7–9.9)	$\propto \mathrm{We}_0^{-1/2}$	increase
Simulations, fixed t/τ_0	Fit, $6 \leqslant t/\tau_0 \leqslant 9.4$	—	—	$\alpha \propto \mathrm{We}_0^{-m}$, $m = 0.37$–0.46	increase

Table 1: Corrections to the growth parameter of the form $\alpha - \alpha_\infty = (\text{coefficient}) \times (\ell_\sigma/h)^\nu$, and the resulting dependence on We_0 at fixed h/λ_0.

with $\ell_\mathrm{in} = \ell_H$. For a smooth interface with $\ell_\mathrm{in} = \ell_\sigma$, (6.3) is (2.3) with $K = \alpha_\infty$: a correction linear in ℓ_σ/h, of the observed sign, and proportional to $\mathrm{We}_0^{-1/2}$ at fixed h/λ_0. Table 1 and figure 2(*b*) summarise the corrections.

7. Comparison with simulations

The simulations of Walker *et al.* (2025, 2026, 2027*a*) solve the compressible multicomponent equations with the high-order compact finite-difference code `Miranda` (Cook 2007), using localised artificial diffusivity with interface sharpening (Brill *et al.* 2025) and a continuum surface force (Brackbill *et al.* 1992), on 1120^3 points at $\mathrm{A_t} = 0.5$ and $\mathrm{Re}_0 = \infty$, for $\mathrm{We}_0 = 50$, 100, 200 and ∞ and a miscible reference case; Walker *et al.* (2027*a*) give the details. Decreasing We_0 delays the transition, the time of maximum growth of the mixedness increasing linearly with σ as $t_\theta/\tau_0 \approx 1.88 + 179/\mathrm{We}_0$ (§ S5 of the supplementary material), and raises the late-time α (figure 1*a*).

Compared at fixed time, the effective exponent in $\alpha \propto \mathrm{We}_0^{-m}$, fitted over $\mathrm{We}_0 = 50$, 100 and 200, drifts from 0.46 at $t = 6\tau_0$ to 0.37–0.38 at $t = 8.5$–$9.4\tau_0$ (figure 3*b* and table S2 of the supplementary material), close at late times to the fit $\alpha \approx 0.49\,\mathrm{We}_0^{-0.35}$ of Walker *et al.* (2026). An exponent obtained at fixed time is therefore not a property of the flow; it depends on when the simulations are compared. The analysis above is formulated in h, and every correction in table 1 is a function of ℓ_σ/h. Plotted against h/λ_0 (figure 1*b*), α without surface tension settles at 0.034–0.036 for $h/\lambda_0 \gtrsim 8$, whereas with surface tension it continues to decrease and remains two to five times larger at equal width. Over $7 \leqslant h/\lambda_0 \leqslant 12.5$, where all four simulations overlap, the excess evaluated at equal h falls on a single line when plotted against ℓ_σ/h (figure 3*a*),

$$\alpha - \alpha_{\mathrm{We}_0=\infty} \approx 8.9\,\frac{\ell_\sigma}{h}\,, \tag{7.1}$$

with the coefficient between 7.7 and 9.9 for the three Weber numbers and a fitted slope of 0.97 on logarithmic axes. This is (2.3) with $K \approx 8.9$. At fixed h/λ_0, the excess is proportional to $\mathrm{We}_0^{-1/2}$, the value that its effective exponent approaches as the $\mathrm{We}_0 = \infty$ case settles (figure 3*b*), and at fixed We_0 it decays as h^{-1}. The exponent agrees with the spectral closure




Figure 3: (*a*) Excess $\alpha - \alpha_{\mathrm{We}_0=\infty}$ at equal h against ℓ_σ/h for $7 \leqslant h/\lambda_0 \leqslant 12.5$, with the fit (7.1), the spectral closure (6.2) with $D_s = 2$, and the magnitude of the direct correction (4.2) with $a_\sigma \approx 3$. (*b*) Effective exponent m in $\alpha \propto \mathrm{We}_0^{-m}$ over $\mathrm{We}_0 = 50$, 100, 200, for α and for $\alpha - \alpha_{\mathrm{We}_0=\infty}$ at fixed t/τ_0 and at fixed h/λ_0, and for α at fixed $t - t_\theta$; x is the corresponding progress variable.

for a smooth interface, but the magnitude is about 250 times $K = \alpha_\infty$; it exceeds the direct capillary correction (4.2) by four orders of magnitude and has the opposite sign.

At fixed h/λ_0, the effective exponent of α itself is only 0.25–0.37, because α includes α_∞, and at fixed time since the transition it rises from about zero to 0.2 (figure 3*b*). The direct capillary term is negligible over the whole late-time range: the capillary group falls below 10^{-2} after the transition and is $(1.5\text{–}3.2) \times 10^{-4}$ at the end of the simulations (figure 2*a*). The capillary length is comparable to the grid spacing. With $\lambda_0 \approx 1.02\,\mathrm{cm}$ and $\Delta x \approx 0.114\,\mathrm{cm}$, $\ell_\sigma/\Delta x = 1.26$, 0.89 and 0.63 for $\mathrm{We}_0 = 50$, 100 and 200, and the late-time effect of surface tension is nevertheless organised by ℓ_σ; the limit of drop breakup at the grid scale in these simulations is analysed by Walker *et al.* (2027*b*).

The mid-plane mass flux indicates where the indirect effect resides (§ S6 of the supplementary material). Without surface tension, the shape factor R_f settles at 8.0, the value for a linear profile, and with surface tension at 9.7–10.7, above the self-similar range. Writing $\langle \rho' w' \rangle_0 = -C_r \rho_{\mathrm{rms}} w_{\mathrm{rms}}$, as in CCM04, with $\rho_{\mathrm{rms}} = (\Delta\rho/2)(1-\theta_0)^{1/2}$ on the mid-plane and $\mathrm{Fr}_0^2 = w_{\mathrm{rms}}^2/\mathrm{A_t} g h$, the definition of R_f gives $\alpha = (R_f^2/16)\, C_r^2\, (1-\theta_0)\, \mathrm{Fr}_0^2$. Averaged over $8 \leqslant t/\tau_0 \leqslant 9.4$, the factor 3.84 in α between $\mathrm{We}_0 = 50$ and $\mathrm{We}_0 = \infty$ splits as 1.76 from R_f^2, 1.49 from the purity $1-\theta_0$ and 1.46 from $C_r^2 \mathrm{Fr}_0^2$, whereas the CCM04 closure, $\alpha \propto (1-\theta_0)^{3/2}$, gives 1.82. Between $\mathrm{We}_0 = \infty$ and the miscible case, the same closure predicts 8.8 for a factor of 1.51, so the purity alone overpredicts the difference between miscible and immiscible growth. Surface tension therefore raises α through the shape factor, the purity and $C_r \mathrm{Fr}_0$ together, and these are the indirect effect required in § 5. The energy budget constrains the sign in the same way: the surface energy lowers α, so an increase with σ must come from a smaller dissipated fraction or from a change in the profile shape.

8. Conclusions

Complete self-similarity leaves no dimensionless group that contains σ, so surface tension can change the late-time growth parameter of immiscible RT turbulence only through corrections in ℓ_σ/h. In the self-similar moment equations extended to immiscible fluids, the direct capillary term is the inverse instantaneous Weber number; it reduces α and scales as We_0^{-1}. The moment balance with fixed constants relaxes approximately as h^{-40}, so the

observed increase of α, which decays only as h^{-1}, requires an indirect effect of surface tension on the flow structure. A correction linear in ℓ_σ/h is the form that this effect takes in simulations: compared at equal mixing width, $\alpha - \alpha_{\mathrm{We}_0=\infty} \approx 8.9\,\ell_\sigma/h$, which is a $\mathrm{We}_0^{-1/2}$ dependence at fixed h/λ_0 and vanishes as $h/\ell_\sigma \to \infty$. A power law in We_0 extracted at fixed time has an exponent that drifts with the time of comparison. A spectral closure in which surface tension sets the inner cut-off of a smooth interface reproduces the exponent and the sign but not the magnitude, and the mass-flux decomposition divides the effect among the shape factor R_f, the purity and $C_r\mathrm{Fr}_0$, while the purity alone overpredicts the difference between miscible and immiscible growth. The coefficient K remains to be determined by a closure for the profile constants χ_0 and $\hat{D}$ in the presence of surface tension.

Supplementary data. Supplementary material is available at https://doi.org/ ....

Acknowledgements. This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the U.S. Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the U.S. Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the U.S. Government or any agency thereof.

Funding. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Department of Energy Computational Science Graduate Fellowship under Award Number DE-SC0024386.

Declaration of interests. The authors report no conflict of interest.

Data availability statement. The data that support the findings of this study are openly available on GitHub at https://github.com/mw6136/alpha-theory.

Author ORCIDs. M. D. Walker, https://orcid.org/0000-0002-4765-6539; S. S. Nixon, https://orcid.org/0009-0003-5313-9081

Author contributions. M. D. Walker: Methodology, Formal analysis, Investigation, Software, Data curation, Writing – original draft. S. S. Nixon: Conceptualization, Methodology, Formal analysis, Writing – review & editing.

REFERENCES

Barenblatt, G. I. 1996 *Scaling, Self-Similarity, and Intermediate Asymptotics*. Cambridge: Cambridge University Press.

Brackbill, J. U., Kothe, D. B. & Zemach, C. 1992 A continuum method for modeling surface tension. *Journal of Computational Physics* **100** (2), 335–354.

Brill, S. R., Olson, B. J. & Bokman, G. T. 2025 An extension of the localized artificial diffusivity method for immiscible and high density ratio flows. *Journal of Computational Physics* **542**, 114366.

Chandrasekhar, S. 1981 *Hydrodynamic and Hydromagnetic Stability*. New York: Dover Publications.

Chertkov, M., Kolokolov, I. & Lebedev, V. 2005 Effects of surface tension on immiscible Rayleigh-Taylor turbulence. *Phys. Rev. E* **71**, 055301.

Cook, A. W. 2007 Artificial fluid properties for large-eddy simulation of compressible turbulent mixing. *Physics of Fluids* **19** (5), 055103.

Cook, A. W., Cabot, W. & Miller, P. L. 2004 The mixing transition in Rayleigh-Taylor instability. *Journal of Fluid Mechanics* **511**, 333–362.

Corrsin, S. 1951 On the spectrum of isotropic temperature fluctuations in an isotropic turbulence. *Journal of Applied Physics* **22** (4), 469–473.

Dimonte, G. & Schneider, M. 2000 Density ratio dependence of Rayleigh-Taylor mixing for sustained and impulsive acceleration histories. *Physics of Fluids* **12** (2), 304–321.

Dimonte, G., Youngs, D. L., Dimits, A., Weber, S., Marinak, M., Wunsch, S., Garasi, C., Robinson, A.,

Andrews, M. J., Ramaprabhu, P., Calder, A. C., Fryxell, B., Biello, J., Dursi, L., MacNeice, P., Olson, K., Ricker, P., Rosner, R., Timmes, F., Tufo, H., Young, Y.-N. & Zingale, M. 2004 A comparative study of the turbulent Rayleigh-Taylor instability using high-resolution three-dimensional numerical simulations: The Alpha-Group collaboration. *Physics of Fluids* **16** (5), 1668–1693.

Hinze, J. O. 1955 Fundamentals of the hydrodynamic mechanism of splitting in dispersion processes. *AIChE Journal* **1** (3), 289–295.

Kolmogorov, A. N. 1949 On the disintegration of drops in a turbulent flow. In *Dokl. Akad. Nauk SSSR*, , vol. 66, pp. 825–828.

Lumley, J. L. 1967 Similarity and the turbulent energy spectrum. *Physics of Fluids* **10** (4), 855–858.

Obukhov, A. M. 1949 Structure of the temperature field in turbulent flows. *Izvestiya Akademii Nauk SSSR, Seriya Geograficheskaya i Geofizicheskaya* **13**, 58–69.

Porod, G. 1951 Die Röntgenkleinwinkelstreuung von dichtgepackten kolloiden Systemen: I. Teil. *Kolloid-Zeitschrift* **124**, 83–114.

Ristorcelli, J. R. & Clark, T. T. 2004 Rayleigh-Taylor turbulence: self-similar analysis and direct numerical simulations. *Journal of Fluid Mechanics* **507**, 213–253.

Roberts, M. S. & Jacobs, J. W. 2016 The effects of forced small-wavelength, finite-bandwidth initial perturbations and miscibility on the turbulent Rayleigh-Taylor instability. *Journal of Fluid Mechanics* **787**, 50–83.

Sohn, S.-I. 2009 Effects of surface tension and viscosity on the growth rates of Rayleigh-Taylor and Richtmyer-Meshkov instabilities. *Phys. Rev. E* **80**, 055302.

Sreenivasan, K. R. 1991 Fractals and multifractals in fluid turbulence. *Annual Review of Fluid Mechanics* **23**, 539–604.

Walker, M. D., Brill, S. R. & Olson, B. J. 2025 Effects of surface tension and viscosity on immiscible Rayleigh-Taylor mixing. In *78th Annual Meeting of the APS Division of Fluid Dynamics*. Houston, TX, session U30, abstract 8.

Walker, M. D., Brill, S. R. & Olson, B. J. 2026 High-fidelity parametric study of surface tension and viscosity effects in immiscible Rayleigh–Taylor mixing. In *19th International Workshop on the Physics of Compressible Turbulent Mixing (IWPCTM19)*. Ann Arbor, MI.

Walker, M. D., Brill, S. R. & Olson, B. J. 2027*a* Immiscible Rayleigh-Taylor turbulence I: Macroscopic behavior and growth dynamics. *Physical Review Fluids* Submitted.

Walker, M. D., Brill, S. R. & Olson, B. J. 2027*b* Immiscible Rayleigh-Taylor turbulence II: Mesoscale mixing and spectral characteristics. *Physical Review Fluids* Submitted.

Youngs, D. L. 1984 Numerical simulation of turbulent mixing by Rayleigh-Taylor instability. *Physica D: Nonlinear Phenomena* **12** (1), 32–44.

Zhao, D., Huang, X. & Li, G. 2026 Interfacial dynamics and energy cascade in immiscible Rayleigh-Taylor turbulence. *Journal of Fluid Mechanics* **1040**, A56.

Zhao, D. & Li, G. 2026 Influence of immiscibility and atwood number on the dynamics of confined Rayleigh-Taylor turbulence. *International Journal of Multiphase Flow* **202**, 105811.

Rapids articles must not exceed this page length