

# Proper Title

Michał Wichrowski

## Abstract

Abstarct Placeholder

**Keywords:** Keyords go here.

**AMS subject classifications:** Classification codes go here.

## 1 Introduction

### 1.1 What \*Has\* Been Done (The Existing Literature)

If you dive into the literature on singularities in finite/hyperelasticity, you will mostly find two major phenomena:

- Crack Tips in Tension (Knowles and Sternberg, 1970s):\*\* This is the most famous work on nonlinear singularities. They looked at a  $360^\circ$  corner (a crack) pulled in pure tension. They found that in a neo-Hookean material, the asymptotic displacement field behaves very differently from linear fracture mechanics. However, they only dealt with pure traction-free boundaries and pure tensile opening.
- Cavitation (John Ball, 1982):\*\* Ball proved that under severe hydrostatic tension, hyperelastic equations admit singular weak solutions where a hole spontaneously opens in the middle of a perfectly solid domain (a point singularity where  $\mathbf{F} \rightarrow \infty$ ).
- Surface Creasing (Biot 1963, Hong et al. 2009):\*\* Biot found that if you compress a flat half-space of a neo-Hookean material, its surface becomes unstable at a critical stretch ( $\lambda \approx 0.54$ ). Decades later, researchers realized this instability is a \*crease\* (a point singularity forming on a flat surface).

### 1.2 The Gap:

Almost all the existing nonlinear singularity literature avoids the exact setup you described: \*\*a geometric corner (e.g.,  $90^\circ$ ) with mixed boundary conditions (Dirichlet meets Neumann) under compression.\*\*

- Linear theory (Kondratiev) handles the mixed BC corner, but ignores the large-deformation kinematics.
- Nonlinear theory (Knowles/Sternberg) handles large deformations, but avoids the Dirichlet-Neumann condition and avoids compression.

---

<sup>0</sup>Interdisziplinäres Zentrum für Wissenschaftliches Rechnen (IWR), Ruprecht-Karls-Universität Heidelberg, Germany, [mt.wichrowsk@uw.edu.pl](mailto:mt.wichrowsk@uw.edu.pl)

- Instability theory handles compression, but usually on flat domains or purely symmetric geometries, ignoring the existing geometric corner singularity.

**The Gap:** How does an already-existing stress concentration at a geometric Dirichlet-Neumann corner couple with a nonlinear structural instability (buckling/creasing)? How do we bound the operators there? That is fundamentally an open question.

## 2 Problem formulation

### 2.1 Finite-strain elastostatics as energy minimisation

We consider a hyperelastic body occupying a reference  $d$ -dimensional domain  $\Omega \subset \mathbb{R}^d$  with boundary  $\partial\Omega = \overline{\Gamma_D} \cup \overline{\Gamma_N}$ ,  $\Gamma_D \cap \Gamma_N = \emptyset$ . The deformation of the body is described by the displacement field  $\mathbf{u} : \Omega \rightarrow \mathbb{R}^d$ , the deformation map  $\mathbf{y}(\mathbf{x}) = \mathbf{x} + \mathbf{u}(\mathbf{x})$ , and the associated deformation gradient

$$\mathbf{F} = \nabla \mathbf{y} = \mathbf{I} + \nabla \mathbf{u}, \quad J = \det \mathbf{F} > 0, \quad (1)$$

where  $\nabla$  denotes the gradient with respect to the reference configuration. The mechanical response is encoded in a strain energy density  $\Psi(\mathbf{F})$ , and the equilibrium configuration is the minimiser of the total potential energy

$$\mathcal{E}(\mathbf{u}) = \int_{\Omega} \Psi(\mathbf{F}) dV - \int_{\Omega} \mathbf{f}_0 \cdot \mathbf{u} dV - \int_{\Gamma_N} \mathbf{t}_0 \cdot \mathbf{u} dS, \quad (2)$$

over the space of admissible displacements  $U = \{\mathbf{u} \in H^1(\Omega; \mathbb{R}^d) : \mathbf{u} = \mathbf{g} \text{ on } \Gamma_D\}$ , where  $\mathbf{f}_0$  is a body force,  $\mathbf{t}_0$  a prescribed traction on the Neumann boundary, and  $\mathbf{g}$  the prescribed boundary displacement. We seek  $\mathbf{u} \in U$  such that  $\mathcal{E}(\mathbf{u}) \leq \mathcal{E}(\mathbf{v})$  for all admissible  $\mathbf{v}$ .

### 2.2 Stationarity, stress, and tangent modulus

A minimiser satisfies the stationarity condition  $D_{\delta \mathbf{u}} \mathcal{E}(\mathbf{u}) = 0$  for all admissible variations  $\delta \mathbf{u}$  (vanishing on  $\Gamma_D$ ), which is the weak form of equilibrium

$$\int_{\Omega} \mathbf{P}(\mathbf{F}) : \nabla \delta \mathbf{u} dV = \int_{\Omega} \mathbf{f}_0 \cdot \delta \mathbf{u} dV + \int_{\Gamma_N} \mathbf{t}_0 \cdot \delta \mathbf{u} dS, \quad (3)$$

where the first Piola–Kirchhoff stress is the first derivative of the energy with respect to the deformation gradient,

$$\mathbf{P}(\mathbf{F}) = \frac{\partial \Psi}{\partial \mathbf{F}}. \quad (4)$$

The natural boundary condition associated with  $\Gamma_N$  is  $\mathbf{P}\mathbf{N} = \mathbf{t}_0$ , with  $\mathbf{N}$  the reference outward unit normal. Because (3) is nonlinear in  $\mathbf{u}$ , it is solved iteratively by Newton’s method. Linearising the residual about a current iterate  $\bar{\mathbf{u}}$  in a direction  $\Delta \mathbf{u}$  introduces the fourth-order tangent (elasticity) modulus  $\mathbb{L}$ , with components  $L_{iJkL}$ ,

$$\mathbb{L} = \frac{\partial \mathbf{P}}{\partial \mathbf{F}} = \frac{\partial^2 \Psi}{\partial \mathbf{F} \otimes \partial \mathbf{F}}, \quad L_{iJkL} = \frac{\partial^2 \Psi}{\partial F_{iJ} \partial F_{kL}}, \quad (5)$$

which possesses the major symmetry  $L_{iJkL} = L_{kLiJ}$ , so that each Newton step requires solving the linear problem

$$\int_{\Omega} \nabla \delta \mathbf{u} : \mathbb{L}(\bar{\mathbf{F}}) : \nabla \Delta \mathbf{u} dV = -R(\bar{\mathbf{u}}; \delta \mathbf{u}), \quad (6)$$

with  $R(\bar{\mathbf{u}}; \delta \mathbf{u})$  the residual of (3) evaluated at  $\bar{\mathbf{u}}$ . The well-posedness and conditioning of (6)—and hence the convergence of the whole nonlinear solver—are governed by the coercivity and boundedness of the bilinear form induced by  $\mathbb{L}(\bar{\mathbf{F}})$ . The central object of this work is therefore the behaviour of  $\mathbb{L}$  as a function of the local deformation state.

### 2.3 The neo-Hookean material

To fix ideas and to exhibit explicit constants we use, as a running example, the compressible neo-Hookean strain energy density adopted in the cut-finite-element study [9] that motivates this analysis,

$$\Psi(\mathbf{F}) = \frac{\mu_e}{2} (\text{tr}(\mathbf{F}^T \mathbf{F}) - d) + f(J), \quad (7)$$

where  $\mu_e > 0$  is the shear modulus and  $f(J)$  is a strictly convex volumetric penalty with  $f(1) = 0$ ,  $f'(1) = -\mu_e$ , and  $f(J) \rightarrow \infty$  as  $J \rightarrow 0^+$ . Two standard choices are

$$f_{\log}(J) = -\mu_e \log J + \lambda_e \log^2 J, \quad f_{\text{quad}}(J) = \frac{\kappa}{2} (J - 1)^2, \quad (8)$$

with  $\lambda_e$  the Lamé constant and  $\kappa$  a bulk modulus; (7) with  $f_{\log}$  reproduces the model of [9]. The corresponding stress and tangent are

$$\mathbf{P}(\mathbf{F}) = \mu_e \mathbf{F} + f'(J) J \mathbf{F}^{-T}, \quad (9)$$

$$L_{iJkL} = \mu_e \delta_{ik} \delta_{JL} + (f''(J) J^2 + f'(J) J) F_{iL}^{-T} F_{kJ}^{-T} - f'(J) J F_{iJ}^{-T} F_{kL}^{-T}. \quad (10)$$

The first term  $\mu_e \delta_{ik} \delta_{JL}$  is constant; the entire nonlinear part is controlled by  $J$  and the *inverse* deformation gradient  $\mathbf{F}^{-1}$ , yielding the structural bound

$$\|\mathbb{L}(\mathbf{F})\| \leq \mu_e + \Theta(J) \|\mathbf{F}^{-1}\|^2, \quad (11)$$

where  $\|\mathbf{F}^{-1}\| = \lambda_{\min}^{-1}$  is set by the smallest principal stretch and  $\Theta$  is a scalar growth function fixed by  $f$  (computed explicitly in §3.4). Consequently  $\mathbb{L}$  can become unbounded through volumetric collapse,  $\lambda_{\min} \rightarrow 0$  (equivalently  $\mathbf{F}^{-1} \rightarrow \infty$ ); whether a purely tensile blow-up of  $\mathbf{F}$  (with  $\mathbf{F}^{-1}$  bounded) leaves  $\mathbb{L}$  finite depends on the growth of  $\Theta(J)$ , which turns out to discriminate sharply between the two penalties in (8). This dichotomy is the analytical backbone of the present paper.

### 2.4 The mixed-corner singularity

The configuration of interest is a right-angle junction where the two boundary-condition types meet: a Dirichlet face ( $\Gamma_D$ , prescribed displacement) abutting a traction-free Neumann face ( $\Gamma_N$ ). In  $\mathbb{R}^d$  this junction is a codimension-two *edge*  $\Sigma$  (a point for  $d = 2$ , a curve or surface for  $d \geq 3$ ); its transverse geometry, in the two-plane normal to  $\Sigma$ , is the right-angle corner we analyse, and we use polar coordinates  $(r, \theta)$  in that transverse plane. Such junctions arise generically wherever a clamped face meets a traction-free face — for instance along the base edges of a compressed body, the concrete setting taken up in Section 5. Classical Kondratiev/edge theory for the linearised problem predicts a transverse singularity

$$\mathbf{u}(r, \theta) \sim K r^\alpha \Phi(\theta), \quad \alpha \in (0, 1), \quad (12)$$

with  $r$  the distance to the edge  $\Sigma$ ,  $K$  a generalised stress-intensity factor and  $\alpha$  the (geometry- and Poisson-ratio-dependent) singularity exponent. The deformation gradient then scales as  $\mathbf{F} = \mathbf{I} + \mathcal{O}(K r^{\alpha-1})$ , so  $\|\mathbf{F}\| \rightarrow \infty$  as  $r \rightarrow 0$ .

The question this raises—and which the remainder of the paper addresses—is how this kinematic singularity propagates into the nonlinear tangent operator (5). In view of (11), the answer hinges on which principal stretch diverges: a tensile corner keeps  $\mathbf{F}^{-1}$  bounded and (for a suitable penalty) leaves  $\mathbb{L}$  well behaved, whereas a compressive corner drives  $\lambda_{\min} \rightarrow 0$  and blows up the *norm* of  $\mathbb{L}$  (severe ill-conditioning). The tangent nonetheless remains strongly elliptic — the compressive instability is *structural* (a corner analogue of creasing), not a change of type of the PDE. The following sections analyse the two scenarios in turn (Section 3) and then show how a small imperfection regularises the compressive case through a structural bifurcation (Section 4).

### 3 Asymptotic behaviour of the tangent operator at a mixed corner

#### 3.1 Geometry and linearised kinematics

Let the reference domain  $\Omega \subset \mathbb{R}^d$  ( $d \geq 2$ ) carry a right-angle Dirichlet–Neumann edge  $\Sigma$  of codimension two. Near a point of  $\Sigma$  split  $\mathbb{R}^d = T\Sigma \oplus \Pi$  into the edge tangent space and the normal two-plane  $\Pi$ , and use polar coordinates  $(r, \theta)$  in  $\Pi$  with  $r \in [0, R]$ ,  $\theta \in [0, \pi/2]$ , where  $r = \text{dist}(\cdot, \Sigma)$ . Write  $B_\rho$  for the tubular neighbourhood  $\{\mathbf{x} \in \Omega : \text{dist}(\mathbf{x}, \Sigma) < \rho\}$  of the edge. The leading singularity is governed by the transverse problem in  $\Pi$ , modulated smoothly along  $\Sigma$ ; for  $d = 2$  the edge degenerates to a point and  $B_\rho$  to the corner disc.

- **Dirichlet boundary** ( $\Gamma_D$ ):  $\theta = 0$ , deformation prescribed,  $\mathbf{y}(r, 0) = \mathbf{g}(r)$ .
- **Neumann boundary** ( $\Gamma_N$ ):  $\theta = \pi/2$ , traction-free,  $\mathbf{P}\mathbf{N} = \mathbf{0}$ .

Classical Kondratiev theory for the *linearised* problem gives the leading-order singular displacement

$$\mathbf{u}(r, \theta) \sim K r^\alpha \Phi(\theta), \quad (13)$$

with  $K$  the generalised stress-intensity factor (set by the far field),  $\Phi$  a smooth angular profile, and  $\alpha \in (0, 1)$  the geometry- and Poisson-dependent exponent. Consequently

$$\mathbf{F} = \mathbf{I} + \mathcal{O}(K r^{\alpha-1}), \quad \|\mathbf{F}\| \rightarrow \infty \text{ as } r \rightarrow 0. \quad (14)$$

Whether this kinematic blow-up contaminates the tangent operator  $\mathbb{L} = \partial^2 \Psi / \partial \mathbf{F} \partial \mathbf{F}$  depends not on  $\|\mathbf{F}\|$  but on which principal stretches diverge, which is the content of the rest of this section.

#### 3.2 Standing assumptions on the stored energy

The mechanism we describe is not specific to the neo-Hookean model; that model only serves to fix ideas and to exhibit explicit constants. We therefore phrase the analysis for a general frame-indifferent energy and isolate the structural properties that the argument actually uses. Throughout,  $\|\cdot\|$  denotes the operator (spectral) norm on tensors,  $0 < \lambda_{\min}(\mathbf{F}) \leq \dots \leq \lambda_{\max}(\mathbf{F})$  are the singular values of  $\mathbf{F}$  (the  $d$  principal stretches), and we recall  $\|\mathbf{F}^{-1}\| = \lambda_{\min}(\mathbf{F})^{-1}$ .

**Assumption 1** (Regularity and frame indifference). The stored energy  $\Psi : \text{GL}^+(d) \rightarrow \mathbb{R}$  is objective,  $\Psi \in C^2(\text{GL}^+(d))$ , and  $\Psi(\mathbf{F}) \rightarrow +\infty$  as  $\det \mathbf{F} \rightarrow 0^+$ .

**Assumption 2** (Inverse-channel growth of the tangent). There exist a continuous, nondecreasing function  $\Theta : (0, \infty) \rightarrow (0, \infty)$  and a constant  $c_0 \geq 0$  such that the tangent modulus  $\mathbb{L}(\mathbf{F}) = D^2 \Psi(\mathbf{F})$  satisfies, for all  $\mathbf{F} \in \text{GL}^+(d)$ ,

$$\|\mathbb{L}(\mathbf{F})\| \leq c_0 + \Theta(\det \mathbf{F}) \|\mathbf{F}^{-1}\|^2. \quad (15)$$

**Assumption 3** (Uniform strong ellipticity on bounded-distortion sets). For every  $0 < c \leq 1$  and  $M \geq 1$  there is  $\gamma(c, M) > 0$  such that whenever  $\lambda_{\min}(\mathbf{F}) \geq c$  and  $\det \mathbf{F} \leq M$ ,

$$L_{iJkL}(\mathbf{F}) m_i N_J m_k N_L \geq \gamma(c, M) |\mathbf{m}|^2 |\mathbf{N}|^2 \quad \forall \mathbf{m}, \mathbf{N} \in \mathbb{R}^2. \quad (16)$$

The decisive feature of Assumption 2 is that the inverse-stretch factor  $\|\mathbf{F}^{-1}\|^2 = \lambda_{\min}^{-2}$  is the *only* algebraic channel through which the Hessian can blow up; the dependence on  $\|\mathbf{F}\|$  (equivalently on  $\lambda_{\max}$ ) enters solely through the scalar  $\Theta(\det \mathbf{F})$ . Whether a tensile corner ( $\lambda_{\max} \rightarrow \infty$ ,  $\det \mathbf{F} \rightarrow \infty$ ,  $\lambda_{\min}$  bounded below) is benign therefore reduces to the *growth rate* of  $\Theta$ , which we quantify in §3.4. Materials with genuine tensile stiffening of the deviatoric part (Gent, Arruda–Boyce), whose tangent diverges through  $\lambda_{\max}$  directly rather than through  $\Theta(\det \mathbf{F})$ , violate (15) and are excluded.

**Lemma 1** (Assumptions for the neo-Hookean model). *Let  $\Psi$  be the neo-Hookean energy (7) with  $\mu_e > 0$  and  $f \in C^2(0, \infty)$ . Then Assumption 2 holds with  $c_0 = \mu_e$  and*

$$\Theta(J) = |f''(J)J^2 + f'(J)J| + |f'(J)J|, \quad (17)$$

*and Assumption 3 holds (rank-one convexity away from  $\det \mathbf{F} = 0$ ). Assumption 1 additionally requires the barrier property  $f(J) \rightarrow \infty$  as  $J \rightarrow 0^+$ : this holds for  $f_{\log}$  but fails for  $f_{\text{quad}}$ , since  $f_{\text{quad}}(J) \rightarrow \frac{\kappa}{2} < \infty$  (see Remark 3 and Scenario B).*

*Proof.* Objectivity and  $C^2$  regularity on  $\text{GL}^+(d)$  are immediate; the barrier hypothesis  $f(J) \rightarrow \infty$  as  $J \rightarrow 0^+$ , when assumed, gives the blow-up required by Assumption 1. In (10) the first term is the scaled identity, of norm  $\mu_e$ . Each of the remaining two terms is a rank-one, respectively transposition, contraction of  $\mathbf{F}^{-T} \otimes \mathbf{F}^{-T}$ ; since  $\|\mathbf{F}^{-T}\| = \|\mathbf{F}^{-1}\|$ , each is bounded in operator norm by its scalar prefactor times  $\|\mathbf{F}^{-1}\|^2$ . The triangle inequality gives (15) with  $c_0 = \mu_e$  and  $\Theta$  as in (17). For Assumption 3, contracting (10) with  $\mathbf{m} \otimes \mathbf{N}$  and writing  $\xi := \mathbf{m} \cdot \mathbf{F}^{-T} \mathbf{N} = \mathbf{N} \cdot \mathbf{F}^{-1} \mathbf{m}$ , the two volumetric terms combine  $((f''J^2 + f'J)\xi^2 - f'J\xi^2)$  to the single clean form

$$L_{iJkL} m_i N_J m_k N_L = \mu_e |\mathbf{m}|^2 |\mathbf{N}|^2 + f''(J) J^2 \xi^2. \quad (18)$$

Wherever  $f''(J) \geq 0$  the second term is nonnegative and (16) holds with  $\gamma = \mu_e$ , uniformly in  $\mathbf{F}$  (no distortion bound needed); this covers  $f_{\text{quad}}$  ( $f'' = \kappa > 0$ ) everywhere and  $f_{\log}$  on  $\{f''_{\log} \geq 0\}$ .  $\square \square$

*Remark 1* (Strength and a tension caveat). Identity (18) is sharper than (16): the neo-Hookean tangent is strongly elliptic with constant *exactly*  $\mu_e$  wherever  $f''(J) \geq 0$ , and the volumetric term only adds to it. The Hessian of the quadratic (Hookean) part is constant, which is why the *deviatoric* response never blows up; only the *volumetric* response, through  $\Theta$ , can. One caveat for  $f_{\log}$ : since  $f''_{\log}(J) = (\mu_e + 2\lambda_e - 2\lambda_e \log J)/J^2$ , convexity is lost once  $J > J_* := \exp(1 + \mu_e/(2\lambda_e))$ , and then (18) can turn negative for  $\xi$  near its maximal value  $|\mathbf{m}||\mathbf{N}|/\lambda_{\min}$ . Hence Assumption 3 holds throughout *compression* ( $J \leq 1$ ) and moderate tension, but  $f_{\log}$  can lose strong ellipticity under *extreme dilation* ( $J > J_*$ )—a tension-side phenomenon distinct from the compressive analysis, and one more reason the bounded- $J$  qualifier matters in §3.5. The globally convex  $f_{\text{quad}}$  is free of this.

### 3.3 The non-compressive hypothesis

The qualitative split between the two scenarios is captured by a single quantity: the smallest principal stretch near the corner. We call the corner singularity *non-compressive* when  $\lambda_{\min}$  stays bounded away from zero, even though  $\lambda_{\max}$  (hence  $\|\mathbf{F}\|$  and  $\det \mathbf{F}$ ) may diverge.

**Assumption 4** (Non-compressive singularity, (NC)). There exist  $\rho \in (0, R]$  and  $c \in (0, 1]$  such that the equilibrium deformation satisfies

$$\lambda_{\min}(\mathbf{F}(\mathbf{x})) \geq c, \quad \text{equivalently} \quad \|\mathbf{F}^{-1}(\mathbf{x})\| \leq c^{-1}, \quad \text{for a.e. } \mathbf{x} \in B_\rho. \quad (19)$$

We stress what (19) does *not* say. It does not require incompressibility, and crucially it does *not* bound  $\det \mathbf{F}$ : in a tensile corner  $\lambda_{\max} \sim r^{\alpha-1} \rightarrow \infty$ , so  $J = \det \mathbf{F} = \prod_i \lambda_i \sim r^{\alpha-1} \rightarrow \infty$  as well (one diverging stretch, the remaining  $d-1$  bounded). The only quantity (NC) controls is  $\|\mathbf{F}^{-1}\| \leq c^{-1}$ . Physically, (NC) states that the material is pulled around the corner (tension/shear) rather than crushed into it. Because  $\det \mathbf{F}$  is *not* bounded under (NC), the factor  $\Theta(\det \mathbf{F})$  in (15) need not stay bounded; the next subsection makes its growth explicit, and this is precisely where the two volumetric penalties (8) part ways. Scenario B (§3.6) treats the complementary, compressive case in which (19) fails.

### 3.4 Explicit volumetric growth functions

The bound (15) exposes *two independent channels* through which the tangent can diverge, and they are kinematically decoupled:

$$\|\mathbb{L}(\mathbf{F})\| \leq \mu_e + \underbrace{\Theta(\det \mathbf{F})}_{\substack{\text{dilation channel} \\ J \rightarrow \infty}} \cdot \underbrace{\|\mathbf{F}^{-1}\|^2}_{\substack{\text{inverse channel} \\ \lambda_{\min} \rightarrow 0}}. \quad (20)$$

The *inverse channel* is the one usually blamed for trouble: it requires  $\lambda_{\min} \rightarrow 0$ , i.e. compression to zero volume, and is the subject of Scenario B. The *dilation channel*,  $\Theta(\det \mathbf{F}) \rightarrow \infty$  as  $J \rightarrow \infty$ , is activated by *expansion* and is precisely what a tensile corner switches on, since  $\lambda_{\max} \sim r^{\alpha-1} \rightarrow \infty$  forces  $J \rightarrow \infty$  even with  $\lambda_{\min}$  bounded below by (NC). Whether this matters is therefore not a property of the corner or of the sign of the loading, but of the *growth rate of  $\Theta$* —a property of the volumetric penalty  $f$  alone. We now evaluate it for the two penalties (8); the contrast shows that “tension is benign” is a model-dependent statement.

**Logarithmic penalty.** For  $f_{\log}(J) = -\mu_e \log J + \lambda_e \log^2 J$ ,

$$f'(J)J = -\mu_e + 2\lambda_e \log J, \quad f''(J)J^2 = \mu_e + 2\lambda_e - 2\lambda_e \log J, \quad (21)$$

so that the combination collapses to a constant,

$$f''(J)J^2 + f'(J)J = 2\lambda_e, \quad (22)$$

and therefore

$$\Theta_{\log}(J) = 2\lambda_e + |2\lambda_e \log J - \mu_e| = \mathcal{O}(\log J) \quad \text{as } J \rightarrow \infty. \quad (23)$$

The volumetric stiffening grows only *logarithmically* in  $J$ .

**Quadratic penalty.** For  $f_{\text{quad}}(J) = \frac{\kappa}{2}(J-1)^2$ ,

$$f'(J)J = \kappa J(J-1), \quad f''(J)J^2 = \kappa J^2, \quad f''(J)J^2 + f'(J)J = \kappa J(2J-1), \quad (24)$$

so

$$\Theta_{\text{quad}}(J) = \kappa J |2J-1| + \kappa J |J-1| = \mathcal{O}(\kappa J^2) \quad \text{as } J \rightarrow \infty. \quad (25)$$

The volumetric stiffening grows *quadratically* in  $J$ .

**Consequence at the corner.** We combine (15) with (NC) ( $\|\mathbf{F}^{-1}\|^2 \leq c^{-2}$ ) and the tensile kinematics. In the non-compressive regime exactly one stretch diverges:  $\lambda_{\max} \sim |K| r^{\alpha-1}$  while the remaining  $d-1$  stretches stay bounded (in particular  $\lambda_{\min}$  bounded below), so  $J = \det \mathbf{F} = \prod_i \lambda_i \sim r^{\alpha-1}$ . (If instead  $\nabla \mathbf{u}$  had two diverging eigenvalues, the corresponding subdeterminant would give  $J \sim r^{2(\alpha-1)}$ ; we work in the one-large-stretch regime characteristic of a Dirichlet–Neumann opening, where  $J \sim r^{\alpha-1}$ .) The tangent norm near the corner then obeys

$$\|\mathbb{L}(\mathbf{x})\| \leq \mu_e + c^{-2} \Theta(\det \mathbf{F}(\mathbf{x})) \sim \begin{cases} \mu_e + \mathcal{O}(\log \frac{1}{r}), & f = f_{\log}, \\ \mathcal{O}(r^{-2(1-\alpha)}), & f = f_{\text{quad}}. \end{cases} \quad (26)$$

Thus the logarithmic penalty keeps the tangent bounded up to a logarithm, whereas the quadratic penalty produces a genuine *algebraic* blow-up of the tangent *even in pure tension*. This is the dilation channel of (20) acting alone, with the inverse channel inert ( $\|\mathbf{F}^{-1}\|$  bounded); it is algebraically singular of a severity comparable to—though mechanistically distinct from—the compressive blow-up of Scenario B, which instead runs through  $\|\mathbf{F}^{-1}\|^2$ . The often-repeated statement that “tension is harmless for neo-Hookean” is therefore an artefact of the logarithmic volumetric model and is false for the quadratic one.

**Physical reading: the large-dilation response.** The dichotomy is, at bottom, a statement about how the bulk energy behaves under unbounded *expansion*, the loading a tensile corner imposes:

| volumetric model          | energy as $J \rightarrow \infty$ | tangent $\Theta(J)$   |
|---------------------------|----------------------------------|-----------------------|
| $\lambda_e \log^2 J$      | $\sim \log^2 J$ (soft)           | $\mathcal{O}(\log J)$ |
| $\frac{\kappa}{2}(J-1)^2$ | $\sim J^2$ (stiff)               | $\mathcal{O}(J^2)$    |

A penalty that is polynomial in  $J$  makes the material resist further dilation ever more stiffly, and the linearised modulus faithfully reports that stiffness as a blow-up; the logarithmic penalty is gentle about expansion and does not. Both penalties are designed to diverge correctly as  $J \rightarrow 0$  (resisting collapse, the inverse channel); their large- $J$  behaviour, which the dilation channel probes, was never physically constrained, and that latitude is exactly where they part. The exact cancellation (22) is the analytic signature of this softness. We make the benign case precise in Theorem 1 and flag the algebraic case in Remark 3.

### 3.5 Scenario A: the tensile (non-compressive) singularity

We work under the mild-growth condition isolated above.

**Assumption 5** (Mild volumetric growth). The growth function satisfies  $\Theta(J) = \mathcal{O}((\log J)^p)$  as  $J \rightarrow \infty$  for some  $p \geq 0$ .

By (23),  $f_{\log}$  satisfies Assumption 5 with  $p = 1$ ; by (25),  $f_{\text{quad}}$  does not.

**Lemma 2** (Subordinate tangent). *Under Assumptions 1, 2, 5 and (NC), for every  $\varepsilon > 0$  there is  $C_\varepsilon$  with*

$$\|\mathbb{L}(\mathbf{x})\| \leq C_\varepsilon r^{-\varepsilon} \quad \text{for a.e. } \mathbf{x} \in B_\rho. \quad (27)$$

*In particular  $\mathbb{L} \in L^q(B_\rho)$  for every  $q < \infty$ , and  $\|\mathbb{L}\|$  is dominated by any negative power of  $r$ .*

*Proof.* By (15) and (NC),  $\|\mathbb{L}\| \leq \mu_e + c^{-2}\Theta(\det \mathbf{F})$ . Assumption 5 and  $\det \mathbf{F} = \mathcal{O}(r^{\alpha-1})$  give  $\Theta(\det \mathbf{F}) = \mathcal{O}((\log \frac{1}{r})^p)$ , and  $(\log \frac{1}{r})^p \leq C_\varepsilon r^{-\varepsilon}$  on  $(0, \rho]$  for every  $\varepsilon > 0$ .  $\square$   $\square$

**Lemma 3** (Uniform ellipticity). *Under Assumptions 1, 3 and (NC), and assuming the volumetric prefactors in (18) remain controlled (automatic for  $f_{\log}$ , where  $f''J^2 + f'J = 2\lambda_e$  is constant and  $|f'J| = \mathcal{O}(\log J)$  is subordinate), the tangent satisfies the Legendre–Hadamard condition on  $B_\rho$  with a slowly varying constant: there are  $\gamma_0 > 0$  and  $\varepsilon > 0$  arbitrarily small with*

$$L_{iJkL}(\mathbf{x}) m_i N_J m_k N_L \geq (\gamma_0 - C_\varepsilon r^\varepsilon) |\mathbf{m}|^2 |\mathbf{N}|^2 \geq \frac{1}{2} \gamma_0 |\mathbf{m}|^2 |\mathbf{N}|^2 \quad (28)$$

for a.e.  $\mathbf{x} \in B_\rho$  after shrinking  $\rho$ , for all  $\mathbf{m}, \mathbf{N} \in \mathbb{R}^2$ .

*Proof.* From (18),  $L_{iJkL} m_i N_J m_k N_L \geq \mu_e |\mathbf{m}|^2 |\mathbf{N}|^2 - |f''J^2 + f'J| |\mathbf{a}|^2 |\mathbf{m}|^2 - |f'J| |\mathbf{a}| |\mathbf{b}| |\mathbf{m}| |\mathbf{N}|$ , where the volumetric correction is bounded using  $|\mathbf{a}| \leq c^{-1} |\mathbf{N}|$ ,  $|\mathbf{b}| \leq c^{-1} |\mathbf{m}|$  by  $c^{-2}(|f''J^2 + f'J| + |f'J|) |\mathbf{m}|^2 |\mathbf{N}|^2 = c^{-2} \Theta(J) |\mathbf{m}|^2 |\mathbf{N}|^2$ . For  $f_{\log}$ ,  $\Theta(J) = \mathcal{O}(\log \frac{1}{r}) \leq C_\varepsilon r^\varepsilon$  on  $(0, \rho]$ ; the volumetric correction is thus  $o(1)$  as  $r \rightarrow 0$ , and after shrinking  $\rho$  it is below  $\frac{1}{2} \mu_e$ , giving (28) with  $\gamma_0 = \mu_e$ .  $\square$   $\square$

For the weighted well-posedness we recall two standard inequalities in the Kondratiev space  $V_\beta^{1,2}(B_\rho)$ —the closure of smooth functions vanishing on  $\Gamma_D \cap \partial B_\rho$  under  $\|\mathbf{v}\|_{V_\beta^{1,2}}^2 = \int_{B_\rho} (r^{2\beta} |\nabla \mathbf{v}|^2 + r^{2\beta-2} |\mathbf{v}|^2) dV$ .

**Lemma 4** (Weighted Hardy and Korn inequalities [6, 8]). *Let  $\beta \in \mathbb{R}$  avoid the (discrete) set of exceptional exponents of the corner. There are constants  $C_H, C_K$  depending only on  $\beta$  and the opening angle such that, for all  $\mathbf{v} \in V_\beta^{1,2}(B_\rho)$  vanishing on  $\Gamma_D \cap \partial B_\rho$ ,*

$$\int_{B_\rho} r^{2\beta-2} |\mathbf{v}|^2 dV \leq C_H \int_{B_\rho} r^{2\beta} |\nabla \mathbf{v}|^2 dV, \quad (29)$$

$$\int_{B_\rho} r^{2\beta} |\nabla \mathbf{v}|^2 dV \leq C_K \int_{B_\rho} r^{2\beta} |\varepsilon(\mathbf{v})|^2 dV, \quad \varepsilon(\mathbf{v}) = \frac{1}{2} (\nabla \mathbf{v} + \nabla \mathbf{v}^T). \quad (30)$$

Let  $\beta_\star$  be the weight exponent for which the linear elasticity corner problem at this Dirichlet–Neumann junction is Fredholm (equivalently  $\beta_\star$  avoids the singular exponents  $\alpha$ ). We may write the Newton-step bilinear form as

$$a(\Delta \mathbf{u}, \delta \mathbf{u}) := \int_{B_\rho} \nabla \delta \mathbf{u} : \mathbb{L}(\bar{\mathbf{F}}) : \nabla \Delta \mathbf{u} dV. \quad (31)$$

**Theorem 1** (Well-posedness of the linearised step, tensile corner). *Under Assumptions 1–5 and the non-compressive hypothesis (NC), the form (31) is bounded and coercive on  $V_{\beta_\star}^{1,2}(B_\rho)$  modulo a compact perturbation: there exist constants  $\Lambda < \infty$ ,  $\gamma_1 > 0$  and a compact form  $k(\cdot, \cdot)$  such that, for all  $\mathbf{u}, \mathbf{v} \in V_{\beta_\star}^{1,2}(B_\rho)$ ,*

$$|a(\mathbf{u}, \mathbf{v})| \leq \Lambda \|\mathbf{u}\|_{V_{\beta_\star}^{1,2}} \|\mathbf{v}\|_{V_{\beta_\star}^{1,2}}, \quad (32)$$

$$a(\mathbf{v}, \mathbf{v}) \geq \gamma_1 \|\mathbf{v}\|_{V_{\beta_\star}^{1,2}}^2 - k(\mathbf{v}, \mathbf{v}). \quad (33)$$

Consequently the linearised corner operator is Fredholm of index zero in  $V_{\beta_\star}^{1,2}$ ; for  $\beta_\star$  avoiding the corner spectrum it is an isomorphism, its solutions admit the asymptotic expansion (13), and the governing singular exponents coincide with those of the linearised elasticity problem.



*Proof. Boundedness.* By Cauchy–Schwarz and Lemma 2, for any  $\varepsilon \in (0, 1)$ ,

$$\begin{aligned} |a(\mathbf{u}, \mathbf{v})| &\leq \int_{B_\rho} \|\mathbb{L}\| |\nabla \mathbf{u}| |\nabla \mathbf{v}| dV \leq C_\varepsilon \int_{B_\rho} r^{-\varepsilon} |\nabla \mathbf{u}| |\nabla \mathbf{v}| dV \\ &\leq C_\varepsilon \left( \int_{B_\rho} r^{2\beta_\star} |\nabla \mathbf{u}|^2 \right)^{1/2} \left( \int_{B_\rho} r^{-2\varepsilon-2\beta_\star} |\nabla \mathbf{v}|^2 \right)^{1/2}. \end{aligned}$$

Since  $\varepsilon$  is arbitrary and  $\beta_\star$  is interior to an interval of admissible weights, the loss  $r^{-2\varepsilon}$  can be absorbed by a weight chosen within that interval; this gives (32) with  $\Lambda = \Lambda(\varepsilon)$ . The key point is that the mild (logarithmic) growth of  $\|\mathbb{L}\|$  is subordinate to every algebraic weight, so it does not change the admissible weight interval.

*Gårding inequality.* By Lemma 3 the integrand symbol  $\mathbb{L}(\mathbf{x})$  is uniformly strongly elliptic on  $B_\rho$  with constant  $\geq \frac{1}{2}\gamma_0$ . Freezing the (measurable, uniformly elliptic) coefficients on a dyadic partition  $\{r \sim 2^{-j}\}$  and applying the constant-coefficient Gårding inequality on each annulus—valid because the principal symbol is positive definite on the Legendre–Hadamard cone—yields, after summation against the weight  $r^{2\beta_\star}$ ,

$$a(\mathbf{v}, \mathbf{v}) \geq \frac{1}{2}\gamma_0 c_K^{-1} \int_{B_\rho} r^{2\beta_\star} |\nabla \mathbf{v}|^2 dV - C \int_{B_\rho} r^{2\beta_\star} |\mathbf{v}|^2 dV,$$

where the Korn inequality (30) converts ellipticity of the symmetric part into control of the full gradient. The remainder term has a strictly higher power of  $r$  than the Hardy weight  $r^{2\beta_\star-2}$ , hence by (29) it is relatively compact with respect to  $\|\cdot\|_{V_{\beta_\star}^{1,2}}$ ; this is the compact form  $k$ , and we obtain (33) with  $\gamma_1 = \frac{1}{2}\gamma_0 c_K^{-1}$ .

*Fredholm property and exponents.* Estimates (32)–(33) place the operator in the Kondratiev–Maz’ya framework [6]: it is Fredholm of index zero in  $V_{\beta_\star}^{1,2}$  whenever  $\beta_\star$  avoids the spectrum of the corner operator pencil. Because the nonlinear correction to the principal symbol is subordinate (it is  $o(r^{-\varepsilon})$  for every  $\varepsilon$ , by Lemma 2), the pencil coincides with that of linearised elasticity; its eigenvalues are exactly the exponents  $\alpha$  of (13), which are therefore unchanged by the nonlinearity.  $\square$   $\square$

*Remark 2* (Stress is singular, operator is not). The first Piola–Kirchhoff stress still blows up,  $\mathbf{P} \sim \mu_e \mathbf{F} \rightarrow \infty$  as  $r \rightarrow 0$ , so the tensile corner is a genuine *stress* singularity. Theorem 1 asserts only that this singularity lives entirely in the *right-hand side* (the residual) of the Newton step, not in its operator: under a mild volumetric penalty the nonlinear problem inherits the regularity theory of the linear one, standard graded meshes restore optimal convergence, and no structural instability occurs. This is the precise sense in which a non-compressive corner “passes like linear elasticity.”

*Remark 3* (The quadratic penalty does *not* pass like linear elasticity). For  $f_{\text{quad}}$ , Assumption 5 fails: by (25) and (26) the tangent grows algebraically,  $\|\mathbb{L}\| = \mathcal{O}(r^{-2(1-\alpha)})$ , *even in pure tension*. The coefficients of (31) are then genuinely singular, of severity comparable to the compressive Scenario B, the boundedness step above fails for  $\beta_\star$ , and the admissible weight interval—hence the effective corner exponents seen by the nonlinear operator—shifts relative to linear elasticity. A sharp analysis of this case (the correct weight, and whether coercivity survives) is governed by the volumetric blow-up rather than by the deviatoric singularity and is left open here.

We emphasise that this serves as a *cautionary comparator* rather than a model we advocate: as written,  $f'_{\text{quad}}(1) = 0 \neq -\mu_e$ , so  $\frac{\kappa}{2}(J-1)^2$  does not combine with the deviatoric term  $\frac{\mu_e}{2}(\text{tr}(\mathbf{F}^T \mathbf{F}) - d)$  to leave the reference configuration stress-free, and in practice it is used together with an isochoric (deviatoric/volumetric) split of  $\mathbf{F}$ . The point is that a standard-looking polynomial penalty already forfeits the benign tensile behaviour enjoyed by the logarithmic model (8) actually used in [9].

*Remark 4* (Self-consistency of (NC)). Hypothesis (NC) is compatible with the singular kinematics (14): an angular profile  $\Phi$  corresponding to opening/shear produces  $\nabla \mathbf{u}$  whose dominant eigenvalue is positive (radial extension), so  $\lambda_{\max} \rightarrow \infty$  while  $\lambda_{\min} = \det \mathbf{F} / \lambda_{\max}$  need not approach zero. The complementary profile, in which the dominant eigenvalue of  $\nabla \mathbf{u}$  is negative, drives  $\lambda_{\min} \rightarrow 0$  and violates (NC); that is exactly Scenario B below.

### 3.6 Scenario B: the compressive singularity (the pathological case)

Now assume the far-field loading pushes the material *into* the corner ( $K < 0$ , radial compression).

**Kinematic consequence.** The naive linear prediction gives  $\nabla \mathbf{u}$  a negative dominant eigenvalue scaling as  $-|K|r^{\alpha-1}$ . Because  $\mathbf{F} = \mathbf{I} + \nabla \mathbf{u}$ , as  $r \rightarrow 0$  (or as the pseudo-time load factor  $t$  increases) there is a critical threshold where the local contraction annihilates the material volume:

$$\det \mathbf{F} \rightarrow 0 \implies J \rightarrow 0, \quad \lambda_{\min} \rightarrow 0. \quad (34)$$

**Operator blow-up (and its model dependence).** Here it is the *inverse channel* of (20) that fires:  $\|\mathbf{F}^{-1}\| = \lambda_{\min}^{-1} \rightarrow \infty$ . The dilation channel  $\Theta(J)$  is now probed at  $J \rightarrow 0^+$  rather than  $J \rightarrow \infty$ , and—exactly as in tension, but with the roles reversed—its behaviour discriminates between the two penalties. Writing  $J = \prod_i \lambda_i \sim \lambda_{\min}$  near the corner (one collapsing stretch, the remaining  $d-1$  bounded, cf. §3.4),

$$\Theta_{\log}(J) \sim 2\lambda_e |\log J| \rightarrow \infty, \quad \Theta_{\text{quad}}(J) \sim 2\kappa J \rightarrow 0 \quad (J \rightarrow 0^+), \quad (35)$$

so that the tangent norm  $\|\mathbb{L}\| \leq \mu_e + \Theta(J)\|\mathbf{F}^{-1}\|^2$  behaves as

$$\|\mathbb{L}(\mathbf{x})\| \sim \begin{cases} \mathcal{O}(\lambda_{\min}^{-2} \log \frac{1}{\lambda_{\min}}), & f = f_{\log}, \\ \mathcal{O}(\lambda_{\min}^{-1}), & f = f_{\text{quad}}. \end{cases} \quad (36)$$

Thus the logarithmic penalty—benign under tension—produces here the *stronger* blow-up,  $\lambda_{\min}^{-2}$  amplified by a logarithm, whereas the quadratic penalty gives the milder rate  $\lambda_{\min}^{-1}$ . This reversal is not a redemption of the quadratic model:  $f_{\text{quad}}(J) \rightarrow \frac{\kappa}{2} < \infty$  as  $J \rightarrow 0^+$ , so it provides *no energy barrier against volume collapse* and violates Assumption 1; the milder tangent growth merely reflects this missing barrier (the material is free to interpenetrate at finite energy), which is a worse pathology, not a better one. The logarithmic penalty's stronger blow-up is, conversely, the analytic price of a genuine barrier  $f_{\log}(J) \rightarrow +\infty$ —and it is precisely that barrier which forces, and gives physical meaning to, the bifurcation rescue of Section 4.

**Ellipticity is *not* lost — the instability is structural.** It is tempting to attribute the breakdown to a loss of strong ellipticity (an indefinite acoustic tensor). For this neo-Hookean model that is *false*, and it is worth stating the fact precisely, as it shapes the rest of the analysis.

**Lemma 5** (Compressive strong ellipticity of the neo-Hookean tangent). *For the energy (7), the rank-one form of the tangent is exactly (18),  $L_{iJkL}m_iN_Jm_kN_L = \mu_e|\mathbf{m}|^2|\mathbf{N}|^2 + f''(J)J^2(\mathbf{m} \cdot \mathbf{F}^{-T}\mathbf{N})^2$ . Wherever  $f''(J) \geq 0$  — in particular throughout compression,  $J \leq 1$ , for both penalties (8) — the tangent is strongly elliptic with constant  $\geq \mu_e > 0$ , uniformly in  $\mathbf{F}$ , and the acoustic tensor  $Q_{ik}(\mathbf{N}) = L_{iJkL}N_JN_L$  is positive definite and never degenerates under compression.*

*Proof.* The identity is (18) (Lemma 1). As  $J \rightarrow 0^+$ ,  $f''(J)J^2 \rightarrow +\infty$  for  $f_{\log}$  (cf. (52)) and  $f''(J)J^2 = \kappa J^2 \geq 0$  for  $f_{\text{quad}}$ ; in both cases the second term only *adds* to the constant  $\mu_e$ .  $\square$   $\square$

Thus the compressive corner is *not* a material instability: the volumetric stiffening that blows up  $\|\mathbb{L}\|$  is exactly the term  $f''(J)J^2$  in (18), and it is positive — it strengthens ellipticity while destroying conditioning. What fails is the *boundedness* of the operator, not its positivity.

**Conclusion for Scenario B.** The assumption that the deformation smoothly follows the fundamental path into the corner therefore produces, not a loss of ellipticity, but two distinct effects: (i) a blow-up of the tangent *norm*,  $\|\mathbb{L}\| \rightarrow \infty$  (eq. (36)), which renders the Newton step catastrophically ill-conditioned even though it remains elliptic; and (ii) for a barrier penalty  $f_{\log}$ , an unphysical infinite accumulation of energy ( $f(J) \rightarrow \infty$  as  $J \rightarrow 0$ ), signalling that the smooth fundamental path is not the physical solution. The genuine mechanism that intervenes is a *structural* bifurcation of the compressed state — the corner analogue of *creasing*, an instability of a material that remains everywhere strongly elliptic (Lemma 5), driven by the geometry and boundary conditions rather than by a change of type of the PDE. To capture it we introduce in Section 4 an imperfect perturbation that regularises the bifurcation and bounds the singularity; we work from here on with the barrier model  $f_{\log}$ , for which collapse is genuinely resisted.

## 4 Imperfect bifurcation analysis and asymptotic bounds

The previous section showed that the perfect compressive path drives  $\lambda_{\min} \rightarrow 0$  and blows up the tangent norm while the material remains strongly elliptic (Lemma 5), so the breakdown is a structural instability rather than a loss of type. The premise of this section is that this scenario is an idealisation: a real system always carries small disturbances—geometric imperfections, load eccentricities, material inhomogeneity—and these select a nearby *regular* solution branch on which the collapse never quite happens. We model the disturbance by a single scalar amplitude  $\epsilon$ , carry out an imperfect-bifurcation analysis, and quantify how the tangent blows up as  $\epsilon \rightarrow 0$ . The punchline is that the blow-up is confined to a vanishingly small neighbourhood of the corner and proceeds at a controlled algebraic rate, so that for every  $\epsilon > 0$  the problem is regular.

### 4.1 The regularised problem and the fundamental path

We localise to the corner neighbourhood  $B_\rho$  of §3, regard the loading as transmitted through data on the outer arc  $\partial B_\rho \setminus (\Gamma_D \cup \Gamma_N)$ , and introduce a pseudo-time load parameter  $t \in [0, 1]$  that monotonically scales the compressive part of that data. Following the answer to the question posed in Scenario B, we add a small, fixed volumetric body force  $\epsilon \mathbf{p}(\mathbf{x})$  with  $\|\mathbf{p}\|_{L^2(B_\rho)} = 1$  and  $\epsilon \ll 1$  as the imperfection (we write the field as  $\mathbf{p}$  to avoid collision with the penalty  $f(J)$ ). The regularised equilibrium is  $G(\mathbf{u}, t, \epsilon) = 0$ ,

$$\langle G(\mathbf{u}, t, \epsilon), \mathbf{v} \rangle = \int_{B_\rho} \mathbf{P}(\mathbf{F}) : \nabla \mathbf{v} \, dV - \epsilon \int_{B_\rho} \mathbf{p} \cdot \mathbf{v} \, dV - \ell_t(\mathbf{v}) = 0 \quad \forall \mathbf{v} \in V, \quad (37)$$

with  $\mathbf{P} = \partial \Psi / \partial \mathbf{F}$ ,  $V$  the admissible test space, and  $\ell_t$  the (linear) load functional at pseudo-time  $t$ . Let  $\mathbf{u}_0(t)$  denote the *fundamental path*, i.e. the branch of (37) with  $\epsilon = 0$  that continues the low-load compressive solution. We use the barrier penalty  $f_{\log}$  throughout, so the collapse  $J \rightarrow 0$  is genuinely resisted (Assumption 1); along  $\mathbf{u}_0(t)$  the corner stretch  $\lambda_c(t) := \lambda_{\min}(\mathbf{F}_0(t))$  decreases as  $t \uparrow 1$  but, on the bifurcating branch we construct, never reaches zero.

*Remark 5* (The instability is structural, not material). By Lemma 5 the tangent stays strongly elliptic throughout compression, so the bifurcation here is a *structural* one (a buckling of the boundary-value problem), not a material change of type. We therefore work with the global form on the bounded domain  $B_\rho$ : the operator  $\mathcal{L}(t) = \partial_{\mathbf{u}} G(\mathbf{u}_0(t), t, 0)$  is uniformly elliptic with compact resolvent, hence has discrete spectrum, and we assume (Assumption 6) that a simple eigenvalue first crosses zero at  $t_c$ . What makes the problem delicate is not a loss of ellipticity but the *norm* blow-up  $\|\mathbb{L}\| \rightarrow \infty$  at the corner (eq. (36)): the coercivity constant stays positive while the boundedness constant diverges, so the crossing eigenvalue  $\mu_1(t)$  and the conditioning are controlled by different mechanisms.

## 4.2 What is provable, and what must be assumed

It is worth separating the genuinely provable facts about the critical point from the irreducible hypotheses. Let  $Q_t(\mathbf{v}) := \int_{B_\rho} \nabla \mathbf{v} : \mathbb{L}(\mathbf{F}_0(t)) : \nabla \mathbf{v} dV$  be the second-variation form on the fundamental path and  $\mu_1(t) := \inf_{\|\mathbf{v}\|_V=1} Q_t(\mathbf{v})$ .

**The critical load exists (conditionally provable).** A clarification is essential here. By Lemma 5 the neo-Hookean material remains strongly elliptic throughout compression, so  $\mu_1(t)$  *cannot* be driven negative by a rank-one (material) instability: any crossing must be a *structural* loss of positivity of the global form  $Q_t$ , i.e. a buckling of the boundary-value problem in which the destabilising test field is not rank-one. This is the creasing mechanism.

**Lemma 6** (Existence of the critical load). *Assume (i) coercivity at zero load,  $\mu_1(0) > 0$ , and (ii) a structural instability of the compressed state: there exist  $t_* \in (0, 1)$  and  $\mathbf{v}_* \in V$  with  $Q_{t_*}(\mathbf{v}_*) < 0$ . Then  $\mu_1$  is continuous on  $[0, t_*]$  and there is a first  $t_c \in (0, t_*]$  with  $\mu_1(t_c) = 0$  and  $\mu_1(t) > 0$  for  $t < t_c$ .*

*Proof.* At  $t = 0$ ,  $\mathbf{F}_0 = \mathbf{I}$  and  $\mathbb{L} = \mu_e \mathbb{I} + (\text{volumetric at } J=1)$  is positive definite, so with Korn's inequality  $\mu_1(0) > 0$ . Hypothesis (ii) gives  $\mu_1(t_*) \leq Q_{t_*}(\mathbf{v}_*) / \|\mathbf{v}_*\|_V^2 < 0$ . As  $t \mapsto \mathbb{L}(\mathbf{F}_0(t))$  is continuous, so is  $\mu_1$ , and the intermediate value theorem gives the first zero  $t_c$ .  $\square$   $\square$

Hypothesis (i) is provable for neo-Hookean (positive-definiteness at  $\mathbf{F} = \mathbf{I}$  plus Korn). Hypothesis (ii) — the existence of a structurally destabilising field — is the genuine content and is *not* elementary: it is the corner counterpart of the Biot/creasing instability [1, 3] and we do not establish it here.

### The critical mode (assumed).

**Assumption 6** (Critical mode). At  $t_c$  the infimum  $\mu_1(t_c) = 0$  is attained at a mode  $\phi \in V$ ,  $\|\phi\|_V = 1$ , spanning a one-dimensional, isolated kernel of  $\mathcal{L}_c$  (the rest of the spectrum bounded away from 0), with  $\phi$  localised near the corner.

This is the irreducible hypothesis: *attainment* (the minimising sequence does not escape into the corner as essential spectrum) and *simplicity*. Simplicity is generic and, for a symmetric corner, can be argued from a reflection symmetry about the bisector; discreteness is the borderline-elliptic difficulty already flagged in Remark 5. We do not prove it.

**The crossing is transversal (provable under monotone loading).**

**Lemma 7** (Transversality). *Suppose the loading is monotonically destabilising at  $t_c$ ,*

$$\langle \phi, \dot{\mathbb{L}}(t_c)\phi \rangle := \int_{B_\rho} \nabla \phi : \partial_t \mathbb{L}(\mathbf{F}_0(t_c)) : \nabla \phi \, dV < 0. \quad (\text{M})$$

*Then, under Assumption 6,  $\mu_1$  is differentiable at  $t_c$  with  $a := -\mu_1'(t_c) = -\langle \phi, \dot{\mathbb{L}}(t_c)\phi \rangle > 0$ .*

*Proof.* For a simple eigenvalue, first-order perturbation theory gives  $\mu_1'(t_c) = \langle \phi, \dot{\mathbb{L}}(t_c)\phi \rangle$ ; hypothesis (M) fixes the sign.  $\square$   $\square$

Condition (M) is the precise meaning of “more compression  $\Rightarrow$  less stiffness in the buckling direction” and holds for any monotone compressive load path; it is an input on the *loading*, not on the solution.

**The imperfection is active (provable, generic).**

**Lemma 8** (Generic activation). *For every  $\mathbf{p}$  outside the closed codimension-one subspace  $\phi^\perp$ , the projection  $\eta := \int_{B_\rho} \mathbf{p} \cdot \phi \, dV \neq 0$ . Hence  $\eta \neq 0$  for generic imperfection.*

**The bifurcation is subcritical (assumed).** The one ingredient genuinely beyond reach of soft arguments is the *sign* of the nonlinearity. Guided by the creasing analogy [1, 3]—surface instabilities of compressed elastomers are well known to be subcritical—we assume it here rather than pretend to derive it.

**Assumption 7** (Subcritical, non-degenerate reduced potential). The reduced bifurcation equation is subcritical with quintic saturation,

$$b := \frac{1}{6} \langle \phi, \partial_{\mathbf{u}}^3 G[\phi, \phi, \phi] \rangle < 0, \quad d := \frac{1}{120} \langle \phi, \partial_{\mathbf{u}}^5 G[\phi, \dots, \phi] \rangle > 0, \quad (38)$$

and the quadratic coefficient vanishes by the reflection symmetry  $w \mapsto -w$  of the perfect compressive state about the corner bisector.

### 4.3 Lyapunov–Schmidt reduction

**Proposition 1** (Normal form). *Under Assumption 6, Lemma 7 and Assumption 7, there is a neighbourhood of  $(\mathbf{u}_0(t_c), t_c, 0)$  in which the solutions of (37) are exactly*

$$\mathbf{u}(t, \epsilon) = \mathbf{u}_0(t) + w \phi + \psi(w, t, \epsilon), \quad \psi \in V^\perp := \{\phi\}^\perp, \quad (39)$$

*where  $\psi$  is smooth with  $\psi = \mathcal{O}(w^2 + \epsilon)$ , and the amplitude  $w$  solves the scalar imperfect bifurcation equation, retained to quintic order because the cubic is destabilising ( $b < 0$ ),*

$$a(t_c - t)w + bw^3 + dw^5 = \epsilon\eta + \mathcal{O}(\epsilon^2 + \epsilon w + |t_c - t|w^2 + w^6), \quad (40)$$

*with  $a > 0$  (Lemma 7),  $b < 0 < d$  (Assumption 7) and  $\eta$  as in Lemma 8.*

*Proof.* Let  $P$  be the  $V$ -orthogonal projection onto  $\text{span}\{\phi\}$  and  $Q = I - P$ . Write  $\mathbf{u} = \mathbf{u}_0(t) + w\phi + \psi$  with  $\psi \in V^\perp$ . The range equation  $QG = 0$  has invertible linearisation  $Q\mathcal{L}_cQ$  on  $V^\perp$  by Assumption 6, so the implicit function theorem yields a unique smooth  $\psi = \psi(w, t, \epsilon)$  with  $\psi(0, t_c, 0) = 0$  and  $\psi = \mathcal{O}(w^2 + \epsilon + |t_c - t|)$ . Substituting into the kernel equation  $\langle G, \phi \rangle = 0$  and Taylor-expanding  $G$  about  $(\mathbf{u}_0(t_c), t_c, 0)$  gives (40): the linear term  $a(t_c - t)w$  from Lemma 7, the cubic  $bw^3$  and quintic  $dw^5$  from Assumption 7 (the quadratic vanishing by symmetry), and the inhomogeneity  $\epsilon\eta$  from  $-\epsilon\langle \mathbf{p}, \phi \rangle$ . Because  $b < 0$ , the cubic alone does not saturate the amplitude, which is why the quintic  $dw^5$  ( $d > 0$ ) is retained.  $\square$   $\square$

#### 4.4 Subcritical branch structure and imperfection sensitivity

**Perfect system.** Set  $\epsilon = 0$  and  $t < t_c$ . Dividing (40) by  $w$  gives  $a(t_c - t) + bw^2 + dw^4 = 0$ , i.e.

$$w^2 = \frac{-b \pm \sqrt{b^2 - 4ad(t_c - t)}}{2d}, \quad (41)$$

which (with  $a > 0$ ,  $b < 0$ ,  $d > 0$ ) admits real nontrivial branches only for  $t \geq t_L$ , where

$$t_L := t_c - \frac{b^2}{4ad} < t_c, \quad w_L^2 = \frac{-b}{2d} = \mathcal{O}(1). \quad (42)$$

At  $t_L$  the two branches meet in a limit point (fold) at the *finite* amplitude  $w_L$ . The fundamental branch  $w = 0$  is stable for  $t < t_c$  and unstable for  $t > t_c$ ; the upper nontrivial branch is stable. Hence on  $[t_L, t_c)$  the compressed state and a finite-amplitude creased state coexist: the bifurcation is subcritical, with hysteresis and a *snap* at  $t_c$ .

**Imperfect system.**

**Proposition 2** (Imperfection sensitivity, Koiter  $\frac{2}{3}$  law). *For  $\epsilon > 0$  the compressed branch terminates at a limit point that issues from the bifurcation point  $(0, t_c)$  and is depressed below  $t_c$  according to*

$$t_s(\epsilon) = t_c - C |\eta \epsilon|^{2/3} + o(\epsilon^{2/3}), \quad C = \frac{3|b|^{1/3}}{2^{2/3}a} > 0, \quad (43)$$

*so the load at which the compressed branch ceases to exist is reduced by  $\mathcal{O}(\epsilon^{2/3})$ . The turning amplitude there is itself small,  $w_* = \mathcal{O}(\epsilon^{1/3})$ ; the  $\mathcal{O}(1)$  amplitude is that of the creased branch onto which the structure then snaps.*

*Proof.* Write (40) as  $g(w, t) = 0$  with

$$g(w, t) := a(t_c - t)w + bw^3 + dw^5 - \epsilon\eta,$$

the analytic remainder of (40) being subdominant at the scaling found below (we verify this at the end). A limit point in the load is a point of the solution set at which the branch turns, i.e. at which  $t$  is stationary along the curve; differentiating  $g = 0$  implicitly, this is the pair

$$g(w, t) = 0, \quad g_w(w, t) = a(t_c - t) + 3bw^2 + 5dw^4 = 0. \quad (44)$$

We follow the limit point that issues from the bifurcation point  $(0, t_c)$  as  $\epsilon \rightarrow 0$ ; this is the one that caps the slightly deflected compressed branch and hence sets the snap load.

*Turning amplitude.* Using the second equation of (44) to eliminate the load,  $a(t_c - t) = -(3bw^2 + 5dw^4)$ , and substituting into the first gives a single equation for the turning amplitude,

$$-2bw^3 - 4dw^5 = \epsilon\eta. \quad (45)$$

Because  $b < 0$ , the small root of (45) as  $\epsilon \rightarrow 0$  obeys  $-2bw^3 = \epsilon\eta(1 + o(1))$ , the quintic entering only at relative order  $w^2$ ; hence

$$w_*(\epsilon) = \left(\frac{|\eta|\epsilon}{2|b|}\right)^{1/3} (1 + o(1)) = \mathcal{O}(\epsilon^{1/3}), \quad (46)$$

in agreement with the unstable-branch scaling (47) evaluated at the turn. In particular  $w_* \rightarrow 0$ , so at this limit point the cubic dominates the quintic,  $dw_*^4/(bw_*^2) = \mathcal{O}(\epsilon^{2/3})$ .

*Snap load.* Inserting (46) into  $g_w = 0$  and discarding the quintic, a relative  $\mathcal{O}(\epsilon^{2/3})$  correction,

$$a(t_c - t_s(\epsilon)) = -3b w_*^2 (1 + o(1)) = 3|b| \left( \frac{|\eta|\epsilon}{2|b|} \right)^{2/3} (1 + o(1)),$$

which is (43) with  $C = 3|b|^{1/3}/(2^{2/3}a) > 0$ , the classical Koiter  $\frac{2}{3}$  power law [5, 2].

*Remainder.* At the limit point  $w = \mathcal{O}(\epsilon^{1/3})$  and  $t_c - t = \mathcal{O}(\epsilon^{2/3})$ , so the omitted terms  $\mathcal{O}(\epsilon^2 + \epsilon w + |t_c - t| w^2 + w^6)$  of (40) are  $\mathcal{O}(\epsilon^{4/3})$ , a relative  $\mathcal{O}(\epsilon^{1/3})$  against the dominant balance  $a(t_c - t)w \sim b w^3 \sim \epsilon \eta = \mathcal{O}(\epsilon)$ . They perturb  $w_*$  and  $t_s(\epsilon)$  only within the stated  $o(\epsilon^{2/3})$ , leaving the exponent and  $C$  unchanged.  $\square$   $\square$

**The unstable continuation branch.** Separately, near  $t_c$  the small-amplitude (backward-bending, unstable) branch of (40) satisfies, at  $t = t_c$ ,  $b w^3 = \epsilon \eta (1 + o(1))$  with the quintic subdominant, so

$$|w(\epsilon)| = \left( \frac{|\eta|}{|b|} \right)^{1/3} \epsilon^{1/3} (1 + o(1)), \quad \epsilon \rightarrow 0. \quad (47)$$

This is the branch a displacement- or arc-length-controlled solver tracks through the instability, and it is the one that feeds the tangent estimate below; it is *not* the branch a load-controlled physical system follows.

#### 4.5 Kinematic relief: lower bound on $J$

The heart of the regularisation is geometric: the buckling mode trades pure compression for rotation and shear, which lifts the determinant off zero. The reason the relief is second order in  $w$ —and hence why the scaling exponents below are what they are—is that the mode is, to leading order, a *rotation*, and rotations do not change volume at first order.

**Lemma 9** (Determinant relief). *Let  $\mathbf{F} = \mathbf{F}_0 + w \nabla \phi + \mathcal{O}(w^2)$  at the corner, with  $\mathbf{F}_0 = \mathbf{F}_0(t_c)$ , and let the rotation condition*

$$\text{cof } \mathbf{F}_0 : \nabla \phi = J_0 \mathbf{F}_0^{-T} : \nabla \phi = 0 \quad (48)$$

*hold (the critical mode is volume-preserving at first order; this is established from the energy barrier in Lemma 10 below, exactly rather than assumed). Let  $d_\phi := \frac{1}{2} \frac{d^2}{dw^2} \big|_0 \det(\mathbf{F}_0 + w \nabla \phi) > 0$  be the second-order coefficient of the determinant — the second mixed discriminant of  $\mathbf{F}_0$  ( $d - 2$  copies) with  $\nabla \phi$  (two copies), which for  $d = 2$  reduces to  $\det(\nabla \phi)$ . Then*

$$J = \det \mathbf{F} = J_0 + w^2 d_\phi + \mathcal{O}(w^3), \quad (49)$$

*so that, with  $\lambda_c = \lambda_{\min}(\mathbf{F}_0)$  and the remaining  $d - 1$  stretches of  $\mathbf{F}$  bounded,  $\prod_{i \geq 2} \lambda_i(\mathbf{F}) = \mathcal{O}(1)$ ,*

$$\lambda_{\min}(\mathbf{F}) = \frac{J}{\prod_{i \geq 2} \lambda_i(\mathbf{F})} \gtrsim C_1 \lambda_c + C_2 w^2, \quad C_1, C_2 > 0. \quad (50)$$

*Proof.* Expanding the determinant in  $w$ ,  $\det(\mathbf{F}_0 + w \nabla \phi) = \sum_{k=0}^d w^k D_k(\mathbf{F}_0, \nabla \phi)$  with  $D_k$  the  $k$ -th mixed discriminant;  $D_0 = \det \mathbf{F}_0 = J_0$ ,  $D_1 = \text{cof } \mathbf{F}_0 : \nabla \phi$ , and  $D_2 = d_\phi$ . The first-order term  $D_1$  vanishes by (48), leaving (49) (the  $\mathcal{O}(w^3)$  collecting  $D_3, \dots, D_d$ ). Since  $J = \prod_i \lambda_i$  and the  $d - 1$  largest stretches are  $\mathcal{O}(1)$  on the relieved branch (the buckling redirects, but does not unboundedly stretch, the material near the corner), (49) gives (50).  $\square$   $\square$

As the load approaches the instability the perfect corner stretch  $\lambda_c \rightarrow 0$ , so the  $w^2$  term dominates the lower bound. On the *unstable continuation branch* (47) (the one a solver tracks through the instability), combining (50) with  $w \sim \epsilon^{1/3}$ ,

$$\lambda_{\min}(\epsilon) \gtrsim C_2 w(\epsilon)^2 \sim \mathcal{O}(\epsilon^{2/3}), \quad \|\mathbf{F}^{-1}\| = \lambda_{\min}^{-1} \sim \mathcal{O}(\epsilon^{-2/3}). \quad (51)$$

On the stable physical branch the picture is different and milder; we contrast the two after Theorem 2.

The rotation condition (48) is not an extra assumption: it is *forced* by the energy barrier, as we now show. This is the precise content of the informal statement “buckling relieves compression”—the barrier makes the volume-preserving mode the only one of finite energy.

**Lemma 10** (The barrier forces the rotation condition). *Let  $f = f_{\log}$ . Then the bulk modulus diverges at collapse,*

$$f''(J) = \frac{\mu_e + 2\lambda_e - 2\lambda_e \log J}{J^2} \sim \frac{2\lambda_e |\log J|}{J^2} \rightarrow +\infty \quad (J \rightarrow 0^+), \quad (52)$$

and consequently any mode  $\phi$  of bounded second-variation energy  $\langle \phi, \mathcal{L}\phi \rangle \leq C$  on the near-collapse fundamental state  $\mathbf{F}_0$  satisfies

$$\text{cof } \mathbf{F}_0 : \nabla \phi = \mathcal{O}\left(\frac{J_0}{\sqrt{|\log J_0|}}\right) \rightarrow 0 \quad (J_0 \rightarrow 0^+). \quad (53)$$

In particular the critical buckling mode is first-order volume-preserving in the collapse limit, so (48) holds asymptotically along the loaded path, and the residual first-order term in (49) is dominated by the relief term  $w^2 d_\phi$ .

*Proof.* Differentiating  $f'_{\log}(J) = -\mu_e/J + 2\lambda_e \log J/J$  gives (52). The volumetric part of the second variation of  $\Psi$  in the direction  $\phi$  contains the term  $f''(J_0)(\delta J)^2$ , where  $\delta J = \text{cof } \mathbf{F}_0 : \nabla \phi = J_0 \mathbf{F}_0^{-T} : \nabla \phi$  is the first variation of the determinant. Since every term other than this one is bounded on the fundamental state, boundedness of  $\langle \phi, \mathcal{L}\phi \rangle$  forces  $f''(J_0)(\delta J)^2 \leq C'$ , hence  $(\delta J)^2 \leq C'/f''(J_0) \sim C' J_0^2/(2\lambda_e |\log J_0|)$ . Taking square roots yields (53).  $\square$   $\square$

## 4.6 Regularised tangent bound and spatial localisation

**Proposition 3** (Stable physical branch: bounded tangent). *Under load control the structure follows the compressed fundamental branch up to the snap at  $t_s(\epsilon) \leq t_c$  (Proposition 2) and then jumps to the creased branch of amplitude  $w = \mathcal{O}(1)$ . On the compressed branch  $\lambda_{\min} \geq \lambda_c(t) \geq \lambda_c(t_c) > 0$ ; after the snap the  $\mathcal{O}(1)$  amplitude relieves the compression to  $\lambda_{\min} = \mathcal{O}(1)$  via Lemma 9. Hence on the physical branch*

$$\max_{\mathbf{x} \in B_\rho} \|\mathbb{L}(\mathbf{x}, \epsilon)\| = \mathcal{O}(1) \quad \text{uniformly in } \epsilon, \quad (54)$$

and the sole effect of the imperfection is the  $\mathcal{O}(\epsilon^{2/3})$  reduction (43) of the snap load. There is no tangent blow-up on the physical branch.

**Theorem 2** (Unstable continuation branch: regularised blow-up). *Under Assumptions 1, 6, 7 and the conclusions (51) of Lemma 9, with the barrier penalty  $f_{\log}$ , the tangent along the unstable continuation branch (47)—the branch tracked by a displacement- or arc-length-controlled solver—satisfies*

$$\max_{\mathbf{x} \in B_\rho} \|\mathbb{L}(\mathbf{x}, \epsilon)\| \leq C \Theta(J) \|\mathbf{F}^{-1}\|^2 \sim \mathcal{O}(\epsilon^{-4/3} \log \frac{1}{\epsilon}), \quad \epsilon \rightarrow 0. \quad (55)$$



*Proof.* By (11),  $\|\mathbb{L}\| \leq \mu_\epsilon + \Theta(J)\|\mathbf{F}^{-1}\|^2$ . For  $f_{\log}$ ,  $\Theta(J) = \mathcal{O}(\log \frac{1}{J})$  as  $J \rightarrow 0$  by (35), and on the unstable branch  $J \sim \lambda_{\min} \sim \epsilon^{2/3}$  by (51), so  $\Theta(J) = \mathcal{O}(\log \frac{1}{\epsilon})$ ; multiplying by  $\|\mathbf{F}^{-1}\|^2 \sim \epsilon^{-4/3}$  gives (55).  $\square$

The two results together are the precise sense in which “the singularity never occurs in reality”: a load-controlled physical system snaps and keeps a bounded tangent (Proposition 3), and the divergent  $\epsilon^{-4/3}$  rate is felt only by an algorithm that suppresses the snap and tracks the unstable branch through the instability (Theorem 2). The spatial localisation below applies to that branch.

The maximum is, however, achieved only in a shrinking zone, and we can now identify the localisation exponent exactly. The relevant geometry is the *collapse locus* of the perfect field.

**Lemma 11** (Transversal collapse:  $s = 1$ ). *On the perfect (unphysical) continuation of the fundamental path beyond  $t_c$ , let  $\Sigma_c \subset B_\rho$  be the collapse locus on which the smallest stretch vanishes,  $\lambda_{\min} = 0$  (equivalently  $J = 0$ ); it is generically of codimension one. At any point of  $\Sigma_c$  where  $\lambda_{\min}$  is a simple singular value of  $\mathbf{F}$ ,  $\lambda_{\min}$  is a smooth function of  $\mathbf{x}$ , and generically it vanishes transversally,*

$$\lambda_{\min}^{\text{field}}(\mathbf{x}) = \kappa_* \text{dist}(\mathbf{x}, \Sigma_c) + \mathcal{O}(\text{dist}^2), \quad \kappa_* > 0. \quad (56)$$

*Hence the localisation exponent is  $s = 1$ . (A non-generic tangential contact,  $\nabla \lambda_{\min} = 0$  on  $\Sigma_c$ , would give  $s \geq 2$ .)*

*Proof.* Where the smallest singular value is simple it is an analytic function of the entries of  $\mathbf{F}$ , hence smooth in  $\mathbf{x}$ . A smooth scalar field vanishing on  $\Sigma_c$  has, by Taylor expansion,  $\lambda_{\min} = \nabla \lambda_{\min} \cdot \mathbf{n} \text{dist}(\mathbf{x}, \Sigma_c) + \mathcal{O}(\text{dist}^2)$  with  $\mathbf{n}$  the unit normal; transversality  $\nabla \lambda_{\min} \neq 0$  (the generic case, by Sard’s theorem) gives (56) with  $\kappa_* = |\nabla \lambda_{\min}| > 0$ .  $\square$

On the imperfect branch the relief of (51) imposes the floor  $\lambda_{\min}(\mathbf{x}, \epsilon) \gtrsim \max\{\kappa_* \text{dist}(\mathbf{x}, \Sigma_c), c \epsilon^{2/3}\}$ , so the high-stiffness zone is the tube of half-width  $\ell(\epsilon)$  around  $\Sigma_c$  in which the floor is active, with (using  $s = 1$ )

$$\ell(\epsilon) \sim \frac{c}{\kappa_*} \epsilon^{2/3} \xrightarrow{\epsilon \rightarrow 0} 0. \quad (57)$$

**Proposition 4** (Narrowness of the singular zone). *With  $\ell(\epsilon)$  as in (57) and  $s = 1$  (Lemma 11), the set on which the tangent is near its maximum is a tube about  $\Sigma_c$  of area  $\mathcal{O}(|\Sigma_c| \ell(\epsilon)) = \mathcal{O}(\epsilon^{2/3}) \rightarrow 0$ , and the integrated stiffness obeys*

$$\int_{B_\rho} \|\mathbb{L}(\mathbf{x}, \epsilon)\|^q dV \leq C \left(1 + \epsilon^{-4q/3} \ell(\epsilon) \log^q \frac{1}{\epsilon}\right) = C \left(1 + \epsilon^{\frac{2}{3}(1-2q)} \log^q \frac{1}{\epsilon}\right), \quad (58)$$

*which stays bounded as  $\epsilon \rightarrow 0$  for every  $q < \frac{1}{2}$ . Thus, although the pointwise tangent diverges like  $\epsilon^{-4/3}$ , its  $L^q$ -mass over the corner remains finite for all  $q < \frac{1}{2}$ : the blow-up is genuinely localised to a tube of vanishing area, and for every  $\epsilon > 0$  the regularised problem is well posed.*

*Proof.* Split  $B_\rho$  into the tube  $T_\epsilon = \{\text{dist}(\cdot, \Sigma_c) < \ell(\epsilon)\}$  and its complement. On  $T_\epsilon$ ,  $\|\mathbb{L}\| \sim \epsilon^{-4/3} \log \frac{1}{\epsilon}$  (Theorem 2) and  $|T_\epsilon| \sim |\Sigma_c| \ell(\epsilon) \sim \epsilon^{2/3}$ , giving the displayed contribution. Off  $T_\epsilon$ , using the transversal coordinate  $\varrho = \text{dist}(\cdot, \Sigma_c)$  and  $\lambda_{\min} \sim \kappa_* \varrho$ , one has  $\|\mathbb{L}\| \sim \varrho^{-2} \log \frac{1}{\varrho}$ , and  $\int_{\ell(\epsilon)}^\rho \varrho^{-2q} \log^q \frac{1}{\varrho} d\varrho$  converges as  $\ell(\epsilon) \rightarrow 0$  precisely when  $2q < 1$ . Both regimes are controlled iff  $q < \frac{1}{2}$ .  $\square$

## 4.7 Discussion

The structure of the result is the one anticipated physically, and it splits cleanly along the stability of the branch followed. The idealised noiseless problem ( $\epsilon = 0$ ) is singular—the tangent is unbounded and ellipticity is lost—but it is a measure-zero idealisation. Because the corner instability is subcritical (the creasing analogy, Assumption 7), a load-controlled physical system never traverses the singular continuation: it stays on the compressed branch and *snaps* to a finite-amplitude crease at a load reduced by only  $\mathcal{O}(\epsilon^{2/3})$  (Proposition 2), keeping a bounded tangent throughout (Proposition 3). The much-discussed divergence is therefore not seen by the physical system at all; it is the worst case for an algorithm—displacement or arc-length control, numerical continuation—that suppresses the snap and tracks the *unstable* branch through the instability. On that branch the determinant is still lifted off zero (Lemma 9, with the volume-preserving character of the mode *forced* by the energy barrier, Lemma 10), the tangent diverges only algebraically,  $\epsilon^{-4/3}$  up to a logarithm (Theorem 2), and the singular character is confined to a tube of width  $\ell(\epsilon) \sim \epsilon^{2/3} \rightarrow 0$  about the collapse locus, remaining  $L^q$ -integrable for all  $q < \frac{1}{2}$  (Lemma 11, Proposition 4). That algebraic rate is the mirror image—through the inverse channel—of the benign tensile growth of §3.4; the same logarithm, traceable to the barrier  $f_{\log}$ , appears in both. Two practical consequences follow. First, the bifurcation is the corner-localised analogue of the classical *creasing* instability of compressed elastomers [1, 3], now seated at a geometric Dirichlet–Neumann junction rather than on a flat surface. Second, the dichotomy is a concrete warning for computation: a solver that continues through the instability will see the conditioning degrade as  $\epsilon^{-4/3}$  and should either admit the snap or refine inside the tube (57); the load-control reduction (43) tells it when the snap is due. The reduction to scalar imperfect-bifurcation form follows the classical theory [2, 4, 5].

## 5 A concrete realization: the clamped–free corner under compression

The preceding analysis is deliberately abstract: it isolates the structural properties of the energy (Assumptions 1–5) and of the loading (the hypotheses of §4) that the argument actually uses, without committing to a domain. We now fix a concrete material and a concrete family of geometries and ask how much of that hypothesis list can be *discharged* rather than assumed. The material is the compressible neo-Hookean energy (7) with the barrier penalty  $f_{\log}$ . The geometry is a body whose boundary carries a clamped (Dirichlet) face meeting a traction-free (Neumann) face at a right angle, under a compressive load.

The abstract development of §3–§4 is already dimension-independent: it analyses the transverse problem at the codimension-two Dirichlet–Neumann edge  $\Sigma$  in  $\mathbb{R}^d$ . Here we instantiate it on concrete bodies and ask how much of that hypothesis list can be *discharged* rather than assumed.

### 5.1 Geometry and the singular set

We consider two model bodies in  $\Omega \subset \mathbb{R}^d$ .

- **(H) Half-ball.**  $\Omega = \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}| < 1, x_d > 0\}$ . The flat base  $\Gamma_D = \{x_d = 0\} \cap \partial\Omega$  is clamped; the spherical cap  $\Gamma_N = \{|\mathbf{x}| = 1\} \cap \partial\Omega$  is traction-free.
- **(C) Box.**  $\Omega = (-1, 1)^{d-1} \times (0, 1)$ . The base  $\Gamma_D = \{x_d = 0\}$  is clamped; the remaining faces are traction-free.

In both cases the load is a monotone compression transmitted through  $\Gamma_N$  (a pressure or an inward normal displacement on the cap, respectively the lateral/top faces), scaled by the pseudo-time  $t \in [0, 1]$  of §4.

The *singular set*  $\Sigma \subset \partial\Omega$  is the locus where  $\overline{\Gamma_D}$  meets  $\overline{\Gamma_N}$ :

- for (H),  $\Sigma = \{|\mathbf{x}| = 1, x_d = 0\} \cong S^{d-2}$ , a *smooth* codimension-two submanifold; the cap is tangent to the vertical at the equator, so it meets the horizontal base at a right angle;
- for (C),  $\Sigma$  is the union of the bottom edges  $\{x_d = 0, x_i = \pm 1\}$ , again right-angle Dirichlet–Neumann edges, *together with* the bottom vertices, where  $d$  faces meet.

Fix  $\mathbf{x}_0 \in \Sigma$  away from any vertex and split  $\mathbb{R}^d = T_{\mathbf{x}_0}\Sigma \oplus \Pi$ , with  $\Pi$  the two-plane normal to  $\Sigma$ . In the slice  $\Pi$  the boundary reduces to a Dirichlet ray and a Neumann ray meeting at  $90^\circ$ : *exactly* the planar junction of §3.5. We write  $B_\rho$  for a tubular neighbourhood of  $\Sigma$  of radius  $\rho$  and use  $(r, \theta)$  for polar coordinates in  $\Pi$  as before.

*Remark 6* (Why the half-ball is the clean model). For (H) the singular set  $\Sigma$  is a smooth closed manifold with no vertices, so the singularity is a *pure edge* everywhere on  $\Sigma$  and the transverse problem is the planar right-angle junction at every point. For (C) the edge interiors behave identically, but the bottom vertices carry an additional, generally stronger, polyhedral-vertex singularity [7] that is outside the scope of the edge analysis; one either rounds the vertices or restricts  $B_\rho$  to the edge interior. We therefore state the dimension-independent results for (H) and note that they hold verbatim on the edge interiors of (C).

## 5.2 Tier 1: the material assumptions are already discharged

For the neo-Hookean energy (7) with  $f_{\log}$ , the standing assumptions of §3 hold with explicit constants and require no information about the domain: Assumption 1 (regularity, objectivity, and the collapse barrier  $f_{\log}(J) \rightarrow \infty$ ), Assumption 2 (the inverse-channel bound, with  $\Theta = \Theta_{\log}$  of (23)), Assumption 3 and the sharper Lemma 5 (strong ellipticity with constant  $\geq \mu_e$  throughout compression), and Assumption 5 (logarithmic growth,  $p = 1$ ) are all established in Lemma 1, Remark 1 and §3.4. None of these proofs uses  $d = 2$ : the tangent formula (10), the rank-one identity (18), and the growth functions (23) are algebraic identities in  $\text{GL}^+(d)$ . Thus the entire material side of the hypothesis list is closed for the concrete model, in every dimension. What remains are the hypotheses on the *solution and loading*, which is where the geometry enters.

## 5.3 The edge exponent and the scenario it selects

By the edge-singularity calculus [6, 7], the leading singular exponents of the linearised elasticity operator along the smooth edge  $\Sigma$  coincide with those of the transverse two-dimensional pencil in  $\Pi$ , modulated smoothly along  $\Sigma$ . That transverse pencil is the planar right-angle Dirichlet–Neumann corner of (13); its leading exponent  $\alpha \in (0, 1)$  is the smallest root of the corresponding clamped-free Williams secular equation and depends only on the Poisson ratio. Consequently the kinematic blow-up (14),  $\mathbf{F} = \mathbf{I} + \mathcal{O}(r^{\alpha-1})$ , holds with a *computable* exponent, and by Theorem 1 the nonlinear tangent inherits exactly this  $\alpha$  in the benign regime.

Which scenario of §3 is realized is governed by the sign of the dominant eigenvalue of  $\nabla \mathbf{u}$  at the edge, equivalently by the sign of the stress-intensity factor  $K$  in (13). This is a global quantity, set by the far field, and is not determined by the local pencil alone. The compressive load we impose selects  $K < 0$ : the material is pushed into the corner, the dominant eigenvalue of  $\nabla \mathbf{u}$  is negative, and Scenario B (§3.6) is the relevant branch. The quantitative form of “compressive enough” is the

hypothesis (SC) below; it simultaneously selects Scenario B and drives the structural instability of §5.4. The complementary tensile load ( $K > 0$ ) selects Scenario A, where (NC) holds and Theorem 1 applies; for the present geometry under compression that case does not arise near the edge.

#### 5.4 Existence of the critical load via a localized surface instability

The one genuinely non-elementary ingredient of §4 was hypothesis (ii) of Lemma 6: the existence of a structurally destabilising field,  $Q_{t_*}(\mathbf{v}_*) < 0$ , which we there declined to establish and flagged as the corner counterpart of the Biot/creasing instability. For the concrete clamped–free geometry we now *prove* it, by reduction to the surface instability of the compressed half-space.

The free face  $\Gamma_N$  under compression is, locally, a traction-free surface compressed parallel to itself — the exact configuration of Biot’s classical analysis [1]. A homogeneously compressed neo-Hookean half-space loses stability to a surface mode once the surface-parallel stretch falls below a threshold  $\lambda_* \in (0, 1)$  (for the incompressible model  $\lambda_* \approx 0.544$ ; for the compressible energy (7) the threshold is the corresponding root of the surface secular equation, still in  $(0, 1)$  [3]). We package the loading requirement as a single hypothesis.

**Assumption 8** (Surface compression reaches threshold, (SC)). There exist  $t_* \in (0, 1]$  and an interior point  $P \in \Gamma_N \cap B_\rho$ , at positive distance from both  $\Sigma$  and  $\overline{\Gamma_D}$ , such that the fundamental state compresses the free face below the half-space threshold there: the surface-parallel principal stretch of  $\mathbf{F}_0(t_*)$  at  $P$  satisfies  $\lambda_{\parallel}(\mathbf{F}_0(t_*); P) < \lambda_*$ .

(SC) is an input on the *loading*, in the same spirit as the monotonicity condition (M) of Lemma 7: it says only that the prescribed compression is strong enough that the free surface dips below its own buckling threshold somewhere near the edge. Since  $t$  merely normalises the load amplitude, (SC) can always be met by loading hard enough, provided the material does not fail elsewhere first.

**Lemma 12** (Localized surface instability  $\Rightarrow$  structural instability). *Under Assumption 8, hypothesis (ii) of Lemma 6 holds: there is  $\mathbf{v}_* \in V$ , supported in  $B_\rho$  and vanishing on  $\Gamma_D$ , with*

$$Q_{t_*}(\mathbf{v}_*) = \int_{B_\rho} \nabla \mathbf{v}_* : \mathbb{L}(\mathbf{F}_0(t_*)) : \nabla \mathbf{v}_* dV < 0. \quad (59)$$

Consequently, with  $\mu_1(0) > 0$  from Lemma 6(i), there is a first critical load  $t_c \in (0, t_*]$  with  $\mu_1(t_c) = 0$ .

*Proof.* Work in coordinates centred at  $P$ : let  $n \geq 0$  be the depth into the body along the inward normal to  $\Gamma_N$  and  $\mathbf{s} \in \mathbb{R}^{d-1}$  the surface-tangent coordinates, so that near  $P$  the body is the half-space  $\{n > 0\}$  and  $\mathbf{F}_0(t_*)$  is  $C^1$  with value  $\mathbf{F}_P := \mathbf{F}_0(t_*; P)$  whose surface-parallel stretch is  $\lambda_{\parallel} < \lambda_*$ .

*Step 1: the homogeneous unstable mode.* Freeze the coefficients at  $\mathbf{F}_P$  and consider the constant-tensor form  $Q_0(\mathbf{v}) = \int_{\{n>0\}} \nabla \mathbf{v} : \mathbb{L}(\mathbf{F}_P) : \nabla \mathbf{v} dV$  on the half-space, with  $\mathbf{v}$  traction-free at  $n = 0$ . Because  $\lambda_{\parallel} < \lambda_*$ , the half-space is past its surface-instability threshold [1, 3]: for each surface wavevector  $\mathbf{k} \in \mathbb{R}^{d-1}$ ,  $k := |\mathbf{k}|$ , there is a surface mode

$$\mathbf{v}_{\mathbf{k}}(\mathbf{s}, n) = \text{Re} [\hat{\mathbf{v}}(kn) e^{i\mathbf{k} \cdot \mathbf{s}}], \quad |\hat{\mathbf{v}}(\zeta)| \leq C e^{-\beta \zeta} \quad (\beta > 0), \quad (60)$$

decaying exponentially into the body on the depth scale  $1/k$ . Since  $\mathbb{L}(\mathbf{F}_P)$  is constant, the rescaling  $(\mathbf{s}, n) \mapsto k^{-1}(\mathbf{s}, n)$  shows that over a surface window  $W$  of area  $|W|$ ,

$$\int_{\{n>0\}} \mathbf{1}_W \nabla \mathbf{v}_{\mathbf{k}} : \mathbb{L}(\mathbf{F}_P) : \nabla \mathbf{v}_{\mathbf{k}} dV = q(\lambda_{\parallel}) k |W| (1 + o(1)), \quad q(\lambda_{\parallel}) < 0, \quad (61)$$

where  $q$  is the Biot secular form, strictly negative past threshold. Likewise  $\int_{\{n>0\}} \mathbf{1}_W |\nabla \mathbf{v}_k|^2 \sim k |W|$  and  $\int_{\{n>0\}} \mathbf{1}_W |\mathbf{v}_k|^2 \sim k^{-1} |W|$ , the latter two by the same scaling.

*Step 2: cut-off.* Let  $\chi$  be smooth,  $\chi \equiv 1$  on  $B(P, \delta/2)$ ,  $\text{supp } \chi \subset B(P, \delta)$ ,  $|\nabla \chi| \leq C/\delta$ , with  $\delta$  small enough that  $B(P, \delta) \cap \bar{\Omega}$  avoids both  $\Sigma$  and  $\bar{\Gamma}_D$  and lies in  $B_\rho$  (possible since  $P$  is at positive distance from both). Set  $\mathbf{v}_* := \chi \mathbf{v}_k$ . Then  $\mathbf{v}_* \in V$  and vanishes on  $\Gamma_D$  (its support avoids  $\bar{\Gamma}_D$ ); it is unconstrained on the free face, as required.

*Step 3: error control.* Write  $\nabla \mathbf{v}_* = \chi \nabla \mathbf{v}_k + (\nabla \chi) \otimes \mathbf{v}_k$  and expand (59). The effective window is  $W \sim B(P, \delta) \cap \Gamma_N$ , of area  $|W| \sim \delta^{d-1}$ , and the decay confines the mass to depth  $1/k$ . Three errors arise.

- (a) *Coefficient variation:* replacing  $\mathbb{L}(\mathbf{F}_0(\mathbf{x}))$  by  $\mathbb{L}(\mathbf{F}_P)$  costs  $\leq \sup_{B(P, \delta)} \|\mathbb{L}(\mathbf{F}_0) - \mathbb{L}(\mathbf{F}_P)\| \int \chi^2 |\nabla \mathbf{v}_k|^2 \leq C\delta \cdot k\delta^{d-1} = Ck\delta^d$  (by  $C^1$  regularity of  $\mathbf{F}_0$  at  $P$ ).
- (b) *Gradient commutator:* the cross term and  $|\nabla \chi|^2 |\mathbf{v}_k|^2$  term are bounded by  $C(\delta^{-1} \int_W |\mathbf{v}_k| |\nabla \mathbf{v}_k| + \delta^{-2} \int_W |\mathbf{v}_k|^2) \leq C(\delta^{d-2} + \delta^{d-3} k^{-1})$ .
- (c) *Window vs. cut-off:* replacing  $\mathbf{1}_W$  by  $\chi^2$  in (61) changes the leading constant by  $1 + o(1)$  only.

The leading term is  $q(\lambda_{\parallel}) k \delta^{d-1} < 0$ . The errors (a)–(b), relative to it, are  $\mathcal{O}(\delta)$ ,  $\mathcal{O}((k\delta)^{-1})$  and  $\mathcal{O}((k\delta)^{-2})$ . Choosing  $\delta = k^{-1/2}$  sends  $\delta \rightarrow 0$  and  $k\delta = k^{1/2} \rightarrow \infty$ , so all relative errors vanish as  $k \rightarrow \infty$ . Hence for  $k$  large enough,

$$Q_{t_*}(\mathbf{v}_*) = q(\lambda_{\parallel}) k \delta^{d-1} (1 + o(1)) < 0,$$

which is (59). The final statement is Lemma 6 with hypothesis (ii) now supplied.  $\square$   $\square$

*Remark 7* (What this closes). Lemma 12 converts the existence of the critical load  $t_c$  — previously conditional on the unproved hypothesis (ii) — into a *theorem* for the concrete clamped–free geometry, contingent only on the loading hypothesis (SC). The mechanism is made explicit: the corner instability *is* the half-space surface (creasing) instability, localised to the free face near the edge by the cut-off (60)–(61). This is the precise sense in which the corner bifurcation is “the corner analogue of creasing.”

*Remark 8* (On the threshold  $\lambda_*$ : compressible vs. incompressible). The proof uses only the *existence* of a surface-instability threshold  $\lambda_* \in (0, 1)$ , not its value. The familiar number  $\lambda_* \approx 0.544$  is Biot’s result for the *incompressible* neo-Hookean half-space [1]; the present material (7) is compressible, and its threshold is the corresponding root of the compressible surface secular equation, which depends on the ratio  $\lambda_e/\mu_e$  (equivalently the Poisson ratio) and tends to 0.544 in the incompressible limit  $\lambda_e/\mu_e \rightarrow \infty$  [3]. This is the same instability with a material-dependent threshold, not a discrepancy: Lemma 12 *reduces* the corner instability to the half-space one and thereby agrees with the creasing literature rather than competing with it. Accordingly we keep  $\lambda_*$  symbolic throughout.

## 5.5 Symmetry: simplicity of the mode and the vanishing quadratic

Both model bodies of §5.1 possess reflection symmetries that the fundamental compressive state inherits: (H) is invariant under every orthogonal map fixing the  $x_d$ -axis, and (C) under each coordinate reflection  $x_i \mapsto -x_i$  ( $i < d$ ). Let  $R$  be such a reflection and  $\mathcal{R}$  its action on displacement fields. Because  $\mathbf{F}_0(t)$  and hence  $\mathbb{L}(\mathbf{F}_0(t))$  are  $R$ -invariant, the second-variation operator  $\mathcal{L}(t)$  commutes with  $\mathcal{R}$ , so its eigenspaces split into the symmetric ( $\mathcal{R} = +1$ ) and antisymmetric ( $\mathcal{R} = -1$ ) sectors. This yields, for the concrete geometry, two of the ingredients that §4 had to posit:

- *Simplicity (part of Assumption 6)*. The lowest eigenvalue within a fixed symmetry sector is generically simple, and the critical mode  $\phi$  is the ground state of one sector; the genericity is the standard one for self-adjoint operators commuting with a finite symmetry group. Thus simplicity need not be assumed separately — only the spectral *attainment* (the borderline-elliptic point of Remark 5) remains genuinely open.
- *Vanishing quadratic coefficient (part of Assumption 7)*. The reduced potential of Proposition 1 inherits the  $\mathcal{R}$ -symmetry of the perfect state. If  $\phi$  lies in the antisymmetric sector then  $w \mapsto -w$  is a symmetry of the reduced equation, forcing the quadratic coefficient to vanish exactly, as Assumption 7 stipulated “by the reflection symmetry about the corner bisector.” Here that bisector symmetry is realized concretely by  $R$ .

What symmetry does *not* give is the sign of the cubic,  $b < 0$ ; that is the irreducible core addressed below.

## 5.6 Kinematic relief

The kinematic relief is already dimension-independent: Lemma 9 gives  $J = J_0 + w^2 d_\phi + \mathcal{O}(w^3)$  in  $\mathbb{R}^d$  with  $d_\phi$  the second mixed discriminant of  $\mathbf{F}_0$  and  $\nabla \phi$  (the second-order coefficient of the determinant,  $= \det \nabla \phi$  for  $d = 2$ ), and the rotation condition (48) that suppresses the first-order term is *forced* by the barrier in every dimension by Lemma 10 (its proof uses only  $f''(J)(\delta J)^2 \leq C$ , valid for all  $d$ ). For the concrete model the only points to add are that the genericity  $d_\phi > 0$  holds off a measure-zero set of modes, by the transversality argument of Lemma 11, and that the collapse locus  $\Sigma_c$  is the codimension-one set inside  $B_\rho$  along which the relieved branch would still collapse; the regularised bound of Theorem 2 and the localisation of Proposition 4 then apply verbatim.

## 5.7 The irreducible core

Table 1 summarises the effect of fixing the concrete model. Of the hypotheses of §3–§4, the material assumptions are discharged by the neo-Hookean structure (§5.2), the existence of the critical load by the localized-creasing Lemma 12 under the loading hypothesis (SC), and mode simplicity together with the vanishing quadratic by reflection symmetry (§5.5). Two hypotheses genuinely resist soft arguments and remain assumptions:

1. *Spectral attainment* (the remaining half of Assumption 6): that the critical infimum is attained by an  $L^2$  mode rather than escaping into the essential spectrum associated with the corner. This is the borderline-elliptic difficulty of Remark 5; the norm blow-up (36) is exactly what makes it delicate, and we do not resolve it.
2. *Subcriticality*,  $b < 0$  (the sign in Assumption 7): the direction in which the cubic bends the branch. The creasing literature [1, 3] establishes subcriticality for the flat surface, which makes  $b < 0$  overwhelmingly likely here, but the sign of the reduced cubic at a *corner* is not delivered by symmetry and we verify it only numerically.

The upshot is that for the clamped-free neo-Hookean corner under compression the abstract picture of §3–§4 is, but for two clearly delimited hypotheses, a sequence of theorems: the material bounds are explicit, the corner is genuinely compressive, the instability is a localized creasing event with a constructed destabilising mode, the critical mode is simple and the bifurcation has no quadratic term by symmetry, and the kinematic relief that regularises the tangent holds in every dimension. The two residual hypotheses — attainment of the critical mode and the sign of the

| Hypothesis                                  | Abstract status          | Status for the concrete model                    |
|---------------------------------------------|--------------------------|--------------------------------------------------|
| Reg./growth/ellip./mild (Ass. 1–5)          | standing assumptions     | <b>proved</b> (Lem. 1, §5.2)                     |
| Edge exponent $\alpha$ , scenario selection | abstract pencil          | <b>computed</b> ; $B$ selected by $K < 0$ (§5.3) |
| Critical load $t_c$ exists (Lem. 6(ii))     | not established          | <b>proved</b> under (SC) (Lem. 12)               |
| Transversality (M) (Lem. 7)                 | input on loading         | <b>holds</b> for monotone compression            |
| Mode simplicity (Ass. 6)                    | assumed                  | <b>generic</b> per symmetry sector (§5.5)        |
| Quadratic coeff. vanishes (Ass. 7)          | assumed by symmetry      | <b>forced</b> by reflection $R$ (§5.5)           |
| Rotation condition / relief (Lem. 10, 9)    | <b>proved</b> (all $d$ ) | genericity $d_\phi > 0$ (§5.6)                   |
| Spectral attainment (Ass. 6)                | assumed                  | <i>open</i> (irreducible core)                   |
| Subcriticality $b < 0$ (Ass. 7)             | assumed                  | <i>numerical</i> (irreducible core)              |

Table 1: Hypothesis ledger for the concrete clamped–free neo-Hookean model. Fixing the material and geometry discharges all but two hypotheses; those two form the irreducible analytical core.

cubic — are precisely the ones that the creasing problem on a flat surface also leaves to spectral theory and to computation, and they are inherited here rather than created by the corner.

## 6 Conclusions

## References

- [1] M. A. BIOT, *Surface instability of rubber in compression*, Applied Scientific Research, Section A, 12 (1963), pp. 168–182.
- [2] M. GOLUBITSKY AND D. G. SCHAEFFER, *Singularities and Groups in Bifurcation Theory, Volume I*, vol. 51 of Applied Mathematical Sciences, Springer, 1985.
- [3] W. HONG, X. ZHAO, AND Z. SUO, *Formation of creases on the surfaces of elastomers and gels*, Applied Physics Letters, 95 (2009), p. 111901.
- [4] J. P. KEENER AND H. B. KELLER, *Perturbed bifurcation theory*, Archive for Rational Mechanics and Analysis, 50 (1973), pp. 159–175.
- [5] W. T. KOITER, *Over de stabiliteit van het elastisch evenwicht (On the stability of elastic equilibrium)*, PhD thesis, Delft University of Technology, 1945.
- [6] V. A. KOZLOV, V. G. MAZ’YA, AND J. ROSSMANN, *Elliptic Boundary Value Problems in Domains with Point Singularities*, vol. 52 of Mathematical Surveys and Monographs, American Mathematical Society, 2001.
- [7] V. G. MAZ’YA AND J. ROSSMANN, *Elliptic Equations in Polyhedral Domains*, vol. 162 of Mathematical Surveys and Monographs, American Mathematical Society, 2010.
- [8] S. A. NAZAROV AND B. A. PLAMENEVSKY, *Elliptic Problems in Domains with Piecewise Smooth Boundaries*, vol. 13 of De Gruyter Expositions in Mathematics, Walter de Gruyter, 1994.
- [9] M. WICHROWSKI AND E. G. NOOMEN, *A Unified CutFEM Formulation for Finite-Strain Elasticity: Energy Minimization and Corner Singularities*, arXiv preprint, (2026).