

MATH211: Linear Methods I

Lecture 2

Matthew Burke

Tuesday 11th September, 2018

Lecture 2

Last time

Reduced row echelon form

Examples

Terminology

Examples

Last time

- ▶ Linear equations in one variable
- ▶ Linear equations in two variables
 - ▶ graphical interpretation
- ▶ What elementary row operations are allowed?
 - ▶ graphical interpretation
- ▶ Linear equations in three variables

Reduced row echelon form

Row echelon form

A matrix is in *row echelon form* iff:-

- ▶ All rows consisting entirely of zeros are at the bottom.
- ▶ The first nonzero entry in each nonzero row is a 1 (called the **leading 1** for that row).
- ▶ Each leading 1 is to the right of all leading 1's in rows above it.

For instance:

$$\begin{bmatrix} 0 & 1 & * & * & * & * & * & * \\ 0 & 0 & 0 & 1 & * & * & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

where * can be any number.

Reduced row echelon form

A matrix is in *reduced row echelon form* iff:-

- ▶ It is a row-echelon matrix.
- ▶ Each leading 1 is the only nonzero entry in its column.

$$\begin{bmatrix} 0 & 1 & * & 0 & 0 & * & * & 0 \\ 0 & 0 & 0 & 1 & 0 & * & * & 0 \\ 0 & 0 & 0 & 0 & 1 & * & * & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

where * can be any number.

Advantages of reduced row echelon form

- ▶ Row echelon form \leftrightarrow ready for back-substitution.
- ▶ Reduced row echelon form \leftrightarrow read off solutions.

Another advantage of reduced row echelon form is that:

Theorem

Reduced row echelon form is unique.

Reading off solutions

- ▶ The augmented matrix

$$\left[\begin{array}{ccccc|c} 1 & 0 & 5 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

is in reduced row echelon form.

- ▶ If there are any rows of the form $(0 \ 0 \dots 0 | 1)$ then we know immediately that there are no solutions.

Identify leading 1s and parameter blocks

- ▶ The leading 1s are shown in green

$$\left[\begin{array}{ccccc|c} 1 & 0 & 5 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

and the other non-zero numbers are shown in blue and red.

- ▶ Variables associated to the leading 1s are the *leading variables*.
- ▶ In this case x_1 , x_2 and x_4 are leading and x_3 and x_5 are not.

The non-leading variables can take any value so we assign parameters to them.

In this case:

$$x_3 = s$$

$$x_5 = t$$

and then we solve for the leading variables in terms of these parameters:

$$x_1 = 1 - 5s - t$$

$$x_2 = -s - 2t$$

$$x_4 = -4t$$

Last time
○

Reduced row echelon form
○○○○○○○●

Examples
○○○○○○○

Terminology
○○○○○○○○○

Examples
○○○○○

Questions?

Examples

Worked example

Solve

$$\left[\begin{array}{ccccc|c} 0 & 0 & 0 & -2 & -8 & 4 \\ -3 & 6 & -4 & -9 & 3 & -1 \\ -1 & 2 & -2 & -4 & -3 & 3 \\ 1 & -2 & 1 & 3 & -1 & 1 \end{array} \right]$$

Infinitely many solutions

Solve

$$\left[\begin{array}{cccc|c} 1 & 3 & 4 & 9 & 1 \\ 5 & 6 & 7 & 8 & 5 \\ 6 & 18 & 24 & 54 & 12 \\ 1 & 1 & 1 & 1 & 1 \end{array} \right]$$

Inconsistent

Solve

$$\left[\begin{array}{cccc|c} 1 & 3 & 1 & 2 & 1 \\ 5 & 6 & 7 & 8 & 5 \\ 6 & 9 & 8 & 10 & 10 \\ 1 & 1 & 1 & 1 & 1 \end{array} \right]$$

Unique solution

Solve

$$\left[\begin{array}{cccc|c} 1 & 2 & 3 & 4 & 1 \\ 5 & 6 & 7 & 8 & 5 \\ 11 & 10 & 10 & 12 & 6 \\ 1 & 1 & 1 & 4 & 1 \end{array} \right]$$

Last time
○

Reduced row echelon form
○○○○○○○○

Examples
○○○○○●

Terminology
○○○○○○○○

Examples
○○○○○

Questions?

Terminology

Gaussian elimination

An *elementary row operation* is one of:-

- ▶ swap two rows
- ▶ multiply a row by a non-zero scalar
- ▶ add a multiple of one row to another

The method of *Gaussian elimination* consists of:-

- ▶ using elementary row operations to reduce a matrix to reduced row echelon form
- ▶ make the non-leading variables parameters
- ▶ read off the solutions

Definition of rank

Definition

The *rank* of a matrix is the number of leading 1s in its reduced row echelon form.

Example

Therefore the rank of the following matrix is 3.

$$\begin{bmatrix} 1 & 0 & 5 & 0 & 1 \\ 0 & 1 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The rank is a property of a matrix without augmentation.

Relationship to solution

For a *consistent* system in n variables of rank r :

- ▶ If $n = r$ there is a unique solution.
- ▶ If $r < n$ there are infinitely many solutions.

Homogeneous systems

A system of equations

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1m} & b_1 \\ a_{21} & a_{22} & \dots & a_{2m} & b_2 \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nm} & b_n \end{array} \right]$$

is *homogeneous* iff for all i we have $b_i = 0$. I.e. one that looks like:

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1m} & 0 \\ a_{21} & a_{22} & \dots & a_{2m} & 0 \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nm} & 0 \end{array} \right]$$

Linear combinations

Definition

A *linear combination* of the vectors v_1, v_2, \dots, v_n is a sum of the form

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n$$

where the a_i are scalars.

Example linear combinations

Example

Suppose that we found that the solution to a system of equations was

$$x_1 = 1 - 5s - t,$$

$$x_2 = -s - 2t$$

$$x_3 = s$$

$$x_4 = -4t$$

$$x_5 = t$$

in parameters s and t . Then we can write:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -5 \\ -1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ -2 \\ 0 \\ -4 \\ 1 \end{bmatrix}$$

Questions?

Reading off example

Consider the system:

$$\left[\begin{array}{ccccc|c} 1 & 0 & -3 & 0 & 6 & 8 \\ 0 & 1 & 1 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 5 & -5 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

What is the rank of the corresponding matrix? What is the number of parameters? Is the system homogeneous?

Worked example

Consider the system

$$\left[\begin{array}{cccc|c} 1 & 0 & -1 & 0 & 0 \\ 2 & 0 & 0 & -1 & 0 \\ 0 & 2 & 0 & -2 & 0 \end{array} \right]$$

What is the rank of the corresponding matrix? What is the number of parameters? Is the system homogeneous?

Computer example

Consider the system

$$\left[\begin{array}{ccc|c} 0 & 1 & 3 & 2 \\ 0 & 0 & 5 & 6 \\ 1 & 5 & 1 & -5 \end{array} \right]$$

What is the rank of the corresponding matrix? What is the number of parameters? Is the system homogeneous?

Computer example

Consider the system

$$\left[\begin{array}{ccc|c} 9 & -1 & 0 & -24 \\ -1 & 7 & -4 & 17 \\ 0 & -4 & 8 & -14 \end{array} \right]$$

What is the rank of the corresponding matrix? What is the number of parameters? Is the system homogeneous?

Questions?