

# 26

MART  
MARCH  
PAZARTESİ  
MONDAY

MART 2012						
H	P	S	Ç	P	C	Ct Pz
9					1	2 3 4
10	5	6	7	8	9	10 11
11	12	13	14	15	16	17 18
12	19	20	21	22	23	24 25
13	26	27	28	29	30	31

M A T H E M A T I C S



MART 2012								
H	P	S	C	P	C	Ct	Pz	
9				1	2	3	4	
10	5	6	7	8	9	10	11	
11	12	13	14	15	16	17	18	
12	19	20	21	22	23	24	25	
13	26	27	28	29	30	31		

MART  
MARCH  
SALI  
TUESDAY

27

## Belirsiz Integral

Tanım:  $f(x)$  genelde olup da  $f'(x) = f(x)$  logaritimsal sağlayıcı bir  $F$  fonksiyonu varsa bu  $F$  fonksiyonu  $f'$  nin antiderivedir.

Tanım:  $f(x)$  herhangi bir fonksiyon olursa  $f$  nin tüm entegrallerinin sınıfında  $f$  fonksiyonun  $x$  değişkenine göre belirsiz integrali denir.

$$\boxed{F(x) = \int f(x) dx}$$

## Temel Integral Formüller


$$\textcircled{1} \int x^m dx = \frac{x^{m+1}}{m+1} + C \quad (m \neq -1)$$

$$\textcircled{2} \int x^{-1} dx = \int \frac{1}{x} dx = \ln|x| + C$$

$$\textcircled{3} \int a^x dx = \frac{a^x}{\ln a} + C$$

$$\textcircled{4} \int e^x dx = e^x + C$$

$$\textcircled{5} \int \sin x dx = -\cos x + C$$



$$\textcircled{6} \int \csc^2 x dx = \int \frac{1}{\sin^2 x} dx = -\cot x + C = \int (1 + \cot^2 x) dx$$

$$\textcircled{7} \int \sec^2 x dx = \int \frac{1}{\cos^2 x} dx = \tan x + C = \int (1 + \tan^2 x) dx$$

$$\textcircled{8} \int \frac{1}{1+x^2} dx = \arctan x + C$$

# 28

MART  
MARCH

ÇARŞAMBA  
WEDNESDAY

MART							2012
H	P	S	Ç	P	C	Ct	Pz
9				1	2	3	4
10	5	6	7	8	9	10	11
11	12	13	14	15	16	17	18
12	19	20	21	22	23	24	25
13	26	27	28	29	30	31	

$$\textcircled{10} \int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C$$

$$\textcircled{11} \int \sinh x dx = \cosh x + C$$

$$\textcircled{12} \int \cosh x dx = \sinh x + C$$

$$\textcircled{13} \int \frac{dx}{\sqrt{1+x^2}} = \ln \left[ x + \sqrt{1+x^2} \right] + C$$

$$\textcircled{14} \int \frac{dx}{\sqrt{x^2-1}} = \ln \left[ x + \sqrt{x^2-1} \right] + C$$

Integralin Öznellikleri

$$\textcircled{1} \int \alpha f(x) dx = \alpha \int f(x) dx$$

$$\textcircled{2} \int [f(x) + g(x)] dx = \int f(x) dx + \int g(x) dx$$

$$\textcircled{3} \int (6x^2 - 4x + 3) dx = 2x^3 - 2x^2 + 3x + C$$

$$\textcircled{4} \int \left( 3\sqrt{x} + \frac{1}{2x^4} \right) dx = \int \left( x^{1/3} + \frac{1}{2} x^{-4} \right) dx$$

$$= \frac{x^{4/3}}{4/3} + \frac{1}{2} \frac{x^{-3}}{-3} + C$$

$$= \frac{3}{4} \sqrt[3]{x^4} - \frac{1}{6x^3} + C$$



MART 2012						
H	P	S	C	P	C	Pz
9			1	2	3	4
10	5	6	7	8	9	10 11
11	12	13	14	15	16	17 18
12	19	20	21	22	23	24 25
13	26	27	28	29	30	31

MART  
MARCH

PERŞEMBE  
THURSDAY

29

$$\cancel{\int} \left( \frac{5}{x} + \frac{6}{\cos^2 x} + 3e^x \right) dx = 5 \ln|x| + 6 \tan x + 3e^x + C$$

$$\begin{aligned}\cancel{\int} \frac{5x^2 - 6x + 3}{\sqrt{x}} dx &= \int \left( \frac{5x^2}{\sqrt{x}} \right) dx - \int \frac{6x}{\sqrt{x}} dx + \int \frac{3}{\sqrt{x}} dx \\ &= \int 5x^{3/2} dx - \int 6x^{1/2} dx + \int 3x^{-1/2} dx \\ &= \frac{5x^{5/2}}{5/2} - \frac{6x^{3/2}}{3/2} + \frac{3x^{1/2}}{1/2} \\ &= 2x^{5/2} - 4x^{3/2} + 6x^{1/2} \\ &= 2\sqrt{x^5} - 4\sqrt{x^3} + 6\sqrt{x} + C\end{aligned}$$

$$\begin{aligned}\cancel{\int} \left( 3 \sinhx + 6 \cosh x + \frac{5}{\sinh^2 x} \right) dx \\ &= 3 \cosh x + 6 \sinhx - 5 \coth x + C\end{aligned}$$

$$\begin{aligned}\cancel{\int} \left( x^3 + 3^x + \frac{3}{\sqrt{x^2 - 1}} \right) dx &=? \\ &= \frac{x^4}{4} + \frac{3^x}{\ln 3} + 3 \ln(x + \sqrt{x^2 - 1}) + C\end{aligned}$$



# 30

MART  
MARCH  
CUMA  
FRIDAY

*Süleyman formüller*

MART 2012						
H	P	S	C	P	Ct	Pz
9				1	2	3
10	5	6	7	8	9	10
11	12	13	14	15	16	17
12	19	20	21	22	23	24
13	26	27	28	29	30	31

$$\begin{aligned}
 (\tan x)^l &= \frac{1 + \tan^2 x}{\tan x} \\
 &= \frac{1}{\cos^2 x} \\
 &= \frac{1}{\cos x}
 \end{aligned}$$

$$\begin{aligned}
 (\cot x)^l &= \frac{\sin^2 x + \cos^2 x}{\sin x} \\
 &= \frac{1 + \cos^2 x}{\sin x} \\
 &= \frac{1}{\sin x}
 \end{aligned}$$



MART			2012				
H	P	S	Ç	P	C	Ct	Pz
9				1	2	3	4
10	5	6	7	8	9	10	11
11	12	13	14	15	16	17	18
12	19	20	21	22	23	24	25
13	26	27	28	29	30	31	

sin  
cos

hyperbolik  
+ - kuralı  
yok

MART  
MARCH  
CUMARTESİ  
SATURDAY

31

## Integral Alımlı Kuralları

### Degrısker degrıstırma

Tanım:  $\phi$  sürekli türnevlere sahip bir fonksiyon olmak üzere  $x = \phi(t)$  doğrusal yapislıdırında  $x'$  in diferansiyeli  $\frac{dx}{dt} = \phi'(t)$ ,  $dx = \phi'(t)dt$  olur. İnden,

$$\int f(x) dx = \int f(\phi(t)) \phi'(t) dt$$

Bu adan integral计算ları daha sonra  $t = \phi^{-1}(x)$  yazılıarak sona gelir.

NİSAN  
APRIL

PAZAR  
SUNDAY

01

$$\star \int (1-6x)^6 dx = ?$$

$$1-6x=t$$

$$-6dx = dt$$

$$\int t^6 \cdot \frac{dt}{-6} = -\frac{1}{24} t^7 + C$$

$$= -\frac{(1-6x)^7}{24} + C$$



# 02

NİSAN

APRIL

PAZARTESİ  
MONDAY

NİSAN

2012

H	P	S	C	P	C	Ct	Pz
13							1
14	2	3	4	5	6	7	8
15	9	10	11	12	13	14	15
16	16	17	18	19	20	21	22
17	23	24	25	26	27	28	29
18	30						

\*  $\int \sin(7x) dx = ?$

$$7x = \alpha$$

$$7dx = d\alpha$$

$$\int \sin \alpha \frac{d\alpha}{7} = -\frac{1}{7} \cdot \cos \alpha + C$$

$$= -\frac{\cos(7x)}{7} + C$$

\*  $\int \frac{(4+\ln x)^5}{x} dx = ?$

$$4+\ln x = t$$

$$\frac{1}{x} dx = dt$$

$$\int t^5 dt = \frac{t^6}{6} + C$$

$$= \frac{(4+\ln x)^6}{6} + C$$

\*  $\int \frac{e^{\sqrt{x}}}{\sqrt{x}} dx = ?$

$$\sqrt{x} = t$$

$$x = t^2$$

$$dx = 2t dt$$

$$\int \frac{e^t}{t} 2t dt = 2e^t + C$$

$$= 2e^{\sqrt{x}} + C$$

\*  $\int \tan x dx = ?$

$$\cos x = t$$

$$-\sin x dx = dt$$

$$\int \frac{\sin x}{\cos x} dx = \int -\frac{dt}{t}$$

$$= -\ln|t| + C$$

$$= -\ln|\cos x| + C$$



NİSAN 2012						
H	P	S	C	P	C	Pz
13	14	2	3	4	5	6
	15	9	10	11	12	13
	16	16	17	18	19	20
	17	23	24	25	26	27
	18	30				28

$$\int e^{sx^2} dx$$

$$\int \cos x dx$$

NİSAN  
APRIL  
SALI  
TUESDAY

03

$$\cancel{\star} \int \cos x dx = ?$$

$$\int \frac{\cos x}{\sin x} dx = \int \frac{dt}{t}$$

$$= \ln |t| + C$$

$$= \ln |\sin x| + C$$

$$\cancel{\star} \int \frac{x^4}{1+x^{10}} dx = ?$$

~~$x^{10} = t$~~ 
 ~~$x^4 dx = dt$~~

$$\begin{array}{ccc} 4 & 10 & 2 \\ 2 & 5 & 2 \\ 2 & -5 & 5 \end{array}$$

$$\int \frac{dt}{5(1+t^2)} = \frac{1}{5} \int \frac{dt}{1+t^2}$$

$$= \frac{1}{5} \arctan(t) + C$$

$$= \frac{1}{5} \arctan x^5 + C$$

$$\cancel{\star} \int e^{sx} dx = ?$$

~~$sx = t$~~ 
 ~~$sdx = dt$~~

$$\int e^t \cdot \frac{dt}{s} = \frac{1}{s} \int e^t dt$$

$$= \frac{1}{s} e^t + C$$

$$= \frac{1}{s} e^{sx} + C$$

osney

$$e^{ax} = ae^{ax}$$

$$\int e^{dx} dx = \frac{1}{d} e^{dx} + C$$

Integral



# 04

NİSAN  
APRIL  
ÇARŞAMBA  
WEDNESDAY

Taner  
der cos ve arc tan  
SEN BUL !!!

NİSAN 2012							
H	P	S	G	P	C	Ct	Pz
13	2	3	4	5	6	7	1
14	9	10	11	12	13	14	8
15	16	17	18	19	20	21	15
16	23	24	25	26	27	28	29
17	30						

~~$\int e^{\frac{x}{\alpha}} dx = se^{\frac{x}{\alpha}} + C$~~

~~$\int \frac{dx}{x^2 + \alpha^2} = ?$~~

$$\begin{aligned} x &= \alpha t \\ dx &= \alpha dt \end{aligned}$$

$$\int \frac{\alpha dt}{\alpha^2 t^2 + \alpha^2} = \int \frac{\alpha dt}{\alpha^2 (t^2 + 1)}$$

$$= \frac{1}{\alpha} \int \frac{dt}{t^2 + 1} = \frac{1}{\alpha} \arctan t + C.$$

$$= \frac{1}{\alpha} \arctan \frac{x}{\alpha} + C$$

~~$\int \frac{dx}{x^2 + s^2} = \frac{1}{\sqrt{s}} \arctan \frac{x}{\sqrt{s}} + C$~~

$$s^2 = \sqrt{s}$$

~~$\int \frac{dx}{\sqrt{\alpha^2 - x^2}}$~~

$$\begin{aligned} x &= \alpha t \\ dx &= \alpha dt \end{aligned}$$

$$\boxed{\int \frac{dx}{\sqrt{1-x^2}} = \arcsin x + C}$$

$$\int \frac{\alpha dt}{\sqrt{\alpha^2 - \alpha^2 t^2}} = \int \frac{\alpha dt}{\sqrt{\alpha^2 (1-t^2)}} = \int \frac{dt}{\sqrt{1-t^2}}$$

$$= \arcsin t + C$$

$$= \alpha \arcsin \frac{x}{\alpha} + C$$



NİSAN 2012						
H	P	S	C	P	C	Pz
13						1
14	2	3	4	5	6	7
15	9	10	11	12	13	14 15
16	16	17	18	19	20	21 22
17	23	24	25	26	27	28 29
18	30					

(D)  $\cos^2 x = 1 - \sin^2 x$  NİSAN  
 $\theta^2 - \theta^2 \sin^2 x$  APRIL  
 PERŞEMBE  
 THURSDAY

05

~~X~~  $\int \frac{dx}{\sqrt{3-2x-x^2}} = ?$

$$3-2x-x^2 \stackrel{?}{=} \theta^2-t^2$$

$$\stackrel{?}{=} 7-2x$$

$$= -[x^2 + 2x - 3]$$

$$\int \frac{dt}{\sqrt{2^2-t^2}} = \arcsin \frac{t}{2} + C$$

$$= -[(x+1)^2 - 4]$$

$$= \arcsin \left( \frac{x+1}{2} \right) + C$$

$$= 4 - (x+1)^2$$

$$= 2^2 - t^2$$

$$x+1=t$$

$$dx=dt$$

~~X~~  $\int x^2 \sqrt{x-2} dx = ?$

$$x-2=t^2$$

$$dx=2t dt$$

$$\int (t^2+2)^2 t \cdot 2t dt$$

$$(t^6 + 4t^4 + 4t^2)^{1/2}$$

$$2 \int t^2 [t^4 + 4t^2 + 4] dt$$

$$2t^6 + 8t^4 + 8t^2$$

$$2 \int (t^6 + 4t^4 + 4t^2) dt$$

$$2 \left[ \frac{t^7}{7} + \frac{4t^5}{5} + \frac{4t^3}{3} \right] + C = 2 \left[ \frac{(x-2)^{7/2}}{7} + \frac{4(x-2)^{5/2}}{5} + \frac{4(x-2)^{3/2}}{3} \right] + C$$

(1) İntegralin herhangi bir yerinde  $\sqrt{\theta^2 - x^2}$  böyle bir ifade versa  $x = \theta \sin t$  dönüşüm yoluyla

~~X~~  $\int \frac{x+8}{\sqrt{9-x^2}} dx$

$$\begin{cases} x = 3 \sin t \\ dx = 3 \cos t dt \end{cases} \rightarrow$$

$$\int \frac{3 \sin t + 8}{\sqrt{9 - 9 \sin^2 t}} 3 \cos t dt$$

$$= \int (3 \sin t + 8) dt = -3 \cos t + 8t + C$$

$$3 \cos t$$



06

NİSAN  
APRILCUMA  
FRIDAY

$$\cos^2 x = 1 - \sin^2 x$$

$\downarrow$   
 $\delta^2 - x^2$

$$1 + \tan^2 x = \sec^2 x \Rightarrow \frac{1}{\cos^2 x}$$

$$\tan^2 x = \sec^2 x - 1$$


Tüm  
formüller  
!!!!

NİSAN 2012						
H	P	S	C	P	C	Ct Pz
13						1
14	2	3	4	5	6	7 8
15	9	10	11	12	13	14 15
16	16	17	18	19	20	21 22
17	23	24	25	26	27	28 29
18	30					

$$\left[ x = 3 \sin t \rightarrow \frac{x}{3} = \sin t \rightarrow t = \arcsin \frac{x}{3} \right]$$

$$= -3 \cos t + 8t + C$$

~~$$= -3 \cos(\arcsin \frac{x}{3}) + 8 \arcsin \frac{x}{3} + C$$~~



$$= -\sqrt{8-x^2} + 8 \arcsin \frac{x}{3} + C$$

② İntegrallerin herhangi bir yemde  $\sqrt{x^2 - a^2}$   
Vorsa  $x = a \sec t$  donusum yapılırı.

~~$$\int \frac{dx}{x \sqrt{x^2 - 9}} = ?$$~~

$$x = 3 \sec t$$

$$dx = 3 \sec t \cdot \sec t \tan t dt$$

~~$$\int \frac{3 \sec t \tan t dt}{3 \sec t \sqrt{9 \sec^2 t - 9}}$$~~

$$\int \frac{\tan t dt}{3 \sqrt{\sec^2 t - 1}} = \int \frac{1}{3} dt = \frac{t}{3} + C$$

$$= \frac{1}{3} \operatorname{arcsec} \frac{x}{3} + C$$



NİSAN 2012							
H	P	S	C	P	C	Ct	Pz
13						1	
14	2	3	4	5	6	7	8
15	9	10	11	12	13	14	15
16	16	17	18	19	20	21	22
17	23	24	25	26	27	28	29
18	30						

$$\int \frac{dx}{\sqrt{1+x^2}} = \operatorname{arcose} x$$

NİSAN  
APRIL  
SALI  
TUESDAY

17

③ İntegratörün herhangi bir yerinde  $\sqrt{a^2+x^2}$ '  
Vorsa  $x = a \tan t$  denkleme yapılı.

$$\int \frac{dx}{x^2 \sqrt{x^2+a^2}}$$

$\rightarrow 2 \sec^2 t$

$$\int \frac{(a \sec^2 t) dt}{a^2 \tan^2 t \sqrt{a^2 \tan^2 t + a^2}}$$

$\cancel{a^2 \sec^2 t}$

$\cancel{a^2 \tan^2 t}$

$$\int \frac{1}{2} \frac{\sec^2 t}{\tan^2 t \cdot 2 \sec t} dt$$

$$\frac{1}{2} \int \frac{\sec t}{\tan^2 t} dt$$

$$\frac{1}{2} \int \frac{\cos t}{\sin^2 t} dt$$


$$\frac{1}{2} \int \frac{du}{v^2} = \frac{1}{2} \int v^{-2} dv$$

$$= \frac{1}{2} \frac{v^{-1}}{-1} + C$$

$$= \frac{1}{2} \frac{1}{uv} + C$$

$$= -\frac{1}{2 \sin t} + C$$

$$= -\frac{\sqrt{a^2+x^2}}{2x} + C$$



# 18

NİSAN  
APRIL  
ÇARŞAMBA  
WEDNESDAY

$$\sqrt{\frac{6}{4}(4x^2 - 1)}$$

$$2x \sqrt{x^2 - \frac{1}{4}}$$

H	P	S	Ç	P	C	Ct	Pz
13							1
14	2	3	4	5	6	7	8
15	9	10	11	12	13	14	15
16	16	17	18	19	20	21	22
17	23	24	25	26	27	28	29
18	30						

~~$\int \frac{dx}{x\sqrt{6x^2-1}}$~~  = ?

$x = \frac{1}{2} \sec t$   
 $dx = \frac{1}{2} \sec t \tan t dt$

~~$\int \frac{1}{2} \sec t \tan t dt$~~   
 ~~$\int \frac{1}{2} \sec t dt$~~

$x = \frac{1}{2} \sec t$   
 $dx = \frac{1}{2} \sec t \tan t dt$

$\int \frac{1}{2} \sec t \tan t dt$   
 ~~$\int \frac{1}{2} \sec t \sqrt{\frac{1}{4} \sec^2 t - \frac{1}{4}}$~~   
 ~~$\frac{1}{2} \tan t$~~

$t + C$

$\arcsin 2x + C$

Q) İntegrallerin herhangi bir yerinde  $\sqrt{ax+b}$   
ve  $\sqrt{ax^2+b}$  olsun;  $\int dx = t$  olsun;  
değerlerin donanıma uygulanır. (Kök konu aynı olmalıdır)

~~$\int \frac{\sqrt{x+1} + 2}{\sqrt{x+8}} dx$~~  = ?

$$\begin{array}{r|l} 6 & 6 \\ 2 & 2 \\ 2 & 1 \end{array}$$

$\int \frac{t^3 + 2}{t^2} 12t^{11} dt$

$x+1 = t^{12}$   
 $dx = 12t^{11} dt$

$$\begin{aligned} \int \left(t + \frac{2}{t^2}\right) 12t^{11} dt &= 12 \int (t^{12} + 2t^9) dt \\ &= 12 \left[ \frac{t^{13}}{13} + \frac{2t^{10}}{10} \right] + C \end{aligned}$$



NİSAN							2012	
H	P	S	C	P	C	Ç	Pz	
13						1		
14	2	3	4	5	6	7	8	
15	9	10	11	12	13	14	15	
16	16	17	18	19	20	21	22	
17	23	24	25	26	27	28	29	
18	30							

NİSAN  
APRIL

PERŞEMBE  
THURSDAY

19

(5) Mekanikla ıçinde trigonometrik haneler ve  
Bir yarım açı yanlarında döşeler  $\tan \frac{x}{2} = t$   
düşükken döşüm 1'e göre olur.

$$\begin{aligned} \text{arc tan } t &= \frac{x}{2} \\ 2 \text{arc tan } t &= x \\ 2 \frac{1}{1+t^2} dt &= dx \end{aligned}$$

$$\tan \frac{x}{2} = t$$

$$\sin \frac{x}{2} = \frac{t}{\sqrt{1+t^2}}$$

$$\cos \frac{x}{2} = \frac{1}{\sqrt{1+t^2}}$$

$$\sin 2x = 2 \sin x \cos x$$

$$\sin x = 2 \sin \frac{x}{2} \cos \frac{x}{2}$$

$$\sin x = \frac{2t}{1+t^2}$$

$$\cos 2x = 2 \cos^2 x - 1$$

$$\cos x = 2 \cos^2 \left( \frac{x}{2} \right) - 1$$

$$\cos x = \frac{1-t^2}{1+t^2} - 1$$

$$\cos x = \frac{1-t^2}{1+t^2}$$

$$\tan x = \frac{\sin x}{\cos x} = \frac{2t}{1-t^2}$$

$$\cot x = \frac{\cos x}{\sin x} = \frac{1-t^2}{2t}$$



# 20

NİSAN  
APRIL  
CUMA  
FRIDAY

$$\begin{aligned} & t+t^2+t-t^3 \quad 2 \quad \frac{dt}{1+t^2} \\ & 1+t^2+2t \\ & 2+2t^2+4t \\ & t^2+2t+1 \quad (t+1)^2 \quad 2 \end{aligned}$$

NİSAN 2012						
H	P	S	C	P	C	Pz
13						1
14	2	3	4	5	6	7 8
15	9	10	11	12	13	14 15
16	16	17	18	19	20	21 22
17	23	24	25	26	27	28 29
18	30					

K  $\int \frac{1+\sin x}{(1+\cos x) \sin x} dx = ?$

$$\tan \frac{x}{2} = t$$

$$\int \frac{\left[1 + \frac{2t}{1+t^2}\right] \frac{2t}{1+t^2}}{\left[1 + \frac{1-t^2}{1+t^2}\right] \frac{2t}{1+t^2}} dt$$

$$\boxed{\frac{2}{1+t^2} dt = dx}$$

$$\begin{cases} \sin x = \frac{2t}{1+t^2} \\ \cos x = \frac{1-t^2}{1+t^2} \end{cases}$$

$$\int \frac{t^2+2t+1}{2t} dt$$

$$\frac{1}{2} \int t dt + \int 1 dt + \frac{1}{2} \int \frac{1}{t} dt$$

$$\frac{t^2}{4} + t + \frac{1}{2} \ln |t| + C$$

K  $\int \frac{\sin x - \cos x}{\sin x + \cos x} dx$

$$\cos x + \sin x = t$$

$$(-\sin x + \cos x) dx = dt$$



NİSAN 2012

H	P	S	C	P	C	Ct	Pz
13							1
14	2	3	4	5	6	7	8
15	9	10	11	12	13	14	15
16	16	17	18	19	20	21	22
17	23	24	25	26	27	28	29
18	30						

NİSAN

APRIL

CUMARTESİ

SATURDAY

21

$$\cancel{\star} \int \cos 3x dx = ?$$

$$3x = t \\ 3dx = dt$$

$$\int \cos ax dx = \frac{1}{a} \sin ax + C$$

$$\cancel{\star} \int \sin 5x dx = ?$$

$$5x = t \\ 5dx = dt$$

NİSAN

APRIL

PAZAR

SUNDAY

22

$$\int \sin ax dx = -\frac{1}{a} \cos ax + C$$

$$\cancel{\star} \int \frac{dx}{5+3 \cos x} = ?$$

$$\tan \frac{x}{2} = t$$

$$dx = \frac{2dt}{1+t^2}$$

$$\cos x = \frac{1-t^2}{1+t^2}$$

$$= \int \frac{dt}{2^2 + t^2} = \frac{1}{2} \arctan \frac{t}{2} + C$$

$$= \frac{1}{2} \arctan \left( \frac{\tan \frac{x}{2}}{2} \right) + C$$



# 23

NİSAN  
APRIL

PAZARTESİ  
MONDAY


NİSAN 2012						
H	P	S	C	P	Ct	Pz
13						1
14	2	3	4	5	6	7 8
15	9	10	11	12	13	14 15
16	16	17	18	19	20	21 22
17	23	24	25	26	27	28 29
18	30					

## 2. Kısım İntegral Yorumu

Tanım:  $v$  ve  $v'$ ,  $x$ 'e bağlı,

$$\int f(x) dx = v \cdot v' - \int v dv$$

$\underbrace{v}_{\text{v}}$      $\underbrace{dv}_{\text{v'}}$



$$\cancel{\int} \int x e^x dx = ?$$

$\underbrace{x}_{\text{v}}$      $\underbrace{e^x dx}_{\text{v'}}$

$$\begin{aligned} x &= v \\ dx &= dv \\ e^x dx &= dv \\ \int e^x dx &= \int dv \\ e^x &= v \end{aligned}$$

$$= x e^x - \int e^x dx$$

$$= x e^x - e^x + C$$

$$\cancel{\int} \int x e^{3x} dx = ?$$

$$\begin{aligned} x &= v \\ dx &= dv \end{aligned}$$

$$\begin{aligned} e^{3x} dx &= dv \\ \frac{e^{3x}}{3} &= v \end{aligned}$$

$$= x \cdot \frac{e^{3x}}{3} - \int \frac{e^{3x}}{3} dx$$

$$= \frac{x e^{3x}}{3} - \frac{1}{3} \int e^{3x} dx$$

$$= \frac{x e^{3x}}{3} - \frac{1}{3} e^{3x} + C$$



NİSAN							2012	
H	P	S	C	P	C	Ct	Pz	
13							1	
14	2	3	4	5	6	7	8	
15	9	10	11	12	13	14	15	
16	16	17	18	19	20	21	22	
17	23	24	25	26	27	28	29	
18	30							

NİSAN  
APRIL  
SALI  
TUESDAY

24

$$\cancel{x} \int x \sin 2x dx = ?$$

$$x = u \\ dx = du$$

$$\sin 2x dx = du \\ -\frac{1}{2} \cos 2x + C$$

$$= x \cancel{\frac{1}{2}} \cos 2x - \int -\frac{1}{2} \cos 2x dx$$

$$\approx -\frac{x \cos 2x}{2} + \frac{1}{4} \sin 2x + C$$

$$\cancel{x} \int \frac{x}{\sin^2 x} dx = ?$$

$$x = u \\ dx = du \\ \sin^2 x \\ -\cot x = v$$

$$= -x \cot x - \int -\cot x dx$$

$$\int -\cot x dx = \int \frac{\cos x}{\sin x} dx \\ \left| \begin{array}{l} \sin x = t \\ \cos x dx = dt \end{array} \right. \\ = \int \frac{dt}{t} \\ = \ln |\sin x|$$

$$\cancel{x} \int \frac{x}{\cos^2 x} dx = ?$$

$$x = u \\ dx = du \\ \cos^2 x \\ \tan x = v$$

$$= x \tan x - \int \tan x dx$$

$$\int \tan x dx = \int \frac{\sin x}{\cos x} dx \\ \left| \begin{array}{l} \cos x = t \\ -\sin x dx = dt \end{array} \right. \\ = -\ln |\cos x| + C$$

$$= x \tan x + \ln |\cos x| + C$$



# 25

NİSAN  
APRIL  
ÇARŞAMBA  
WEDNESDAY

LAPTU

NİSAN 2012							
H	P	S	C	P	C	Ct	Pz
13							1
14	2	3	4	5	6	7	8
15	9	10	11	12	13	14	15
16	16	17	18	19	20	21	22
17	23	24	25	26	27	28	29
18	30						

$$\cancel{x} \int x^4 \ln x dx$$

$$\ln x = u$$

$$\frac{1}{x} dx = du$$

$$x^4 dx = du$$

$$\frac{x^5}{5} = u$$

$$= (\ln x) \frac{x^5}{5} - \int \frac{x^5}{5} \cdot \frac{1}{x} dx$$

$$= (\ln x) \frac{x^5}{5} - \frac{1}{5} \int x^4 dx$$

$$= (\ln x) \frac{x^5}{5} - \frac{x^5}{25} + C$$

$$\cancel{x} \int \arctan x dx$$

$$= \arctan x \cdot x - \int x \frac{1}{1+x^2} dx$$

~~$$\arctan x \cdot x - \int x \frac{1}{1+x^2} dx$$~~

$$\cancel{\arctan x \cdot x - \int x \frac{1}{1+x^2} dx}$$

$$\begin{cases} \arctan x = u & dx = du \\ \frac{dx}{1+x^2} = du & x = v \end{cases}$$

$$\int \frac{x dx}{1+x^2}$$

$$\begin{aligned} 1+x^2 &= t \\ 2x dx &= dt \end{aligned}$$

$$\int \frac{1}{t} \frac{dt}{2}$$

$$= \frac{1}{2} \ln |t| + C$$



NİSAN 2012							
H	P	S	C	P	C	Ct	Pz
13						1	
14	2	3	4	5	6	7	8
15	9	10	11	12	13	14	15
16	16	17	18	19	20	21	22
17	23	24	25	26	27	28	29
18	30						

tert laper

NİSAN  
APRIL

PERŞEMBE  
THURSDAY

26

$$\int e^{ax} \cos bx dx = ?$$

$$\begin{aligned} u &= e^{ax} \\ du &= a e^{ax} dx \end{aligned}$$

$$= e^{ax} \frac{1}{b} \sin bx - \int \frac{1}{b} \sin bx \cdot a e^{ax} dx$$

$$\begin{aligned} \cos bx dx &= dv \\ \frac{1}{b} \sin bx &= v \end{aligned}$$

$$= \frac{1}{b} e^{ax} \sin bx - \frac{a}{b} \int e^{ax} \sin bx dx$$

$$\begin{aligned} u &= e^{ax} \\ du &= a e^{ax} dx \\ \sin bx dx &= dv \\ -\frac{1}{b} \cos bx &= v \end{aligned}$$

$$= -\frac{1}{b} e^{ax} \cos bx + \frac{a}{b} \int \cos bx e^{ax} dx$$

$$= \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx - \frac{a^2}{b^2} \int e^{ax} \cos bx dx$$

$$I = \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx - \frac{a^2}{b^2} I$$

$$\frac{a^2 + b^2}{b^2} I = \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx$$

$$I = \frac{b^2}{a^2 + b^2} \left[ \frac{1}{b} e^{ax} \sin bx + \frac{a}{b^2} e^{ax} \cos bx \right] + C$$



# 27

NİSAN  
APRIL  
CUMA  
FRIDAY

NİSAN 2012						
H	P	S	C	P	C	Cl
13						1
14	2	3	4	5	6	7
15	9	10	11	12	13	14
16	16	17	18	19	20	21
17	23	24	25	26	27	28
18	30					

2X.

~~2X.~~  $\int \frac{xe^x}{(x+1)^2} dx$

$v = xe^x$ 
 $dv = (e^x + xe^x) dx$

$\frac{dx}{(x+1)^2} = dv$

$= xe^x - \frac{t}{x+1} - \int \frac{-1}{x+1} e^x (x+1) dx$

$\int \frac{dx}{(x+1)^2} = dv$ 
 $\int \frac{dt}{t^2} = v$ 
 $x+1=t$ 
 $dx=dt$

$= \frac{-xe^x}{x+1} + \int e^x dx$

$\int t^{-2} dt = -t^{-1}$

$= \frac{-xe^x}{x+1} + e^x + C$

$= -\frac{1}{t}$ 
 $= -\frac{1}{x+1}$

$v = \frac{-1}{x+1}$

~~2X.~~  $\int \sqrt{x^2 + 5} dx$

$= \sqrt{x^2 + 5} \cdot x - \int x \frac{x}{\sqrt{x^2 + 5}} dx$

$\sqrt{x^2 + 5} = u$ 
 $dx = du$ 
 $\frac{1}{2} (x^2 + 5)^{-\frac{1}{2}} \cdot 2x dx = du$ 
 $x = v$ 
 $\frac{x}{\sqrt{x^2 + 5}} dx = du$

$\int \frac{x^2 + 5 - 5}{\sqrt{x^2 + 5}} dx = \int \frac{(x^2 + 5) - 5}{\sqrt{x^2 + 5}} dx$

$= \int \sqrt{x^2 + 5} dx - 5 \int \frac{1}{\sqrt{x^2 + 5}} dx$

Formül 13

$x = \sqrt{5}t$ 
 $dx = \sqrt{5}dt$ 
 $\int \frac{1}{\sqrt{5+t^2}} dt$

$x\sqrt{x^2 + 5} + 5 \int \frac{1}{\sqrt{x^2 + 5}} dx \rightarrow I$

$x\sqrt{x^2 + 5} + 5 \ln \left[ x + \sqrt{5+x^2} \right] - I$

$I = x\sqrt{x^2 + 5} + 5 \ln \left[ x + \sqrt{5+x^2} \right]$



MAYIS	2012					
H	P	S	C	P	Ct	Pz
18		1	2	3	4	5 6
19	7	8	9	10	11	12 13
20	14	15	16	17	18	19 20
21	21	22	23	24	25	26 27
22	28	29	30	31		

HAFTA

MAYIS  
MAY  
CUMARTESİ  
SATURDAY

05

$$\int x^2 e^{-x} dx = ?$$

$$\begin{aligned} & \cancel{x^2} \\ & \cancel{\int x^2 e^{-x} dx = 0} \\ & \cancel{x^3} \\ & \cancel{\int x^3 e^{-x} dx = 0} \\ & \cancel{x^3} = v \\ & \cancel{x^2} = u \\ & \cancel{dx = du} \\ & \cancel{-e^{-x}} = v \end{aligned}$$

$$\begin{aligned} &= -\cancel{e^{-x} \cdot \frac{x^3}{3}} - \cancel{\int \frac{x^3}{3} \cdot (-e^{-x}) dx} \\ &= -e^{-x} \cancel{\frac{x^3}{3}} + \cancel{\left( \frac{1}{3} \right)} \int x^3 e^{-x} dx \end{aligned}$$

$$\begin{aligned} & x^2 = u & e^{-x} dx = dv \\ & 2x dx = du & -e^{-x} = v \end{aligned}$$

$$= x^2 \cdot (-e^{-x}) - \int f(e^{-x}), 2x dx$$

$$= -x^2 e^{-x} + \int 2x e^{-x} dx$$

$$= -x^2 e^{-x} + 2 \left[ -xe^{-x} - e^{-x} \right] + C$$

$$\begin{aligned} & x = u & e^{-x} dx = dv \\ & dx = du & -e^{-x} = v \\ & -xe^{-x} + \int e^{-x} dx \end{aligned}$$

MAYIS  
MAY

06

PAZAR  
SUNDAY

$$-xe^{-x} - e^{-x} + C$$

Not:

$$\int x^{10} e^x dx = e^x \left[ x^{10} + 10x^9 + \dots \right] + C$$

Kendisi Tersi

Tersi *(Tersi)*

$$\int x^3 e^x dx = e^x \left[ x^3 + 3x^2 + 6x + 6 \right] + C$$

J 782 *(yeni)*

30.05.10

20 MAYIS 2012



07

MAYIS

MAY

PAZARTESİ  
MONDAY

Béziers Curve

H	P	S	C	P	C	Ct	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

## (3) İNTEGRALİ RİTMLİLER

$$\text{①} \int \cos^n x dx = \int \underbrace{\cos^{n-1} x}_{U} \underbrace{\cos x dx}_{dv}$$

$$U = \cos^{n-1} x$$

$$dU = (n-1) \cos^{n-2} x \cdot (-\sin x) dx$$

$$dv = \cos x dx$$

$$V = \sin x$$

$$= (\cos^{n-1} x) (\sin x) + \int \underbrace{(n-1) \sin^2 x \cos^{n-2} x dx}_{(1-\cos^2 x)}$$

$$= \cos^{n-1} x \sin x + (n-1) \left[ \int \cos^{n-2} x dx - \int \cos^n x dx \right]$$

$$I = \cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x dx - (n-1) I$$

$$I = \frac{\cos^{n-1} x \sin x + (n-1) \int \cos^{n-2} x dx}{n}$$

$\int \sin^n x dx$  sea bu (!!)



MAYIS 2012						
H	P	S	Ç	P	C	Cl Pz
18	1	2	3	4	5	6
19	7	8	9	10	11	12 13
20	14	15	16	17	18	19 20
21	21	22	23	24	25	26 27
22	28	29	30	31		

木亭

$$\cos 2x = 2\cos^2 x - 1$$

MAYIS  
MAY  
SALI  
TUESDAY

08

$$\text{A) } \int \cos^4 x dx = \frac{1}{4} \cos^3 x \sin x + \frac{3}{4} \int \cos^2 x dx$$

$$\int \cos^2 x dx = \frac{1}{2} \int (\cos 2x + 1) dx$$

$$= \frac{1}{2} \left[ \frac{1}{2} \sin 2x + x \right].$$

$$= \frac{1}{4} \cos^3 x \sin x + \frac{3}{16} \sin 2x + \frac{3x}{8} + C$$

~~Gelişmeyen  
Düzenli~~

$$\int \cos^4 x dx = \int \underbrace{\cos^3 x}_{u} \underbrace{\cos x dx}_{dv}$$

$$\int \sin^5 x dx = ?$$

$$\textcircled{2} \quad \int \csc^n x dx = \int -\frac{1}{\sin^n x} dx$$

$$= \int \underbrace{\frac{1}{\sin^{n-2} x}}_{u} \cdot \underbrace{\frac{1}{\sin^2 x} dx}_{dv}$$

$$u = \sin^{2-n} x$$

$$du = (2-n) \sin^{1-n} x (\cos x) dx$$

$$\frac{dx}{\sin^2 x} = dv$$

$$-\cot x = v \Rightarrow -\frac{\cos x}{\sin x} = v$$



# 09

MAYIS

MAY

ÇARŞAMBA  
WEDNESDAY

MAYIS		2012					
H	P	S	Ç	P	C	Ct	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

$$= \sin^{2-n} x \left( \frac{-\cos x}{\sin^n x} \right) - \int \frac{-\cos x}{\sin^n x} (2-n) \sin^{1-n} x \cos x dx$$

$$= \frac{-\cos x}{\sin^{n-1} x} + (2-n) \int \frac{\cos^2 x}{\sin^n x} dx$$

~~$$\frac{1 - \sin^2 x}{\sin^n x}$$~~

$$= \frac{-\cos x}{\sin^{n-1} x} + \boxed{(2-n) \int \frac{1}{\sin^n x} dx} - (2-n) \int \frac{1}{\sin^{n-2} x} dx$$

I

$$I = \frac{-\cos x}{\sin^{n-1} x} + (n-2) \int \frac{1}{\sin^{n-2} x} dx + (2-n) I$$

$$I = \frac{1}{(n-1)} \cdot \left[ \frac{-\cos x}{\sin^{n-1} x} + (n-2) \int \frac{1}{\sin^{n-2} x} dx \right]$$

~~$$\int \operatorname{cosec}^6 x dx = \int \frac{1}{\sin^6 x} dx.$$~~

$$= \frac{-\cos x}{\sin^5 x} + \frac{4}{5} \int \frac{1}{\sin^4 x} dx$$

$$\int \frac{1}{\sin^4 x} dx = \frac{-\cos x}{3 \sin^3 x} + \frac{2}{3} \int \frac{1}{\sin^2 x} dx$$

$$= \boxed{\frac{-\cos x}{3 \sin^3 x} - \frac{2}{3} \cot x}$$



$\int \sec^6 x dx$  bitti!!

MAYIS	2012						
H	P	S	Ç	P	C	Ct	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

MAYIS  
MAY

PERŞEMBE  
THURSDAY

10

$$(3) \int \tan^n x dx = \int \tan^{n-2} x \tan^2 x dx$$

$$= \int \tan^{n-2} x (\tan^2 x + 1 - 1) dx$$

$$= \int \tan^{n-2} x (\tan^2 x + 1) dx - \int \tan^{n-2} x dx$$

$$\int \tan^{n-2} x (\tan^2 x + 1) dx$$

$$\int t^{n-2} dt = \frac{t^{n-1}}{n-1} + C$$

$$= \frac{\tan^{n-1} x}{n-1} + C$$

$$= \frac{\tan^{n-1} x}{n-1} - \int \tan^{n-2} x dx$$

$$\cancel{\int \tan^4 x dx = ?}$$

$$= \frac{\tan^3 x}{3} - \int \tan^2 x dx$$

$$\begin{aligned} \tan x &= t \\ (1+\tan^2 x) dx &= dt \end{aligned}$$

$$\boxed{\int \tan^2 x dx = \int (\tan^2 x + 1) - 1 dx = \tan x - x}$$

$$= \frac{\tan^3 x}{3} - \tan x + x + C$$

$$\int \cot^4 x dx \text{ butlu?}$$



# 11

MAYIS

MAY

CUMA

FRIDAY

MAYIS 2012						
H	P	S	Ç	P	C	Cl
18		1	2	3	4	5
19	7	8	9	10	11	12
20	14	15	16	17	18	19
21	21	22	23	24	25	26
22	28	29	30	31		27

$$\textcircled{4} \quad \int \frac{dx}{(\alpha^2 + x^2)^n} = ?$$

Bir deşte derece  
azalır. Gözükse  
derece kendisinden  
gözelik:

$$\int \frac{1}{(\alpha^2 + x^2)^{n-1}} dx$$

$v$        $dv$

$v = (\alpha^2 + x^2)^{1-n}$   
 $dv = (1-n)(\alpha^2 + x^2)^{-n} (2x) dx$   
 $dx = dv$   
 $x = v$

$$= \frac{1}{(\alpha^2 + x^2)^{n-1}} \cdot x - \int x \cdot (1-n)(\alpha^2 + x^2)^{-n} (2x) dx$$

$$x^2 = \alpha^2 - a^2$$

$$\int \frac{1}{(\alpha^2 + x^2)^{n-1}} dx = \frac{x}{(\alpha^2 + x^2)^{n-1}} - 2(1-n) \int \frac{x^2}{(\alpha^2 + x^2)^n} dx$$

$$= \frac{x}{(\alpha^2 + x^2)^{n-2}} - 2(1-n) \int \frac{1}{(\alpha^2 + x^2)^{n-1}} dx + 2(n-1)\alpha^2 \int \frac{1}{(\alpha^2 + x^2)^n} dx$$

$$2(1-n)\alpha^2 \int \frac{1}{(\alpha^2 + x^2)^n} dx = \frac{x}{(\alpha^2 + x^2)^{n-1}} + (1-2n) \int \frac{1}{(\alpha^2 + x^2)^{n-1}} dx$$

$$= \frac{x}{(\alpha^2)^{n-1}}$$



MAYIS	2012						
H	P	S	Ç	P	C	Çt	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

Yıldız P.M.P'

MAYIS  
CUMARTESİ  
SATURDAY

12

$$\int \frac{dx}{(\alpha^2+x^2)^n} = \frac{1}{2(1-n)\alpha^2} \left[ \frac{x}{(\alpha^2+x^2)^{n-1}} + (1-2n) \int \frac{1}{(\alpha^2+x^2)^{n-1}} dx \right]$$

\*  $\int \frac{dx}{(6x^2+6x+10)^2} = \int \frac{dx}{((2x+3)^2+3^2)^2}$

$$\int \frac{dt/2}{(t^2+3^2)^2} = \frac{1}{2} \int \frac{dt}{(3^2+t^2)^2} \quad \begin{aligned} 2x+1 &= t \\ 2dx &= dt \end{aligned}$$

$$d=3$$

$$n=2$$

$$= \frac{1}{2} \left[ \frac{-1}{18} \left( \frac{t}{(9+t^2)} \right) + 3 \int \frac{1}{(9+t^2)} dt \right] \quad \begin{array}{l} \text{MAYIS} \\ \text{MAY} \\ \text{PAZAR} \\ \text{SUNDAY} \end{array}$$

$$= \frac{-1}{36} \left[ \frac{t}{9+t^2} - \arctan \frac{t}{3} \right] + C \quad \int \frac{dx}{\alpha^2+x^2} = \frac{1}{\alpha} \arctan \frac{x}{\alpha}$$

$$= \frac{-1}{36} \left[ \frac{2x+1}{6x^2+6x+10} - \arctan \frac{2x+1}{3} \right] + C$$

\*  $\int \frac{dx}{(1+x^2)^2} = ?$

$$x = \tan t \quad dx = (1+\tan^2 t) dt$$

$$\int \frac{(1+\tan^2 t) dt}{(1+\tan^2 t)^2} = \int \frac{dt}{1+\tan^2 t} = \int \cos^2 t dt$$

$$\int \frac{(1+\tan^2 t) dt}{(1+\tan^2 t)^2} = \int \frac{dt}{1+\tan^2 t}$$

$$\int \frac{dt}{1+\tan^2 t} = \int \frac{dt}{\cos^2 t} = \int \cos^2 t dt$$




# 14

MAYIS  
MAY  
PAZARTESİ  
MONDAY

MAYIS		2012	
H	P	S	C
18		1	2 3 4 5 6
19	7	8	9 10 11 12 13
20	14	15 16 17 18 19 20	
21	21	22 23 24 25 26 27	
22	28	29 30 31	

$$= \int \frac{1}{2} (\cos 2t + 1) dt = \frac{1}{2} \left[ \frac{1}{2} \sin 2t + t \right] + C$$

$$x = \tan t$$



$$\sin 2t = 2 \sin t \cos t$$

$$= 2 \frac{x}{\sqrt{x^2+1}} \cdot \frac{1}{\sqrt{x^2+1}}$$

$$= \frac{1}{2} \cdot \frac{2x}{(x^2+1)} + \frac{1}{2} \operatorname{arctan} x + C$$

## BASIT KESİRLERDE AYRMA YÖNTEMİ

$$\frac{P(x)}{Q(x)} = K(x) + \frac{R(x)}{Q(x)}$$

$$\int \frac{P(x)}{Q(x)} dx = \int K(x) dx + \int \frac{R(x)}{Q(x)} dx ?$$

$$\frac{R(x)}{Q(x)} = \frac{M}{px+q} + \frac{N}{(px+q)^n} + \frac{S}{ax^2+bx+c} + \frac{Z}{(ax^2+bx+c)^n}$$



MAYIS 2012						
H	P	S	C	P	C	Ct Pz
18		1	2	3	4	5 6
19	7	8	9	10	11	12 13
20	14	15	16	17	18	19 20
21	21	22	23	24	25	26 27
22	28	29	30	31		

MAYIS  
MAY  
SALI  
TUESDAY

15

K

$$\int \frac{3x^2+2x+3}{3x+2} dx = \left( x + \frac{3}{3x+2} \right) dx$$

$$= \int x dx + 3 \int \frac{1}{3x+2} dx$$

$$3x+2=t$$

$$= \frac{x^2}{2} + \ln|3x+2| + C$$

$$\int \frac{dx}{3x+2} = \int \frac{dt/3}{t}$$

$$= \frac{1}{3} \ln|t| + C$$

$$= \frac{1}{3} \ln|3x+2|$$

K

$$\int \frac{dx}{2-3x} = \int -\frac{dt/3}{t} = -\frac{1}{3} \ln|t| + C$$

$$2-3x=t$$

$$-3dx=dt$$

$$= -\frac{1}{3} \ln|2-3x| + C$$

FÖRÜMLÜ:

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln|ax+b| + C$$

$$\int \frac{1}{2-\sqrt{2}x} dx = -\frac{1}{\sqrt{2}} \ln|2-\sqrt{2}x| + C$$

$$\int \frac{dx}{x+t} = \ln|x+t| + C$$



# 16

MAYIS  
MAY

ÇARŞAMBA  
WEDNESDAY

MAYIS 2012						
H	P	S	C	P	C	Ct Pz
18		1	2	3	4	5 6
19	7	8	9	10	11	12 13
20	14	15	16	17	18	19 20
21	21	22	23	24	25	26 27
22	28	29	30	31		

$$\cancel{X} \int \frac{1}{(5x+2)^3} dx = \int \frac{1}{t^3} \frac{dt}{5} = \frac{1}{5} \int t^{-3} dt$$

$$\begin{aligned} 5x+2 &= t \\ 5dx &= dt \\ \Rightarrow &= \frac{1}{5} \frac{t^{-2}}{-2} = \frac{-1}{10t^2} + C \\ &= \frac{-1}{10(5x+2)^2} + C \end{aligned}$$

$$\cancel{X} \int \frac{2x+5+1-1}{x^2+5x+13} dx = \int \frac{2x+5}{x^2+5x+13} dx + \int \frac{-1}{x^2+5x+13} dx$$

$$\boxed{\begin{aligned} x^2+5x+13 &= t \\ (2x+5)dx &= dt \end{aligned}}$$

$$\textcircled{a} \int \frac{dt}{t} = \ln|t| = \ln|x^2+5x+13|$$

$$\textcircled{b} \int \frac{-1}{x^2+5x+13} dx = \int \frac{-1}{\left(x+\frac{5}{2}\right)^2 + \frac{27}{4}} dx$$

$$\int \frac{dx}{a^2+x^2} = \frac{1}{a} \arctan \frac{x}{a}$$

$$\boxed{\begin{aligned} x+\frac{5}{2} &= t \\ dx &= dt \end{aligned}}$$

$$= \int \frac{-dt}{t^2 + \left(\frac{3\sqrt{3}}{2}\right)^2} = \frac{-1}{3\sqrt{3}} \operatorname{arctan} \frac{t}{\frac{3\sqrt{3}}{2}}$$

$$2x+5=2t$$

$$= \frac{-2}{3\sqrt{3}} \operatorname{arctan} \left( \frac{2x+5}{3\sqrt{3}} \right)$$

cevap

$$I = \ln|x^2+5x+13| - \frac{2}{3\sqrt{3}} \operatorname{arctan} \left( \frac{2x+5}{3\sqrt{3}} \right) + C$$



MAYIS 2012							
H	P	S	Ç	P	C	Çt	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

MAYIS  
MAY

PERŞEMBE  
THURSDAY

17

$$\cancel{X} \int \frac{dx}{(4+x^2)^2} = I \quad \text{indirgenen bəzintis,}$$

$$\int \frac{1}{(4+x^2)} dx$$

$$\left[ \begin{array}{l} (4+x^2)^{-1} = v \\ -\frac{2x}{(4+x^2)^2} = dv \end{array} \right] \quad \begin{array}{l} dx = dv \\ x = v \end{array}$$

$$= \frac{1}{4+x^2} x - \int x \cdot \frac{-2x}{(4+x^2)^2} dx$$

$$= \frac{x}{4+x^2} + 2 \int \frac{x^2+4}{(4+x^2)^2} dx$$

$$= \frac{x}{4+x^2} + 2 \int \frac{\frac{1}{x^2+4}}{(4+x^2)^2} dx - 8 \int \frac{dx}{(4+x^2)^2}$$

$$8 \int \frac{1}{(4+x^2)^2} dx = \frac{x}{4+x^2} + \int \frac{dx}{x^2+4}$$

$$= \frac{x}{8(4+x^2)} + \frac{1}{16} \arctan \frac{x}{2} + C$$

$$\cancel{X} \cancel{v} \quad x = 2 \tan t$$

# 18

MAYIS

MAY

CUMA

FRIDAY

Fayda Garip olursa oysa yersa gittir

MAYIS							2012	
H	P	S	C	P	C	Ct	Pz	
18		1	2	3	4	5	6	
19	7	8	9	10	11	12	13	
20	14	15	16	17	18	19	20	
21	21	22	23	24	25	26	27	
22	28	29	30	31				

$$\cancel{\int} \int \frac{4}{x^2-4} dx \Rightarrow \frac{A}{x-2} + \frac{B}{x+2} = \frac{4}{x^2-4}$$

$$Ax + 2A + Bx - 2B = 4$$

$$A+B=0$$

$$2A - 2B = 4$$

$$A=1, B=-1$$

$$\int \frac{dx}{x-2} + \int \frac{-dx}{x+2}$$

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln(ax+b)$$

$$\ln|x-2| + \ln|x+2| + C = \ln\left|\frac{x-2}{x+2}\right| + C$$

Tümler

2

$$(x+1)(x-1)^3(3x^2+4)(x^2-4)(x^2+5x+13)^2 \cancel{(x^3+13)}$$

$$= \frac{A}{x+1} + \frac{B}{x-1} + \frac{C}{(x-1)^2} + \frac{D}{(x-1)^3} + \frac{Ex+F}{3x^2+4} + \frac{H}{x-2} + \frac{I}{x+2}$$

$$+ \frac{Jx+K}{x^2+5x+13} + \frac{Mx+N}{(x^2+5x+13)^2} + \frac{Zx^2+Yx+S}{x^3+13}$$

$$\cancel{\int} \int \frac{2x+1}{(x-1)^2} dx \quad \frac{2x+1}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

$$Ax - A + B = 2x + 1$$

$$A=2 \quad B=3$$

$$\int \frac{2}{x-1} dx + \int \frac{3}{(x-1)^2} dx$$

$$x-1=t \\ dx=dt$$

$$2 \ln|x-1| - \frac{3}{(x-1)} + C$$

$$\int \frac{3}{t^2} dt \\ = 3t^{-1}$$

$$= \frac{3}{(x-1)}$$



MAYIS							2012	
H	P	S	C	P	C	Ct	Pz	
18		1	2	3	4	5	6	
19	7	8	9	10	11	12	13	
20	14	15	16	17	18	19	20	
21	21	22	23	24	25	26	27	
22	28	29	30	31				

MAYIS

MAY

CUMARTESİ

SATURDAY

19

$$x^2 - 2x + 1$$

$$(x-1)x^2 + 1$$

$$x^3 + x - x^2 - 1$$

X

$$\int \frac{-2x+6}{(1+x^2)(x-1)^2} dx$$

$$\frac{-2x+6}{(x^2+1)(x-1)^2} = \frac{Ax+B}{x^2+1} + \frac{C}{x-1} + \frac{D}{(x-1)^2}$$

$$\begin{aligned} & (Ax^3 - Ax^2 + Ax) + (Bx^2 - 2Bx + B) \\ & (Cx^3 - Cx^2 + Cx) + (Dx^2 + D) \end{aligned}$$

$$A + B \neq 0$$

$$C = -2$$

$$-A + B - C + D = 0$$

$$D = 1$$

$$A - 2B + C = -2$$

$$A = 2$$

$$B - C + D = 4$$

$$B = 1$$

$$\int \frac{-2}{x-1} dx = -2 \ln|x-1|$$

MAYIS  
MAY

20

$$\int \frac{1}{(x-1)^2} dx = \int \frac{1}{t^2} dt = t^{-2} dt = \frac{t^{-1}}{-1} = -\frac{1}{t} = \frac{1}{x-1}$$

PAZAR

SUNDAY

$$\boxed{\begin{aligned} x-1 &= t \\ dx &= dt \end{aligned}}$$

$$\int \frac{2x+1}{1+x^2} dx = \int \frac{2x}{1+x^2} dx + \int \frac{1}{1+x^2} dx$$

$$\begin{aligned} & \boxed{\begin{aligned} 1+x^2 &= t \\ 2x dx &= dt \end{aligned}} \\ & = \int \frac{dt}{t} + \arctan x \\ & = \ln|t| + \arctan x \end{aligned}$$

$$= -2 \ln|x-1| - \frac{1}{x-1} + \ln|1+x^2| + \arctan x + C$$



# 21

MAYIS  
MAY

PAZARTESİ  
MONDAY

$$\frac{2x^4 - 6x^3 + 7x^2 - 2x - 2}{x^3 - 3x^2 + 3x - 1} \quad | \quad x^3 - 3x^2 + 3x - 1$$

MAYIS		2012						
H	P	S	C	P	C	Ct	Pz	
18		1	2	3	4	5	6	
19	7	8	9	10	11	12	13	
20	14	15	16	17	18	19	20	
21	21	22	23	24	25	26	27	
22	28	29	30	31				

$$\text{A} \int \frac{2x^4 - 6x^3 + 7x^2 - 2x - 2}{x^3 - 3x^2 + 3x - 1} dx$$

$$= \int \left[ 2x + \frac{x^2 - 2}{(x-1)^3} \right] dx$$

$$= \underbrace{\int 2x dx}_{x^2} + \underbrace{\int \frac{x^2 - 2}{(x-1)^3} dx}$$

$$\frac{x^2 - 2}{(x-1)^3} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{(x-1)^3}$$

$$Ax^2 - 2Ax + A + Bx - B + C = x^2 - 2$$

$$A = 1$$

$$B = 2$$

$$C = -1$$

$$\int \frac{dx}{x-1} + \int \frac{2dx}{(x-1)^2} - \int \frac{dx}{(x-1)^3}$$

$$x-1 \neq 0 \\ dx \neq 0$$

~~$$\int \frac{1}{x-1} dx + 2 \int \frac{1}{(x-1)^2} dx - \int \frac{1}{(x-1)^3} dx$$~~

$$\ln|x-1| - \frac{2}{x-1} + \frac{1}{2(x-1)^2}$$

$$= x^2 + \ln|x-1| - \frac{2}{x-1} + \frac{1}{2(x-1)^2} + C$$



MAYIS 2012

H	P	S	Ç	P	C	Cl	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

MAYIS

MAY

SALI

TUESDAY

22

$$\text{A} \int \frac{2}{x^2-1} dx$$

$$\frac{2}{x^2-1} = \frac{A}{x-1} + \frac{B}{x+1}$$

$$Ax + A + Bx - B = 2$$

$$A+B=0$$

$$A-B=2$$

$$A=1 \quad B=-1$$

$$\int \frac{dx}{x-1} - \int \frac{dx}{x+1}$$

$$\ln|x-1| - \ln|x+1| + C$$

$$\text{A} \int \frac{e^y dy}{e^{2y} + 3e^y + 2}$$

$$\frac{e^y}{(e^y+2)(e^y+1)} = \frac{A}{e^y+2} + \frac{B}{e^y+1}$$

$$A+B=1$$

$$2A+2B=0$$

$$A=-1$$

$$B=2$$

$$\cancel{\int \frac{e^y}{e^y+2} dy} + 2 \cancel{\int \frac{e^y}{e^y+1} dy} = \cancel{B} e^{-y} - A e^y$$

$$e^y dy = dx$$

$$\int \frac{1}{v^2 + 3v + 2} dv$$

$$\boxed{v = e^y}$$

$$\boxed{dv = e^y dy}$$

$$= \ln|v+1| - \ln|v+2| + C$$

$$= \ln \left| \frac{e^y+1}{e^y+2} \right| + C$$



# 23

MAYIS

MAY

ÇARŞAMBA  
WEDNESDAY

MAYIS		2012					
H	P	S	Ç	P	C	Ct	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

$$\int \frac{\sin \theta d\theta}{\cos^2 \theta + \cos \theta - 2}$$

$$= \int \frac{-du}{u^2 + u - 2}$$

$$\begin{matrix} +2 \\ -1 \end{matrix}$$

$$\left[ \begin{array}{l} \cos \theta = u \\ -\sin \theta d\theta = du \end{array} \right]$$

$$\frac{-1}{u^2 + u - 2} = \frac{A}{u+2} + \frac{B}{u-1}$$

$$Au - A + Bu + 2B = -1$$

$$A + B = 0$$

$$-A + 2B = -1$$

$$B = -\frac{1}{3}$$

$$A = \frac{1}{3}$$

$$= \frac{1}{3} \int \frac{du}{u+2} - \frac{1}{3} \int \frac{du}{u-1}$$

$$= \frac{1}{3} \ln |\cos \theta + 2| - \frac{1}{3} \ln |\cos \theta - 1| + C$$

$$= \frac{1}{3} \ln \left| \frac{\cos \theta + 2}{\cos \theta - 1} \right| + C$$

## TRİGONOMETRİK İNTEGRALLER

$$\sin ax \sin bx = -\frac{1}{2} [\cos(a+b)x - \cos(a-b)x]$$

$$\sin ax \cos bx = \frac{1}{2} [\sin(a+b)x + \sin(a-b)x]$$

$$\cos ax \cos bx = \frac{1}{2} [\cos(a+b)x + \cos(a-b)x]$$



MAYIS	2012						
H	P	S	Ç	P	C	Cl	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

(1) (3)  $\sin \alpha x \rightarrow (2)$

MAYIS  
PERŞEMBE  
THURSDAY

24

- $\cos(\alpha+b)x = \cos(\alpha x + b x)$   
 $= \cos(\alpha x) \cos(b x) - \sin(\alpha x) \sin(b x)$
- $\cos(\alpha-b)x = \cos(\alpha x - b x)$   
 $= \cos(\alpha x) \cos(b x) + \sin(\alpha x) \sin(b x)$
- $\cos(\alpha-b)x - \cos(\alpha+b)x = 2 \sin(\alpha x) \sin(b x)$

$$\frac{1}{2} [\cos(\alpha-b)x - \cos(\alpha+b)x] = \sin(\alpha x) \sin(b x)$$

$$\cancel{\int \sin 7x \sin 3x dx} = \int -\frac{1}{2} [\cos 10x - \cos 4x] dx$$

$$= -\frac{1}{2} \left[ \frac{1}{10} \sin 10x - \frac{1}{4} \sin 4x \right] + C$$

$$\int \cos(\alpha x) dx = \frac{1}{\alpha} \sin(\alpha x) + C$$

$$\cancel{\int \sin 5x \cos 3x dx} = \int \frac{1}{2} [\sin 8x + \sin 2x] dx$$

$$= \frac{1}{2} \left[ -\frac{1}{8} \cos 8x + \frac{1}{2} \cos 2x \right] + C$$

$$\int \sin \alpha x dx = -\frac{1}{\alpha} \cos \alpha x + C$$

$$\cancel{\int \cos 4x \cos 3x dx} = \int \frac{1}{2} [\cos 7x + \cos x] dx$$

$$= \frac{1}{2} \left[ \frac{1}{7} \sin 7x + \sin x \right] + C$$



# 25

MAYIS  
MAY  
CUMA  
FRIDAY

$\sin \rightarrow a$   
 $\cos \rightarrow b$

MAYIS 2012											
H	P	S	C	P	C	Ct	Pz				
18		1	2	3	4	5	6				
19	7	8	9	10	11	12	13				
20	14	15	16	17	18	19	20				
21	21	22	23	24	25	26	27				
22	28	29	30	31							

$$\star \int \sin 3x \cos 5x \, dx = \frac{1}{2} \int [\sin 8x + \sin(-2x)] \, dx$$

$$= \frac{1}{2} \int [\sin 8x - \sin 2x] \, dx$$

$$= \frac{1}{2} \left[ -\frac{1}{8} \cos 8x + \frac{1}{2} \cos 2x \right] + C$$

$$\textcircled{1} \int \sin^m x \cos^n x \, dx$$

İki rolle tek

(a)  $m \rightarrow \text{tek}$  ( $n \text{ çift}$ )  $\rightarrow \cos x = t$   $[m > n \quad \sin x = t]$

(b)  $n \rightarrow \text{tek}$  ( $m \text{ çift}$ )  $\rightarrow \sin x = t$   $[n > m \quad \cos x = t]$

(c)  $m$  çift ve  $n$  çift

$$\cos 2x = 1 - 2 \sin^2 x$$

$$\cos 2x = 2 \cos^2 x - 1$$

$$2 \sin^2 x = 1 - \cos 2x$$

$$\cos 2x + 1 = 2 \cos^2 x$$

$$\boxed{\sin^2 x = \frac{1 - \cos 2x}{2}}$$

$$\boxed{\cos^2 x = \frac{\cos 2x + 1}{2}}$$

$$\star \int \sin^5 x \cos^2 x \, dx$$

$m=5, n=2$

$$\boxed{\begin{aligned} \cos x &= t \\ -\sin x \, dx &= dt \end{aligned}}$$

$$= \int \sin^4 x \cdot \sin x \cos^2 x \, dx$$

$$= \int (1 - \cos^2 x)^2 \cos^2 x \sin x \, dx$$

$$= \int (1 - t^2)^2 t^2 (-dt)$$



MAYIS							2012	
H	P	S	C	P	C	Ct	Pz	
18		1	2	3	4	5	6	
19	7	8	9	10	11	12	13	
20	14	15	16	17	18	19	20	
21	21	22	23	24	25	26	27	
22	28	29	30	31				

MAYIS  
MAY  
CUMARTESİ  
SATURDAY

26

$$\begin{aligned}
 &= - \int (1 - 2t^2 + t^4) t^2 dt \\
 &= - \int (t^2 - 2t^4 + t^6) dt \\
 &= - \left[ \frac{t^3}{3} - 2 \frac{t^5}{5} + \frac{t^7}{7} \right] + C \\
 &= - \frac{1}{3} \cos^3 x + \frac{2}{5} \cos^5 x - \frac{1}{7} \cos^7 x + C
 \end{aligned}$$

~~A~~  $\int \sin^4 x \cos^3 x dx$   $m=4$   $n=3$

$$\begin{cases} t = \sin x \\ dt = \cos x dx \end{cases}$$

tek olur  
haricinde  
t de

$$= \int \sin^4 x \cos^2 x \cos x dx$$

$$= \int \sin^4 x (1 - \sin^2 x) \cos x dx$$

$$= \int t^4 (1 - t^2) dt$$

$$= \int (t^4 - t^6) dt = \frac{t^5}{5} - \frac{t^7}{7} + C = \frac{\sin^5 x}{5} - \frac{\sin^7 x}{7} + C$$

~~A~~  $\int \sin^3 x \cos^3 x dx$   $m=3$   $n=3$

$$\begin{cases} \sin x = t \\ \cos x dx = dt \end{cases}$$

Birçok olur t de

$$= \int \sin^3 x \cos^2 x \cos x dx$$

$$= \int t^3 (1 - t^2) dt$$

$$= \int (t^3 - t^5) dt$$

$$= \frac{t^{10}}{10} - \frac{t^{12}}{12} + C$$

$$= \frac{\sin^{10} x}{10} - \frac{\sin^{12} x}{12} + C$$



$(\cos 5x)^7 / (3x^2 + 5x + 6)^3$

**28**

MAYIS  
MAY  
PAZARTESİ  
MONDAY

$$1 - 2\cos 2x + \cos^2 2x$$

$$\cos 2x - 2\cos^2 2x + \cos^3 2x$$

**MAYIS**

**2012**

H	P	S	C	P	C	Ct	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

~~dx~~  $\int \sin^6 x \cos^2 x dx$

$$= \int \left[ \frac{1}{2} (1 - \cos 2x) \right]^2 \times \frac{1}{2} [1 + \cos 2x] dx$$

$$= \frac{1}{8} \int (1 - 2\cos 2x + \cos^2 2x)(1 + \cos 2x) dx$$

$$= \frac{1}{8} \int (\cancel{\cos^3 2x} - \cancel{\cos^2 2x} - \cancel{\cos 2x} + 1) dx$$

④  ~~$\int \cos 3x dx = x$~~

③  ~~$\int -\cos 2x dx = -\frac{1}{2} \sin 2x$~~

$$\begin{aligned} \textcircled{2} \int -\cos 2x dx &= \int -\frac{1}{2} [1 + \cos 4x] dx \\ &= -\frac{1}{2} \left[ x + \frac{1}{4} \sin 4x \right] \\ &= \cancel{-\frac{1}{2} x} - \frac{1}{8} \sin 4x \end{aligned}$$

$$\begin{aligned} \cos 2x &= 2\cos^2 x - 1 \\ \cos 4x &= 2\cos^2 2x - 1 \end{aligned}$$

①  $\int \cos^3 2x dx = \int \cos^3 2x \cdot \sin^0 2x dx$

$$\boxed{\begin{aligned} \sin 2x &= t \\ 2\cos 2x dx &= dt \end{aligned}}$$

$$= \int \cos^2 2x \cos 2x dx$$

$$= \int (-\sin^2 2x) \cos 2x dx$$

$$= \int (1 - t^2) dt / 2$$

$$= \frac{1}{2} \left[ t - \frac{t^3}{3} \right] + C = \cancel{\frac{1}{2} \left[ \sin 2x - \frac{\sin^3 2x}{3} \right]}$$



MAYIS 2012						
H	P	S	C	P	C	Pz
18		1 2 3 4	5 6			
19	7	8 9 10 11	12 13			
20	14	15 16 17	18 19 20			
21	21	22 23 24	25 26 27			
22	28	29 30 31				

MAYIS  
MAY  
SALI  
TUESDAY

29

$$= \frac{1}{8} \left[ \frac{x}{2} - \frac{1}{8} \sin 6x - \frac{\sin 32x}{6} \right] + C$$

②

$$\int \tan^m x \sec^n x dx$$

a)

n çift

r se

$$\boxed{\tan x = t}$$

b)

m tek

r se

$$\boxed{\sec x = t}$$

c)

m çift

n tek

r se

$$\boxed{\tan^2 x = \sec^2 x - 1}$$

deği. kalkınan cüz



$$\int \tan^6 x \sec^4 x dx$$

$$\boxed{\tan x = t}$$

$$\sec^2 x dx = dt$$

$$= \int \tan^6 x \sec^2 x \sec^2 x dx$$

$$= \int \tan^6 x (1 + \tan^2 x) \sec^2 x dx$$

$$= \int t^6 (1 + t^2) dt$$

$$= \int (t^6 + t^8) dt = \frac{t^7}{7} + \frac{t^9}{9} + C$$

$$= \frac{\tan^7 x}{7} + \frac{\tan^9 x}{9} + C$$



$t^4 - 2t^2 + 1$

# 30

MAYIS  
MAY  
ÇARŞAMBA  
WEDNESDAY

$$\tan^2 x = \sec^2 x - 1$$

MAYIS		2012					
H	P	S	Ç	P	C	Ct	Pz
18		1	2	3	4	5	6
19	7	8	9	10	11	12	13
20	14	15	16	17	18	19	20
21	21	22	23	24	25	26	27
22	28	29	30	31			

~~A~~  $\int \tan^5 x \sec^3 x dx$

$$\sec x = t$$

$$\sec x \tan x dx = dt$$

$$= \int \tan^4 x \sec^2 x \sec x \tan x dx$$

$$= \int (\sec^2 x - 1)^2 \sec^2 x \sec x \tan x dx$$

$$= \int (t^2 - 1)^2 t^2 dt$$

$$= \int (t^6 - 2t^4 + t^2) dt$$

$$= \frac{t^7}{7} - 2 \frac{t^5}{5} + \frac{t^3}{3} + C$$

$$= \frac{\sec^7 x}{7} - 2 \frac{\sec^5 x}{5} + \frac{\sec^3 x}{3} + C$$

~~A~~  $\int \tan^2 x \sec x dx$

$$\tan^2 x = \sec^2 x - 1$$

$$= \int (\sec^2 x - 1) \sec x dx$$

$$= \int (\sec^3 x - \sec x) dx$$

$$= \underbrace{\int \sec^3 x dx}_b - \underbrace{\int \sec x dx}_a$$

$\sec^3 x$  in tek konuvtelennin integrali  
öynü yellow le yapılsı

①  $\int \sec x dx = \int \frac{(\sec^2 x + \sec x \tan x)}{\sec x + \tan x} dx$

$$(\sec x + \tan x)$$



(A'ye 3' dere)

$\int \cosec x dx$

$(\cosec x + \cot x)$

MAYIS							2012	
H	P	S	C	P	C	Ct	Pz	
18		1	2	3	4	5	6	
19	7	8	9	10	11	12	13	
20	14	15	16	17	18	19	20	
21	21	22	23	24	25	26	27	
22	28	29	30	31				

MAYIS  
MAY  
PERŞEMBE  
THURSDAY

31

$$= \int \frac{(\sec^2 x + \sec x \tan x) dx}{\sec x + \tan x}$$

$$\sec x + \tan x = t$$

$$(\sec^2 x + \sec x \tan x) dx$$

$$= \int \frac{dt}{t} = \ln |t| + C = \ln |\sec x + \tan x|$$

(b)  $\int \sec^3 x dx - I$

(2'inci kuvvet ayrılr.)

$$\int \underbrace{\sec x}_{u} \underbrace{\sec^2 x dx}_{dv}$$

$$\begin{cases} \sec x = u \\ \sec x \tan x dx = dv \\ \tan x \neq v \end{cases}$$

$$= \sec x \tan x - \int \tan^2 x \sec x dx$$

$$\boxed{\tan^2 x = \sec^2 x - 1}$$

$$= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx$$

$$= \sec x \tan x - \int (\sec^3 x - \sec x) dx$$

$$I = \sec x \tan x - I + \int \sec x dx$$

$$I = \frac{1}{2} \left[ \sec x \tan x + \ln |\sec x + \tan x| \right]$$

Çözüm

$$= \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| + C$$



# 01

HAZİRAN  
JUNE  
CUMA  
FRIDAY

HAZİRAN 2012						
H	P	S	Ç	P	C	Cl
22					1	2 3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

(3)  $\int \cot^n x \csc^m x dx$

(a)  $n$  çift ise  $\cot x = t$

(b)  $m$  tek ise  $\csc x = t$

(c)  $m$  çift  
 $n$  tek ise  $\cot^2 x = \csc^2 x - 1$

~~(3)~~  $\int \cot^2 x \csc^4 x dx$

$$\begin{cases} \cot x = t \\ -\csc^2 x dx = dt \end{cases}$$

$$= \int \cot^2 x (\cot^2 x + 1) \csc^3 x dx \quad (\cot x)^2 = -(1+dt)$$

$$= \int t^2 (t^2 + 1) dt$$

$$= - \int (t^6 + t^2) dt = - \left[ \frac{t^7}{7} + \frac{t^3}{3} \right] + C \quad \begin{cases} \cot^2 x = \csc^2 x - 1 \\ \csc^2 x dx = dt \end{cases}$$

$$= - \left[ \frac{\cot^7 x}{7} + \frac{\cot^5 x}{5} \right] + C$$

~~(3)~~  $\int \cot^5 x \csc^5 x dx$

$$\begin{cases} \csc x = t \\ -\csc x \cot x dx = dt \end{cases}$$

$$= \int \cot^4 x \csc^4 x \cot x \csc x dx$$

$$= \int (\csc^2 x - 1)^2 \csc^4 x \cot x \csc x dx$$

$$= \int (t^2 - 1)^2 \cdot t^4 dt$$

$$= \int (t^8 + 2t^6 - t^4) dt$$



HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22					1	2 3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

$$\frac{x^2 - 1}{x^4 - 2x^2 + 1}$$

HAZİRAN  
JUNE  
CUMARTESİ  
SATURDAY

02

$$\int \cot^4 x \cosec x dx$$

$$\cot^2 x = \cosec^2 x - 1$$

$$\int (\cosec^2 x - 1)^2 \cosec x dx$$

$$\int (\cosec^5 x - 2\cosec^3 x + \cosec x) dx$$

c

b

a

$$\textcircled{a} \int \cosec x dx = \int \frac{(\cosec^2 x + \cosec x \cot x) dx}{\cosec x + \cot x}$$

$$= \int -\frac{dt}{t} = -\ln|t|$$

$$= -\ln|\cosec x + \cot x|$$

$$(\cosec x + \cot x) = t$$

$$(-\cosec x \cot x - \cosec^2 x) dx = dt$$

$$\textcircled{b} \int \cosec x \cosec^2 x dx$$

HAZİRAN  
JUNE  
PAZAR  
SUNDAY

03

$$= -\cosec x \cot x + \int \cot^2 x \cosec x dx$$

$$\cosec x = u$$

$$\cosec^2 x dx = du$$

$$-\cot x = v$$

$$= -\cosec x \cot x - \int (\cosec^2 x - 1) \cosec x dx$$

$$= -\cosec x \cot x - \int \cosec^3 x + \int \cosec x dx$$

$$I = \frac{1}{2} \left[ -\cosec x \cot x - \ln|\cosec x + \cot x| \right]$$

$$2 \int \cosec^2 x dx = -\cosec x \cot x - \ln|\cosec x + \cot x|$$



# 04

HAZİRAN

JUNE

PAZARTESİ  
MONDAY

HAZİRAN 2012					
H	P	S	Ç	P	C
22					1 2 3
23	4	5	6	7	8 9 10
24	11	12	13	14	15 16 17
25	18	19	20	21	22 23 24
26	25	26	27	28	29 30

$$\textcircled{2} \int \csc^5 x dx = \int \underbrace{\csc^3 x}_{v} \underbrace{\csc^2 x dx}_{du}$$

$$-\csc^3 x \cot x - \int (-\cot x) (-3 \csc^3 x \cot x) dx$$

$$-\csc^3 x \cot x - 3 \int \cot^2 x \csc^3 x dx$$

$$\csc^2 x - 1$$

$$\csc^3 x du$$

$$3 \csc^2 x (-\csc x \cot x) dx$$

$$\csc^3 x dx = du$$

$$-\cot x = v$$

$$-\csc^3 x \cot x - 3 \int (\csc^5 x - \csc^3 x) dx$$

$$-\csc^3 x \cot x - 3 \int \csc^5 x dx + 3 \int \csc^3 x dx$$

$$\int \csc^5 x dx = \frac{1}{6} \left[ -\csc^3 x \cot x + \frac{3}{2} \int -\csc x \cot x - \ln |\csc x + \cot x| \right]$$

$$= \frac{1}{6} \csc^3 x \cot x - \frac{3}{8} \csc x \cot x - \frac{3}{8} \ln |\csc x + \cot x|$$

Cevap

$$= -\frac{1}{6} \csc^3 x \cot x - \frac{3}{8} \csc x \cot x - \frac{3}{8} \ln |\csc x + \cot x|$$

$$+ \csc x \cot x + \ln |\csc x + \cot x| - \ln |\csc x + \cot x| + C$$

$$= -\frac{1}{6} \csc^3 x \cot x + \frac{5}{8} \csc x \cot x - \frac{3}{8} \ln |\csc x + \cot x| + C$$



HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22				1	2	3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

HAZİRAN  
JUNE  
SALI  
TUESDAY

05

(4)  $\int R(\sin x, \cos x) dx$

①  $R(\sin x, \cos x) = -R(\sin x, \cos x)$   
 $\boxed{\sin x = t}$

②  $R(-\sin x, \cos x) = -R(\sin x, \cos x)$   
 $\boxed{\cos x = t}$

③  $R(-\sin x, -\cos x) = R(\sin x, \cos x)$   
 $\boxed{\sin x = t}$

A  $\int \frac{\sin^3 x}{\cos^2 x} dx$        $\cos x = t$   
 $-\sin x dx = dt$

 $= \int \frac{\sin^2 x \sin x}{\cos^2 x} dx$ 
 $= \int \frac{(1 - \cos^2 x) \sin x}{\cos^2 x} dx = \int \frac{(1 - t^2) \cdot (-dt)}{t^2} = \int \left(1 - \frac{1}{t^2}\right) dt$ 
 $= \int dt - \int t^{-2} dt$ 
 $= t - \frac{t^{-1}}{-1} + C$ 
 $= t + \frac{1}{t} + C$ 
 $= \cos x + \frac{1}{\cos x} + C$



# 06

HAZİRAN

JUNE

ÇARŞAMBA  
WEDNESDAY

HAZİRAN 2012						
H	P	S	Ç	P	C	Cı
22					1	2 3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

$$\begin{aligned}
 & \text{K} \int \sin^3 x \cot^5 x dx = \int \sin^3 x \frac{\cos^5 x}{\sin^5 x} dx \\
 & = \int \frac{\cos^5 x}{\sin^2 x} dx \\
 & = \int \frac{(1 - \sin^2 x)^2 \cos x dx}{\sin^2 x} \\
 & = \int \frac{(1 - t^2)^2 dt}{t^2} = \int \frac{(1 - 2t^2 + t^4) dt}{t^2} \\
 & = \int \frac{1}{t^2} dt - 2 \int dt + \int t^2 dt \\
 & = -\frac{1}{t} - 2t + \frac{t^3}{3} + C \\
 & = -\frac{1}{\sin x} - 2\sin x + \frac{\sin^3 x}{3} + C
 \end{aligned}$$

$$\text{K} \int \frac{1}{\sin^2 x \cos^n x} dx$$

$$\tan x = t$$

$$\begin{aligned}
 & \cancel{\int dx} = dt \\
 & \cancel{\cos^2 x}
 \end{aligned}$$

$$(1 + \tan^2 x) dt = dt$$

$$dx = \frac{dt}{(1 + \tan^2 x)}$$

$$dx = \frac{dt}{1 + t^2}$$

$$\cos x = \frac{1}{\sqrt{1+t^2}}$$

$$\sin x = \frac{t}{\sqrt{1+t^2}}$$



HAZİRAN 2012

H	P	S	Ç	P	C	Ct	Pz
22					1	2	3
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN

JUNE

PERŞEMBE  
THURSDAY

07

ABD 758

$$\begin{aligned}
 &= \int \frac{1}{\frac{t^2}{1+t^2} \cdot \frac{1}{(1+t^2)^2}} \cdot \frac{dt}{1+t^2} \\
 &= \int \frac{1+2t^2+t^4}{t^2} dt \\
 &= \int (t^{-2} + 2 + t^2) dt \\
 &= -\frac{1}{t} + 2t + \frac{t^3}{3} + C
 \end{aligned}$$

~~$$\int \sin \frac{7x}{2} \sin \frac{x}{2} dx$$~~

$$\begin{aligned}
 &= \int -\frac{1}{2} [\cos 6x - \cos 3x] dx \\
 &= -\frac{1}{2} \int \cos 6x dx + \frac{1}{2} \int \cos 3x dx \\
 &= -\frac{1}{8} \sin 6x - \frac{1}{6} \sin 3x + C
 \end{aligned}$$

~~$$\int \sin^3 x \cos^2 x dx$$~~

$\cos x = t$   
 $-\sin x dx = dt$

$$\int \sin^2 x \cos^2 x \sin x dx$$

$$\int (1 - \cos^2 x) \cos^2 x \sin x dx$$

$$\int (1 - t^2) t^2 (-dt)$$



# 08

**HAZİRAN**

JUNE

**CUMA**

FRIDAY

**HAZİRAN 2012**

H	P	S	Ç	P	C	Ct	Pz
22						1	2 3
23	4	5	6	7	8	9 10	
24	11	12	13	14	15	16 17	
25	18	19	20	21	22	23 24	
26	25	26	27	28	29	30	

$$\cancel{\star} \int \tan^3 x \sec^4 x dx$$

$$\left. \begin{array}{l} \sec x = t \\ \sec x \tan x dx = dt \end{array} \right\}$$

$$= \int \tan^2 x \sec^3 x \sec x \tan x \sec x dx$$

$$= \int (\cancel{\sec^2 x} - 1) \sec^3 x \sec x \tan x \sec x dx$$

$$= \int (t^2 - 1) t^3 dt$$

$$= \int (t^5 - t^3) dt$$

$$= \frac{t^6}{6} - \frac{t^4}{4} + C = \frac{\sec^6 x}{6} - \frac{\sec^4 x}{4} + C$$

$$\cancel{\star} \int \cot^4 x \cosec^6 x dx$$

$$\begin{aligned} \cot x &= t \\ -\cosec^2 x dx &= dt \end{aligned}$$

$$= \int \cot^4 x \cosec^4 x \cosec^2 x dx$$

$$= \int \cot^4 x (\cot^2 x + 1)^2 \cosec^2 x dx$$

$$= \int t^4 (t^2 + 1)^2 (-dt)$$

$$= - \int (t^8 + 2t^6 + t^4) dt$$

$$= - \left[ \frac{t^9}{9} + 2 \frac{t^7}{7} + \frac{t^5}{5} \right] + C$$

$$= -\frac{1}{9} \cot^9 x + \frac{2}{7} \cot^7 x + \frac{1}{5} \cot^5 x + C$$



**HAZİRAN 2012**


H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

$$\frac{t^2(5+4t^2)}{1+t^2} \cdot \frac{dt}{1+t^2}$$

$5t^2 + 4t^4$   
**HAZİRAN**  
 $1 + 2t^2 + t^4$   
**JUNE**  
**CUMARTESİ**  
**SATURDAY**

**09**

$$\int \frac{\sin^4 x}{\cos^2 x} dx$$



$$\sin x = \frac{t}{\sqrt{1+t^2}}$$

$$\cos x = \frac{1}{\sqrt{1+t^2}}$$

$$= \int \frac{\frac{t^4}{(1+t^2)^2}}{4 + \frac{1}{1+t^2}} \cdot \frac{dt}{1+t^2}$$

$$= \int \frac{t^4}{(1+t^2)^3} \cdot \frac{1+t^2}{5+4t^2} dt$$

$$= \int \frac{t^4}{(1+t^2)^2 (5+4t^2)} dt$$

$$= \int \frac{t^4}{4t^6 + 13t^4 + 16t^2 + 5} dt$$

$$\begin{aligned} \tan x &= t \\ (1+\tan^2 x) dx &= dt \\ dx &= \frac{dt}{1+\tan^2 x} \\ dx &= \frac{dt}{1+t^2} \end{aligned}$$

**HAZİRAN**  
**JUNE**  
**PAZAR**  
**SUNDAY**

**10**



# 11

HAZİRAN  
JUNE

PAZARTESİ  
MONDAY

HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22					1	2 3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

$$\sin^4 x = (1 - \cos^2 x)^2 = 1 - 2\cos^2 x + \cos^4 x$$

$$\frac{d(-2\cos^2 x + \cos^4 x)}{\cos^2 x - 6} \Big|_{25}$$

$$\int \left( (\cos^2 x - 6) + \frac{25}{6 + \cos^2 x} \right) dx$$

$$\cos 2x = 2\cos^2 x - 1$$

$$\int \underbrace{\frac{1}{2} [\cos 2x + 1]}_1 dx - \underbrace{\int 6 dx}_2 + \underbrace{\int \frac{25}{4 + \cos^2 x} dx}_3$$

$$\textcircled{1} \quad \frac{1}{2} \left[ \frac{1}{2} \sin 2x + x \right]$$

$$\textcircled{2} \quad -6x$$

$$\textcircled{3} \quad \int \frac{25}{6t + \frac{1}{1+t^2}} \cdot \frac{dt}{1+t^2} = \int \frac{25}{5+6t^2} dt = \int \frac{\frac{25}{2} du}{5+u^2}$$

$$= \frac{25}{2} \left[ \frac{1}{\sqrt{5}} \arctan \frac{u}{\sqrt{5}} \right] = \frac{25}{2} \left[ \frac{1}{\sqrt{5}} \arctan \frac{2t}{\sqrt{3}} \right]$$

$$\text{Cevap} \quad = \frac{1}{5} \sin 2x + \frac{x}{2} - 6x + \frac{25}{2\sqrt{2}} \arctan \frac{2(\tan x)}{\sqrt{3}} + C$$



HAZİRAN 2012

H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN

JUNE

SALI

TUESDAY

12

## İBRAS YÖNTEL FONKSİYONLARIN İNTEGRALİ

(1)

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}}$$

(a)  $ax^2 + bx + c = k^2 - v^2$   $k = \sqrt{-c}$   $v = x'$   $f(x)$

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}} = \int \frac{dx}{\sqrt{k^2 - v^2}} = \arcsin \frac{v}{k} + C$$

(b)  $ax^2 + bx + c = v^2 - p$   $s = \sqrt{-p}$

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}} = \int \frac{dv}{\sqrt{v^2 - p}} = \ln |v + \sqrt{v^2 - p}| + C$$

~~$\int \frac{dx}{\sqrt{-x^2 + 2x + 3}}$~~

$$-x^2 + 2x + 3 = -(x^2 - 2x - 3) = -(x-1)^2 + 4$$

$$= \int \frac{dx}{\sqrt{2^2 - (x-1)^2}}$$

$$\begin{cases} x-1 = u \\ dx = du \end{cases}$$

$$= \int \frac{du}{\sqrt{2^2 - u^2}}$$

$$= \arcsin \frac{u}{2} + C$$

$$= \arcsin \left( \frac{x-1}{2} \right) + C$$



# 13

HAZİRAN  
JUNE  
ÇARŞAMBA  
WEDNESDAY

HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22						1 2 3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

$$\int \frac{dx}{\sqrt{6x^2 + 6x + 3}}$$

$$6x^2 + 6x + 3 = (2x+1)^2 + 2$$

$$= \int \frac{dx}{\sqrt{(2x+1)^2 + 2}}$$

$$\begin{cases} 2x+1 = u \\ 2dx = du \end{cases}$$

$$= \int \frac{du/2}{\sqrt{u^2+2}} = \frac{1}{2} \ln |u + \sqrt{u^2+2}| + C$$

$$= \frac{1}{2} \ln |2x+1 + \sqrt{(2x+1)^2+2}| + C$$

$$\int \frac{dx}{\sqrt{x^2 - 2x + 3}}$$

$$x^2 - 2x + 3 = (x-1)^2 + 2$$

$$= \int \frac{dx}{\sqrt{(x-1)^2 + 2}}$$

$$\begin{cases} x-1 = u \\ dx = du \end{cases}$$

$$= \int \frac{du}{\sqrt{u^2+2}} = \frac{1}{2} \ln |u + \sqrt{u^2+2}| + C$$

$$= \frac{1}{2} \ln |x-1 + \sqrt{x^2 - 2x + 3}| + C$$



HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22				1	2	3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

HAZİRAN  
JUNE  
PERŞEMBE  
THURSDAY

14

$$\int \frac{dx}{\sqrt{2x^2 - 8x - 6x^2}}$$

$$-4x^2 + 8x - 21 = (2x+2)^2$$

-25

$$= \int \frac{dx}{\sqrt{s^2 - (2x+2)^2}}$$

$$\begin{cases} 2x+2 = u \\ 2dx = du \end{cases}$$

$$= \int \frac{du/2}{\sqrt{s^2 - u^2}} = \frac{1}{2} \operatorname{arc sm} \left( \frac{u}{s} \right) + C$$

$$\frac{1}{2} \operatorname{arc sm} \left( \frac{2x+2}{s} \right) + C$$

$$\int \frac{dx}{\sqrt{x^2 - 6x + 5}}$$

$$x^2 - 6x + 5 = (x-2)^2 + 1$$

$$= \int \frac{dx}{\sqrt{(x-2)^2 + 1}}$$

$$\begin{cases} x-2 = u \\ dx = du \end{cases}$$

$$= \int \frac{du}{\sqrt{u^2 + 1}} = \ln |u + \sqrt{u^2 + 1}| + C$$

$$= \ln |x-2 + \sqrt{x^2 - 6x + 5}| + C$$

$$\int \frac{dx}{\sqrt{6x^2 - 6x + 5}}$$

$$6x^2 - 6x + 5 = (2x-1)^2 + 4$$

$$= \int \frac{dx}{\sqrt{(2x-1)^2 + 4}}$$

$$\begin{cases} 2x-1 = u \\ 2dx = du \end{cases}$$

$$= \int \frac{du/2}{\sqrt{u^2 + 4}}$$

$$= \frac{1}{2} \ln |u + \sqrt{u^2 + 4}| + C$$

$$= \frac{1}{2} \ln |2x-1 + \sqrt{6x^2 - 6x + 5}| + C$$



# 15

HAZİRAN  
JUNE  
CUMA  
FRIDAY

HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22					1	2 3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

$$\textcircled{2} \int \frac{mx+n}{\sqrt{ax^2+bx+c}} dx$$

$$\boxed{dx^2+bx+c=t}$$

$$(2ax+b)dx=dt$$

ile  
 $(mx+n)dx$  sonrası bire bir

~~$\int \frac{3x+2}{\sqrt{x^2+bx+c}} dx$~~

$$x^2+bx+c=t$$

$$(2x+b)dx=dt$$

$$\begin{aligned} 3x+2 &= 3\left[x + \frac{b}{3}\right] \\ &= \frac{3}{2}\left[2x + \frac{b}{3}\right] \\ &= \frac{3}{2}\left[2x + b + \frac{b}{3} - b\right] \\ &= \frac{3}{2}\left[2x + b - \frac{8}{3}\right] \\ &= \frac{3}{2}\left[2x + b\right] - 4 \end{aligned}$$

$$= \int \frac{\frac{3}{2}(2x+b)-4}{\sqrt{x^2+bx+c}} dx = \frac{3}{2} \int \frac{2x+b}{\sqrt{x^2+bx+c}} dx - 4 \int \frac{1}{\sqrt{x^2+bx+c}} dx$$

$$\textcircled{2} \frac{3}{2} \int \frac{2x+b}{\sqrt{x^2+bx+c}} dx = \frac{3}{2} \int \frac{dt}{\sqrt{t}} + \frac{3}{2} \frac{t^{1/2}}{1/2}$$

$$= 3\sqrt{t} = 3\sqrt{x^2+bx+c}$$



HAZİRAN 2012

H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN

JUNE

CUMARTESİ

SATURDAY

16

$$(2) -6 \int \frac{dx}{\sqrt{x^2 + 6x + 1}}$$

$$x^2 + 6x + 1 = (x+2)^2 - 3$$

$$-6 \int \frac{1}{\sqrt{(x+2)^2 - 3}} dx$$

$$\begin{cases} x+2 = u \\ dx = du \end{cases}$$

$$-6 \int \frac{du}{\sqrt{u^2 - 3^2}} = -6 \left( \ln \left| u + \sqrt{u^2 - 3^2} \right| \right) = -6 \left( \ln \left| x+2 + \sqrt{x^2 + 6x + 1} \right| \right)$$

~~$$= 3 \sqrt{x^2 + 6x + 1} - 6 \ln \left| x+2 + \sqrt{x^2 + 6x + 1} \right| + C$$~~

~~$$\int \frac{x+3}{\sqrt{x^2 + 2x + 2}} dx$$~~

$$x^2 + 2x + 2 = t$$

$$(2x+2)dx = 2(x+1)dx$$

~~$$\begin{aligned} x+3 &= \frac{1}{2}[2x+6] \\ &= \frac{1}{2}[2x+2+4] \end{aligned}$$~~

HAZİRAN

JUNE

PAZAR

SUNDAY

17

$$-\int \frac{(x+1)+2}{\sqrt{x^2 + 2x + 2}} dx = \int \frac{(x+1)dx}{\sqrt{x^2 + 2x + 2}} + 2 \int \frac{dx}{\sqrt{x^2 + 2x + 2}}$$

①                            ②

$$\textcircled{1} \quad \int \frac{dt/2}{\sqrt{t^1}} = \frac{1}{2} \int t^{-1/2} dt = \frac{1}{2} \frac{t^{1/2}}{\frac{1}{2}} = \sqrt{t} = \sqrt{x^2 + 2x + 2}$$



# 18

HAZİRAN

JUNE

PAZARTESİ  
MONDAY

HAZİRAN

2012

H	P	S	Ç	P	C	Ct	Pz
22						1	2 3
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

②

$$2 \int \frac{dx}{\sqrt{x^2 + 2x + 2}} = 2 \int \frac{dx}{\sqrt{(x+1)^2 + 1}}$$

$$\begin{cases} x+1 = u \\ dx = du \end{cases}$$

$$= 2 \int \frac{du}{\sqrt{u^2 + 1}} = 2 \left[ \ln \left| u + \sqrt{u^2 + 1} \right| \right] = 2 \left[ \ln \left| x+1 + \sqrt{x^2 + 2x + 1} \right| \right]$$

Larv

$$\sqrt{x^2 + 2x + 1} + 2 \left[ \ln \left| x+1 + \sqrt{x^2 + 2x + 1} \right| \right] + C$$

③

$$\int \frac{dx}{(px+q)\sqrt{ax^2 + bx + c}}$$

$$\boxed{\frac{1}{px+q} = t}$$

$$\cancel{K} \int \frac{dx}{(x-1)\sqrt{x^2 + 3}}$$

$$\boxed{\frac{1}{x-1} = t}$$

$$\frac{1}{t} = x-1$$

$$\frac{t}{t} + 1 = x$$

$$-\frac{t+1}{t} = x$$

$$\boxed{\frac{-1}{t^2} dt = dx}$$

$$x^2 = \frac{t^2 + 2t + 1}{t^2}$$

$$x^2 + 3 = \frac{t^2 + 2t + 1}{t^2}$$

$$= \int \frac{t - \frac{1}{t^2} dt}{\sqrt{\frac{at^2 + 2bt + c}{t^2}}}$$

$$= \int \frac{-dt}{t + \sqrt{\frac{at^2 + 2bt + c}{t^2}}}$$

$$= \int \frac{-dt}{\sqrt{\frac{at^2 + 2bt + c}{t^2}}}$$



HAZİRAN 2012

H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN

JUNE

SALI

TUESDAY

19

$$2t^2 - 2t + t^2$$

$$= \int \frac{-dt}{\sqrt{\left(2t+\frac{1}{2}\right)^2 + \frac{3}{4}}} = \int \frac{-dt/2}{\sqrt{v^2 + 3/h^2}} = -\frac{1}{2} h \left[ v + \sqrt{v^2 + \frac{3}{h^2}} \right] + C$$

$$\boxed{2t + \frac{1}{2} = v \\ 2dt = dv}$$

$$= -\frac{1}{2} h \ln \left[ 2t + \frac{1}{2} + \sqrt{h^2 + 2t + 1} \right] + C$$

$$= -\frac{1}{2} h \ln \left[ \frac{2}{x-1} + \frac{1}{2} + \sqrt{h \left( \frac{1}{x-1} \right)^2 + \frac{2}{x-1} + 1} \right] + C$$

$$\cancel{\int \frac{dx}{(x+1)\sqrt{x^2 - 2x + 2}}}$$

$$\frac{1}{x+1} = t$$

$$\frac{1}{t} = x+1$$

$$\frac{1}{t} - 1 = x$$

$$\frac{1-t}{t} = x$$

$$\boxed{-\frac{1}{t^2} dt = dx}$$

$$x^2 = \frac{t^2 + 2t + 1}{t^2} = \frac{t - 2t + t^2}{t^2}$$

$$-2x = \frac{2t - 2}{t}$$

$$+2$$

$$= \int t \frac{-dt}{\sqrt{\frac{t^2 + 2t + 1}{t^2} + \frac{2t - 2}{t} + 2}} = \int \frac{-dt}{\sqrt{5t^2 - 6t + 1}}$$

$$= \int \frac{-dt}{\sqrt{5t^2 - 6t + 1}}$$

$$= \int \frac{-dt}{\sqrt{\left(\sqrt{5}t - \frac{2}{\sqrt{5}}\right)^2 + \frac{1}{5}}} = \int \frac{-dt/\sqrt{5}}{\sqrt{v^2 + 1/5}}$$

$$= \int \frac{-dv/\sqrt{5}}{\sqrt{v^2 + 1/5}}$$

$$\boxed{\sqrt{5}t - \frac{2}{\sqrt{5}} = v \\ \sqrt{5}dt = dv}$$

$$= -\frac{1}{\sqrt{5}} \ln \left[ v + \sqrt{v^2 + 1/5} \right] + C$$

$$= -\sqrt{5} h \left[ \sqrt{5}t - \frac{2}{\sqrt{5}} + \sqrt{5t^2 - 6t + 1} \right] + C$$



1  
X

# 20

HAZİRAN  
JUNE  
ÇARŞAMBA  
WEDNESDAY

$x^2, x^3, \dots$

$x^2+bx+c$

BN dence  
dogru

Poyda

H	P	S	C	P	C	Ct	Pz
22						1	2 3
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

$$\textcircled{1} \int \frac{P_n(x)}{\sqrt{dx^2 + bx + c}} dx = Q_{n-1}(x) \sqrt{dx^2 + bx + c} + \lambda \int \frac{1}{\sqrt{dx^2 + bx + c}} dx$$

Bu formü geneldir, sonda her bir fonksiyon formesi olinsotk bilmeceler bulur. A) Bilmeceler sağ formata yerine yofur.

$$\star \int \frac{x^2}{\sqrt{1-x^2}} dx = (Ax+B)\sqrt{1-x^2} + \lambda \int \frac{1}{\sqrt{1-x^2}} dx$$

$$\frac{x^2}{\sqrt{1-x^2}} = A\sqrt{1-x^2} + (Ax+B) \frac{-x}{\sqrt{1-x^2}} + \frac{\lambda}{\sqrt{1-x^2}}$$

$$x^2 = A(1-x^2) - x(Ax+B) + \lambda$$

$$x^2 = A - Ax^2 - Ax^2 + Bx + \lambda$$

$$x^2 = x^2[-2A] + x[-B] + [A+\lambda]$$

$$A = -\frac{1}{2}, \quad B = 0, \quad \lambda = \frac{1}{2}$$

$$= \left(-\frac{1}{2}x\right)\sqrt{1-x^2} + \frac{1}{2} \int \frac{dx}{\sqrt{1-x^2}}$$

$$= \left(-\frac{1}{2}x\right)\sqrt{1-x^2} + \frac{1}{2} \arcsin x + C$$



HAZİRAN		2012					
H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

$$(x^2+4)^{1/2} = \frac{1}{2} (x^2+4)^{-1/2} \cdot (2x)$$

HAZİRAN

JUNE

PERŞEMBE

THURSDAY

21

$$\int \frac{x^4 + 4x^2}{\sqrt{x^2+4}} dx = (Ax^3 + Bx^2 + Cx + D)\sqrt{x^2+4} + \lambda \int \frac{1}{\sqrt{x^2+4}} dx$$

$$\frac{x^4 + 4x^2}{\sqrt{x^2+4}} = (3Ax^2 + 2Bx + C)\sqrt{x^2+4} + (Ax^3 + Bx^2 + Cx + D) \frac{x}{\sqrt{x^2+4}}$$

$$x^4 + 4x^2 = (3Ax^2 + 2Bx + C)(x^2 + 4) + (Ax^3 + Bx^2 + Cx + D)x + \lambda$$

$$x^4 + 4x^2 = 3Ax^4 + 2Bx^3 + Cx^2 + 12Ax^2 + 8Bx + 4C + Ax^4 + Bx^3 + Cx^2 + Dx + \lambda$$

$$x^4 + 4x^2 = x^4[4A] + x^3[3B] + x^2[12A + 2C] + x[8B + D] + 4C + \lambda$$

$$A = 1/4 \quad B = 0 \quad C = 1/2 \quad D = 0 \quad \lambda = -2$$

$$= \left( \frac{x^3}{4} + \frac{x}{2} \right) \sqrt{x^2+4} - 2 \int \frac{1}{\sqrt{x^2+4}} dx$$

$$= \left( \frac{x^3}{4} + \frac{x}{2} \right) \sqrt{x^2+4} - 2 \ln \left[ x + \sqrt{x^2+4} \right] + C$$



# 22

HAZİRAN  
JUNE  
CUMA  
FRIDAY

H	P	S	Ç	P	C	Ct	Pz
22						1	2 3
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

(5)

$$\int \frac{dx}{(x-p)^n \sqrt{ax^2 + bx + c}}$$

$$\left[ \frac{1}{(x-p)} = t \right]$$

X

$$\int \frac{dx}{(x-1)^2 \sqrt{x^2 + 2x - 2}}$$

$$x^2 + 2x - 2$$

$$\frac{t^2 + 2t + 1}{t^2} + \frac{2t + 2}{t} - 2 = \frac{t^2 + 4t + 1}{t^2}$$

$$= \int \frac{t^2 - dt}{\sqrt{\frac{t^2 + 4t + 1}{t^2}}}$$

$$\frac{1}{x-1} = t$$

$$\frac{1}{t} = x-1$$

$$\frac{1}{t} + 1 = x$$

$$\frac{t+1}{t} = x$$

$$\left[ -\frac{1}{t^2} dt = dx \right]$$

$$= \int \frac{-t dt}{\sqrt{t^2 + 4t + 1}}$$

2. yöntem

$$t^2 + 4t + 1 = u$$

$$(2t+4)dt = du$$

$$-t = -\frac{1}{2}(2t+4)$$

$$= -\frac{1}{2}(2t+4-4)$$

$$= -\frac{1}{2}(2t+4) + 2$$

$$= \int \frac{-\frac{1}{2}(2t+4) + 2}{\sqrt{t^2 + 4t + 1}} dt$$

$$= -\frac{1}{2} \int \frac{2t+4}{\sqrt{t^2 + 4t + 1}} dt + 2 \int \frac{dt}{\sqrt{t^2 + 4t + 1}}$$



HAZİRAN 2012

H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN

JUNE

CUMARTESİ

SATURDAY

23

$$\textcircled{1} -\frac{1}{2} \int \frac{2t+4}{\sqrt{t^2+4t+1}} dt = -\frac{1}{2} \int \frac{dt}{\sqrt{u}} = -\frac{1}{2} \int u^{-\frac{1}{2}} du = -\frac{1}{2} \cdot \frac{u^{\frac{1}{2}}}{\frac{1}{2}} = -\sqrt{u} \\ = -\sqrt{t^2+4t+1}$$

$$\textcircled{2} 2 \int \frac{dt}{\sqrt{t^2+4t+1}} \stackrel{1. \text{ yöntem}}{=} 2 \int \frac{dt}{\sqrt{(t+2)^2-3}} \\ = 2 \ln \left[ t+2 + \sqrt{t^2+4t+1} \right] \\ \cancel{= -\sqrt{t^2+4t+1} + 2 \ln \left[ t+2 + \sqrt{t^2+4t+1} \right] + C} \\ = -\sqrt{\frac{1}{(x-1)^2} + \frac{4}{x-1} + 1} + 2 \cancel{\ln \left[ \frac{1}{x-1} + 2 + \sqrt{\frac{1}{(x-1)^2} + \frac{4}{x-1} + 1} \right]} \quad \begin{array}{l} \text{HAZİRAN} \\ \text{JUNE} \\ \text{PAZAR} \\ \text{SUNDAY} \end{array} \quad 24$$

~~$$\int \frac{dx}{(x+1)^2 \sqrt{x^2-2x}}$$~~

~~zam~~

$$= \int \frac{t^2 \frac{-1}{t^2} dt}{\sqrt{\frac{3t^2-4t+1}{t^2}}} \\ = \int \frac{-1}{\sqrt{3t^2-4t+1}} dt$$

$$\frac{t}{x+1} = t$$

$$\frac{1}{t} = x+1$$

$$\frac{1}{t} - 1 = x$$

$$\frac{1-t}{t} = x$$

$$-\frac{1}{t^2} dt = dx$$

$$x^2 - 2x \\ \frac{1-2t+t^2}{t^2} - 2 \frac{1+t}{t} \\ = \frac{3t^2-4t+1}{t^2}$$



# 25

HAZİRAN  
JUNE  
PAZARTESİ  
MONDAY

HAZİRAN 2012						
H	P	S	Ç	P	C	Ct Pz
22				1	2	3
23	4	5	6	7	8	9 10
24	11	12	13	14	15	16 17
25	18	19	20	21	22	23 24
26	25	26	27	28	29	30

$$= \int \frac{t dt}{\sqrt{3t^2 - 6t + 1}}$$

$$\begin{aligned} 3t^2 - 6t + 1 &= 0 \\ (6t - 1)dt &= dt \\ -t &= -\frac{1}{6}[6t - 1] \\ &= -\frac{1}{6}[6t - 1 + 1] \\ &= -\frac{1}{6}(6t - 1) - \frac{2}{3} \end{aligned}$$

$$= \int \frac{-\frac{1}{6}(6t - 1) - \frac{2}{3}}{\sqrt{3t^2 - 6t + 1}} dt$$

$$= -\frac{1}{6} \int \frac{6t - 1}{\sqrt{3t^2 - 6t + 1}} dt - \frac{2}{3} \int \frac{1}{\sqrt{3t^2 - 6t + 1}} dt$$

$$\textcircled{1} \quad -\frac{1}{6} \int \frac{6t - 1}{\sqrt{3t^2 - 6t + 1}} dt = -\frac{1}{6} \int \frac{dv}{\sqrt{v^2}} = -\frac{1}{6} \int v^{-\frac{1}{2}} dv = \frac{1}{6} v^{\frac{1}{2}} = \frac{1}{6} \sqrt{v} = \frac{1}{6} \sqrt{3t^2 - 6t + 1}$$

$$= -\frac{1}{6} \sqrt{3t^2 - 6t + 1}$$

$$= -\frac{1}{6} \sqrt{3(t^2 - 2t + \frac{1}{3})} = -\frac{1}{6} \sqrt{3((t - \frac{1}{3})^2 - \frac{1}{3})}$$

$$\textcircled{2} \quad -\frac{2}{3} \int \frac{dt}{\sqrt{3t^2 - 6t + 1}} = -\frac{2}{3} \int \frac{1}{\sqrt{(3t - \frac{2}{\sqrt{3}})^2 - \frac{1}{3}}} dt$$

$$\begin{cases} \sqrt{3t - \frac{2}{\sqrt{3}}} = u \\ \sqrt{3}dt = du \end{cases}$$

$$= -\frac{2}{3} \int \frac{du}{\sqrt{u^2 - \frac{1}{3}}}$$



H	P	S	Ç	P	C	Ct	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN  
JUNE  
SALI  
TUESDAY

26

$$= -\frac{2}{3\sqrt{3}} \ln \left[ t + \sqrt{t^2 - \frac{1}{3}} \right]$$

$$= -\frac{2}{3\sqrt{3}} \ln \left[ \sqrt{3}t - \frac{2}{\sqrt{3}} + \sqrt{3t^2 - 6t + 1} \right]$$

*carp*

$$= -\frac{1}{3} \sqrt{3t^2 - 6t + 1} - \frac{2}{3\sqrt{3}} \ln \left[ \sqrt{3}t - \frac{2}{\sqrt{3}} + \sqrt{3t^2 - 6t + 1} \right] + C$$

$$\left| t = \frac{1}{x+1} \right.$$

### ⑥ BINOM INT.

$$\int x^n (a + bx^p)^q dx$$

a)  $q = \text{tam sayı ise}$  olkek  $(p \text{ doðar}, p \text{ deðer}) = k$  olur

$$\boxed{x = t^k}$$

b)  $q \neq \text{tam sayı ise}$  ~~deðer~~  $q$ 'nın paydası  $n$  olur  $\frac{p+1}{p} = \text{tam sayı}$   
~~deðer~~

$$\boxed{(a + bx^p) = t^n}$$

c)  $q \neq \text{karsayı}, \frac{p+1}{p} \neq \text{Tamsayı}, \frac{p+1}{p} + q = \text{karsayı ve } q \text{nun paydası } n \text{ ise}$

$$\boxed{ax^{-p} + b = t^n}$$



# 27

HAZİRAN

JUNE

ÇARŞAMBA  
WEDNESDAY

HAZİRAN

2012

H	P	S	Ç	P	C	Ct	Pz
22						1	2 3
23	4	5	6	7	8	9 10	
24	11	12	13	14	15	16 17	
25	18	19	20	21	22	23 24	
26	25	26	27	28	29	30	

$$\cancel{\int} \sqrt[3]{x} (1+2\sqrt{x})^2 dx = \int x^{1/3} (1+2x^{1/2})^2 dx$$

$$r = \frac{1}{3}, \quad p = \frac{1}{2}, \quad q = 2$$

düştü de geceler ( $q = \text{konsan}$ )

$$\text{deki } (2, 3) = 6 = L$$

$$\boxed{x = t^L = t^6}$$

$$dx = 6t^5 dt$$

$$= \int t^2 (1+2t^3)^2 6t^5 dt$$

$$= 6 \int t^7 (1+2t^3)^2 dt$$

$$= 6 \int t^7 (1+4t^3+4t^6) dt$$

$$= 6 \int (t^7 + 4t^{10} + 4t^{13}) dt$$

$$= 6 \left[ \frac{t^8}{8} + \frac{4t^{11}}{11} + \frac{4t^{14}}{14} \right] + C$$

$$= 6 \left[ \frac{x^{8/6}}{8} + \frac{4x^{11/6}}{11} + \frac{4x^{14/6}}{14} \right] + C$$



HAZİRAN 2012

H	P	S	Ç	P	C	C1	Pz
22				1	2	3	
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

HAZİRAN

JUNE

PERŞEMBE

THURSDAY

28

$$\cancel{\int \frac{\sqrt{1+3\sqrt{x^2}}}{3\sqrt{x^2}} dx} = \int x^{-2/3} [1+x^{1/3}]^{1/2} dx$$

$$r = -\frac{2}{3} \quad p = \frac{1}{3} \quad q = \frac{1}{2}$$

$q \neq \text{Tanses!}$   $\frac{r+1}{p} = \frac{1/3}{1/3} = 1 = \text{Tanses!}$

bşibbi ke sebzeler  $n=2 \rightarrow q' \text{nun şeripte}$

pontez sine  $t^n$  dekar

$$1+x^{1/3} = t^2$$

$$\frac{1}{3} x^{-2/3} dx = 2t dt$$

$$t = \sqrt{1+3\sqrt{x}}$$

$$= \int 6t dt (t^2)^{1/2} = 6 \int t^2 dt = 6 \frac{t^3}{3} + C = 2t^3 + C$$

$$= 2(\sqrt{1+3\sqrt{x}})^3 + C$$

$$\cancel{\int \sqrt{\frac{x}{1-x^3}} dx} = \int \frac{\sqrt{x}}{\sqrt{1-x^3}} dx = \int x^{1/2} (1-x^3)^{-1/2} dx$$

$$r = 1/2 \quad p = 3 \quad q = -1/2$$

$$\frac{r+1}{p} = \frac{3/2}{3} = 1/2$$

$$\frac{r+1}{p} + q = 0 \rightarrow \text{Tanses!}$$

$$n=2$$

C'sizliklara giden yollar



# 29

HAZİRAN  
JUNE  
CUMA  
FRIDAY

H	P	S	Ç	P	C	Ct	Pz
22						1	2 3
23	4	5	6	7	8	9	10
24	11	12	13	14	15	16	17
25	18	19	20	21	22	23	24
26	25	26	27	28	29	30	

$$x^{-3} - 1 = t^2 \rightarrow x^{-3} = t^2 + 1$$

$$-3x^{-4} dx = 2t dt$$

$$dx = -\frac{2}{3} x^4 t dt$$

$$dx = -\frac{2}{3} \left( \frac{1}{t^2+1} \right)^{1/3} t dt$$

$$\frac{1}{x^3} = t^2 + 1 \rightarrow \frac{1}{x^3} - 1 = t^2$$

$$\frac{1}{t^2+1} = x^3 \quad t = \sqrt[3]{1-x^3}$$

$$\left( \frac{1}{t^2+1} \right)^{1/3} = x$$

$$= \int \frac{1}{(t^2+1)^{4/3}} \left[ 1 - \frac{1}{t^2+1} \right]^{-1/2} \cdot \left( \frac{-2}{3} \right) \frac{t}{(t^2+1)^{1/3}} dt$$

$$= -\frac{2}{3} \int \frac{1}{(t^2+1)^{3/2}} \frac{t^{-1}, t}{(t^2+1)^{-1/2}} dt$$

$$= -\frac{2}{3} \int \frac{1}{t^2+1} dt = -\frac{2}{3} \arctan t + C$$

$$= -\frac{2}{3} \arctan \sqrt{\frac{1-x^3}{x^3}} + C$$



TEMMUZ 2012

H	P	S	C	P	C	Ct	Pz
26						1	
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

4 - P - %65-Kutuz  
 5 - S - %35-Nur TEMMUZ  
 6 - P - %5-Kutuz  
 7 - P - 9.5 - 10 CUMARTESİ  
 8 ? - 9.50 - FNA SATURDAY

07

$$\int \frac{2x^4 - 3x^3 + 5}{3x^2} dx$$

$$\frac{2x^4 - 3x^2 + 5}{3x^2}$$

$$\int \left( \frac{2}{3}x^2 + \frac{5}{3}\frac{1}{x^2} \right) dx = \frac{2}{3} \frac{x^3}{3} + \frac{5}{3} \frac{x^{-1}}{-1} - \frac{x^2}{2}$$

$$\frac{2}{9}x^3 - \frac{5}{3}x^{-1} - \frac{x^2}{2} + C$$

$$\int x^5 \ln x dx = \frac{x^6}{36} (A \ln x + B) + C. \quad A, B, C?$$

$$\begin{aligned} \ln x &= u \\ x^5 &= v \end{aligned}$$

$$x^5 dx = v du$$

f(x)

TEMMUZ

JULY

PAZAR

SUNDAY

08

$$\left( \frac{x^6}{36} [A \ln x + B] + C \right)' = x^5 \ln x$$

$$\frac{6x^5}{36} (A \ln x + B) + \frac{x^6}{36} \frac{A}{x} + 0 = x^5 \ln x$$

$$\ln x = \frac{A \ln x + B}{6} + \frac{A}{36}$$

$$\ln x = \frac{A}{6} \ln x + \frac{B}{6} + \frac{A}{36}$$

$$A = 6$$

$$B = -1$$

$$C \in \mathbb{R}$$



09

TEMMUZ

JULY

PAZARTESİ

MONDAY

TEMMUZ

2012

H	P	S	Ç	P	C	Ct	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

~~T~~ Tercüde kendişinin  $f'$  katmanı eşit olur pozitif  
bir  $f$  fonksiyonu için  $f(0)=1$  oldugu için  $f(x)=?$

$$f'(x) = 2f(x)$$

$$\frac{f'(x)}{f(x)} = 2$$

$$= \int \frac{f'(x)}{f(x)} dx = \int 2 dx$$

$$\left| \begin{array}{l} f(x) = u \\ f'(x)dx = du \end{array} \right.$$

$$= \int \frac{du}{u} = 2x + c$$

$$\ln|u| = 2x + c$$

$$\ln|f(x)| = 2x + c$$

$$\ln|f(x)| = 2x + c \rightarrow \text{pozitif}$$

$$f(x) = e^{2x+c}$$

$$f(0) = 1 = e^c \Rightarrow c = 0$$

$$\boxed{f(x) = e^{2x}}$$



TEMMUZ 2012						
H	P	S	C	P	C	Ç
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

TEMMUZ  
JULY  
SALI  
TUESDAY

10

$$\mathcal{X} \int \frac{x^3}{1+x^8} dx$$

okey (3,8)

$$\begin{aligned} & \cancel{dx} \cancel{x^8} \cancel{x^{20}} \\ & \cancel{8x^7 dx} \cancel{24x^{23} dt} \\ & \boxed{x^4 = t} \\ & \boxed{4x^3 dx = dt} \end{aligned}$$

$$2^3 \cdot 3$$

$$= \frac{1}{4} \int \frac{dt}{1+t^2}$$

$$= \frac{1}{4} \arctan t + C$$

$$= \frac{1}{4} \arctan x^4 + C$$

$$\mathcal{X} \int \sin((2x+1)\pi) dx$$

tahtas

$$(2x+1)\pi = t$$

$$2\pi dx = dt$$

$$= -\frac{1}{2\pi} \cos t + C$$

$$= -\frac{1}{2\pi} \cos((2x+1)\pi) + C$$

$$\mathcal{X} \int \frac{x dx}{\sqrt{x+1}}$$

$$x+1 = t^2$$

$$= 2 \int (t^2 - 1) dt$$

$$\approx 2 \frac{t^3}{3} - 2t + C$$

$$= \frac{2}{3} \frac{(x+1)^{3/2}}{3} - 2(x+1)^{1/2} + C$$



# 11

TEMMUZ  
JULY  
ÇARŞAMBA  
WEDNESDAY

$$e^x - e^{-x}/2$$

$$\sqrt{a^2 - x^2} \rightarrow \text{osmt}$$

$$\sqrt{x^2 - a^2} \rightarrow \text{osct}$$

$$\sqrt{a^2 + x^2} \rightarrow \text{otant}$$

TEMMUZ 2012						
H	P	S	C	P	C	Ct Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

~~$$\int \frac{dx}{x^2 \sqrt{16-x^2}}$$~~

~~$$\int t^{3/2} (-2) dt$$~~


$$x = 4 \sin t$$

$$dx = 4 \cos t dt$$

$$\begin{aligned} \frac{x}{4} &= t \rightarrow x = 4t \\ x^2 &= t \\ -2x^3 dx &= dt \\ -2x^3 dt &= dx \\ -2x^3 &= dt \\ \left(\frac{1}{t}\right)^3 &= dt \end{aligned}$$

$$\begin{aligned} 1 \int \frac{4 \cos t}{16 \sin^2 t + \sqrt{16 - 16 \sin^2 t}} dt \\ &= \frac{1}{16} \cos t + C = \frac{1}{16} \end{aligned}$$

$$= -\frac{1}{16} \cot t + C = -\frac{1}{16} \cancel{\cos t} \frac{\sqrt{16-x^2}}{x} + C$$



~~$$\int \frac{2^x dx}{\sqrt{8-4^x}}$$~~

$$\begin{aligned} 2^x &= t \\ 2^x \ln 2 dx &= dt \end{aligned}$$

$$= \frac{1}{\ln 2} \int \frac{dt}{\sqrt{1-t^2}}$$

$$= \frac{1}{\ln 2} \arcsin t + C$$

$$= \frac{1}{\ln 2} \arcsin(2^x) + C$$



TEMMUZ 2012

H	P	S	C	P	C	Cl	Pz
26						1	
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

TEMMUZ

JULY

PERŞEMBE

THURSDAY

12

$$\int \frac{e^{2x}}{\sqrt{e^x + 1}} dx$$

$$1+t^2 = t^2$$

$$e^x dx = 2t dt$$

$$= \int \frac{(t^2 - 1) 2t dt}{\sqrt{t^2 + 1}} = 2 \int (t^2 - 1) dt = 2 \left( \frac{t^3}{3} - t \right) + C$$

$$= \frac{2}{3} (e^x + 1)^{3/2} - \sqrt{e^x + 1} + C$$

$$\int \frac{1}{1 + \cos x} dx$$

$$\tan \frac{x}{2} = t$$

$$\frac{x}{2} = \arctant$$

$$x = 2 \arctant$$

$$dx = 2 \frac{1}{1+t^2} dt$$

$$\cos x = \frac{1-t^2}{1+t^2}$$

$$= \int \frac{1}{1 + \frac{1-t^2}{1+t^2}} \cdot \frac{1}{1+t^2} dt$$

$$= \int \frac{1+t^2}{2} \cdot \frac{dt}{1+t^2}$$

$$= \tan \frac{x}{2} + C$$

$$\frac{1-t^2 + 1+t^2}{1+t^2}$$

$$\int \frac{1}{1 + [2\cos^2(\frac{x}{2}) - 1]} dx = \int \frac{dx}{2\cos^2(\frac{x}{2})}$$

$$\frac{x}{2} = t$$

$$\frac{dx}{2} = dt$$



# 13

TEMMUZ  
JULY  
CUMA  
FRIDAY

3.

$$2+2t^2$$

TEMMUZ 2012						
H	P	S	C	P	C	Ct Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

~~$$\int \frac{dx}{2+\sin x - \cos x}$$~~

$$\left| \text{for } \frac{x}{2} = t \right)$$

$$= \int \frac{\frac{2}{1+t^2} dt}{2 + \frac{2t}{1+t^2} + \frac{1-t^2}{1+t^2}} = \int \frac{\frac{2}{1+t^2}}{\frac{3t^2+2t+1}{1+t^2}} dt$$

$$= \int \frac{2}{3t^2+2t+1} dt$$

~~$$t^2+2t+3=0$$~~

~~$$= \int \left( \frac{2}{t+1} + \frac{2}{t+2} \right) dt$$~~

~~$$\frac{2}{t^2+2t+3} = \frac{A}{t+1} + \frac{B}{t+2}$$~~

~~$$At+A+Bt+B=2$$~~

~~$$At+B=0$$~~

~~$$2A+B=2$$~~

~~$$A=2, B=-2$$~~

~~$$= \ln |t+1| - \ln |t+2| + C$$~~

$$\int \frac{A dx}{e^x+x^2} = A \frac{1}{\alpha} \operatorname{arctan} \frac{x}{\alpha} + C$$

~~$$= 2 \int \frac{dt}{(\sqrt{3}t + \frac{1}{\sqrt{3}})^2 + \frac{2}{3}}$$~~

$$\sqrt{3}t + \frac{1}{\sqrt{3}} = u$$

$$(3dt = du)$$

$$= 2 \int \frac{du}{u^2 + (\frac{\sqrt{2}}{\sqrt{3}})^2} = \frac{2}{\sqrt{3}} \cdot \frac{1}{\frac{\sqrt{2}}{\sqrt{3}}} \operatorname{arctan} \frac{u}{\frac{\sqrt{2}}{\sqrt{3}}} + C$$

$$= \sqrt{2} \operatorname{arctan} \frac{\sqrt{3}t + \frac{1}{\sqrt{3}}}{\sqrt{2}}$$



TEMMUZ 2012					
H	P	S	C	P	Ct
26					1
27	2	3	4	5	6
28	9	10	11	12	13
29	16	17	18	19	20
30	23	24	25	26	27
31	30	31			28 29

TEMMUZ  
JULY  
CUMARTESİ  
SATURDAY

14

$$= \sqrt{2} \operatorname{arctan} \frac{3t+1}{\sqrt{2}} + C$$

$$= \sqrt{2} \operatorname{arctan} \frac{3(\tan \frac{x}{2})+1}{\sqrt{2}} + C$$

~~$$\int x^2 \cos x dx$$~~

$$= x^2 \sin x - \int (\sin x) \cdot 2x dx$$

$$= x^2 \sin x - 2 \left[ x \cos x + \int \cos x dx \right]$$

$$= x^2 \sin x + 2x \cos x - 2 \sin x + C$$

~~$$\int \cos(\ln x) dx$$~~

~~$$\begin{aligned} \ln x &= t \\ dt &= dx \end{aligned}$$~~

TEMMUZ  
JULY  
PAZAR  
SUNDAY

15

$$= (\cos(\ln x))x - \int \frac{1}{x} (-\sin(\ln x))dx$$

$$= x \cos(\ln x) + \int \sin(\ln x) dx$$

$$= x \cos(\ln x) + x \sin(\ln x) - \int x \frac{1}{x} \cos(\ln x) dx$$

$$I = \frac{1}{2} \left[ x(\cos(\ln x) + \sin(\ln x)) \right] + C$$

$$\cos(\ln x) = v$$

$$\frac{1}{x} (-\sin(\ln x)) dx = dv$$

$$dx = dv$$

$$x = y$$

$$\sin(\ln x) = v$$

$$\frac{1}{x} (\cos(\ln x)) dx = dv$$



# 16

TEMMUZ  
JULY

PAZARTESİ  
MONDAY

$$(\log a^v) = \frac{v}{v} \log a e$$

TEMMUZ 2012							
H	P	S	C	P	C	Çt	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

~~\*~~   $\int \log_5 x \, dx$

$$\begin{cases} \log_5 x = v \\ \left( \frac{1}{x} \log_5 e \right) dx = du \end{cases}$$

$$dx = dv$$

$$x = v$$

$$= (\log_5 x)x - \int x \frac{1}{x} \log_5 e \, dx$$

$$= x \log_5 x - x \log_5 e + C$$

~~\*~~   $\int \sin^n x \, dx = \int \underbrace{\sin^{n-1} x}_{v} \underbrace{\sin x \, dx}_{du}$

$$\begin{cases} \sin^{n-1} x = v \\ (n-1) \cos^{n-2} x \, dx = du \end{cases}$$

$$\sin x \, dx = dv$$

$$-\cos x = u$$

$$-\sin^{n-1} x \cos x + \int \cos x (n-1) \cos^{n-2} x \, dx$$



TEMMUZ 2012

H	P	S	Ç	P	C	Ct	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

TEMMUZ

JULY

SALI

TUESDAY

17

~~$\int \frac{dx}{x^2+x-6}$~~

$$\frac{1}{x^2+x-6} = \frac{A}{x+3} + \frac{B}{x-2}$$

$$Ax - 2A + Bx + BB = 1$$

$$A+B=0$$

$$-2A+BB=1$$

$$-\frac{1}{5} \ln|x+3| + \frac{1}{5} \ln|x-2| + C$$

~~$\int \frac{x^4}{x^4-1} dx$~~

$$\frac{x^4}{x^4-1} = \frac{Ax+B}{x^2+1} + \frac{C}{x+1} + \frac{D}{x-1}$$

$$(x^2+1)(x+1)(x-1)$$

$$D = 1/4 \quad C = -1/4$$

$$A = 0 \quad B = 1/2$$

$$\int \left( 1 + \frac{1}{x^4-1} \right) dx$$

$$= x + \frac{1}{4} \ln|x-1| + \frac{1}{4} \ln|x+1| + \frac{1}{2} \arctan x + C$$

~~$\int \cos 2x \cos 3x dx$~~

$$\cos ax \cos bx = \frac{1}{2} [\cos(a+b)x + \cos(a-b)x]$$

$$= \frac{1}{2} \int (\cos 5x + \cos x) dx$$

$$= \frac{1}{2} \left[ \frac{1}{5} \sin 5x + \sin x \right] + C$$



# 18

TEMMUZ

JULY

ÇARŞAMBA  
WEDNESDAY

TEMMUZ 2012						
H	P	S	C	P	C	Cl Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

$$\cancel{\int} \tan^3 x \sec^4 x dx$$

$$\begin{cases} \tan x = t \\ \sec^2 x dx = dt \\ 1 + \tan^2 x = \sec^2 x \end{cases}$$

$$= \int \tan^3 x \sec^2 x \sec^2 x dx$$

$$= \int t^3 (1+t^2) dt$$

$$= \int (t^3 + t^5) dt = \frac{t^4}{4} + \frac{t^6}{6} + C = \frac{\tan^4 x}{4} + \frac{\tan^6 x}{6} + C$$

$$\cancel{\int} \frac{dx}{(sin x + cos x)^2}$$

$$\left( \tan x = t \text{ ve } \frac{\tan x}{\sqrt{2}} = t \right)$$

derecelendir

$$= \int \frac{1}{(1+t^2)^2} dt$$

$$\left( \frac{t}{\sqrt{1+t^2}} + \frac{1}{\sqrt{1+t^2}} \right)^2$$

$$\cancel{dt} = \sec t \tan t dt$$

$$dt = \frac{1}{1+t^2} dt$$

$$t \quad \sqrt{1+t^2}$$

$$= \int \frac{1}{(1+u^2)^2} du$$

$$u = \tan x$$

$$du = \sec^2 x dx$$

$$= -\frac{1}{1+u^2} + C = -\frac{1}{1+\tan^2 x} + C = -\frac{1}{\sec^2 x} + C$$



TEMMUZ 2012

H	P	S	Ç	P	C	Cl	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

TEMMUZ

JULY

PERŞEMBE  
THURSDAY

19

$$\int \frac{3\sqrt{1+\sqrt[3]{x}}}{\sqrt{x}} dx = \int x^{-\frac{1}{2}} (1+x^{\frac{1}{3}})^{\frac{3}{2}} dx$$

$$(1+x^{\frac{1}{3}})^{\frac{3}{2}} = t^3$$

$$\frac{1}{6} x^{-\frac{3}{2}} dx = 3t^2 dt$$




$$dx = 18t^2 x^{\frac{3}{2}} dt$$

$$dx = (t^3 - 1)^3 18t^2 dt$$

$$x^{\frac{1}{3}} = t^3 - 1$$

$$x = (t^3 - 1)^3, x^{-\frac{1}{2}} = (t^3 - 1)^{-\frac{1}{2}}$$

## BELİRLİ İNTEGRAL



Tanım:  $f: [a, b] \rightarrow \mathbb{R}$  fonksiyon  
sayıları olsun  $[a, b]$  aralığının

$P = \{x_0, \dots, x_n\}$  partisyonu  $\mathcal{P}$

$M_k = \max\{f(x), x_{k-1} \leq x \leq x_k\}$

$m_k = \min\{f(x), x_{k-1} \leq x \leq x_k\}$



olsun →

# 20

TEMMUZ  
JULY  
CUMA  
FRIDAY

TEMMUZ 2012						
H	P	S	Ç	P	C	Ct Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

$$A(f, P) = \sum_{i=1}^n \Delta x_i M_i$$

$$\Delta x_5 = x_5 - x_3$$

$$\bar{V}(f, P) = \sum_{i=1}^n \Delta x_i M_i$$

$$R(f, P) = \sum_{i=1}^n \Delta x_i f(x_i)$$

$f$  fonksiyonun  $P$  partisi üzerindeki  $f$  fonksiyonun toplamı dem

Tanım:  $f$ ,  $[a, b]$  aralığında sürekli ve  $b-a$  partisi  $n$  tane alt aralığa bölünse de  $f$  fonksiyonu olsun ve

$$x_k^* \in [x_{k-1}, x_k]$$

$$\lim_{\|P\| \rightarrow 0} \sum_{i=1}^n f(x_i^*) \Delta x_i = I$$

(mots versi, bu limite  $a$  da  $b$  ye kadar  $f$  fonksiyonun integrali dem ve

$$\int_a^b f(x) dx$$



TEMMUZ 2012

H	P	S	C	P	C	CL	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

TEMMUZ

JULY

CUMARTESİ

SATURDAY

21

Teorem:  $f [a, b]$  aralığı integrallenebilir.

$\forall x \in (a, b)$  için  $F'(x) = f(x)$  ise

$$\int_a^b f(x) dx = f(x) \Big|_a^b = f(b) - f(a)$$

(Newton-Leibnitz)

~~1~~  $\int_1^2 2x dx = x^2 \Big|_1^2 = 2^2 - 1^2 = 3$

~~2~~  $\int_0^{\pi/2} \cos x dx = \sin x \Big|_0^{\pi/2} = 1$

## Bölüm: İntegral Uyeler

TEMMUZ

JULY

PAZAR

SUNDAY

22

①  $f$  farklı sürekli ise integrallenebilir.

② Borulu sürekli ise integrallenebilir.

③ Monoton ise integrallenebilir

④  $\int_a^a f(x) dx = 0$

⑤  $\int_a^b f(x) dx = - \int_b^a f(x) dx$

⑥  $\int_a^b f(x) dx = \int_a^b f(t) dt = \dots = \int_a^b f(u) du$



# 23

TEMMUZ

JULY

PAZARTESİ  
MONDAY

TEMMUZ 2012						
H	P	S	C	P	C	Ct Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

$$\textcircled{A} \quad c \in [a, b]$$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

$$\textcircled{B} \quad \int_0^{\pi/2} \sin^{3n} x \cos x dx$$

$$\sin x = t \\ \cos x dx = dt$$

$$= \int_0^{\pi/2} t^{3n} dt$$

$$= \frac{t^{3n+1}}{3n+1} \Big|_0^{\pi/2} = \frac{2}{3} t^{3n+1} \Big|_0^{\pi/2} = \frac{2}{3} \sin^{3n+1} x \Big|_0^{\pi/2} = \frac{2}{3}$$

$$\textcircled{C} \quad \int_a^b f(x) dx = uv \Big|_a^b - \int_a^b v du$$

$$\int_0^{\pi/2} \frac{\cos x}{1 + \sin^2 x} dx$$

$$\begin{cases} \sin x = t \\ \cos x dx = dt \end{cases}$$

$$= \int_0^{\pi/2} \frac{dt}{1+t^2} = \arctan t \Big|_0^{\pi/2}$$

$$= \arctan t \Big|_0^{\pi/2}$$

$$x=0 \rightarrow \sin 0 = 0 \\ x=\pi/2 \rightarrow \sin \pi/2 = 1$$

$$= \arctan 1 - \arctan 0$$

$$= \pi/4 - 0 = \pi/4$$



TEMMUZ 2012						
H	P	S	C	P	C	Ct
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

TEMMUZ  
JULY  
SALI  
TUESDAY

24

Lecture

$$\int_0^{\pi} x \cos x dx$$

$$x = v \\ dx = dv$$

$$\cos x dx = dv \\ \sin x = v$$

$$x \sin x - \int_0^{\pi} \sin x dx$$

$$(\pi \sin \pi) - [1 - 1] = -2$$

$$\textcircled{1} \quad f(x) \geq 0 \quad \Rightarrow \quad \int_0^b f(x) dx \geq 0$$

$$\textcircled{2} \quad \left| \int_a^b f(x) dx \right| \leq \int_0^b |f(x)| dx$$

$$\textcircled{3} \quad \begin{array}{l} \cancel{f \text{ tek}} \text{ net se} \\ \cancel{f \text{ çift}} \text{ net se} \end{array} \quad \begin{array}{l} f(-x) = -f(x) \text{ tek} \\ f(-x) = +f(x) \text{ çift} \end{array}$$

$$\begin{array}{l} f(-x) = x^2 \text{ çift} \\ f(-x) = -x^3 \text{ tek} \end{array} \quad \begin{array}{l} f(x) = \frac{x^2}{1+x^4} \\ f(-x) = \frac{x^2}{1+x^4} \end{array} \quad \begin{array}{l} f(x) = x^3 + 1 \\ f(-x) = -x^3 + 1 \end{array}$$

gitir

ne tek ne çift

$$\text{if } f \text{ tek} \text{ se} \quad \int_{-a}^a f(x) dx = 0$$

$$\text{if } f \text{ çift se} \quad \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$



# 25

TEMMUZ

JULY

ÇARŞAMBA  
WEDNESDAY

TEMMUZ 2012

H	P	S	Ç	P	C	Ct	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

X

$$\int_{-\pi/2}^{\pi/2} \cos^2 x dx$$

$-\pi/2$

$$= 2 \int_0^{\pi/2} \cos^2 x dx$$

$$= 2 \int_0^{\pi/2} \frac{1}{2} [\cos 2x + 1] dx$$

$$= \frac{1}{2} \sin 2x + x \Big|_0^{\pi/2} = \left( \frac{1}{2} \sin \pi + \frac{\pi}{2} \right) - \left( \frac{1}{2} \sin 0 + 0 \right)$$

$$\approx \pi/2$$

X

$$\int_{-\pi}^{\pi} \frac{x^3 \cos x}{1 + \sin^{10} x} dx = 0$$

## (B) integrallerin Törəmə

$f, [a, b]$  integrallenebilər funks

$$F(x) = \int_a^x f(t) dt$$

$$F'(x) = \frac{d}{dx} \int_a^x f(t) dt = f(v(x)) v'(x) - f(v(x)) v'(x)$$

(Leibnitz)



TEMMUZ							2012	
H	P	S	C	P	C	Ct	Pz	
26								1
27	2	3	4	5	6	7	8	
28	9	10	11	12	13	14	15	
29	16	17	18	19	20	21	22	
30	23	24	25	26	27	28	29	
31	30	31						

TEMMUZ

JULY

PERSEMBE

THURSDAY

26

$$\begin{array}{|c|c|c|} \hline 31 & 30 & 31 \\ \hline \end{array} \quad \text{THUR} \quad \frac{x^2}{2}$$

~~A~~  $F(x) = \int_{-x}^x \sin(t^2) dt$        $F'(x) = ?$

$$= \sin(x^4) \cdot 2x - \sin(x^3) \cdot 1$$

$$\lim_{x \rightarrow 0} \frac{1}{x^2} \int_0^x \frac{t^2}{t^4+1} dt = ?$$

$\infty \cdot 0$  Beharrlichkeit!

$\frac{0}{0}$  wegen  $\frac{\infty}{\infty}$ : sparsam

L'Hospital rule oft

$$= \lim_{x \rightarrow 0} \int_0^x \frac{t^2}{t^4+1} dt = \frac{0}{0}$$

↑ *Indefinite Integral*

$$= \lim_{x \rightarrow \infty} \frac{x^2}{x^4 + 1} = 0$$

$$= \lim_{x \rightarrow \infty} \frac{x}{2(x^4+1)} = \frac{0}{2} = 0$$




# 27

TEMMUZ  
JULY  
CUMA  
FRIDAY

TEMMUZ 2012						
H	P	S	C	P	C	Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

Fresik


Tümnic (Ortalama Değer)  $\equiv$  Ağırlık Merkezi



$f$ ,  $[a, b]$  aralığında  
integrali varsa olsun  
 $y = f(x)$  fonksiyonu  
 $[a, b]$  aralığında,  
ortalama

$$\bar{y} = \frac{1}{b-a} \int_a^b f(x) dx, \quad f(\bar{x}) = \bar{y}$$

yazın oradan  
diğerlerse denetle  
fazla (.)



$$\bar{y} = \frac{1}{2-(-1)} \int_{-1}^2 (3x+4) dx$$

$$\bar{y} = 3 \frac{x^2}{2} + 4x \Big|_{-1}^2$$

$$\bar{y} = (6+8) - \left( \frac{-3}{2} + 4 \right)$$

$$\bar{y} = 17/2 \approx 8,5$$

$$\bar{y} = 3\bar{x} + 4$$

$$17/2 = 3\bar{x} + 4$$

$$\bar{x} = 3/2$$



TEMMUZ 2012						
H	P	S	C	P	C	Cl Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

$\frac{12}{3}$   $\frac{12 \cdot 12^2}{3} = 144 \cdot 4$  TEMMUZ  
JULY

CUMARTESİ  
SATURDAY

28


~~Q~~  $y = x^2$  ,  $[0, 3]$  cross ort

$$g = \frac{1}{3-0} \int_0^3 x^2 dx$$


$$\bar{y} = \frac{1}{3} \left( \frac{x^3}{3} \right) \Big|_0^3$$

$$\bar{y} = 3$$

$$\bar{x} = \sqrt{3} \approx 1,71$$



~~Q~~ Taban yarıçapı = 3 yüzseklik = 12 dik  
dairesel koni - tepeinden paralel y eksenine  
uzunluksız kesikyes. Dairesel kesitin TEMMUZ  
ortalama = ? JULY



$$\bar{y} = \frac{1}{12} \int_0^{12} \pi r^2 dy$$

$$\bar{y} = \frac{\pi}{12} \left( \frac{r^3}{3} \right) \Big|_0^{12}$$

29

$$A = \pi r^2$$

$$\frac{r}{3} = \frac{y}{12}$$

$$r = \frac{y}{4}$$

$$A = \pi \frac{y^2}{16}$$

$$\bar{y} = \frac{1}{12} \int_0^{12} \frac{1}{16} \pi y^2 dy$$

$$= \frac{1}{12} \frac{1}{16} \pi \left[ \frac{y^3}{3} \right] \Big|_0^{12}$$

$$= 3\pi$$

# 30

TEMMUZ

JULY

PAZARTESİ

MONDAY

TEMMUZ 2012							
H	P	S	Ç	P	C	Ct	Pz
26							1
27	2	3	4	5	6	7	8
28	9	10	11	12	13	14	15
29	16	17	18	19	20	21	22
30	23	24	25	26	27	28	29
31	30	31					

$$\int_{-2}^2 |x| dx = \int_0^2 x dx + \int_{-1}^1 x dx$$

$$|x| = \begin{cases} -x & , x \leq 0 \\ x & , x > 0 \end{cases}$$

$$= \int_{-1}^0 -x dx + \int_0^2 x dx$$

$$= -\frac{x^2}{2} \Big|_{-1}^0 + \frac{x^2}{2} \Big|_0^2$$

$$= (0) - \left(-\frac{1}{2}\right) + (2) - (0)$$

$$= 5/2$$

$$\int_0^3 [x] dx$$

Kurdistan  
boyut olmaz  
en boyut  
sayı  
(Tüm sayılar)

$$[3,9] = 3$$

$$[3,2] = 3$$

$$[3,3] = 3$$

$$[-2,4] = 3$$

$$[-2,7] = -3$$

$$[4,2] = 4$$

$$= \int_0^1 0 dx + \int_1^2 1 dx + \int_2^3 2 dx$$

$$= x \Big|_1^2 + 2x \Big|_2^3$$

$$= -1 + 2 = \boxed{1}$$



TEMMUZ 2012						
H	P	S	C	P	C	Pz
26						1
27	2	3	4	5	6	7 8
28	9	10	11	12	13	14 15
29	16	17	18	19	20	21 22
30	23	24	25	26	27	28 29
31	30	31				

TEMMUZ

JULY

SALI

TUESDAY


31

$$\text{A} \quad \int_0^2 x[0x] dx$$

$$= \int_0^1 x \cdot 0 dx + \int_1^2 x \cdot 1 dx = 3/2$$

$$\text{B} \quad \int_0^4 \operatorname{sgn}(x^2 - 4x + 3) dx$$

$$\operatorname{sgn}(f) = \begin{cases} 1, & f > 0 \\ 0, & f = 0 \\ -1, & f < 0 \end{cases}$$



$$= \int_0^1 1 dx + \int_1^3 -1 dx + \int_3^4 1 dx$$

$$\text{C} \quad f(x) = 2|x|, \quad [-1, 1] \text{ arası ort bul.}$$

$$\bar{y} = \frac{1}{1-(-1)} \int_{-1}^1 2|x| dx$$

$$\bar{y} = \int_{-1}^1 |x| dx$$

$$\bar{y} = \int_{-1}^0 -x dx + \int_0^1 x dx$$

$$\bar{y} = 1$$



# 01


AĞUSTOS

AUGUST

ÇARŞAMBA  
WEDNESDAY

AĞUSTOS 2012						
H	P	S	Ç	P	C	Ct Pz
31				1	2	3
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	

## ALAN HESABI




$$y = f(x) \geq 0$$

$$x=a$$

$$x=b$$

$Ox$ - ekseni ( $y=0$ )

$$A = \int_a^b f(x) dx$$




$$y = f(x) < 0$$

$$x=a$$


$$x=b$$

$Ox$ - ekseni ( $y=0$ )

$$A = - \int_a^b f(x) dx$$



$$A = \int_a^b f(x) dx - \int_b^c f(x) dx$$



$$x = g(y) \geq 0$$

$$y=c$$

$$y=d$$

$Oy$ - ekseni ( $x=0$ )

$$A = \int_c^d g(y) dy$$




AĞUSTOS 2012						
H	P	S	Ç	P	C	Ct Pz
31			1	2	3	4 5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	

$$26\frac{26^3}{3} - \frac{3^3}{3}$$

AĞUSTOS  
AUGUST  
9.(8-1)2.3.2.2.3 PERŞEMBE  
THURSDAY

02



$$x = g(y) \leq 0$$

$$y = c \\ y = d$$


Oy-eleşik soru

$$A = - \int_c^d g(y) dy$$



$$A = A_1 + A_2$$

$$A = \int_c^e g(y) dy - \int_e^d g(y) dy$$




$$x = 3 \quad x = 6, \text{ OX-eleşik soru}$$

$$A = \int_3^6 x^2 dx$$

$$= \frac{x^3}{3} \Big|_3^6 \\ = 6^3 - 3^3$$

$$y = x^3 - 3x \quad OX-eleşik$$



$$y = 0$$

$$0 = x(x^2 - 3)$$

OS

$$x_1 = 0$$

$$x_2 = \sqrt{3}$$

$$x_3 = -\sqrt{3}$$




# 03

AGUSTOS  
AUGUST  
CUMA  
FRIDAY

$$\frac{dy}{dx}^2 - 2x^2 + 3x - \frac{4x^2 + 3x}{2}$$


AGUSTOS		2012		
S	C	P	C	P
31	1	2	3	4 5
6	7	8	9 10	11 12
13	14	15 16	17	18 19
20	21	22 23	24	25 26
27	28	29	30	31



$$A = \int_{-5}^5 (x^3 - 3x) dx$$

$$A = 2 \int_0^5 (x^3 - 3x) dx$$

$$\cancel{x} y = x^2 - 6x + 3 \quad x=2 \quad x=6 \quad 0x=0$$




$$A = - \int_2^3 (x^2 - 6x + 3) dx + \int_3^6 (x^2 - 6x + 3) dx$$

$$A = - \left( \frac{x^3}{3} - \frac{6x^2}{2} + 3x \right) \Big|_2^3 + \left( \frac{x^3}{3} - \frac{6x^2}{2} + 3x \right) \Big|_3^6$$

$$A = \frac{2}{3} + \left( \frac{6}{3} - 0 \right)$$

$$A = 2 \text{ br}^2$$



$$A = \int_1^8 x^3 dy$$

$$A = \int_1^8 y^{1/3} dy$$

$$\frac{16}{3} \cdot 3 = \frac{16}{3} b r^2$$

$$-\frac{3}{9} y^{\frac{4}{3}} \Big|_1^8$$



AĞUSTOS 2012

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

AĞUSTOS

AUGUST

CUMARTESİ

SATURDAY


04

$$y = \frac{1}{x}$$

$$x = 1$$

$$x = e$$


$$\partial x - \partial c$$



$$A = \int_1^e \frac{1}{x} dx$$

$$= \ln x \Big|_1^e$$

$$= 1 \ln e$$



$$\ln 0^+ = -\infty$$

$$\ln 1 = 0$$

$$\ln e = 1$$

$$\ln 100 = +\infty$$

AĞUSTOS

AUGUST


05

Nes egri alan



$$A = A_1 + A_2$$

$$A_2 = \int_a^b (f(x) - g(x)) + \int_c^b (g(x) - f(x))$$



$$A = \int_c^d (f(x) - g(x)) + \int_e^d (g(x) - f(x))$$



üsten alt toplu yada sağdan soldan  
gilas

# 06

AĞUSTOS  
AUGUST  
PAZARTESİ  
MONDAY


$$x^2 + 2x \\ x(2-x) \quad 0 \quad 13$$

10  
15  
20  
25  
30

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		



$$y = x^2 + 2 \\ y = 2x - x^2$$




$$\text{Og} = e^t$$

$$A = \int_0^3 [(x^2 + 2) - (2x - x^2)] dx$$

$$A = \int_0^3 (2x^2 - 2x + 2) dx$$

$$A = 2x^3/3 - 2x^2/2 + 2x \Big|_0^3 = 15 \text{ br}^2$$

$y = x^2$  ile  $x = y^2$  arası alan



$$A = \int_0^1 (x - x^2) dx$$

$$1/3 \text{ br}^2$$



AGUSTOS 2012

H	P	S	Ç	P	C	Cl	Pz
31		1	2	3	4	5	
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

$$x = y^2$$

$$y^2 = \frac{y+4}{3}$$

AGUSTOS  
AUGUST

$$3y^2 - y - 4 = 0$$

SALI


TUESDAY

07

~~A~~

$$y = 3x - 6$$

$$y+1 \quad 3y-9 \quad \frac{y}{3} \quad -1$$



$$A = \int_{-1}^{2} \left( \frac{y+4}{3} - y^2 \right) dy$$

$$A = \int_{-1}^{2} \left( \frac{1}{3}y + \frac{4}{3} - y^2 \right) dy$$

$$= \left[ \frac{1}{3} \frac{y^2}{2} + \frac{4}{3}y - \frac{y^3}{3} \right]_{-1}^{2}$$

$$\left( \frac{1}{3} \cdot \frac{16}{9} \cdot \frac{1}{2} + \frac{4}{3} \cdot \frac{4}{3} - \frac{1}{3} \cdot \frac{64}{27} \right)$$

$$- \left( \frac{1}{3} \cdot \frac{1}{2} + \frac{4}{3}(-1) + \frac{1}{3} \right)$$

$$y = 3x - x^3 \quad , \quad y = x$$


$$x(3-x^2)$$

$$x^3 + 2x$$

$$x^3 - 2x = 0$$

$$x(x^2 - 2) = 0$$

$$0 \quad \sqrt{2} \quad -\sqrt{2}$$



$\int_0^2$

$$A = 2 \int_0^2 (3x - x^3 - x) dx$$

$$= 2 \left( -\frac{x^4}{4} + 3x^2 \right) \Big|_0^2$$

$$= -2 + 12$$

$$= 10 \text{ br}^2$$



# 08

AĞUSTOS  
AUGUST  
ÇARŞAMBA  
WEDNESDAY

$$8y^2 = 2y$$


$$y^2 + 2y - 8 = 0$$

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

~~08~~

$$x = 2y$$

$$x = 8 - y^2$$



$$A = \int_{-2}^2 (8 - y^2 - 2y) dy$$

$$\left( 8y - \frac{y^3}{3} - 2y^2 \right) \Big|_{-2}^2$$

Vite  
Sınırı

Parabolik D. V. E. B.

$$\textcircled{1} \quad \begin{aligned} x &= g(t) \\ y &= h(t) \end{aligned} \quad \rightarrow t_1 \leq t \leq t_2$$

$$\begin{array}{l} x \geq 0 \\ x \geq b \end{array} \quad \forall t \in [t_1, t_2]$$

$$A = \int_{t_1}^{t_2} |h(t)| g'(t) dt$$

$\downarrow$   
 $y$   
 $\downarrow$   
 $x'$



AĞUSTOS 2012						
H	P	S	Ç	P	C	Ct Pz
31			1	2	3	4 5
32	6	7	8	9 10	11 12	
33	13	14	15	16 17	18 19	
34	20	21	22	23 24	25 26	
35	27	28	29	30	31	

AGÜSTOS  
AUGUST  
PERŞEMBE  
THURSDAY

09

$$\textcircled{2} \quad x = g(t) \\ y = h(t)$$

$$t_3 \leq t \leq t_4$$

$$y = c \\ y = d \quad \text{ve} \quad \text{Og eksen}$$

$$A = \int_{t_3}^{t_4} |g(t)| h'(t) dt$$

$|x|$        $y'$


~~A~~

$$x = 3 \cos t \\ y = 2 \sin t$$

$x=0$        $t_1 = ?$   
 $x=3$        $t_2 = ?$

$$x = 3 \cos t \Rightarrow 0 = 3 \cos t$$

$$\boxed{t_1 = \frac{\pi}{2}}$$



$$A = \int_{\pi/2}^0 |2 \sin t|, (-3 \cos t) dt$$

$$= 6 \int_0^{\pi/2} \sin^2 t dt$$

$$= 6 \int_0^{\pi/2} \frac{1}{2} [1 - \cos 2t] dt$$

$$= 3 \left[ t - \frac{\sin 2t}{2} \right] \Big|_0^{\pi/2}$$

$$= 3 \left( \left( \frac{\pi}{2} - 0 \right) - (0 - 0) \right)$$

$$= \frac{3\pi}{2} br^2$$

$$\cos 2x = 1 - 2 \sin^2 x$$

$$2 \sin^2 x = 1 - \cos 2x$$

$$\sin^2 x = \frac{1}{2} [1 - \cos 2x]$$

$$\text{Cevap: } 4A = 6\pi br^2$$




10

AĞUSTOS  
AUGUST  
CUMA  
FRIDAY

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

$x = a(t - \sin t)$   
 $y = a(1 - \cos t)$

cycloid eğrisinin bir çevresi  
 bir yayının  $Ox$  ekseninde  
 yaptığı 16m mesafe



$$x=0 \Rightarrow 0=a(t-\sin t) \Rightarrow t_1=0$$

$$x=2\pi a \Rightarrow 2\pi a=a(t-\sin t) \Rightarrow t_2=2\pi$$

$$A = \int_0^{2\pi} |a(1-\cos t)| \underbrace{a(1-\cos t)}_{|x'|} dt$$

-1 ile 1 arasıdır (pozitif olur)

$$= a^2 \int_0^{2\pi} (1-\cos t)^2 dt$$

$$= a^2 \int_0^{2\pi} [t - 2\cos t + \cos^2 t] dt$$

$$\textcircled{2} \quad \int_0^{2\pi} t dt = t \Big|_0^{2\pi} = 2\pi$$

$$\textcircled{3} \quad \int_0^{2\pi} 2\cos t dt = 2\sin t \Big|_0^{2\pi} = 0$$

(3)



AGUSTOS 2012

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

AĞUSTOS

AUGUST

CUMARTESİ

SATURDAY

11


$$\int_0^{2\pi} \cos^2 t dt = \int_0^{2\pi} \frac{1}{2} [2\cos 2t + 1] dt \\ = \frac{1}{2} \left[ \frac{\sin 2t}{2} + t \right] \Big|_0^{2\pi} = \frac{1}{2} [(0+2\pi) - (0+0)] \\ = \pi$$

cevap

$$A = d^2 [2\pi - 0 + \pi] = 3\pi d^2 \text{ br}^2$$



$$x = a \cos^3 t \\ y = a \sin^3 t$$



AĞUSTOS

AUGUST


PAZAR

SUNDAY


12

## HESAP HESABI

### ① Kesit Yontomu



$[a, b]$  aralığında yerlesitnelerin  $\int_a^b f(x) dx$   
Ox-eksenine dik kesitlerle elde edilen  
kesitlerin alanları  $A(x)$  olsun




# 13

AĞUSTOS  
AUGUST  
PAZARTESİ  
MONDAY

$$\frac{8}{16} \cdot \frac{16}{\beta} r^2$$


H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

$$H = \int_a^b A(x) dx$$



$$H = \int_c^d A(y) dy$$

~~BB~~ Tabanının bir karesi: 12 br<sup>2</sup> kare  
Yerstdakiler 16 br kare prizmeler hanesi



$$\frac{y}{16} = \frac{r}{12}$$

$$r = 3/4 y$$

$$H = \int_0^{16} \frac{3}{16} y^2 dy$$

$$= \frac{3}{16} \left( \frac{y^3}{3} \right) \Big|_0^{16}$$

$$= 768 \text{ br}^3$$

Pacut  $\rightarrow \frac{1}{3} \cdot 6^2 \cdot 16$   
 $\frac{1}{3} \cdot 16 \cdot 16$



AGUSTOS 2012

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

AGUSTOS


AUGUST

SALI

TUESDAY

14

~~yoncağı = 3 br kore hacmi~~



$$r = \sqrt{9 - x^2}$$


$$H = \int_{-3}^3 A(x) dx$$

$$= \int_{-3}^3 \pi (9 - x^2) dx$$

$$= 2\pi \left( 9x - \frac{x^3}{3} \right) \Big|_0^3$$

$$= 36\pi \text{ br}^3$$

## ② Disk Yontem



$y = f(x)$
$x = a$
$x = b$
$Ox - \text{eksen}$

$Ox$  ekseni etrafında  
dönüşüm

$$H = \pi \int_a^b [f(x)]^2 dx$$




# 15

AĞUSTOS  
AUGUST

ÇARŞAMBA  
WEDNESDAY


Lagrenel Eşitlerin hali!!!

AĞUSTOS 2012						
H	P	S	Ç	P	C	
31			1	2	3	4 5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	




$$\begin{cases} x = g(y) \\ y = c \\ y = d \end{cases} \rightarrow Oy\text{-ek} \text{ etrafı}\text{ donsun}$$

$$H = \pi \int_c^d [g(y)]^2 dy$$



$$\begin{cases} y = f(x) \\ y = g(x) \\ x = a \\ x = b \end{cases} \rightarrow Ox\text{-ek} \text{ etrafı}\text{ donsun}$$

$$H = \pi \int_a^b ([f(x)]^2 - [g(x)]^2) dx$$




$$\begin{cases} x = f(y) \\ x = g(y) \\ y = c \\ y = d \end{cases} \rightarrow Oy\text{-ek} \text{ etrafı}\text{ donsun}$$

$$H = \pi \int_c^d ([f(y)]^2 - [g(y)]^2) dy$$

~~$$g = \frac{1}{\sqrt{x}}$$~~

$$x = 1, x = e, \text{ } Ox\text{-ek}$$



AGUSTOS 2012						
H	P	S	Ç	P	C	Ct Pz
31		1	2	3	4	5
32	6	7	8	9	10	11
33	13	14	15	16	17	18
34	20	21	22	23	24	25
35	27	28	29	30	31	

$x^4 = 4x$   
 $x^3 - 4x = 0$   
 AGUSTOS  
 AUGUST  
 PERŞEMBE  
 THURSDAY

16

$$H = \pi \int_{1}^{e} \left( \frac{1}{\sqrt{x}} \right)^2 dx$$

$$= \pi (\ln x) \Big|_1^e$$

$$= \pi b r^3$$

A


$$y = \ln x$$

$$Ox - ek$$

$$Oy - ek$$

$$y = 1$$

Oy-elektif, den  $H = ?$



$$h_0^+ = \infty$$

$$h_1^- = 0$$

$$h_2 = 1$$

$$h_\infty = \infty$$

$$H = \pi \int_0^1 (e^y)^2 dy$$

$$= \pi \frac{e^{2y}}{2} \Big|_0^1$$

$$= \frac{\pi}{2} [e^2 - 1] \quad b r^3$$


A

$$y = 2\sqrt{x}$$

$$y = x^2$$

$$x = 1$$

Ox-elektif, döner



17

AĞUSTOS

AUGUST

CUMA  
FRIDAY

AĞUSTOS 2012							
H	P	S	C	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		


$$H = \pi \int_0^4 ((2\sqrt{x})^2 - (x^2)^2) dx$$

$$= \pi \int_0^4 (4x - x^4) dx$$

$$= \pi \left( 4 \frac{x^2}{2} - \frac{x^5}{5} \right) \Big|_0^4$$

$$= \frac{9\pi}{5} b^3$$

~~K~~ Merkezi  $(0,4)$ ,  $r=2$  eksenler  $Ox$  da



$$H = \pi \int_{-2}^2 \left[ (4 + \sqrt{4-x^2})^2 - (4 - \sqrt{4-x^2})^2 \right] dx$$

$$= 2\pi \int_0^2 (16\sqrt{4-x^2}) dx$$



AĞUSTOS 2012

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

AĞUSTOS

AUGUST

CUMARTESİ

SATURDAY

18

$$\begin{aligned}
 & \int \sqrt{h-x^2} dx \\
 & x = 2smt \\
 & dx = 2cost dt \\
 & = \int \sqrt{h-hsmt^2} 2cost dt \\
 & = \int h \cos^2 t dt \\
 & = \int h \frac{1}{2} [\cos 2t + 1] dt \\
 & = 2 \left[ \frac{1}{2} \sin 2t + t \right] = \sin 2t + t
 \end{aligned}$$


19

AĞUSTOS

AUGUST

PAZAR

SUNDAY



$$\begin{aligned}
 \sin 2t &= 2smt \cos t \\
 &= 2 \frac{x}{2} \frac{\sqrt{h-x^2}}{2} \\
 &= \frac{x\sqrt{h-x^2}}{2}
 \end{aligned}$$

$$H = 32\pi \left[ \frac{x\sqrt{h-x^2}}{2} + 2 \arcsin \frac{x}{2} \right] \Big|_0^2$$

$$\begin{aligned}
 &= 32\pi \left[ \pi - 0 \right] \\
 &= 32\pi^2 b r^3
 \end{aligned}$$




# 20

**AGUSTOS**  
AUGUST  
**PAZARTESİ**  
MONDAY

Disk → y sinir ise y kalkımları ve sağda sonuçlar

AGUSTOS 2012						
H	P	S	C	P	C	Ct Pz
31			1	2	3	4 5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	


## ③ KABUK YONTEMI



$$\begin{cases} y = f(x) \\ y = g(x) \\ x = a \\ x = b \end{cases}$$

oy egrisinden  
don

$$H = 2\pi \int_a^b |x[f(x) - g(x)]| dx$$



$$\begin{cases} x = f(y) \\ x = g(y) \\ y = c \\ y = d \end{cases}$$


Ox egrisinden  
don

$$H = 2\pi \int_c^d |y[f(y) - g(y)]| dy$$

$$\begin{aligned} y &= -x^2 + 6x - 3 \\ y &= x - 3 \end{aligned}$$

Oy-egrisinden don

$$\begin{aligned} x^2 - 4x + 3 \\ -3 \\ -1 \end{aligned}$$



AĞUSTOS 2012						
H	P	S	Ç	P	C	Çt Pz
31			1	2	3	4 5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	

AĞUSTOS  
AUGUST  
SALI  
TUESDAY

21

$$H = 2\pi \int_0^3 |x[-x^2 + 4x - 3 - x + 3]| dx$$

$$= 2\pi \int_0^3 |x[-x^2 + 3x]| dx$$

~~$$= 2\pi \int_0^3 |-x^3 + 3x^2| dx$$~~

$$= 2\pi \int_0^3 |x^2(3-x)| dx$$

↓ pozitif      ↓ pozitif  
 (0'dan 3'e)


$$= 2\pi \int_0^3 (3x^2 - x^3) dx$$

$$= 2\pi \left( x^3 - \frac{x^4}{4} \right) \Big|_0^3 =$$

~~$$y = \ln x$$~~

~~$0x - ek$~~        ~~$0x - ek$~~        $H = ?$   
 ~~$0y - ek$~~        ~~$0y - ek$~~   
 $y = 1$

$$H = 2\pi \int_0^1 |y| [1 - e^y] dy$$



# 22

AĞUSTOS

AUGUST

ÇARŞAMBA

WEDNESDAY

AGÜSTOS

2012

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

$$H = 2\pi \int_0^1 (ye^y) dy$$

$$\int ye^y dy$$

LAPTO

$$y = u$$

$$e^y dy = du$$

$$e^y = v$$

$$ye^y - \int e^y dy$$


$$ye^y - e^y$$

$$e^y [y - 1]$$

$$H = 2\pi \left( e^y [y - 1] \right) \Big|_0^1$$

$$= 2\pi \left( e^1 (1-1) - e^0 (0-1) \right)$$

$$= 2\pi$$




$$H = H_1 + H_2$$

$$= \pi \int_0^1 (1^2 - 0^2) dx + \pi \int_e^1 (e^2 - h_x^2) dx$$

$$= \pi + \pi$$

$$\Rightarrow 2\pi$$



AGUSTOS 2012

H	P	S	Ç	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

$$H = \pi \int_{0}^{\pi/2} (1 - \sin x)^2 dx$$

AGUSTOS

AUGUST

PERŞEMBE  
THURSDAY

23

$$y = \sin x$$


$$y = 1$$

$$Oy - ek$$

$$y = 1 \text{ etrafında döndürse}$$

$$H = ?$$

Dikdörtgen



$$f = \text{uzaklık} = 1 - \sin x$$

$$H = \pi \int_0^{\pi/2} (1 - \sin x)^2 dx$$

$$= \pi \int_0^{\pi/2} (1 - 2\sin x + \sin^2 x) dx$$

$$= \pi \left( x + 2\cos x + \frac{x}{2} - \frac{\sin 2x}{4} \right) \Big|_0^{\pi/2}$$

$$= \frac{3}{4} \pi^2 - 2\pi b r^3$$



# 24

AĞUSTOS  
AUGUST  
CUMA  
FRIDAY

AĞUSTOS 2012						
H	P	S	Ç	P	C	Pz
31				1	2	3
32	6	7	8	9	10	11
33	13	14	15	16	17	18
34	20	21	22	23	24	25
35	27	28	29	30	31	

Pozisyon Değişkenler Vektor Eşitliğinin Formül

$$x = v(t)$$

$$t_1 \leq t \leq t_2$$

$$y = w(t)$$

Sınırlılığı da  
Ox-eksen dan

$$H=?$$

$$H = \pi \int_{t_1}^{t_2} y^2 x' dt$$

$$x = v(t)$$

$$t_3 \leq t \leq t_4$$

$$y = w(t)$$


Sınırlılığı da  
Oy-eksen dan  
H=1

$$H=1$$

$$H = \pi \int_{t_3}^{t_4} x^2 y' dt$$

$$\begin{aligned} x &= (t - \sin t) \\ y &= (1 - \cos t) \end{aligned}$$

cyclloid eğrisinin bir  
bergasıının Ox ekseninden  
H=?



$$H = \pi \int_0^{2\pi} (1 - \cos t)^2 (1 - \cos t)' dt$$

$$= \pi \int_0^{2\pi} (1 - 3\cos t + 3\cos^2 t - \cos^3 t) dt$$



AGUSTOS 2012

H	P	S	C	P	C	Ct	Pz
31		1	2	3	4	5	
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

$$\cos^2 t = \frac{1}{2} [1 + \cos 2t]$$

AĞUSTOS

AUGUST

CUMARTESİ


SATURDAY

25

$$H = \pi \left[ t - 3Sm t + \frac{3t}{2} + \frac{3\sin 2t}{4} + Sm t - \frac{\sin 3t}{3} \right] \Big|_0^{2\pi}$$

$$H = 5\pi^2 br^3$$

Eğri Uzunluğu



$$\begin{aligned}
 & \int \cos^3 t dt \\
 & \int \cos^2 t \cos t dt \\
 & \int (1 - \sin^2 t) \cos t dt \\
 & \int (1 - v^2) dv \\
 & = v - \frac{v^3}{3} \\
 & = Sm t - \frac{\sin 3t}{3}
 \end{aligned}$$

AĞUSTOS

AUGUST

PAZAR

26

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY

SUNDAY


SUNDAY

# 27

AĞUSTOS  
AUGUST  
PAZARTESİ  
MONDAY

AĞUSTOS 2012						
H	P	S	C	P	C	Ct Pz
31			1	2	3	4 5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	

~~A~~  $r=1$  olur conform çevre yüzüğü



$$x^2 + y^2 = 1$$

$$y = \sqrt{1-x^2}$$

$$y = (1-x^2)^{1/2}$$

$$y' = \frac{1}{2} (1-x^2) \cdot (-2x)$$

$$y' = \frac{-x}{\sqrt{1-x^2}}$$

$$(y')^2 = \frac{x^2}{1-x^2}$$

$$l = 4 \int_0^1 \sqrt{1 + \frac{x^2}{1-x^2}} dx$$

$$= 4 \int_0^1 \frac{1}{\sqrt{1-x^2}} dx$$

$$= 4 \arcsin l_0$$

$$= 4 \frac{\pi}{2}$$

$$= 2\pi$$



AĞUSTOS 2012						
H	P	S	Ç	P	C	Cl
31		1	2	3	4	5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	

$$\frac{1 - \cancel{3x^2} + x^4}{\cancel{16x^2}}$$

AĞUSTOS  
AUGUST  
SALI  
TUESDAY

28

$$y = \ln x - \frac{x^2}{8}, [2, 4] \text{ da } l = ?$$

$$\begin{aligned} y &= \frac{1}{x} - \frac{1}{6}x \Rightarrow (y')^2 = \frac{1}{x^2} - \frac{1}{2} + \frac{x^2}{16} \\ l &= \int_2^4 \sqrt{1 + \left( \frac{1}{x^2} - \frac{1}{2} + \frac{x^2}{16} \right)} dx \\ &\quad \boxed{\frac{x^4 + 8x^2 + 1}{16x^2}} \end{aligned}$$

$$(a-b)^2 = a^2 - 2ab + b^2$$

$$\begin{aligned} &+ 2ab \\ &= (a+b)^2 \end{aligned}$$

$$\begin{aligned} 1 + (y')^2 &= 1 + \left( \frac{1}{x} - \frac{x}{6} \right)^2 \\ &= 1 + \left( \frac{1}{x} \right)^2 - \frac{1}{2} + \left( \frac{x}{6} \right)^2 \end{aligned}$$

$$= \left( \frac{1}{x} \right)^2 + \frac{1}{2} + \left( \frac{x}{6} \right)^2$$

$$= \left( \frac{1}{x} + \frac{x}{6} \right)^2$$

$$\begin{aligned} l &= \int_2^4 \left( \frac{1}{x} + \frac{x}{6} \right) dx = \ln x + \frac{x^2}{8} \Big|_2^4 \\ &= \left( \ln 2 + \frac{3}{2} \right) \text{ br} \end{aligned}$$



# 29

AĞUSTOS

AUGUST

ÇARŞAMBA  
WEDNESDAY

AĞUSTOS 2012							
H	P	S	C	P	C	Ct	Pz
31			1	2	3	4	5
32	6	7	8	9	10	11	12
33	13	14	15	16	17	18	19
34	20	21	22	23	24	25	26
35	27	28	29	30	31		

Parametrik Eşitsizlik Vanligi

$$x = v(t)$$

$$t_1 \leq t \leq t_2$$


$$y = \varphi(t)$$

$$l = \int_{t_1}^{t_2} \sqrt{(x')^2 + (y')^2} dt$$

~~K~~  $x = (t - \sin t)$   
 $y = (1 - \cos t)$


$$0 \leq t \leq 2\pi$$

parametrik  
eşitsizlik  
başlangıç  
eşitliklerin



$$x' = 1 - \cos t$$

$$y' = \sin t$$



$$l = \int_0^{2\pi} \sqrt{(1 - \cos t)^2 + (\sin t)^2} dt$$

$$(x')^2 + (y')^2 = 1 - 2\cos t + \cos^2 t + \sin^2 t$$


$$= 2(1 - \cos t)$$

$$= 2[1 - (1 - 2\sin^2 \frac{t}{2})]$$

$$= 4 \sin^2 \frac{t}{2}$$



AĞUSTOS 2012						
H	P	S	C	P	C	Çt Pz
31		1	2	3	4	5
32	6	7	8	9	10	11 12
33	13	14	15	16	17	18 19
34	20	21	22	23	24	25 26
35	27	28	29	30	31	



AĞUSTOS

AUGUST

PERŞEMBE

THURSDAY

30

$$\begin{aligned}
 l &= \int_0^{2\pi} \sqrt{4 \sin^2 \frac{t}{2}} \, dt \\
 &= \int_0^{2\pi} 2 \sin \frac{t}{2} \, dt \\
 &= -4 \cos \frac{t}{2} \Big|_0^{2\pi} \\
 &= 8b
 \end{aligned}$$

APT  
Geliştiğimiz  
Sıra  
Tarih

~~$x = \cos t$~~

$$-\pi \leq t \leq \pi$$

$$y = t + \sin t$$

efem  
isikli bolu  
 $\sqrt{2}$  kg ✓

$$\begin{aligned}
 (x')^2 + (y')^2 &= \sin^2 t + 1 + 2\cos t + \cos^2 t \\
 &= 2(1 + \cos t) \\
 &= 2\left(1 + \left(2\cos^2 \frac{t}{2} - 1\right)\right) \\
 &= 4\cos^2 \frac{t}{2}
 \end{aligned}$$

$$\begin{aligned}
 l &= \int_{-\pi}^{\pi} \sqrt{4 \cos^2 \frac{t}{2}} \, dt \\
 &= \int_{-\pi}^{\pi} 2 \cos \frac{t}{2} \, dt = 8 \int_0^{\pi/2} \cos \frac{t}{2} \, dt = 8\sqrt{2}
 \end{aligned}$$

$\sqrt{2}$  kg ✓




31

AGÜSTOS  
AUGUST  
CUMA  
FRIDAY

AGÜSTOS 2012						
H	P	S	C	P	C	Ct
31			1	2	3	4
32	6	7	8	9	10	11
33	13	14	15	16	17	18
34	20	21	22	23	24	25
35	27	28	29	30	31	

## Dönel Yüzeylerin Alanı

(a)



$y = f(x)$  fonksiyonu  $[a, b]$


Ox eksen etrafında

Dönel yüzeyin yüzey alanı

$$S = 2\pi \int_a^b |y| \sqrt{1 + (y')^2} dx$$

$$= 2\pi \int_a^b |f(x)| \sqrt{1 + [f'(x)]^2} dx$$

(b)



$x = f(y)$  fonksiyonu  $[a, b]$


Oy eksen etrafında

Dönel yüzey alanı

$$S = 2\pi \int_c^d |x| \sqrt{1 + (x')^2} dy$$

$$= 2\pi \int_c^d |f(y)| \sqrt{1 + [f'(y)]^2} dy$$

(c)



$y = f(x)$  fonksiyonu  $[a, b]$


$y = k$  ekstra eksen etrafında

Dönel yüzey alanı

$$S = 2\pi \int_a^b |f(x) - k| \sqrt{1 + [f'(x)]^2} dx$$



EYLÜL						2012		
H	P	S	Ç	P	C	Ct	Pz	
35						1	2	
36	3	4	5	6	7	8	9	
37	10	11	12	13	14	15	16	
38	17	18	19	20	21	22	23	
39	24	25	26	27	28	29	30	



EYLÜL  
SEPTEMBER  
CUMARTESİ  
SATURDAY

# 01




$$\left\{ \begin{array}{l} x=h \\ x=f(y) \end{array} \right.$$

$$x=f(y) \text{ form } \{x,y\}$$

$x=h$  et den  
 $y$  sey ols

$$S = 2\pi \int |f(y) - h| \sqrt{1 + [f'(y)]^2} dy$$

~~$\oint$~~   $r=3$  korem yozey olsu



$$x^2 + y^2 = 9$$

$$y = \pm \sqrt{9 - x^2} \Rightarrow y' = \frac{-x}{\sqrt{9 - x^2}}$$

$$S = 2\pi \int_{-3}^3 \sqrt{|9 - x^2|} \sqrt{1 + \frac{x^2}{9 - x^2}} dx$$

$$= 2\pi \int_0^3 \sqrt{9 - x^2} \frac{3}{\sqrt{9 - x^2}} dx$$

$$= 12\pi \times \int_0^3 1 = 36\pi b r^2$$


EYLÜL  
SEPTEMBER  
PAZAR  
SUNDAY

# 02

~~$\oint$~~   $y = x^3$

$$0 \leq x \leq 1$$

Ox-etsel  $S = ?$



$$y' = 3x^2$$

$$(y')^2 = 9x^4$$

$$S = 2\pi \int_0^1 |x^3| \sqrt{1 + 9x^4} dx$$

$$= 2\pi \int_0^1 x^3 \sqrt{1 + 9x^4} dx$$

$$1 + 9x^4 = 0$$

$$36x^3 dx = 0$$



# 03

EYLÜL  
SEPTEMBER  
PAZARTESİ  
MONDAY

EYLÜL 2012						
H	P	S	Ç	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

$$S = 2\pi \int_0^1 \sqrt{1 + \frac{dy}{dx}^2} dx$$

$$= \frac{\pi}{18} \frac{u^{3/2}}{3/2} \Big|_0^1$$

$$= \frac{\pi}{18} (1 + 9x^2)^{3/2} \Big|_0^1$$


$$= \frac{\pi}{18} [90^{3/2} - 1] b r^2$$

~~$x = \sqrt{y}$~~

$0 \leq y \leq 2$

Oy-efekt den

Free slot



$$S = 2\pi \int_0^2 |\sqrt{y}| \sqrt{1 + \frac{1}{4y}} dy$$

$$S = 2\pi \int_0^2 \sqrt{y} \frac{\sqrt{4y+1}}{2\sqrt{y}} dy$$

$$= \pi \int_0^2 \sqrt{4y+1} dy$$

$ayt+1=0$   
 $ayt=2k$

$$= \pi \int_0^2 u^{1/2} \frac{du}{2}$$

$$= \frac{\pi}{4} \cdot \frac{u^{3/2}}{3/2} \Big|_0^2$$

$$= \frac{\pi}{6} (4y+1)^{3/2} \Big|_0^2$$

$$= \frac{\pi}{6} (27-1) = \frac{13}{3} br^2$$




EYLÜL 2012					
H	P	S	Ç	P	C
35				1	2
36	3	4	5	6	7
37	10	11	12	13	14
38	17	18	19	20	21
39	24	25	26	27	28
	29	30			

EYLÜL  
SEPTEMBER  
SALI  
TUESDAY


04

$$3y^2 = x(3-x)^2$$

İlmigirin ox-eksen et  $\delta\sigma$   $s=?$



$$y = \sqrt{\frac{x(3-x)^2}{3}} = \frac{1}{3} \sqrt{x} (3-x)$$



$$y' = \frac{1}{3} \left[ \frac{1}{2\sqrt{x}}(3-x) + \sqrt{x}(-1) \right]$$

$$= \frac{3-x}{6\sqrt{x}} - \frac{\sqrt{x}}{3}$$

$$(y')^2 = \frac{9-6x+x^2}{36x} = \frac{6\cancel{x^2}-2x\cancel{x}}{18\cancel{x}} + \frac{x}{36x}$$

$$= \frac{9-6x+x^2-12x+4x^2+4x^2}{36x}$$

$$= \frac{9x^2-18x+9}{36x}$$

$$1+(y')^2 = \frac{9x^2+18x+9}{36x} = \frac{x^2+2x+1}{4x}$$




# 05

EYLÜL  
SEPTEMBER  
ÇARŞAMBA  
WEDNESDAY

EYLÜL 2012						
H	P	S	Ç	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

$$\begin{aligned}
 S &= 2\pi \int_0^3 \left| \frac{1}{3} \sqrt{x} (3-x) \right| \sqrt{\frac{(x+1)^2}{6x}} dx \\
 &= 2\pi \int_0^3 \frac{1}{3} \sqrt{x} (3-x) \frac{(x+1)}{2\sqrt{x}} dx \\
 &= \frac{\pi}{3} \int_0^3 (-x^2 + 2x + 3) dx \\
 &= \frac{\pi}{3} \left[ \frac{-x^3}{3} + x^2 + 3x \right] \Big|_0^3 \\
 &= \frac{\pi}{3} [-9 + 9 + 9] = 3\pi b r^2
 \end{aligned}$$

$x^2 + (y-1)^2 = 1$ ,  $y = -1$  et doğrular  $S = ?$



$$y = 1 \pm \sqrt{1-x^2}$$

$$S = 2\pi \int_0^b |f(x) - k| \sqrt{1 + [f'(x)]^2} dx$$

$$y = -1$$



EYLÜL 2012						
H	P	S	Ç	P	C	Ct Pz
35					1	2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

EYLÜL  
SEPTEMBER  
PERŞEMBE  
THURSDAY

06

(a)  $y = 1 + \sqrt{1-x^2}$  (ösgen kisim)

$$S_1 = 2\pi \int_{-1}^1 \left| 1 + \sqrt{1-x^2} - (-1) \right| \sqrt{\frac{1}{1-x^2}} dx$$

$$y' = \frac{-x}{\sqrt{1-x^2}} \Rightarrow (y')^2 = \frac{x^2}{1-x^2}, \Rightarrow 1+(y')^2 = \frac{1}{1-x^2}$$

(b)  $y = 1 - \sqrt{1-x^2}$  (alt kisim)

$$1+(y')^2 = \frac{1}{1-x^2}$$

$$S_1 = 2\pi \int_{-1}^1 \left| 1 + \sqrt{1-x^2} - (-1) \right| \sqrt{\frac{1}{1-x^2}} dx$$

$$S_2 = 2\pi \int_{-1}^1 \left| 1 - \sqrt{1-x^2} - (-1) \right| \sqrt{\frac{1}{1-x^2}} dx$$

$$S_1 + S_2 = S = 2\pi \int_{-1}^1 |4| \frac{1}{\sqrt{1-x^2}} dx$$

$$= 16\pi \int_0^1 \frac{1}{\sqrt{1-x^2}} dx$$

$$= 16\pi \arcsin x \Big|_0^1$$

$$= 16\pi (\arcsin 1 - \arcsin 0)$$

$$= 16\pi \left( \frac{\pi}{2} - 0 \right) = 8\pi^2 b r^2$$



07

EYLÜL  
SEPTEMBER  
CUMA  
FRIDAY

EYLÜL 2012						
H	P	S	Ç	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

# Parametrik Eşitlerin Öğrenildiği Yüzey Alanı

(a)

$$x = u(t)$$

$$y = v(t)$$

$$t_1 \leq t \leq t_2$$

Parametrikin Ox-de  
efektör hanesi ile  
alan统筹推进  
yazdırılır

$$S = 2\pi \int_{t_1}^{t_2} |y| \sqrt{(x')^2 + (y')^2} dt$$

(b)

$$x = u(t)$$

$$y = v(t)$$

$$t_3 \leq t \leq t_4$$

Par. eğrisi Oy-ele  
etr dor ile  
alan统筹推进  
yazdırılır

$$S = 2\pi \int_{t_3}^{t_4} |x| \sqrt{(x')^2 + (y')^2} dt$$


(c)

$$x = t - \sin t$$

$$y = 1 - \cos t$$

$$0 \leq t \leq 2\pi \quad \text{Ox-ele dor}$$

$$S = ?$$



$$x' + y' = 1 - \cos t + \sin t$$

$$\times \quad (x')^2 + (y')^2 = 1 - 2\cos t + \cos^2 t + \sin^2 t \\ = 2(1 - \cos t)$$

$$S = 2\pi \int_0^{2\pi} |1 - \cos t| \sqrt{2 - 2\cos t} dt$$

$$= 2\sqrt{2} \pi \int_0^{2\pi} (1 - \cos t) \sqrt{1 - \cos t} dt$$



EYLÜL 2012									
H	P	S	Ç	P	C	Ct	Pz		
35						1	2		
36	3	4	5	6	7	8	9		
37	10	11	12	13	14	15	16		
38	17	18	19	20	21	22	23		
39	24	25	26	27	28	29	30		

$$1 - \cos t = 0$$

(Sabit de -du)

EYLÜL  
SEPTEMBER  
CUMARTESİ  
SATURDAY

08

$$1 - \cos t = 1 - \left\{ 1 - 2 \sin^2 \frac{t}{2} \right\} = 2 \sin^2 \frac{t}{2}$$

$$S = 252 \pi \int_0^{2\pi} (1 - \cos t) \sqrt{2 \sin^2 \frac{t}{2}} dt$$

$$= 6\pi \int_0^{2\pi} 2 \sin^2 \frac{t}{2} \sin \frac{t}{2} dt$$

$$= 8\pi \int_0^{2\pi} \left( 1 - \cos^2 \frac{t}{2} \right) \sin \frac{t}{2} dt$$

$$= 8\pi \int_0^{2\pi} (1 - v^2) (-2dv)$$

$$= -16\pi \int_0^{2\pi} \left[ v - \frac{v^3}{3} \right] dv$$

$$= -16\pi \int_0^{2\pi} \left[ \cos \frac{t}{2} - \frac{\cos^3 \frac{t}{2}}{3} \right] dt$$

$$= -16\pi \left[ \left( 1 + \frac{1}{3} \right) - \left( 1 - \frac{1}{3} \right) \right]$$

$$= \frac{64}{3} \pi b r^2$$

EYLÜL  
SEPTEMBER  
PAZAR  
SUNDAY

09

$$\begin{cases} \cos \frac{t}{2} = 0 \\ -\frac{1}{2} \sin \frac{t}{2} dt = du \end{cases}$$



10

EYLÜL  
SEPTEMBER  
PAZARTESİ  
MONDAY

EYLÜL 2012						
H	P	S	C	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

## Genel Leisteindmüz (Has Olmayan) İnt.

İntip sınırlıdan er or 611 sonsuz olmaz  
değer sınırları ise yerine yazılındır sonsuz olmazsa  
①  $y = f(x)$  fonk  $[a, t]$  integrallenebilir olsun

$$\int_a^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

②  $y = f(x)$  fonk  $[t, b]$  m+

$$\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

③  $y = f(x)$  fonk  $\Lambda$ m

$$\int_{-\infty}^{+\infty} f(x) dx = \lim_{t \rightarrow -\infty} \int_t^c f(x) dx + \lim_{t \rightarrow +\infty} \int_c^t f(x) dx$$

Bu integralin sonuc versa "yolculuk"  
seyi yoksad sonuc sonsuz veya baslegsos  
"İndirim" demir



EYLÜL		2012	
H	P	S	C
35			1 2
36	3	4	5 6 7
37	10	11	12 13 14 15 16
38	17	18	19 20 21 22 23
39	24	25	26 27 28 29 30

$\frac{e^x}{Tn!}$

EYLÜL  
SEPTEMBER  
SALI  
TUESDAY

11

$$\int_1^{\infty} e^{-x} dx$$

$$= \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx$$

$$= \lim_{t \rightarrow \infty} (-e^{-x}) \Big|_1^t$$

$$= \lim_{t \rightarrow \infty} \left( -e^{-t} + e^{-1} \right)$$

$$= \lim_{t \rightarrow \infty} \left( -\frac{1}{e^t} + \frac{1}{e} \right)$$

$$= -\frac{1}{e^\infty} + \frac{1}{e}$$

$$= -\frac{1}{\infty} + \frac{1}{e}$$

$$= 0 + \frac{1}{e} = \frac{1}{e}$$

serice çıktı yani yoklansık  
yoklansılık oranı ( $1/e$ )dır.

$y = \frac{1}{1+x^2}$  ile  $Ox$ -el enosunda keler alan

$$A = \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$$

$$= \lim_{t \rightarrow \infty} \int_{-t}^t \frac{1}{1+x^2} dx + \lim_{s \rightarrow \infty} \int_s^{\infty} \frac{1}{1+x^2} dx$$

$$= \lim_{t \rightarrow \infty} (\arctan x \Big|_0^t) + \lim_{s \rightarrow \infty} (\arctan x \Big|_0^s)$$



# 12

EYLÜL  
SEPTEMBER  
ÇARŞAMBA  
WEDNESDAY

EYLÜL 2012						
H	P	S	Ç	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

$$\begin{aligned} & \lim_{t \rightarrow -\infty} (\arctan 0 - \arctan t) + \lim_{s \rightarrow \infty} (\arctan s - \arctan 0) \\ &= -\arctan(-\infty) + \arctan(\infty) \end{aligned}$$

$$= \frac{\pi}{2} + \frac{\pi}{2}$$

$$= \pi b r^2$$

yakınsak

~~$\int_0^b x^2 dx$~~

$$= \lim_{t \rightarrow \infty} \int_0^t x^2 dx$$

$$= \lim_{t \rightarrow \infty} \left( \frac{x^3}{3} \right) \Big|_0^t$$

$$= \lim_{t \rightarrow \infty} \left( \frac{t^3}{3} \right) = \infty$$

ırıksız

~~I.TP~~

①  $y = f(x)$ ,  $[a, b]$

SILİNTİ  
Sayı

$$\int_a^b f(x) dx = \lim_{t \rightarrow b} \int_0^t f(x) dx$$

②  $b[y = f(x)]$ ,  $[0, b]$

$$\int_0^b f(x) dx = \lim_{t \rightarrow b} \int_0^t f(x) dx$$



EYLÜL 2012					
H	P	S	Ç	P	Ct Pz
35				1	2
36	3	4	5	6	7
37	10	11	12	13	14
38	17	18	19	20	21
39	24	25	26	27	28
				29	30

EYLÜL  
SEPTEMBER  
PERŞEMBE  
THURSDAY

13

c)  $y = f(x)$ ,  $[a, b]$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_{a+i-1}^{a+i} f(x) dx + \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_{c+i-1}^{c+i} f(x) dx$$

$$\int_0^1 \frac{dx}{\sqrt[4]{1-x}}$$

$$= \lim_{t \rightarrow 1^-} \int_0^t (1-x)^{-1/4} dx$$

$$= \lim_{t \rightarrow 1^-} \left[ -\frac{4}{3} (1-x)^{3/4} \Big|_0^t \right]$$

$$\begin{cases} 1-x=u \\ -dx=du \end{cases}$$

$$\int_0^1 u^{-1/4} (-du)$$

$$= -\frac{4}{3} (1-x)^{3/4}$$

$$= \lim_{t \rightarrow 1^-} \left[ -\frac{4}{3} \left( (1-t)^{3/4} - 1 \right) \right]$$

$$= \frac{4}{3} \quad \underline{\text{yolun sonu}}$$



# 14

EYLÜL  
SEPTEMBER  
CUMA  
FRIDAY

$$\int_{-1}^1 \frac{1}{x^2} dx \rightarrow \textcircled{6} \quad \int_{-1}^1 \frac{1}{x^2} dx + \int_{-1}^1 \frac{1}{x^2} dx \quad \textcircled{0}$$

II. TSP

EYLÜL 2012						
H	P	S	C	P	C	Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

~~K~~  $\int_0^2 \frac{1}{x^2} dx$

$$= \lim_{t \rightarrow 0} \int_t^2 \frac{1}{x^2} dx$$

$$= \lim_{t \rightarrow 0} \int_t^2 x^{-2} dx$$

$$= \lim_{t \rightarrow 0} \left( -\frac{1}{x} \right) \Big|_t^2$$

$$= \lim_{t \rightarrow 0} \left( -\frac{1}{2} + \frac{1}{t} \right)$$

$$= \infty \quad \underline{\text{İfadeler}}$$

~~K~~  $\int_{-1}^0 \frac{1}{1-x^2} dx$

$$= \int_{-1}^0 \frac{1}{1-x^2} dx + \int_0^1 \frac{1}{1-x^2} dx$$

$$= \lim_{t \rightarrow -1} \int_t^0 \frac{1}{1-x^2} dx + \lim_{s \rightarrow 1} \int_0^s \frac{1}{1-x^2} dx$$

$$= \lim_{t \rightarrow -1} (\arcsin t - \arcsin(-t)) + \lim_{s \rightarrow 1} (\arcsin s - \arcsin 0)$$

$$\int \frac{1}{1-x^2} dx = \frac{A}{1+x} + \frac{B}{1-x} = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right|$$



EYLÜL 2012						
H	P	S	Ç	P	C	Pz
35				1	2	
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

EYLÜL  
SEPTEMBER  
CUMARTESİ  
SATURDAY **15**

$$\begin{aligned}
 &= \lim_{t \rightarrow -1} \left[ \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right| \Big|_0^t \right] + \lim_{s \rightarrow 1} \left[ \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right| \Big|_0^s \right] \\
 &= \lim_{t \rightarrow -1} \frac{1}{2} \left[ \ln 1 - \ln \left| \frac{1+t}{1-t} \right| \right] + \lim_{s \rightarrow 1} \left( \frac{1}{2} \left[ \ln \left| \frac{1+s}{1-s} \right| - \ln 1 \right] \right) \\
 &= \frac{\infty}{2} + \frac{\infty}{2} = \infty \quad \text{(räksel)}
 \end{aligned}$$

~~III. Tp~~ seginin bir sonsuz.  
 Seginin olğuleri sabitdir.  
 $\int f(x) dx$  sonsuz  
 $\alpha \rightarrow$  sitedi

EYLÜL  
SEPTEMBER  
PAZAR  
SUNDAY **16**

~~III. Hp~~  $\Rightarrow$  Itp + Itp

$\alpha \rightarrow$  sitedi  
 $\int f(x) dx$   
 $\infty \rightarrow$  sonsuz

$\int_{-\infty}^{\infty} f(x) dx$  orada  $\alpha$  sayısi sitedi.



# 17

EYLÜL  
SEPTEMBER  
PAZARTESİ  
MONDAY

EYLÜL 2012						
H	P	S	Ç	P	C	Cl
35					1	2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

$$\int_0^{\infty} \frac{1}{x^2} dx = \underbrace{\int_1^{\infty} \frac{1}{x^2} dx}_{\text{I. TIP}} + \underbrace{\int_0^1 \frac{1}{x^2} dx}_{\text{II. TIP}}$$

$$= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2} dx + \lim_{s \rightarrow 0} \int_s^1 \frac{1}{x^2} dx$$

Açıklama

$$\int \frac{dx}{x^3} = \lim_{t \rightarrow \infty} \int_1^t \frac{dx}{x^3}$$

$$= \lim_{t \rightarrow \infty} \left[ \frac{-1}{2x^2} \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \left( \frac{-1}{2t^2} + \frac{1}{2} \right)$$

$$= 0 + \frac{1}{2}$$

$$= 1/2 \quad (\text{Yakınsaklı})$$

$$\int_2^{\infty} \frac{dx}{x\sqrt{x^2-1}} = \lim_{t \rightarrow \infty} \int_2^t \frac{dx}{x\sqrt{x^2-1}}$$

$$\int \frac{dx}{x\sqrt{x^2-1}}$$

$$\begin{aligned} \frac{1}{x} &= t \Rightarrow \frac{1}{t} = x \\ \cancel{\frac{1}{x^2} dx} &= dt \\ -\frac{1}{t^2} dt &= dx \end{aligned}$$



EYLÜL 2012

H	P	S	Ç	P	C	Cl	Pz
35					1	2	
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

$$\lim_{t \rightarrow 0} \frac{2}{t} = \infty$$

$$\frac{2}{t} \leftarrow 60 \text{ cm s}^{-2}$$

$x^{-\frac{1}{2}}$   $x^{-\frac{1}{3}}$   $x^{-\frac{1}{4}}$   $x^{-\frac{1}{5}}$   $x^{-\frac{1}{6}}$   
**EYLÜL**  
**SEPTEMBER**  
**SALI**  
**TUESDAY**

**18**

$$\begin{aligned}
 &= \int \frac{t - \frac{1}{t^2} dt}{\sqrt{1-t^2}} = \int \frac{-dt}{t \sqrt{1-t^2}} = - \int \frac{dt}{\sqrt{1-t^2}} \\
 &= \lim_{t \rightarrow 0} \left[ -\arcsin\left(\frac{1}{t}\right) \right] \Big|_0^{\infty} \\
 &= \lim_{t \rightarrow \infty} \left( -\arcsin\frac{1}{t} + \arcsin\frac{1}{2} \right) = 0 + \frac{\pi}{6} = \frac{\pi}{6}
 \end{aligned}$$



$$\begin{aligned}
 &\int_0^{\infty} x e^{-x} dx = \lim_{t \rightarrow 0} \int_0^t \frac{dx}{x \sqrt{x}}
 \end{aligned}$$

$$= \lim_{t \rightarrow 0} \left( -2x^{-\frac{1}{2}} \right) \Big|_0^t$$

$$= \lim_{t \rightarrow 0} \left( -2 + \frac{2}{\sqrt{t}} \right) \Big|_0^{\infty}$$

$$\approx -2 + \infty$$

$$\approx \infty$$

# 19

EYLÜL  
SEPTEMBER  
ÇARŞAMBA  
WEDNESDAY

$\arcsin x =$

EYLÜL							2012	
H	P	S	Ç	P	C	Ct	Pz	
35						1	2	
36	3	4	5	6	7	8	9	
37	10	11	12	13	14	15	16	
38	17	18	19	20	21	22	23	
39	24	25	26	27	28	29	30	

$$\int_0^1 \frac{\arcsin x}{\sqrt{1-x^2}} dx$$

$$= \lim_{t \rightarrow 1} \int_0^t \frac{\arcsin x}{\sqrt{1-x^2}} dx$$

$$\boxed{\arcsin x = v}$$

$$\frac{1}{\sqrt{1-x^2}} dx = dv$$

$$= \lim_{t \rightarrow 1} \left( \frac{(\arcsin x)^2}{2} \right) \Big|_0^t$$

$$= \lim_{t \rightarrow 1} \left( \frac{\arcsin^2 t}{2} - \frac{\arcsin^2 0}{2} \right)$$

$$= \frac{\pi^2}{8} \quad (\text{yol uysa})$$

$$\int_0^1 \frac{dx}{\sqrt{1-x^2}} = \int_0^{1/2} \frac{dx}{\sqrt{1-x^2}} + \int_{1/2}^1 \frac{dx}{\sqrt{1-x^2}}$$

$$\textcircled{a} \quad \int_0^{1/2} \frac{dx}{\sqrt{1-x^2}} = \lim_{t \rightarrow 0} \int_t^{1/2} \frac{dx}{\sqrt{1-x^2}}$$



EYLÜL 2012

H	P	S	Ç	P	C	Cl	Pz
35						1	2
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

EYLÜL  
SEPTEMBER  
PERŞEMBE  
THURSDAY

20

$$\int \frac{dx}{\sqrt{x-x^2}} = \int \frac{dx}{\sqrt{x(x-1)}} = \int \frac{dx}{x\sqrt{1-\frac{1}{x}}}$$

$$\begin{aligned} \frac{1}{x} &= v \\ 1 &= \end{aligned}$$

$$= \int \frac{dx}{\sqrt{\left(\frac{1}{2}\right)^2 - v^2}}$$

$$= dx \operatorname{csch} \frac{v}{\frac{1}{2}}$$

$$= \operatorname{arcsm} 2v$$

$$= \operatorname{arcsm}(2x-1)$$

$$x - x^2 = -\frac{1}{4}$$

$$= -\int (x-\frac{1}{2})^2 - \frac{1}{4}$$

$$= \frac{1}{4} - (x-\frac{1}{2})^2$$

$$= \left(\frac{1}{2}\right)^2 - v^2$$

$$x - \frac{1}{2} = v$$

$$dx = dv$$

$$= \lim_{t \rightarrow 0} \operatorname{arcsm}(2x-1) \Big|_t^{\frac{1}{2}}$$

$$= \lim_{t \rightarrow 0} (\operatorname{arcsm} 0 - \operatorname{arcsm}(2t-1))$$

$$= 0 - \operatorname{arcsm}(-1)$$

$$= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2}$$

$$\textcircled{B} \quad = \lim_{t \rightarrow 1} \operatorname{arcsm}(2x-1) \Big|_0^t$$

$$\Rightarrow \lim_{t \rightarrow 1} (\operatorname{arcsm}(2t-1) - \operatorname{arcsm} 0)$$

$$= \operatorname{arcsm} 1 - \operatorname{arcsm} 0$$

$$= \frac{\pi}{2}$$

$$\textcircled{A} + \textcircled{B} \Rightarrow \frac{\pi}{2} + \frac{\pi}{2} = \pi$$




# 21

EYLÜL  
SEPTEMBER  
CUMA  
FRIDAY

EYLÜL 2012						
H	P	S	Ç	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

~~y = e<sup>-x</sup>~~, Oy-ele, Ox-ele



$$A = \int_0^{\infty} e^{-x} dx$$

$$= \lim_{t \rightarrow \infty} \int_0^t e^{-x} dx$$

$$= \lim_{t \rightarrow \infty} (-e^{-x}) \Big|_0^t$$

$$= \lim_{t \rightarrow \infty} (-e^{-t} + 1)$$

$$= 1$$

~~y = e<sup>-x</sup>~~, Oy-ele, Ox-ele,  $\int e^{-x} dx = -e^{-x}$  H(?)

$$H = \pi \int_0^{\infty} f^2 dx$$

$$H = \pi \int_0^{\infty} e^{-2x} dx$$

$$= \pi \left( \frac{1}{2} e^{-2x} \right) \Big|_0^{\infty}$$



EYLÜL

2012

H	P	S	Ç	P	C	Cl	Pz
35					1		
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

EYLÜL

SEPTEMBER

CUMARTESİ

SATURDAY

22



$$= \pi \lim_{t \rightarrow \infty} \left( -\frac{e^{-2t}}{2} + \frac{e^t}{2} \right)$$

$$= \pi/2 \text{ br}^3$$

~~AA~~  $x^2 + hy^2 = 4$  elipsi just yeri  $Ox$ -ab düz

$$y = \pm \sqrt{4-x^2}$$

$$y' = \frac{1}{2} (-2x) \cdot \frac{1}{2\sqrt{4-x^2}} = \frac{-x}{2\sqrt{4-x^2}}$$



EYLÜL

SEPTEMBER

PAZAR

SUNDAY

23

$$(y')^2 = \frac{x^2}{4(4-x^2)}$$

$$S = 2\pi \int_{-2}^2 \left| \frac{\sqrt{4-x^2}}{2} \right| \sqrt{1 + \frac{x^2}{4(4-x^2)}} dx$$

$$S = 2\pi \int_0^2 \sqrt{4-x^2} \cdot \frac{\sqrt{16-3x^2}}{2\sqrt{4-x^2}} dx$$

$$S = \pi \int_0^2 \sqrt{16-3x^2} dx$$



# 24

EYLÜL  
SEPTEMBER  
PAZARTESİ  
MONDAY

EYLÜL 2012						
H	P	S	C	P	C	Ct Pz
35						1 2
36	3	4	5	6	7	8 9
37	10	11	12	13	14	15 16
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

$$= \pi \int_0^{\frac{\pi}{2}} \sqrt{16 - 3x^2} dx$$

$$\begin{aligned} x^2 &= r^2 \\ r^2 - a^2 & \\ d^2 + x^2 & \end{aligned}$$

$$\begin{cases} x = \frac{4}{\sqrt{3}} \sin t \\ dx = \frac{4}{\sqrt{3}} \cos t dt \end{cases}$$

$$\begin{aligned} 16 - 3x^2 & \\ \left(\frac{4}{\sqrt{3}}\right)^2 - x^2 & \end{aligned}$$

$$= \pi \int_0^{\frac{\pi}{2}} \sqrt{16 - 3 \frac{16}{3} \sin^2 t} \cdot \frac{4}{\sqrt{3}} \cos t dt$$

$$= \frac{16}{\sqrt{3}} \pi \int_0^{\frac{\pi}{2}} 4 \cos t \cos t dt$$

$$= \frac{16}{\sqrt{3}} \pi \int_0^{\frac{\pi}{2}} \cos^2 t dt$$

$$\cos^2 t = \frac{1}{2} [1 + \cos 2t]$$

$$= \frac{16}{\sqrt{3}} \pi \int_0^{\frac{\pi}{2}} \frac{1}{2} [1 + \cos 2t] dt$$

$$= \frac{8}{\sqrt{3}} \pi \left( t + \frac{\sin 2t}{2} \right) \Big|_0^{\frac{\pi}{2}} = \frac{8}{\sqrt{3}} \pi$$

$$= \frac{8}{\sqrt{3}} \pi \left( \frac{\pi}{2} + \frac{1}{2} \right) 6r^2$$

$$\begin{aligned} x=0 &\Rightarrow t=0 \\ x=2 &\Rightarrow t=\pi/3 \end{aligned}$$



EYLÜL

2012

H	P	S	Ç	P	C	Ct	Pz
35				1	2		
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

$$\text{Ox-eşref dön} \Rightarrow S = 2\pi \int_{0}^{2\pi} |y| \sqrt{(x')^2 + (y')^2} dt$$

EYLÜL

SEPTEMBER  
TUESDAY

$$\text{Qiy-eşref dön} \Rightarrow S = 2\pi \int_{0}^{\pi} |x| \sqrt{(x')^2 + (y')^2} dt$$

25

$$x = e^t \sin t \\ y = e^t \cos t$$

$$0 \leq t \leq \pi/2$$

Ox-eşref,  
don  
yüzeyler - ?

$$x' = e^t \sin t + e^t \cos t$$

$$y' = e^t \cos t - e^t \sin t$$

$$(x')^2 = e^{2t} \sin^2 t + 2e^{2t} \sin t \cos t + e^{2t} \cos^2 t$$

$$(y')^2 = e^{2t} \cos^2 t - 2e^{2t} \sin t \cos t + e^{2t} \sin^2 t$$

$$(x')^2 + (y')^2 = 2e^{2t}$$

$$S = 2\pi \int_0^{\pi/2} |e^t \cos t| \sqrt{2e^{2t}} dt$$

$$= 2\sqrt{2}\pi \int_0^{\pi/2} e^{2t} \cos t dt$$

LAPTO



# 26

EYLÜL  
SEPTEMBER  
ÇARŞAMBA  
WEDNESDAY

$\frac{2}{\sqrt{2}}$   $\frac{3}{3}$   
 $\frac{3}{3}$   $\frac{3}{3}$

EYLÜL 2012						
H	P	S	Ç	P	C	Pz
35						1 2
36	3	4	5	6	7	8 9
38	17	18	19	20	21	22 23
39	24	25	26	27	28	29 30

K

$$\begin{aligned}x &= t^3 \\y &= 2t+3\end{aligned}$$

$$-1 \leq t \leq 1$$

Oy-et. son  
5 = ?

$$x' = 3t^2$$

$$y' = \cancel{2t} 2$$

$$(x')^2 + (y')^2 = 9t^4 + 4$$

$$S = 2\pi \int_{-1}^1 |t^3| \sqrt{9t^4 + 4} dt$$

$$= 4\pi \int_0^1 t^3 \sqrt{9t^4 + 4} dt \quad 9t^4 + 4 = u \\ 36t^3 dt = du$$

$$= 4\pi \int_0^1 \sqrt{u} \frac{du}{36}$$

$$= \frac{1}{3}\pi \int_0^1 \sqrt{u} du$$

;

;

;

$$= \frac{2\pi}{27} \left[ \left( \frac{3}{2}u^{\frac{3}{2}} - 8 \right) \Big|_0^1 \right] br^2$$



EYLÜL 2012

H	P	S	Ç	P	C	Cl	Pz
35					1	2	
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

EYLÜL  
SEPTEMBER  
PERŞEMBE  
THURSDAY

27

A  $x = \cos t$   $-\pi \leq t \leq \pi$  egrileri varsa

 $y = t + \sin t$

$$l = \int_{t_1}^{t_2} \sqrt{(x')^2 + (y')^2} dt$$

$x' = -\sin t$

$y' = 1 + \cos t$

$$(x')^2 + (y')^2 = \sin^2 t + 1 + 2\cos t + \cos^2 t$$
 $= 2 + 2\cos t$

$$l = \int_{-\pi}^{\pi} \sqrt{1 + \cos t} dt$$
 $\cos t = 2\cos^2 \frac{t}{2} - 1$

$$= 2\sqrt{2} \int_0^{\pi} \sqrt{2\cos^2 \frac{t}{2}} dt$$

$$= 4 \int_0^{\pi} \cos \frac{t}{2} dt$$

$$= 8 \sin \frac{t}{2} \Big|_0^{\pi}$$

$$= 8 \text{ br}$$



# 28

EYLÜL  
SEPTEMBER  
CUMA  
FRIDAY

SN 21  
2012  
1  
 $\frac{1}{\cos^2 x}$

H	P	S	Ç	P	C	Ct	Pz
35						1	2
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

~~A~~  $x = \ln(\sec y)$  eger  $y=0$ ,  $y = \frac{\pi}{3}$  ~~ve~~  $\tan y$

$$x = \ln r \rightarrow l = \int_0^b \sqrt{1 + (y')^2} dx$$

$$y = \tan \rightarrow l = \int_c^d \sqrt{1 + (x')^2} dy$$

~~$\frac{1}{\sec y} \cdot (\sec y \tan y) = \tan y$~~

$$l = \int_0^{\pi/3} \sqrt{1 + \tan^2 y} dy$$

~~$= \int_0^{\pi/3} \sqrt{\sec^2 y} dy$~~ 

$$= \int_0^{\pi/3} \sec y dy$$
 ~~$\sec y + \tan y dy$~~ 

$$(sec y + \tan y)$$

$$= \int_0^{\pi/3} \frac{\sec^2 y + \sec y \tan y}{\sec y + \tan y} dy$$

$\sec y + \tan y = t$   
 $(\sec^2 y + \sec y \tan y) dy = dt$

$$= \ln |\sec y + \tan y| \Big|_0^{\pi/3}$$

$$= \ln \left| \sec \frac{\pi}{3} + \tan \frac{\pi}{3} \right| - \ln \left| \sec 0 + \tan 0 \right|$$

$$= \ln (2 + \sqrt{3}) - \ln (1 + 0) = \ln (2 + \sqrt{3}) \text{ br}$$



EYLÜL

2012

H	P	S	Ç	P	C	Ct	Pz
35					1	2	
36	3	4	5	6	7	8	9
37	10	11	12	13	14	15	16
38	17	18	19	20	21	22	23
39	24	25	26	27	28	29	30

Kabuk (y-sınır)Disk (x-sınır)

EYLÜL



SEPTEMBER

CUMARTESİ

SATURDAY

29

$$x = y^2, x=0, y=2 \quad \text{Ox. et de} \quad H=?$$

Gelişti

$$H = 2\pi \int_0^2 y [f(y) - g(y)] dy$$

$$= 2\pi \int_0^2 y (y^2 - 0) dy$$

$$= 2\pi \int_0^2 y^3 dy$$

$$= \pi/2 b r^3$$

EYLÜL  
SEPTEMBER

30

PAZAR  
SUNDAYDost

$$H = 2\pi \int_0^1 (f(x))^2 - (g(x))^2 dx$$

$$= \pi \int_0^1 1^2 - (\sqrt{x})^2 dx$$

$$= \pi \int_0^1 (1-x) dx$$

$$= \pi \left[ x - \frac{x^2}{2} \right] \Big|_0^1$$

$$= \pi/2 b r^3$$



# 01


EKİM  
OCTOBER

PAZARTESİ  
MONDAY

H	P	S	Ç	P	C	Ct	Pz
40	1	2	3	4	5	6	7
41	8	9	10	11	12	13	14
42	15	16	17	18	19	20	21
43	22	23	24	25	26	27	28
44	29	30	31				

~~JK~~  $xy = 5 \rightarrow y = \frac{5}{x}$  (yak)  $H = ?$

$y + x = 6 \rightarrow y = 6 - x$



$$\frac{5}{x} = 6 - x$$

$$5 = 6x - x^2$$

$$x^2 - 6x + 5 = 0$$

$$\begin{array}{l|l} x & -5 \\ \hline x & \end{array}$$

$$\begin{array}{l} x=5 \\ x=1 \end{array}$$

$$H = 2\pi \int_1^5 |x[f(x) - g(x)]| dx$$

$$= 2\pi \int_1^5 |x[(6-x) - (\frac{5}{x})]| dx$$

$$= 2\pi \int_1^5 |6x - x^2 - 5| dx$$

$$-(x^2 - 6x + 5)$$

$$= 2\pi \int_1^5 (6x - x^2 - 5) dx$$

$$\begin{array}{r} + \cancel{f} - \cancel{f} + \\ - \cancel{f} - \cancel{f} - \end{array}$$



## \* Parametrik Densit Vektörler Egzersiz Alan

$$\textcircled{1} \quad x = g(t) \\ y = h(t)$$

$$t_1 \leq t \leq t_2$$

$$x=0 \\ x=b$$

$$A = \int_{t_1}^{t_2} |y| x' dt$$


$$\textcircled{2} \quad x = g(t) \\ y = h(t)$$

$$t_3 \leq t \leq t_4$$

$$y=c \\ y=d$$

$$A = \int_{t_3}^{t_4} |x| y' dt$$

$$\cancel{x = a \cos^3 t} \\ y = a \sin^3 t$$



$$x=0 \\ x=d$$

$$0 = d \cos^3 t \\ 0 = a \cos^3 t$$

$$t_1 = \pi/2 \\ t_2 = 0$$

$$A = \int_{\pi/2}^0 |a \sin^3 t| 3a \cos^2 t (-\sin t) dt$$

$$= 3a^2 \int_0^{\pi/2} \sin^4 t \cos^2 t dt$$

$$= 3a^2 \int_0^{\pi/2} \left[ \frac{1}{2}(1 - \cos 2t) \right]^2 \left[ \frac{1}{2}(1 + \cos 2t) \right] dt$$

$$= \frac{3a^2}{8} \int_0^{\pi/2} (1 - 2\cos 2t + \cos^2 2t)(1 + \cos 2t) dt$$

$$= \frac{3a^2}{8} \int_0^{\pi/2} (\overset{\textcircled{4}}{\cos^3 2t} - \overset{\textcircled{3}}{\cos^2 2t} - \overset{\textcircled{2}}{\cos 2t} + \overset{\textcircled{1}}{1}) dt$$

$$\cos 2x = 2\cos^2 x - 1 \\ \cos 2x = 1 - 2\sin^2 x$$

$$\textcircled{1} \quad \int_0^{\pi/2} dt = \pi/2$$

$$\textcircled{2} \quad - \int_0^{\pi/2} \cos 2t dt = -\frac{1}{2} \sin 2t \Big|_0^{\pi/2} = -\frac{1}{2} (\sin \pi - \sin 0) = 0$$

$$\textcircled{3} \quad - \int_0^{\pi/2} \cos^2 2t dt = - \int_0^{\pi/2} \frac{1}{2}(1 + \cos 4t) dt = -\frac{1}{2} \left( t + \frac{1}{4} \sin 4t \right) \Big|_0^{\pi/2} = -\frac{1}{2} \left( \frac{\pi}{2} + \frac{1}{4} \sin 2\pi \right) - \left( 0 + \frac{1}{4} \sin 0 \right) = -\frac{\pi}{4}$$

$$\textcircled{4} \quad \int_0^{\pi/2} \cos^3 2t dt = \int_0^{\pi/2} \cos^2 2t \sin 2t dt = 0 \quad \boxed{\begin{matrix} \sin 2t = 0 \\ 2\cos 2t dt = 0 \end{matrix}}$$

$$AA = \frac{3a^2}{8} \cdot 4 \cdot \left( \frac{\pi}{2} - \frac{\pi}{4} \right) = \frac{3a^2}{4}$$

by now do only 1 question  
we have seen many before  
so you do it yourself


## Hacim Hesabı

① Kıst Yani (Düzgün cisimler)

$$H = \int_a^b A(x) dx ; \quad H = \int_c^d A(y) dy$$




② Disk Yani



$$H = \pi \int_a^b [f(x)]^2 dx \approx \pi r^2$$

(sinirler ile dairesel yarım dağılımları)

③ Kalınlık Yani



$$H = 2\pi \int_a^b x [f(x) - g(x)] dx$$

$$\approx (2\pi d)(r)$$

## \*Parametrik Egrilerin Hacimleri

①  $x = u(t)$        $t_1 \leq t \leq t_2$   
 $y = v(t)$        $Ox$ -ekseni den

$$H = \pi \int_{t_1}^{t_2} y^2 x' dt$$

②  $x = u(t)$        $t_3 \leq t \leq t_4$   
 $y = v(t)$        $Oy$ -ekseni den

$$H = \pi \int_{t_3}^{t_4} x^2 y' dt$$

## EGRİ UZUNLUĞU

①  $y = f(x)$ ,  $[a, b]$ ,  $\ell = \int_a^b \sqrt{1 + (y')^2} dx$


②  $x = g(y)$ ,  $[c, d]$ ,  $\ell = \int_c^d \sqrt{1 + (x')^2} dy$

## \*Parametrik Egrilerin Egrî Uzunluğu


$x = u(t)$        $t_1 \leq t \leq t_2$   
 $y = v(t)$

$$\ell = \int_{t_1}^{t_2} \sqrt{(x')^2 + (y')^2} dt$$

# DANIEL YÖZGEMİN YÖZGEM ALAMI



$$S = 2\pi \int_0^b |f(x)| \sqrt{1 + (f'(x))^2} dx$$



$$S = 2\pi \int_0^b |f(x) - k| \sqrt{1 + (f'(x))^2} dx$$

## \* Parametrik Parametrik Olususunda Yüzey Alanı

①  $x = u(t)$   
 $y = v(t)$

$$t_1 \leq t \leq t_2$$

Ox-ekseni

$$S = 2\pi \int_{t_1}^{t_2} |y| \sqrt{(x')^2 + (y')^2} dt$$

②  $x = u(t)$   
 $y = v(t)$

$$0 \leq t \leq t_2$$

Oy-ekseni

$$S = 2\pi \int_{0}^{t_2} |x| \sqrt{(x')^2 + (y')^2} dt$$

## GENELLEŞTİRİLMİŞ İNTEGRAL

**I. TIP** Sınırlarımızın en az biri sınırsız, diğerleri ise sınırlı olmalıdır.

$$\int_a^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

$$\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx + \lim_{s \rightarrow -\infty} \int_s^b f(x) dx$$

II.TIP

$$\textcircled{6} \quad \int_a^b f(x) dx = \lim_{t \rightarrow b} \int_a^t f(x) dx$$

$$\textcircled{7} \quad \int_b^a f(x) dx = - \int_a^b f(x) dx$$

$$\textcircled{8} \quad \int_a^c f(x) dx = \textcircled{9} \quad \int_a^c f(x) dx + \textcircled{6} \quad \int_c^c f(x) dx$$

$$= \lim_{t \rightarrow c} \int_a^t f(x) dx + \int_m^c \int_a^x f(x) dx$$

III.TIP

$$\text{II.TIP} \rightarrow \text{II.TIP} + \text{I.TIP}$$

$$\begin{array}{l} \textcircled{10} \\ \int_a^\infty f(x) dx \\ \hline \textcircled{11} \\ \int_{-\infty}^a f(x) dx \\ \textcircled{12} \end{array}$$

$$\int_{-\infty}^\infty f(x) dx \rightarrow \text{orada } 'a' \text{ soyis sifat}$$

$$\int_0^\infty x e^{-x} dx$$