

Problem 1. Consider a one-period market model with no dividends. For the sake of this problem, call an adapted real-valued process $Z = (Z_t)_{t \in \{0,1\}}$ an ‘anti-martingale deflator’ iff

- $Z_0 \geq 0$, $Z_1 \geq 0$ almost surely and $\mathbb{P}(Z_0 = 0 = Z_1) < 1$,
- $Z_1 P_1$ is integrable and $\mathbb{E}(Z_1 P_1) = -Z_0 P_0$

(a) Show that if there exists a numéraire portfolio, then there does not exist an anti-martingale deflator.

The goal for the rest of the problem is to prove the converse of part (a). To that end, let

$$F(h) = e^{-h \cdot P_0} + \mathbb{E}[e^{-h \cdot P_1} \zeta]$$

where $\zeta = e^{-\|P_1\|^2/2}$. Let $(h_k)_k$ be a minimising sequence for F .

(b) Show that if there exists a numéraire, then $\inf_{h \in \mathbb{R}^n} F(h) = 0$.

(c) Show that if $(h_k)_k$ is bounded, then there exists an anti-martingale deflator.

For the rest of the problem, suppose $(h_k)_k$ is unbounded.

(d) Show that there exists a vector $\eta \in \mathbb{R}^n$ with $\|\eta\| = 1$ such that $\eta \cdot P_0 \geq 0$, $\eta \cdot P_1 \geq 0$ almost surely, and that

$$\inf_{h \in \mathbb{R}^n} F(h) = \mathbb{1}_{\{\eta \cdot P_0 = 0\}} + \mathbb{E}[\mathbb{1}_{\{\eta \cdot P_1 = 0\}} \zeta]$$

(e) Now let

$$G(h) = \mathbb{1}_{\{\eta \cdot P_0 = 0\}} e^{-h \cdot P_0} + \mathbb{E}[\mathbb{1}_{\{\eta \cdot P_1 = 0\}} e^{-h \cdot P_1} \zeta].$$

Show that $G(h) = \lim_k F(k\eta + h)$ and conclude that G is maximised at $h = 0$. Assuming $G(0) > 0$, show that there exists an anti-martingale deflator.

Solution 1. (a) Suppose there exists a numéraire portfolio η and suppose $Z_t \geq 0$ for $t = 0, 1$ and $\mathbb{E}(Z_1 P_1) = -Z_0 P_0$. Multiplying by η yields

$$(*) \quad -Z_0 N_0 = \mathbb{E}(Z_1 N_1)$$

where $N_t = \eta \cdot P_t$ for $t = 0, 1$. Since $N_t > 0$ a.s. we have $Z_t N_t \geq 0$ for $t = 0, 1$. Combined with equation (*) we have $Z_0 N_0 = 0$. Since $N_0 \neq 0$ we conclude that $Z_0 = 0$. Equation (*) also says $\mathbb{E}(Z_1 N_1) = 0$, so by the pigeonhole principle $Z_1 N_1 = 0$ a.s. Again, since $N_1 \neq 0$ a.s. we have $Z_1 = 0$ a.s. In particular, Z_0, Z_1 is not an anti-martingale deflator.

(b) Suppose there exists a numéraire η so that $N_t = \eta \cdot P_t > 0$ a.s. for $t = 0, 1$. Let $h_k = k \eta$ for $k = 1, 2, \dots$. Note $e^{-h_k \cdot P_t} = e^{-k N_t} \rightarrow 0$ a.s. for $t = 0, 1$. Hence $F(h_k) \rightarrow 0$ by the dominated convergence theorem.

(c) If $(h_k)_k$ is bounded, we can pass to a convergent subsequence by the Bolzano-Weierstrass theorem. Let h^* be the limit. Note $F(h_k) \rightarrow F(h^*)$ by the dominated convergence theorem. So $F(h^*) = \inf_h F(h)$, i.e. h^* is a minimiser.

(d) We can pass to a subsequence such that $\|h_k\| \uparrow \infty$. Let $\hat{h}_k = h_k / \|h_k\|$. Since $(\hat{h}_k)_k$ is bounded we can pass to a further subsequence where $\hat{h}_k \rightarrow \eta$ where $\|\eta\| = 1$. Note for $t = 0, 1$ that

$$e^{-h_k \cdot P_t} = (e^{-\hat{h}_k \cdot P_t})^{\|h_k\|} \rightarrow \infty \text{ on the event } \{\eta \cdot P_t < 0\}$$

Since $F(h_k) \leq F(0)$ for all k , we see $\eta \cdot P_t \geq 0$ a.s. for $t = 0, 1$ to complete

(e) Since $e^{-k \eta \cdot P_t} \rightarrow \mathbb{1}_{\{\eta \cdot P_t = 0\}}$, we have $G(h) = \lim_k F(k\eta + h)$ by the dominated convergence theorem. Clearly $G(h) \geq \inf_H F(H)$. But by part (d), we have $\inf_H F(H) = G(0)$. Since G is minimised at 0, we have $\nabla G(0) = 0$, or equivalently

$$\mathbb{E}[\mathbb{1}_{\{\eta \cdot P_1 = 0\}} \zeta P_1] = -\mathbb{1}_{\{\eta \cdot P_0 = 0\}} P_0$$

Assuming $G(0) > 0$, an anti-martingale deflator is $Z_0 = \mathbb{1}_{\{\eta \cdot P_0 = 0\}}$ and $Z_1 = \mathbb{1}_{\{\eta \cdot P_1 = 0\}} \zeta$.

Problem 2. What are the economically appropriate definitions of numéraire portfolio and equivalent martingale measure in the case where the assets may pay a dividend?

Solution 2. A numéraire is a previsible process η such that $\eta_1 \cdot P_0 > 0$ and $\eta_{t+1} \cdot P_t = \eta_t \cdot (P_t + \delta_t) > 0$ a.s. for all $t \geq 1$. An equivalent martingale measure relative to the numéraire is a measure \mathbb{Q} under which

$$\frac{P_t}{N_t} = \mathbb{E}^{\mathbb{Q}} \left(\frac{P_{t+1} + \delta_{t+1}}{N_t} \middle| \mathcal{F}_t \right)$$

for all $t \geq 0$. It remains to explain why these definitions are economically appropriate. First note that assuming there is a numéraire according to our revised definition, implies that there is an arbitrage if and only if there is a terminal consumption arbitrage. Next, there is an equivalent martingale measure according to our revised definition if and only if there is a martingale deflator.

Problem 3. Consider a one-period market with three assets. The first asset is a riskless asset with risk-free rate r . The second asset is a stock with prices $(S_t)_{t \in \{0,1\}}$. The third is a contingent claim on the stock with time 1 price $\xi_1 = g(S_1)$, where the function g is convex. Show that if there is no arbitrage, then $\xi_0 \geq \frac{1}{1+r} g[(1+r)S_0]$. Assuming $\xi_0 < \frac{1}{1+r} g[(1+r)S_0]$, find an arbitrage explicitly.

Hint: By the convexity of g , there exists a function λ such that $g(x) \geq g(y) + \lambda(x)(x - y)$ for all $x, y \in \mathbb{R}$.

Solution 3. Suppose $\xi_0 < \frac{1}{1+r} g[(1+r)S_0]$ and let $L = \lambda[(1+r)S_0]$. Consider the portfolio $H = (-\frac{1}{1+r} g[(1+r)S_0] + LS_0, -L, +1)$. Note

$$\begin{aligned} c_0 &= -H \cdot (B_0, S_0, \xi_0) \\ &= \frac{1}{1+r} g[(1+r)S_0] - \xi_0 > 0 \end{aligned}$$

and

$$\begin{aligned} c_1 &= H \cdot (B_1, S_1, \xi_1) \\ &= -g[(1+r)S_0] + LS_0(1+r) - LS_1 + g(S_1) \geq 0 \end{aligned}$$

Problem 4. (Bayes's formula) Let \mathbb{P} and \mathbb{Q} be equivalent probability measures defined on (Ω, \mathcal{F}) with density $Z = \frac{d\mathbb{Q}}{d\mathbb{P}}$. Let $\mathcal{G} \subseteq \mathcal{F}$ be a sigma-field. Prove the identity:


$$\mathbb{E}^{\mathbb{Q}}(X|\mathcal{G}) = \frac{\mathbb{E}^{\mathbb{P}}(ZX|\mathcal{G})}{\mathbb{E}^{\mathbb{P}}(Z|\mathcal{G})}$$

for each random variable X such that X is \mathbb{Q} -integrable.

Solution 4. Let $Y = \frac{\mathbb{E}^{\mathbb{P}}(ZX|\mathcal{G})}{\mathbb{E}^{\mathbb{P}}(Z|\mathcal{G})}$. Note that Y is \mathcal{G} -measurable. Hence, we need only verify the equation $\mathbb{E}^{\mathbb{Q}}(Y\mathbb{1}_G) = \mathbb{E}^{\mathbb{Q}}(X\mathbb{1}_G)$ for all $G \in \mathcal{G}$; equivalently, we must verify $\mathbb{E}^{\mathbb{P}}(ZY\mathbb{1}_G) = \mathbb{E}^{\mathbb{P}}(ZX\mathbb{1}_G)$ for all $G \in \mathcal{G}$.

$$\begin{aligned}\mathbb{E}^{\mathbb{P}}(\mathbb{1}_G Y Z) &= \mathbb{E}^{\mathbb{P}}[\mathbb{E}(\mathbb{1}_G Y Z | \mathcal{G})] \text{ tower property} \\ &= \mathbb{E}^{\mathbb{P}}[\mathbb{1}_G Y \mathbb{E}(Z | \mathcal{G})] \text{ taking out what's known} \\ &= \mathbb{E}^{\mathbb{P}}[\mathbb{1}_G \mathbb{E}^{\mathbb{P}}(XZ | \mathcal{G})] \\ &= \mathbb{E}^{\mathbb{P}}[\mathbb{E}^{\mathbb{P}}(\mathbb{1}_G XZ | \mathcal{G})] \text{ pulling in what's known} \\ &= \mathbb{E}^{\mathbb{P}}(\mathbb{1}_G XZ) \text{ tower property}\end{aligned}$$

Problem 5. * Consider a trinomial two-asset model with prices $P = (B, S)$ where $B_0 = B_1 = 1$ and S is given by



Find all risk-neutral measures for this model. Now introduce a call option with payout $\xi_1 = (S_1 - 2)^+$. Show that there is an open interval I such that the augmented market (B, S, ξ) has no arbitrage if and only if $\xi_0 \in I$.

Solution 5. A risk neutral measure solves $\mathbb{E}^{\mathbb{Q}}(S_1) = S_0$, (no discounting is needed since the numéraire is cash)

$$\begin{aligned}3p + 2q + r &= 2 \\ p + q + r &= 1\end{aligned}$$

and hence $(p, q, r) = (p, 1 - 2p, p)$ for $0 < p < 1/2$, where $p = \mathbb{Q}\{S_1 = 3\}$, etc. By the fundamental theorem of asset pricing, there is no arbitrage if and only if $\xi_0 = \mathbb{E}^{\mathbb{Q}}(\xi_1) = p$. So $I = (0, 1/2)$.

Problem 6. Consider an arbitrage-free bond market. Let P_t^T be the price of the bond of maturity T at time t , where $1 \leq t \leq T$. Let the spot rate be $r_t = \frac{1}{P_{t-1}^t} - 1$ and the bank account be $B_t = \prod_{s=1}^t (1 + r_s)$ for all $t \geq 1$ as usual.

(a) Let \mathbb{Q} be a risk-neutral measure, i.e. an equivalent martingale measure relative to the bank account. Show that

$$P_t^T = B_t \mathbb{E}^{\mathbb{Q}}(B_T^{-1} | \mathcal{F}_t)$$

for all $0 \leq t \leq T$.

(b) Consider a European contingent claim with maturity T and payout r_T . Show that this claim can be replicated by trading in bonds.

(c) Consider a forward contract initiated at time t for the payout at time T of r_T . The forward interest rate f_t^T at time t for maturity T is defined to be the forward price of this payout. Show that

$$f_t^T = \frac{P_t^{T-1}}{P_t^T} - 1$$

- (d) Show that if the spot rate is not random, then $f_t^T = r_T$.
(e) Let \mathbb{Q}^T be a T -forward measure, i.e. an equivalent martingale measure relative to the bond of maturity T . Show that the forward rate process $(f_t^T)_{0 \leq t < T}$ is a \mathbb{Q}^T martingale.
(f) The quantity

$$y_t^T = (P_t^T)^{-\frac{1}{T-t}} - 1$$

is called the yield at time t of the bond maturing at time T .

Show that the following are equivalent

- (1) $f_t^T \geq y_t^T$ a.s. for all $0 \leq t < T$
- (2) $T \mapsto y_t^T$ is non-decreasing a.s. for all $t \geq 0$.

(g) Show that the following are equivalent:

- (1) $r_t \geq 0$ a.s. for all $t \geq 1$
- (2) $t \mapsto B_t$ is non-decreasing a.s.
- (3) $T \mapsto P_t^T$ is non-increasing a.s. for each $t \geq 0$
- (4) $f_t^T \geq 0$ a.s. for all $0 \leq t < T$.
- (5) $y_t^T \geq 0$ a.s. for all $0 \leq t < T$.
- (6) each martingale deflator is a supermartingale.

Solution 6. (a) The discounted price $P^T/B = (P_t^T/B_t)_{0 \leq t \leq T}$ is a martingale for any risk-neutral measure. The result follows from $P_T^T = 1$ and the martingale property.

(b) Work backwards: Note that the payout r_T is \mathcal{F}_{T-1} measurable, and can be realised by holding r_T bonds of maturity T during the period $(T-1, T]$. The time $T-1$ cost of this strategy is $r_T P_{T-1}^T = 1 - P_{T-1}^T$. Now to replicate this time $T-1$ payout by holding one bond of maturity $T-1$ and selling one bond of maturity T .

In summary, the replication strategy is at time 0 to buy one bond of maturity $T-1$ and to sell one bond of maturity T . At time $T-1$, invest the payout of the bond of maturity $T-1$ into $1/P_{T-1}^T$ bonds of maturity T .

(c) *A dual approach:* From lectures

$$f_t^T = \mathbb{E}^{\mathbb{Q}^T}(r_T | \mathcal{F}_t)$$

where \mathbb{Q}^T is a T -forward measure. Since $r_T = \frac{1}{P_{T-1}^T} - 1$ and by changing to a risk-neutral measure (using part (a)) we have

$$\begin{aligned} P_t^T \mathbb{E}^{\mathbb{Q}^T} \left(\frac{1}{P_{T-1}^T} | \mathcal{F}_t \right) &= B_t \mathbb{E}^{\mathbb{Q}} \left(\frac{B_T^{-1}}{\mathbb{E}^{\mathbb{Q}}(B_T^{-1} | \mathcal{F}_{T-1}) B_{T-1}} | \mathcal{F}_t \right) \text{ cond on } \mathcal{F}_{T-1} \text{ and tower} \\ &= B_t \mathbb{E}^{\mathbb{Q}}(B_{T-1}^{-1} | \mathcal{F}_t) \\ &= P_t^{T-1} \end{aligned}$$

A primal approach: Consider the forward claim initiated at time t with time T payout $\xi_T = r_T - f_t^T$. From part (b), the cost at time t to replicate r_T is $P_t^{T-1} - P_t^T$, and the cost at time t to replicate the \mathcal{F}_t -measurable payout f_t^T is $f_t^T P_t^T$. Hence $\xi_t = P_t^{T-1} - (1 + f_t^T) P_t^T$. But the initial price of a forward is $\xi_t = 0$. Solving for f_t^T yields the formula.

(d) If the spot rate is not random, then from part (a) we have $P_t^T = B_t/B_T$, and hence

$$f_t^T = \frac{P_t^{T-1}}{P_t^T} - 1 = \frac{B_T}{B_{T-1}} - 1 = r_T$$

(e) This follows from Doob's observation that $M_t = \mathbb{E}(\xi|\mathcal{F}_t)$ defines a martingale whenever ξ is integrable.

(f) $f_t^T \geq y_t^T \Leftrightarrow (P_t^{T-1})^{T-t} \geq (P_t^T)^{T-t-1} \Leftrightarrow y_t^{T-1} \leq y_t^T$

(g) (1) $\Leftrightarrow B_t = (1 + r_t)B_{t-1} \geq B_{t-1} \Leftrightarrow$ (2)

(2) $\Rightarrow P_t^{T+1} = B_t \mathbb{E}^{\mathbb{Q}}(B_{T+1}^{-1}|\mathcal{F}_t) \leq B_t \mathbb{E}^{\mathbb{Q}}(B_T^{-1}|\mathcal{F}_t) = P_t^T \Leftrightarrow$ (3).

(3) $\Rightarrow 1 = P_t^t \geq P_t^{t+1} = B_t \mathbb{E}^{\mathbb{Q}}(B_{t+1}^{-1}|\mathcal{F}_t) = B_t/B_{t+1}$ since B is predictable \Leftrightarrow (2).

(4) $\Leftrightarrow P_t^{T-1} \geq P_t^T \Leftrightarrow$ (3).

(5) $\Rightarrow P_t^{t+1} \leq 1 \Leftrightarrow r_{t+1} \geq 0 \Leftrightarrow$ (1)

(3) $\Rightarrow P_t^T \leq 1 \Leftrightarrow$ (5)

(6) $\Leftrightarrow r_t = \frac{Y_{t-1}}{\mathbb{E}(Y_t|\mathcal{F}_{t-1})} - 1 \geq 0 \Leftrightarrow$ (1)

Problem 7. (a) Let X_1, X_2, \dots be a sequence of non-negative random variables such that $\mathbb{E}(X_n) = 1$ for all n . Use the Borel–Cantelli lemma to show

$$\limsup_{n \rightarrow \infty} X_n^{1/n} \leq 1 \text{ a.s.}$$

(b) Consider a bond market as in problem 6. The *long rate* at time t is defined as $\ell_t = \lim_{T \rightarrow \infty} y_t^T$ whenever the limit exists.

Suppose that bonds are priced according to the formula in 6(a) for a fixed risk-neutral measure, and that the long rate exists a.s. at all times. Show that the long rate is non-decreasing, that is

$$\ell_s \leq \ell_t \text{ a.s. for all } 0 \leq s \leq t,$$

a fact first discovered by Dybvig, Ingersoll & Ross in 1996.

Solution 7. (a) For each $\epsilon > 0$ we have

$$\sum_{n=1}^{\infty} \mathbb{P}[X_n > (1 + \epsilon)^n] \leq \sum_{n=1}^{\infty} (1 + \epsilon)^{-n} = \frac{1}{\epsilon} < \infty$$

by Markov's inequality. The first Borel–Cantelli lemma then says

$$\mathbb{P}(X_n^{1/n} > 1 + \epsilon \text{ infinitely often}) = 0$$

This shows $\limsup_{n \rightarrow \infty} X_n^{1/n} \leq 1$ as claimed.

(b) Now, let P_t^T be the bond price, B_t the bank account, and $\tilde{P}_t^T = P_t^T/B_t$ the discounted bond price. Suppose the long rate ℓ_t exists, so that

$$\ell_t = - \lim_{T \rightarrow \infty} \frac{1}{T} \log P_t^T = - \lim_{T \rightarrow \infty} \frac{1}{T} \log \tilde{P}_t^T.$$

By assumption, the discounted bond prices are given by

$$\tilde{P}_t^T = \mathbb{E}^{\mathbb{Q}}(B_T^{-1}|\mathcal{F}_t)$$

each $0 \leq t \leq T$ and a fixed risk-neutral measure \mathbb{Q} , and, in particular, \tilde{P}^T is a martingale for each $T > 0$.

Fix $0 \leq s \leq t$, and let

$$X_n = \frac{\tilde{P}_t^n}{\tilde{P}_s^n}.$$

Note $\mathbb{E}(X_n) = \mathbb{E}[\mathbb{E}(X_n|\mathcal{F}_s)] = 1$ for each n . The first part implies

$$\ell_s - \ell_t = \lim_{n \rightarrow \infty} \frac{1}{n} \log X_n \leq 0 \text{ a.s.}$$

as required.

Problem 8. Let S be a positive supermartingale. Show that there is a positive non-decreasing predictable process A and a positive martingale M such that $A_0 = M_0 = 1$ and $S_t = S_0 M_t / A_t$ for all $t \geq 0$.

Solution 8. Let

$$A_t = \prod_{s=1}^t \frac{S_{s-1}}{\mathbb{E}(S_s|\mathcal{F}_{s-1})}$$

so that A is predictable and non-decreasing since $A_{t+1} = A_t S_t / \mathbb{E}(S_{t+1}|\mathcal{F}_t) \geq A_t$ since S is a supermartingale.

Let

$$M_t = \prod_{s=1}^t \frac{S_s}{\mathbb{E}(S_s|\mathcal{F}_{s-1})}$$

Apply Problem 3 from example sheet 1 to show that M is a martingale. By construction $M/A = S/S_0$.

Problem 9. * Let $(Y_t)_{0 \leq t \leq T}$ be a given adapted, integrable process, and let $(U_t)_{0 \leq t \leq T}$ be its Snell envelope.

(a) Show that if Y is a supermartingale then $U_t = Y_t$ for all t , and if Y is submartingale, then $U_t = \mathbb{E}(Y_T|\mathcal{F}_t)$.

(b) Let τ be any stopping time taking values in $\{0, \dots, T\}$. Show that the process $(U_{t \wedge \tau})_{0 \leq t \leq T}$ is a supermartingale.

(c) Define the random time τ_* by

$$\tau_* = \min \{t \in \{0, \dots, T\} : U_t = Y_t\}.$$

Show that τ_* is a stopping time. Furthermore, show that the process $(U_{t \wedge \tau_*})_{t \in \{0, \dots, T\}}$ is a martingale and, in particular, $U_0 = \mathbb{E}(Y_{\tau_*})$. (That is, τ_* is an optimal stopping time, possibly different than τ^* defined in lectures.)

Solution 9. (a) In both cases we proceed by induction. First suppose that Y is a supermartingale, and that $U_{t+1} = Y_{t+1}$ for some $t < T$. Then

$$U_t = \max\{Y_t, \mathbb{E}(U_{t+1}|\mathcal{F}_t)\} = \max\{Y_t, \mathbb{E}(Y_{t+1}|\mathcal{F}_t)\} = Y_t$$

since $Y_t \geq \mathbb{E}(Y_{t+1}|\mathcal{F}_t)$ by assumption, completing the induction. Similarly, suppose that Y is a submartingale, and that $U_{t+1} = \mathbb{E}(Y_T|\mathcal{F}_{t+1})$. Then

$$U_t = \max\{Y_t, \mathbb{E}(U_{t+1}|\mathcal{F}_t)\} = \max\{Y_t, \mathbb{E}[\mathbb{E}(Y_T|\mathcal{F}_{t+1})|\mathcal{F}_t]\} = \mathbb{E}(Y_T|\mathcal{F}_t)$$

by the tower property and the assumption $Y_t \leq \mathbb{E}(Y_T|\mathcal{F}_t)$, and we're done.

(b) Since U is a supermartingale and the event $\{\tau \geq t+1\} = \{\tau \leq t\}^c$ is in \mathcal{F}_t , we have

$$\begin{aligned} \mathbb{E}[U_{(t+1) \wedge \tau} - U_{t \wedge \tau} | \mathcal{F}_t] &= \mathbb{E}[\mathbb{1}_{\{t+1 \leq \tau\}}(U_{t+1} - U_t) | \mathcal{F}_t] \\ &= \mathbb{1}_{\{t+1 \leq \tau\}} \mathbb{E}[U_{t+1} - U_t | \mathcal{F}_t] \\ &\leq 0 \end{aligned}$$

so the stopped process $(U_{t \wedge \tau})_{0 \leq t \leq T}$ is also supermartingale.

(c) Now, the event

$$\{\tau_* > t\} = \{Y_0 < U_0, \dots, Y_t < U_t\}$$

is \mathcal{F}_t -measurable since both Y and U are adapted, hence τ_* is a stopping time.

Since $U_t = \mathbb{E}(U_{t+1} | \mathcal{F}_t)$ on the event $\{t+1 \leq \tau_*\}$ we have

$$\begin{aligned} U_{(t+1) \wedge \tau_*} - U_{t \wedge \tau_*} &= \mathbb{1}_{\{t+1 \leq \tau_*\}}(U_{t+1} - U_t) \\ &= \mathbb{1}_{\{t+1 \leq \tau_*\}}[U_{t+1} - \mathbb{E}(U_{t+1} | \mathcal{F}_t)]. \end{aligned}$$

In particular $\mathbb{E}[Y_{\tau_*}] = \mathbb{E}[U_{\tau_*}] = \mathbb{E}[U_{T \wedge \tau_*}] = U_0$.

Problem 10. Let $(X_k)_{k \in K}$ be a collection of real-valued random variables, where K is an arbitrary (possibly uncountable) index set. Our aim is to show there exists a random variable Y taking values in $\mathbb{R} \cup \{+\infty\}$ such that

- $Y \geq X_k$ almost surely for all $k \in K$, and
- if $Z \geq X_k$ almost surely for all $k \in K$ then $Z \geq Y$ almost surely.

This will show that the $Y = \text{ess sup}_k X_k$ exists.

(a) Show that there is no loss assuming that $|X_k(\omega)| \leq 1$ for all (k, ω) . Hint: Consider $\hat{X}_k = \tan^{-1}(X_k)$.

From now on, assume $|X_k(\omega)| \leq 1$ for all (k, ω) . Let \mathcal{C} be the collection of all countable subsets of K . Let

$$x = \sup_{A \in \mathcal{C}} \mathbb{E}[\sup_{k \in A} X_k]$$

Let $A_n \in \mathcal{C}$ be such that $\mathbb{E}[\sup_{k \in A_n} X_k] > x - 1/n$ and let $B = \cup_n A_n$. Let $Y = \sup_{k \in B} X_k$.

(b) Why is Y a random variable, i.e. measurable? Show that $\mathbb{E}(Y) = x$.

(c) Pick a $k \in K$, and let $Y_k = \max\{Y, X_k\} = \sup_{h \in B \cup \{k\}} X_h$. Show that $\mathbb{E}(Y_k) = x$. Why does this imply that $X_k \leq Y$ almost surely?

(d) Let Z be a random variable such that $Z \geq X_k$ a.s. for all $k \in K$. Prove that $Z \geq Y$ a.s.

Solution 10. (a) Suppose we know that every family of uniformly bounded random variables (indexed by some arbitrary index set) has an essential supremum. So given the family $(X_k)_{k \in K}$ let $\hat{X}_k = \tan^{-1}(X_k)$. Then the \hat{X}_k are bounded so there exists $\hat{Y} = \text{ess sup}_k \hat{X}_k$. Let $Y = \tan \hat{Y}$. Since the function \tan is strictly increasing it is easy to see that Y has the properties characterising the essential supremum of $(X_k)_k$.

(b) Since A_n is countable, the function $Y_n = \sup_{k \in A_n} X_k$ is measurable for each n . Also $\cup_n A_n = B$ is countable and hence Y is also a random variable. For instance, note

$$\{Y > b\} = \cup_{k \in B} \{X_k > b\}.$$

Now, by replacing A_n with $\cup_{i=1}^n A_i$ we may assume $A_{n-1} \subseteq A_n$ for all $n \geq 1$, and hence we have $Y = \sup_n Y_n = \lim_n Y_n$. The result follows from the bounded convergence theorem.

(c) Since $Y_k \geq Y$ we have $\mathbb{E}(Y_k) \geq \mathbb{E}(Y) = x$. On the other hand,

$$\begin{aligned} x &= \sup_{A \in \mathcal{C}} \mathbb{E}[\sup_{h \in A} X_h] \\ &\geq \mathbb{E}[\sup_{h \in B \cup \{k\}} X_h] \\ &= \mathbb{E}(Y_k). \end{aligned}$$

Since $Y - Y_k \geq 0$ a.s. and $\mathbb{E}(Y - Y_k) = 0$, the pigeonhole principle implies $Y = Y_k$ a.s. Since $\max\{Y, X_k\} = Y$ we have $X_k \leq Y$.

(d) If $\mathbb{P}(Z \geq X_k) = 1$ for all $k \in K$, then

$$\mathbb{P}(Z \geq Y) = \mathbb{P}(\cap_{k \in B} \{Z \geq \hat{X}_k\}) = 1$$

since B is countable.