

1 **Human Movement Variability and Aging**

2 Nicholas Stergiou^{1,2}, Jenny A. Kent¹, and Denise McGrath³

3 ¹Center for Research in Human Movement Variability, Biomechanics Research Building,

4 University of Nebraska at Omaha, Omaha, Nebraska

5 ²College of Public Health, University of Nebraska Medical Center, Omaha, Nebraska

6 ³School of Public Health, Physiotherapy and Population Science, University College Dublin,

7 Dublin, Ireland

8

9Address for correspondence:

10Dr. Nicholas Stergiou

11University of Nebraska at Omaha

12Biomechanics Research Building 214

136160 University Drive, Omaha, NE 68182-0860, USA

14Tel: +1-402-554-3247; Fax: +1-402-554-3693

15Email: nstergiou@unomaha.edu

16

17

18

19

Abstract

20

Keywords: Biomechanics, Gait, Elderly, Complexity, Fractals

21

22 An optimal level of variability enables us to interact adaptively and safely to a
 23 continuously changing environment, where often our movements must be adjusted in a matter of
 24 milliseconds. A large body of research exists that demonstrates natural variability in healthy gait
 25 (along with variability in other, healthy biological signals e.g. heart rate), and a loss of this
 26 variability in ageing and injury, as well as in a variety of neurodegenerative and physiological
 27 disorders. We submit that this field of research is now in pressing need of an innovative “next
 28 step” that goes beyond the many descriptive studies that characterize levels of variability in
 29 various patient populations. We need to devise novel therapies that will harness the existing
 30 knowledge on biological variability and create new possibilities for those in the grip of disease.
 31 We also propose that the nature of the specific physiological limitation present in the
 32 neuromuscular apparatus may be less important in the physiological complexity framework than
 33 the control mechanisms adopted by the older individual in the coordination of the available
 34 degrees of freedom. The theoretical underpinnings of this framework suggest that interventions
 35 designed to restore healthy system dynamics may optimize functional improvements in older
 36 adults. We submit that interventions based on the restoration of optimal variability and
 37 movement complexity could potentially be applied across a range of diseases or dysfunctions as
 38 it addresses the adaptability and coordination of available degrees of freedom, regardless of the
 internal constraints of the individual.

39Human Movement Variability and Aging

40 When we perform the same task multiple times, we can easily observe that we never
41execute it the exact same way. This is obvious even with elite performers, like athletes and
42musicians. When it comes down to simple everyday walking, every one of us is an elite
43performer, but as we observe our steps behind us on the sand we can clearly see that they are
44never identical. These natural fluctuations in motor performance define the presence of human
45movement variability which is ubiquitous in all biological systems. Recently, the role of
46movement variability has attracted significant attention due to its relationship to pathology and
47performance (Harrison & Stergiou, 2015; N. Stergiou & Decker, 2011).

48 Previous theoretical frameworks consider variability as an indicator of noise in the
49control system and it has been quantified using traditional linear statistical measures (e.g.
50standard deviation; (Newell & Corcos, 1993; Schmidt & Lee, 2005; N. Stergiou, Harbourne, &
51Cavanaugh, 2006)). Such measures contain no information about how the motor system responds
52to change over time. Practically, the linear measures are measures of centrality and thus, provide
53a description of the amount or magnitude of the variability around that central point. This is
54accomplished by quantifying the magnitude of variation in a set of values independent of their
55order in the distribution. From this perspective, clinicians and scientists believe that the mean is
56the “gold standard” of healthy behavior. Any deviation from this gold standard is error, or
57undesirable behavior, or the result of instability. However, recent literature from several
58disciplines and medical areas, including brain function and disease dynamics have shown that
59many apparently “noisy” phenomena are the result of nonlinear interactions and have
60deterministic origins (Amato, 1992; Buchman, Cobb, Lapedes, & Kepler, 2001; J. T. Cavanaugh,
61Kochi, & Stergiou, 2010; Garfinkel, Spano, Ditto, & Weiss, 1992; Goldstein, Toweill, Lai,

62Sonnenthal, & Kimberly, 1998; Orsucci, 2006; Slutzky, Cvitanovic, & Mogul, 2001; Toweill &
63Goldstein, 1998; Wagner, Nafz, & Persson, 1996). Based on this information, it has been
64proposed that the natural fluctuations that are present in normal motor tasks (e.g. stride-to-stride
65fluctuations in normal walking) are characterized by an appropriate or optimal state of variability
66(Harrison & Stergiou, 2015; N. Stergiou et al., 2006; N. Stergiou & Decker, 2011). Optimal
67variability is associated with complex interactions across multiple control systems, feedback
68loops and regulatory processes that enable an organism to function and adapt to the demands of
69everyday life ((Harrison & Stergiou, 2015; N. Stergiou et al., 2006; N. Stergiou & Decker, 2011);
70Figure 1). This physiological complexity is recognized as an inherent attribute of healthy
71biological systems, whereas the loss of complexity with aging and disease is thought to reduce
72the adaptive capabilities of the individual (Buzzi, Stergiou, Kurz, Hageman, & Heidel, 2003; A.
73L. Goldberger, 2001; A. L. Goldberger et al., 2002). A loss of complexity can refer to either an
74overly constrained, periodic system, or an overly random, incoherent system (Dossey, 2010).
75Healthy human function thus requires the coexistence of the oppositional factors of coherence
76and chaos (Dossey, 2010). It is important to note that the specialized concept of “chaos”
77discussed here is distinct from, and indeed contrary to, the English language notion of chaos
78which means confusion and disorder. In fact, when we refer to mathematical chaos in a system,
79we are pointedly referring to an underlying order or pattern that is contained within a complex,
80variable system, capable of sudden and marked change.

81 **INSERT FIGURE 1 ABOUT HERE**

82 The physiological complexity paradigm was first introduced almost 30 years ago by
83Goldberger and colleagues, based on heart rate dynamics (A. L. Goldberger, Findley, Blackburn,
84& Mandell, 1984). Since then there has been an abundance of literature that reveals the presence

85 of physiological complexity in healthy systems, and a loss of complexity in ill-health e.g. in
 86 brain activity and pathological cognitive processes (Bystritsky, Nierenberg, Feusner, &
 87 Rabinovich, 2012; A. C. Yang et al., 2013), heart failure (A. L. Goldberger et al., 2002), fetal
 88 distress syndromes (A. L. Goldberger, 1996), respiration (Peng et al., 2002), and ambulatory
 89 behaviors in older adults (J. T. Cavanaugh et al., 2010; J. T. Cavanaugh, Coleman, Gaines, Laing,
 90 & Morey, 2007). Evidence for loss of complexity in the motor system (e.g. locomotor system)
 91 with aging and disease first emerged in the mid- to late-nineties (J. M. Hausdorff et al., 1997; J.
 92 M. Hausdorff, Cudkowicz, Firtion, Wei, & Goldberger, 1998), with subsequent cross-sectional
 93 studies relating levels of motor and specifically locomotor complexity to diseases, conditions or
 94 age categories, many of which were performed by our laboratory (Buzzi et al., 2003; Decker,
 95 Moraiti, Stergiou, & Georgoulis, 2011; Deffeyes et al., 2009). However, despite the enormous
 96 potential that this emerging field presents for a) improving our understanding of aging and
 97 disease/dysfunction, b) providing sensitive biomarkers for evaluating behavioral or
 98 pharmacological interventions, and c) designing novel interventions based on restoration of
 99 complexity, its actual impact in the field of rehabilitative medicine has been minimal.

100 The reason for the relatively slow progress may be due to the fact that no single statistical
 101 measure can be used to assess the complexity of physiologic systems. Instead, an extensive
 102 “toolkit” of evolving metrics is needed to probe different aspects of these complicated
 103 behaviours (A. L. Goldberger, Peng, & Lipsitz, 2002). These metrics must be implemented with
 104 an in-depth appreciation of their purpose, strengths and limitations (N. Stergiou, 2004; N.
 105 Stergiou, 2016). Fractal-based complexity metrics hold great promise in the study of walking.
 106 Multiple studies support the idea that stride-to-stride variations in healthy gait exhibit nonlinear
 107 and fractal-like fluctuations extending over hundreds of steps, reflective of complexity in healthy

108gait (J. M. Hausdorff, Peng, Ladin, Wei, & Goldberger, 1995; J. M. Hausdorff et al., 1996; J. M.
 109Hausdorff, 2007). The classic definition of a fractal, first described by Mandelbrot (Mandelbrot,
 1101977), is a geometric object with “self-similarity” over multiple measurement scales (L. A.
 111Lipsitz, 2002). The outputs of the locomotor system measured over time exhibit such fractal
 112properties (Delignieres & Torre, 2009), demonstrating power-law scaling such that the smaller
 113the frequency of oscillation (f) of these signals, the larger their amplitude (amplitude squared is
 114power) (L. A. Lipsitz, 2002). This power-law relation can be expressed as $1/f$, and is referred to
 115as pink noise, where oscillations appear self-similar when observed over seconds, minutes,
 116hours, or days. Pathological gait observed in the elderly population demonstrates a breakdown of
 117this $1/f$ scaling; in other words, a loss of complexity where their movement dynamics are either
 118too periodic or too random (Herman, Giladi, Gurevich, & Hausdorff, 2005).

119 What is the function of this $1/f$ scaling, commonly seen in well-coordinated behaviours,
 120and less so in non-optimal performance or with aging and disease? Vaillancourt and colleagues
 121submit that $1/f$ scaling confers enhanced connectivity between motor control processes (D. E.
 122Vaillancourt & Newell, 2003), while a breakdown in fractal scaling (as observed in stride interval
 123time series in pathological gait) arises from a gradual deterioration in the number of “functioning
 124elements of a given system and/or a decrease in the interactions between these components” (A.
 125L. Goldberger, 2001; A. L. Goldberger et al., 2002). From this perspective, there is no particular
 126component that causes $1/f$ scaling to occur, or indeed to break down. Instead, it is an emergent
 127property that stems from the interactions across the many spatio-temporal scales of organization
 128of an organism (Van Orden, Holden, & Turvey, 2005), giving rise to a “nimble” but coordinated
 129motor control system. Vaillancourt and colleagues have demonstrated that any behavioural
 130performance results from the coordination of the degrees of freedom available to the individual

with respect to constraints imposed from the neuromuscular system, the task, and the environment (D. E. Vaillancourt, Sosnoff, & Newell, 2004). Drawing from evidence that strongly supports a breakdown of fractal scaling in aging physiological systems (L. A. Lipsitz & Goldberger, 1992; D. E. Vaillancourt & Newell, 2002), they have further shown that older adults have a reduced capacity to adapt to faster time scales in both feedback and feedforward processes (Sosnoff & Newell, 2008). Other support for this hypothesis is provided by Bierbaum *et al.* (Bierbaum, Peper, Karamanidis, & Arampatzis, 2011) who showed that locomotor behaviour in older adults is more conservative compared to the young, leading to disadvantages in the reactive adaptation during disturbed walking. Hsu *et al.* (Hsu, Chou, & Woollacott, 2013) concluded that in normal aging, adults lose the compensatory strategy afforded by the flexible control of multiple joints when stabilizing the center of mass after receiving a balance perturbation. Similarly, our previous work has shown that older adults exhibit reduced adaptive capabilities (Byrne *et al.*, 2002; Kurz & Stergiou, 2003).

Causes of gait and balance disturbances in the elderly are multifactorial. For example, abnormal gait can be due to a single disease or multiple diseases developing simultaneously across all sensorimotor levels (Martens & Almeida, 2012; Vinti, Couillandre, & Thoumie, 2010), increased reliance on central mechanisms (Seidler *et al.*, 2010), atrophy or disease of brain structures involved in these central mechanisms (Seidler *et al.*, 2010), degeneration of neurotransmitter systems (R. Cham, Studenski, Perera, & Bohnen, 2008; R. Cham, Perera, Studenski, & Bohnen, 2007), reduced muscle mass and/or muscle quality (Shin, Valentine, Evans, & Sosnoff, 2012), psychosocial factors (Baik & Lang, 2007), and medication use (Boudreau *et al.*, 2009; Hilmer *et al.*, 2009). Compensatory postures, slower gait speeds and “cautious” gait are all trademarks of abnormal gait in the elderly. The capacity to maintain

154balance when walking through unpredictably changing environments is critical to independent
155living for this population. Increasingly, altered gait variability is being reported in older adults
156and has been associated with a variety of disorders ranging from joint and skeletal problems
157(Kiss, 2011) (lowest-level gait disturbances) to Huntington's disease (J. M. Hausdorff et al.,
1581997), Parkinson's disease (J. M. Hausdorff et al., 1998; Kurz, Markopoulou, & Stergiou, 2010),
159higher level gait disorders (Herman et al., 2005), and falls (J. M. Hausdorff, 2007; Montero-
160Odasso, Muir, & Speechley, 2012; Paterson, Hill, & Lythgo, 2011; Toebe, Hoozemans, Furrer,
161Dekker, & van Dieën, 2012). Reduced adaptive capacity in the locomotor system has been linked
162to falls due to the difficulty that older adults experience in recovering quickly from a loss of
163dynamic balance (Madigan & Lloyd, 2005; Wojcik, Thelen, Schultz, Ashton-Miller, &
164Alexander, 1999).

165 There are compelling findings in both animal and human studies that suggest that the
166complexity of locomotor patterns provide a rich source of information that could be relevant to
167the diagnosis and management of a variety of diseases that affect an aging population. Our
168previous research has shown that highly active older adults exhibit more complex patterns of
169locomotor activity than less active older adults, despite the absence of differences between these
170groups in standard measures of variability of their step counts (J. T. Cavanaugh et al., 2010). Hu
171and colleagues have recently shown that older adults and dementia patients have disrupted fractal
172activity patterns (K. Hu, Van Someren, Shea, & Scheer, 2009) and that the degree of disruption is
173positively related to the burden of amyloid plaques - a marker of Alzheimer's disease severity
174(K. Hu, Harper, Shea, Stopa, & Scheer, 2013). They also found that fractal scaling in activity
175fluctuations is unrelated to the average level of activity as assessed within and between subjects
176(K. Hu et al., 2004). A study of primates suggests that a loss of complexity in locomotor

behaviour that is associated with illness and aging, reduces the efficiency with which an animal is able to cope with heterogeneity in its natural environment (Macintosh, Alados, & Huffman, 2011). Japanese quail became less periodic and more complex in their locomotor behaviour when they were stimulated to explore, without there being commensurate changes in the percentage of total time spent walking, or in the average duration of the walking events (Kembro, Perillo, Pury, Satterlee, & Marin, 2009). Additionally, fractal scaling has been observed in the locomotor activity of young, healthy small mammals, a feature that is less evident in aged animals (Anteneodo & Chialvo, 2009).

While the relationships between fractal patterns of locomotor activity and health are indeed intriguing, Hu and colleagues have recently identified a possible neural site that is responsible for scale-invariant regulation of a neurophysiological system over a range of time scales. They demonstrated that lesioning the suprachiasmatic nucleus (SCN) of the anterior hypothalamus in rats i.e., the neural node responsible for circadian rhythms, led to the disappearance of fractality in both heart rate and locomotor rhythms (K. Hu, Scheer, Buijs, & Shea, 2008; K. Hu, Scheer, Ivanov, Buijs, & Shea, 2007). Additionally, they have recently shown that the degree of disruption to fractal activity in dementia patients is strongly associated with vasopressinergic and neurotensinergic neurons (two major circadian neurotransmitters) in post-mortem SCN, and can better predict changes of the two neurotransmitters than other traditional circadian measures (K. Hu et al., 2013). The authors concluded that the SCN impacts human activity regulation at multiple time scales and that disrupted fractal activity may serve as a non-invasive biomarker of SCN neurodegeneration in dementia. A further study by this group demonstrated that multi-unit neural activity of the SCN in mice and rats exhibited fractal fluctuations in vivo that were abolished in preparations in vitro. These empirical results suggest

200that it is not the activity of the SCN in isolation, but the activity of the SCN in concert with other
201physiological mechanisms that lead to fractal fluctuations in physiological output (K. Hu et al.,
2022011). It has been widely shown that most peripheral organs and tissues, including skeletal
203muscle, can express circadian oscillations in isolation, yet still receive and may require input
204from the SCN in vivo (Mohawk, Green, & Takahashi, 2012). A well-functioning circadian
205system therefore requires SCN interaction with peripheral oscillators. Many authors have
206discussed the feedback loops that are ubiquitous at the molecular, cellular, tissue and systems
207level between the inputs and outputs of the circadian system (X. Yang, 2010).

208 Therapeutic interventions that boost the circadian signal and restore the temporal order of
209a system may act to ameliorate some of the decline seen in aged individuals. It is accepted that
210the age-related attenuation of the central timing signal generated by the SCN is associated with a
211number of health problems such as metabolic syndrome, neurodegenerative disorders, and
212cardiovascular diseases (Kondratova & Kondratov, 2012). Transplantation of a ‘young’ SCN into
213aged animals resulted in improvements in numerous rhythmic functions, including behavioural
214rhythms in locomotion (Li & Satinoff, 1995). Tranah and colleagues have shown that older
215community dwelling adults with weak circadian locomotor activity rhythms have a higher
216mortality risk and increased risk of developing dementia and mild cognitive impairment (Tranah
217et al., 2010; Tranah et al., 2011). It has been shown consistently in animal models that aging does
218not affect the size or the number of neurons in the SCN (Madeira, Sousa, Santer, Paula-Barbosa,
219& Gundersen, 1995); rather, aging brings about significant changes in electrophysiological and
220neurochemical outputs of the SCN (Colwell, 2011). Previous studies suggest that some but not
221all peripheral circadian oscillators exhibit age-related changes in rhythmicity (Yamazaki et al.,
2222002) and that some of the related tissues retain the capacity to oscillate but are not appropriately

driven in vivo by physical activity rhythms (Asai et al., 2001). Locomotor activity can influence SCN function via neuronal feedback loops (Hughes & Piggins, 2012). Information on the precise role that circadian abnormalities play in the aging process is somewhat limited, however it has been hypothesized that the fragmentation of behavioural activity with aging may worsen the age-related defects in the central clock function, leading to a downward spiral (Farajnia, Deboer, Rohling, Meijer, & Michel, 2014). Together, these considerations suggest that interventions to regulate circadian activity rhythm abnormalities, are warranted in older adults (Tranah et al., 2011).

We have recently shown that complexity in the locomotor system is a modifiable property in both young and elderly adults, by walking while listening to an auditory stimulus with a complex structure (Hunt, McGrath, & Stergiou, 2014; Kaipust, McGrath, Mukherjee, & Stergiou, 2013). In other words, we have shown that fractal patterns in gait parameters (i.e. stride intervals) can be restored in older adults using a fractal-based auditory cue in an audio-motor entrainment paradigm. Such an intervention to restore fractal patterns in locomotor activity in older adults, enabled by wearable sensor technology that delivers the relevant cues, could have a positive effect on circadian rhythms and adaptive capability promoting SCN interaction with an effective fractal based peripheral oscillator. It could also revolutionize the current practice of metronomic auditory cueing for the rehabilitation of pathological gait by replacing the metronome model with a novel auditory stimulus with a 1/f structure (i.e. pink noise), embedded into music.

Although walking to a metronomic auditory stimulus has been shown to increase gait tempo and stride length (Ford, Malone, Nyikos, Yelisetty, & Bickel, 2010; Roerdink et al., 2009), its implicit outcome is to reduce the natural stride-to-stride fluctuations to zero, thus destroying

the 1/f scaling (J. M. Hausdorff et al., 1996) that enables the neuromuscular system to adapt to a continuously changing environment. Humans have been coordinating their movement to external rhythms since antiquity. Entrainment to an external rhythm can occur even when there is a high degree of rhythmic complexity and ambiguity in music (Skoe & Kraus, 2009; Snyder, 2003). The dynamical systems approach describes musical rhythmic entrainment as an active, self-sustained, periodic oscillation at multiple time scales, enabling the listener to use predictive timing to maintain a stable, multi-periodic pattern and synchronize movements at the main beat or other metrical levels (Large, 2000). There is convincing evidence in the literature suggesting that both healthy and diseased or injured adults can entrain their gait to a metronome (Delval et al., 2008; Ford et al., 2010; Hayden, Clair, Johnson, & Otto, 2009). Drawing from dynamical systems theory, we have advanced this field by showing that the structure of an auditory stimulus is expressed in the patterns of gait variability produced by both young and elderly adults (Hunt et al., 2014; Kaipust et al., 2013). A recent study by Hove *et al.* (Hove, Suzuki, Uchitomi, Orimo, & Miyake, 2012) showed that an “interactive” auditory stimulus, based on nonlinear oscillators restored 1/f scaling in Parkinson’s disease patients that persisted 5 minutes after the stimulus was removed, indicating stabilization of the internal rhythm generating system and the reintegration of timing networks. These experiments show that complex (rather than periodic) interaction is important for the (re)emergence of 1/f structure in human gait behaviour.

In conclusion, an optimal level of variability enables us to interact adaptively and safely to a continuously changing environment, where often our movements must be adjusted in a matter of milliseconds. A large body of research exists that demonstrates natural variability in healthy gait (along with variability in other, healthy biological signals e.g. heart rate), and a loss of this variability in ageing and injury, as well as in a variety of neurodegenerative and

269physiological disorders. We submit that this field of research is now in pressing need of an
270innovative “next step” that goes beyond the many descriptive studies that characterize levels of
271variability in various patient populations. We need to devise novel therapies that will harness the
272existing knowledge on biological variability and create new possibilities for those in the grip of
273disease.

274 We also propose that the nature of the specific physiological limitation present in the
275neuromuscular apparatus may be less important in the physiological complexity framework than
276the control mechanisms adopted by the older individual in the coordination of the available
277degrees of freedom. The theoretical underpinnings of this framework suggest that interventions
278designed to restore healthy system dynamics may optimize functional improvements in older
279adults (Harrison & Stergiou, 2015; Manor & Lipsitz, 2013; N. Stergiou & Decker, 2011). We
280submit that interventions based on the restoration of optimal variability and movement
281complexity could potentially be applied across a range of diseases or dysfunctions as it addresses
282the adaptability and coordination of available degrees of freedom, regardless of the internal
283constraints of the individual.

284

285

286 **Acknowledgements**

287 This work was supported by the Center for Research in Human Movement Variability of
288the University of Nebraska Omaha and the NIH (P20GM109090).

289

290
291

References

- 292Amato, I. (1992). Chaos breaks out at NIH, but order may come of it. *Science (New York, N.Y.)*,
293 256(5065), 1763-1764.
- 294Anteneodo, C., & Chialvo, D. R. (2009). Unraveling the fluctuations of animal motor activity.
295 *Chaos (Woodbury, N.Y.)*, 19(3), 033123. doi:10.1063/1.3211189 [doi]
- 296Asai, M., Yoshinobu, Y., Kaneko, S., Mori, A., Nikaido, T., Moriya, T., Shibata, S. (2001).
297 Circadian profile of per gene mRNA expression in the suprachiasmatic nucleus,
298 paraventricular nucleus, and pineal body of aged rats. *Journal of Neuroscience Research*,
299 66(6), 1133-1139.
- 300Baik, J. S., & Lang, A. E. (2007). Gait abnormalities in psychogenic movement disorders.
301 *Movement Disorders*, 22(3), 395-399.
- 302Bierbaum, S., Peper, A., Karamanidis, K., & Arampatzis, A. (2011). Adaptive feedback potential
303 in dynamic stability during disturbed walking in the elderly. *Journal of Biomechanics*,
304 44(10), 1921-1926.
- 305Boudreau, R. M., Hanlon, J. T., Roumani, Y. F., Studenski, S. A., Ruby, C. M., Wright, R. M., . . .
306 Simonsick, E. M. (2009). Central nervous system medication use and incident mobility
307 limitation in community elders: The health, aging, and body composition study.
308 *Pharmacoepidemiology and Drug Safety*, 18(10), 916-922.
- 309Buchman, T. G., Cobb, P. J., Lapedes, A. S., & Kepler, T. B. (2001). Complex systems analysis:
310 A tool for shock research. *Shock*, 16(4), 248-251.
- 311Buzzi, U., Stergiou, N., Kurz, M., Hageman, P., & Heidel, J. (2003). Nonlinear dynamics
312 indicates aging affects variability during gait. *Clinical Biomechanics*, 18(5), 435-443.
313 doi:10.1016/S0268-0033(03)00029-9

- 314Byrne, J. E., Stergiou, N., Blanke, D., Houser, J. J., Kurz, M. J., & Hageman, P. A. (2002).
315 Comparison of gait patterns between young and elderly women: An examination of
316 coordination. *Perceptual and Motor Skills*, 94(1), 265-280.
- 317Bystritsky, A., Nierenberg, A. A., Feusner, J. D., & Rabinovich, M. (2012). Computational non-
318 linear dynamical psychiatry: A new methodological paradigm for diagnosis and course of
319 illness. *Journal of Psychiatric Research*, 46(4), 428-435.
- 320Cavanaugh, J. T., Kochi, N., & Stergiou, N. (2010). Nonlinear analysis of ambulatory activity
321 patterns in community-dwelling older adults. *The Journals of Gerontology.Series A,*
322 *Biological Sciences and Medical Sciences*, 65(2), 197-203. doi:10.1093/gerona/glp144 [doi]
- 323Cavanaugh, J. T., Coleman, K. L., Gaines, J. M., Laing, L., & Morey, M. C. (2007). Using step
324 activity monitoring to characterize ambulatory activity in Community-Dwelling older adults.
325 *Journal of the American Geriatrics Society*, 55(1), 120-124.
- 326Cham, R., Studenski, S. A., Perera, S., & Bohnen, N. I. (2008). Striatal dopaminergic
327 denervation and gait in healthy adults. *Experimental Brain Research*, 185(3), 391-398.
- 328Cham, R., Perera, S., Studenski, S. A., & Bohnen, N. I. (2007). Striatal dopamine denervation
329 and sensory integration for balance in middle-aged and older adults. *Gait & Posture*, 26(4),
330 516-525.
- 331Colwell, C. S. (2011). Linking neural activity and molecular oscillations in the SCN. *Nature*
332 *Reviews Neuroscience*, 12(10), 553-569.
- 333Decker, L. M., Moraiti, C., Stergiou, N., & Georgoulis, A. D. (2011). New insights into anterior
334 cruciate ligament deficiency and reconstruction through the assessment of knee kinematic
335 variability in terms of nonlinear dynamics. *Knee Surgery, Sports Traumatology,*
336 *Arthroscopy*, 19(10), 1620-1633.

- 337Deffeyes, J. E., Kochi, N., Harbourne, R. T., Kyvelidou, A., Stuberg, W. A., & Stergiou, N.
338 (2009). Nonlinear detrended fluctuation analysis of sitting center-of-pressure data as an early
339 measure of motor development pathology in infants. *Nonlinear Dynamics, Psychology, and*
340 *Life Sciences*, 13(4), 351-368.
- 341Delignieres, D., & Torre, K. (2009). Fractal dynamics of human gait: A reassessment of the 1996
342 data of hausdorff et al. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 106(4), 1272-
343 1279. doi:10.1152/japplphysiol.90757.2008 [doi]
- 344Delval, A., Krystkowiak, P., Delliaux, M., Blatt, J., Derambure, P., Destée, A., & Defebvre, L.
345 (2008). Effect of external cueing on gait in huntington's disease. *Movement Disorders*,
346 23(10), 1446-1452.
- 347Dossey, L. (2010). Coherence, chaos, and the coincidentia oppositorum. *Explore: The Journal of*
348 *Science and Healing*, 6(6), 339-345.
- 349Farajnia, S., Deboer, T., Rohling, J. H., Meijer, J. H., & Michel, S. (2014). Aging of the
350 suprachiasmatic clock. *The Neuroscientist : A Review Journal Bringing Neurobiology,*
351 *Neurology and Psychiatry*, 20(1), 44-55. doi:10.1177/1073858413498936 [doi]
- 352Ford, M. P., Malone, L. A., Nyikos, I., Yelisetty, R., & Bickel, C. S. (2010). Gait training with
353 progressive external auditory cueing in persons with parkinson's disease. *Archives of*
354 *Physical Medicine and Rehabilitation*, 91(8), 1255-1261.
- 355Garfinkel, A., Spano, M. L., Ditto, W. L., & Weiss, J. N. (1992). Controlling cardiac chaos.
356 *Science (New York, N.Y.)*, 257(5074), 1230-1235.
- 357Goldberger, A. L. (2001). Heartbeats, hormones, and health - is variability the spice of life?
358 *American Journal of Respiratory and Critical Care Medicine*, 163(6), 1289-1290.

- 359Goldberger, A. L., Amaral, L. A. N., Hausdorff, J. M., Ivanov, P. C., Peng, C. K., & Stanley, H.
360 E. (2002). Fractal dynamics in physiology: Alterations with disease and aging. *Proceedings*
361 *of the National Academy of Sciences of the United States of America*, 99, 2466-2472.
362 doi:10.1073/pnas.012579499
- 363Goldberger, A. L. (1996). Non-linear dynamics for clinicians: Chaos theory, fractals, and
364 complexity at the bedside. *The Lancet*, 347(9011), 1312-1314.
- 365Goldberger, A. L., Findley, L. J., Blackburn, M. R., & Mandell, A. J. (1984). Nonlinear dynamics
366 in heart failure: Implications of long-wavelength cardiopulmonary oscillations. *American*
367 *Heart Journal*, 107(3), 612-615.
- 368Goldberger, A. L., Peng, C. -, & Lipsitz, L. A. (2002). What is physiologic complexity and how
369 does it change with aging and disease? *Neurobiology of Aging*, 23(1), 23-26.
- 370Goldstein, B., Toweill, D., Lai, S., Sonnenthal, K., & Kimberly, B. (1998). Uncoupling of the
371 autonomic and cardiovascular systems in acute brain injury. *The American Journal of*
372 *Physiology*, 275(4 Pt 2), R1287-92.
- 373Harrison, S. J., & Stergiou, N. (2015). Complex adaptive behavior and dexterous action.
374 *Nonlinear Dynamics, Psychology, and Life Sciences*, 19(4), 345-394.
- 375Hausdorff, J. M., Mitchell, S. L., Firtion, R., Peng, C. K., Cudkowicz, M. E., Wei, J. Y., &
376 Goldberger, A. L. (1997). Altered fractal dynamics of gait: Reduced stride-interval
377 correlations with aging and huntington's disease. *Journal of Applied Physiology*, 82(1), 262-
378 269.
- 379Hausdorff, J. M., Peng, C. K., Ladin, Z., Wei, J. Y., & Goldberger, A. L. (1995). Is walking a
380 random walk? evidence for long-range correlations in stride interval of human gait. *Journal*
381 *of Applied Physiology (Bethesda, Md.: 1985)*, 78(1), 349-358.

- 382 Hausdorff, J. M., Purdon, P. L., Peng, C. K., Ladin, Z., Wei, J. Y., & Goldberger, A. L. (1996).
383 Fractal dynamics of human gait: Stability of long-range correlations in stride interval
384 fluctuations. *Journal of Applied Physiology*, 80(5), 1448-1457.
- 385 Hausdorff, J. M. (2007). Gait dynamics, fractals and falls: Finding meaning in the stride-to-stride
386 fluctuations of human walking. *Human Movement Science*, 26(4), 555-589.
387 doi:10.1016/j.humov.2007.05.003
- 388 Hausdorff, J. M., Cudkowicz, M. E., Firtion, R., Wei, J. Y., & Goldberger, A. L. (1998). Gait
389 variability and basal ganglia disorders: Stride-to-stride variations of gait cycle timing in
390 parkinson's disease and huntington's disease. *Movement Disorders*, 13(3), 428-437.
- 391 Hayden, R., Clair, A. A., Johnson, G., & Otto, D. (2009). The effect of rhythmic auditory
392 stimulation (RAS) on physical therapy outcomes for patients in gait training following
393 stroke: A feasibility study. *International Journal of Neuroscience*, 119(12), 2183-2195.
- 394 Herman, T., Giladi, N., Gurevich, T., & Hausdorff, J. M. (2005). Gait instability and fractal
395 dynamics of older adults with a "cautious" gait: Why do certain older adults walk fearfully?
396 *Gait & Posture*, 21(2), 178-185.
- 397 Hilmer, S. N., Mager, D. E., Simonsick, E. M., Ling, S. M., Windham, B. G., Harris, T. B., . . .
398 Study, H. A. (2009). Drug burden index score and functional decline in older people. *The*
399 *American Journal of Medicine*, 122(12), 1142-1149. e2.
- 400 Hove, M. J., Suzuki, K., Uchitomi, H., Orimo, S., & Miyake, Y. (2012). Interactive rhythmic
401 auditory stimulation reinstates natural 1/f timing in gait of parkinson's patients. *PLoS One*,
402 7(3)
- 403 Hsu, W., Chou, L., & Woollacott, M. (2013). Age-related changes in joint coordination during
404 balance recovery. *Age*, 35(4), 1299-1309.

- 405Hu, K., Harper, D. G., Shea, S. A., Stopa, E. G., & Scheer, F. A. (2013). Noninvasive fractal
406 biomarker of clock neurotransmitter disturbance in humans with dementia. *Scientific*
407 *Reports*, 3, 2229. doi:10.1038/srep02229 [doi]
- 408Hu, K., Meijer, J. H., Shea, S. A., vanderLeest, H. T., Pittman-Polletta, B., Houben, T., . . .
409 Scheer, F. A. (2011). Fractal patterns of neural activity exist within the suprachiasmatic
410 nucleus and require extrinsic network interactions. *PloS One*, 7(11), e48927-e48927.
- 411Hu, K., Scheer, F. A., Buijs, R. M., & Shea, S. A. (2008). The endogenous circadian pacemaker
412 imparts a scale-invariant pattern of heart rate fluctuations across time scales spanning
413 minutes to 24 hours. *Journal of Biological Rhythms*, 23(3), 265-273.
414 doi:10.1177/0748730408316166 [doi]
- 415Hu, K., Van Someren, E. J., Shea, S. A., & Scheer, F. A. (2009). Reduction of scale invariance of
416 activity fluctuations with aging and alzheimer's disease: Involvement of the circadian
417 pacemaker. *Proceedings of the National Academy of Sciences of the United States of*
418 *America*, 106(8), 2490-2494. doi:10.1073/pnas.0806087106 [doi]
- 419Hu, K., Ivanov, P. C., Chen, Z., Hilton, M. F., Stanley, H. E., & Shea, S. A. (2004). Non-random
420 fluctuations and multi-scale dynamics regulation of human activity. *Physica A: Statistical*
421 *Mechanics and its Applications*, 337(1), 307-318.
- 422Hu, K., Scheer, F. A. J. L., Ivanov, P. C., Buijs, R. M., & Shea, S. A.,. (2007). The
423 suprachiasmatic nucleus functions beyond circadian rhythm generation. *Neuroscience*,
424 149(3), 508-517.
- 425Hughes, A. T., & Piggins, H. D. (2012). Feedback actions of locomotor activity to the circadian
426 clock. *Progress in Brain Research*, 199, 305-336. doi:10.1016/B978-0-444-59427-3.00018-6
427 [doi]

- 428 Hunt, N., McGrath, D., & Stergiou, N. (2014). The influence of auditory-motor coupling on
429 fractal dynamics in human gait. *Scientific Reports*, 4, 5879. doi:doi:10.1038/srep05879
- 430 Kaipust, J. P., McGrath, D., Mukherjee, M., & Stergiou, N. (2013). Gait variability is altered in
431 older adults when listening to auditory stimuli with differing temporal structures. *Annals of*
432 *Biomedical Engineering*, 41(8), 1595-1603.
- 433 Kembro, J. M., Perillo, M. A., Pury, P. A., Satterlee, D. G., & Marin, R. H. (2009). Fractal
434 analysis of the ambulation pattern of japanese quail. *British Poultry Science*, 50(2), 161-170.
- 435 Kiss, R. M. (2011). Effect of severity of knee osteoarthritis on the variability of gait parameters.
436 *Journal of Electromyography and Kinesiology*, 21(5), 695-703.
- 437 Kondratova, A. A., & Kondratov, R. V. (2012). The circadian clock and pathology of the ageing
438 brain. *Nature Reviews Neuroscience*, 13(5), 325-335.
- 439 Kurz, M. J., Markopoulou, K., & Stergiou, N. (2010). Attractor divergence as a metric for
440 assessing walking balance. *Nonlinear Dynamics, Psychology, and Life Sciences*, 14(2), 151-
441 164.
- 442 Kurz, M. J., & Stergiou, N. (2003). The aging humans neuromuscular system expresses less
443 certainty for selecting joint kinematics during gait. *Neuroscience Letters*, 348(3), 155-158.
- 444 Large, E. W. (2000). On synchronizing movements to music. *Human Movement Science*, 19(4),
445 527-566.
- 446 Li, H., & Satinoff, E. (1995). Changes in circadian rhythms of body temperature and sleep in old
447 rats. *The American Journal of Physiology*, 269(1 Pt 2), R208-14.
- 448 Lipsitz, L. A. (2002). Dynamics of stability: The physiologic basis of functional health and
449 frailty. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*,
450 57(3), B115-25.

- 451Lipsitz, L. A., & Goldberger, A. L. (1992). Loss of 'complexity' and aging: Potential applications
452 of fractals and chaos theory to senescence. *Jama*, 267(13), 1806-1809.
- 453Macintosh, A. J., Alados, C. L., & Huffman, M. A. (2011). Fractal analysis of behaviour in a wild
454 primate: Behavioural complexity in health and disease. *Journal of the Royal Society,*
455 *Interface / the Royal Society*, 8(63), 1497-1509. doi:10.1098/rsif.2011.0049 [doi]
- 456Madeira, M. D., Sousa, N., Santer, R. M., Paula-Barbosa, M. M., & Gundersen, H. J. G. (1995).
457 Age and sex do not affect the volume, cell numbers, or cell size of the suprachiasmatic
458 nucleus of the rat: An unbiased stereological study. *Journal of Comparative Neurology*,
459 361(4), 585-601.
- 460Madigan, M. L., & Lloyd, E. M. (2005). Age and stepping limb performance differences during a
461 single-step recovery from a forward fall. *The Journals of Gerontology.Series A, Biological*
462 *Sciences and Medical Sciences*, 60(4), 481-485. doi:60/4/481 [pii]
- 463Mandelbrot, B. B. (1977). *The fractal geometry of nature*. New York: W.H.Freeman and
464 Company.
- 465Manor, B., & Lipsitz, L. A. (2013). Physiologic complexity and aging: Implications for physical
466 function and rehabilitation. *Progress in Neuro-Psychopharmacology and Biological*
467 *Psychiatry*, 45, 287-293.
- 468Martens, K. A. E., & Almeida, Q. J. (2012). Dissociating between sensory and perceptual deficits
469 in PD: More than simply a motor deficit. *Movement Disorders*, 27(3), 387-392.
- 470Mohawk, J. A., Green, C. B., & Takahashi, J. S. (2012). Central and peripheral circadian clocks
471 in mammals. *Annual Review of Neuroscience*, 35, 445-462.

- 472Montero-Odasso, M., Muir, S. W., & Speechley, M. (2012). Dual-task complexity affects gait in
473 people with mild cognitive impairment: The interplay between gait variability, dual tasking,
474 and risk of falls. *Archives of Physical Medicine and Rehabilitation*, 93(2), 293-299.
- 475Newell, K. M., & Corcos, D. M. (Eds.). (1993). *Variability and motor control*. Champaign IL:
476 Human Kinetics Publishers.
- 477Orsucci, F. F. (2006). The paradigm of complexity in clinical neurocognitive science. *The*
478 *Neuroscientist : A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 12(5),
479 390-397. doi:12/5/390 [pii]
- 480Paterson, K., Hill, K., & Lythgo, N. (2011). Stride dynamics, gait variability and prospective
481 falls risk in active community dwelling older women. *Gait & Posture*, 33(2), 251-255.
- 482Peng, C. K., Mietus, J. E., Liu, Y. H., Lee, C., Hausdorff, J. M., Stanley, H. E., . . . Lipsitz, L. A.
483 (2002). Quantifying fractal dynamics of human respiration: Age and gender effects. *Annals*
484 *of Biomedical Engineering*, 30(5), 683-692. doi:10.1114/1.1481053
- 485Roerdink, M., Lamothe, C. J., van Kordelaar, J., Elich, P., Konijnenbelt, M., Kwakkel, G., &
486 Beek, P. J. (2009). Rhythm perturbations in acoustically paced treadmill walking after
487 stroke. *Neurorehabilitation and Neural Repair*, 23(7), 668-678.
488 doi:10.1177/1545968309332879 [doi]
- 489Schmidt, R. A., & Lee, T. D. (2005). *Motor control and learning: A behavioral emphasis* .
490 Champagne, IL.: Human Kinetics.
- 491Seidler, R. D., Bernard, J. A., Burutolu, T. B., Fling, B. W., Gordon, M. T., Gwin, J. T., . . . Lipps,
492 D. B. (2010). Motor control and aging: Links to age-related brain structural, functional, and
493 biochemical effects. *Neuroscience & Biobehavioral Reviews*, 34(5), 721-733.

- 494Shin, S., Valentine, R. J., Evans, E. M., & Sosnoff, J. J. (2012). Lower extremity muscle quality
495 and gait variability in older adults. *Age and Ageing*, 41(5), 595-599.
496 doi:10.1093/ageing/afs032 [doi]
- 497Skoe, E., & Kraus, N. (2009). Hearing it again and again: On-line subcortical plasticity in
498 humans. *PloS One*, 5(10), e13645.
- 499Slutzky, M. W., Cvitanovic, P., & Mogul, D. J. (2001). Deterministic chaos and noise in three in
500 vitro hippocampal models of epilepsy. *Annals of Biomedical Engineering*, 29(7), 607-618.
- 501Snyder, J. S. (2003). Tapping to bach: Resonance-based modeling of pulse. *Music Perception*,
502 21(1), 43-80.
- 503Sosnoff, J. J., & Newell, K. M. (2008). Age-related loss of adaptability to fast time scales in
504 motor variability. *The Journals of Gerontology.Series B, Psychological Sciences and Social*
505 *Sciences*, 63(6), 344-352. doi:63/6/P344 [pii]
- 506Stergiou, N., & Decker, L. M. (2011). Human movement variability, nonlinear dynamics, and
507 pathology: Is there a connection? *Human Movement Science*, 30(5), 869-888.
508 doi:10.1016/j.humov.2011.06.002 [doi]
- 509Stergiou, N., Harbourne, R., & Cavanaugh, J. (2006). Optimal movement variability: A new
510 theoretical perspective for neurologic physical therapy. *Journal of Neurologic Physical*
511 *Therapy : JNPT*, 30(3), 120-129.
- 512Stergiou, N. (Ed.). (2004). *Innovative analyses of human movement*. Champaign, IL: Human
513 Kinetics.
- 514Stergiou, N. (Ed.). (2016). *Nonlinear analysis for human movement variability* CRC Press.

- 515 Toebe, M. J. P., Hoozemans, M. J. M., Furrer, R., Dekker, J., & van Dieën, J. H. (2012). Local
- 516 dynamic stability and variability of gait are associated with fall history in elderly subjects.
- 517 *Gait & Posture*, 36(3), 527-531.
- 518 Toweill, D. L., & Goldstein, B. (1998). Linear and nonlinear dynamics and the pathophysiology
- 519 of shock. *New Horizons (Baltimore, Md.)*, 6(2), 155-168.
- 520 Tranah, G. J., Blackwell, T., Ancoli-Israel, S., Paudel, M. L., Ensrud, K. E., Cauley, J. A., . . .
- 521 Stone, K. L. (2010). Circadian activity rhythms and mortality: The study of osteoporotic
- 522 fractures. *Journal of the American Geriatrics Society*, 58(2), 282-291.
- 523 Tranah, G. J., Blackwell, T., Stone, K. L., Ancoli-Israel, S., Paudel, M. L., Ensrud, K. E., . . .
- 524 Cummings, S. R. (2011). Circadian activity rhythms and risk of incident dementia and mild
- 525 cognitive impairment in older women. *Annals of Neurology*, 70(5), 722-732.
- 526 Vaillancourt, D. E., & Newell, K. M. (2003). Aging and the time and frequency structure of force
- 527 output variability. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 94(3), 903-912.
- 528 doi:10.1152/jappphysiol.00166.2002 [doi]
- 529 Vaillancourt, D. E., Sosnoff, J. J., & Newell, K. M. (2004). Age-related changes in complexity
- 530 depend on task dynamics. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 97(1), 454-
- 531 455. doi:10.1152/jappphysiol.00244.2004 [doi]
- 532 Vaillancourt, D. E., & Newell, K. M. (2002). Changing complexity in human behavior and
- 533 physiology through aging and disease. *Neurobiology of Aging*, 23(1), 1-11.
- 534 Van Orden, G. C., Holden, J. G., & Turvey, M. T. (2005). Human cognition and 1/f scaling.
- 535 *Journal of Experimental Psychology: General*, 134(1), 117-123.
- 536 Vinti, M., Couillandre, A., & Thoumie, P. (2010). Does somatosensory loss induce adaptation of
- 537 the gait initiation process? *Neuroscience Letters*, 480(3), 178-181.

- 538Wagner, C. D., Nafz, B., & Persson, P. B. (1996). Chaos in blood pressure control.
539 *Cardiovascular Research*, 31(3), 380-387. doi:0008636396000077 [pii]
- 540Wojcik, L. A., Thelen, D. G., Schultz, A. B., Ashton-Miller, J. A., & Alexander, N. B. (1999).
541 Age and gender differences in single-step recovery from a forward fall. *The Journals of*
542 *Gerontology Series A: Biological Sciences and Medical Sciences*, 54(1), M44-M50.
- 543Yamazaki, S., Straume, M., Tei, H., Sakaki, Y., Menaker, M., & Block, G. D. (2002). Effects of
544 aging on central and peripheral mammalian clocks. *Proceedings of the National Academy of*
545 *Sciences of the United States of America*, 99(16), 10801-10806.
546 doi:10.1073/pnas.152318499 [doi]
- 547Yang, A. C., Huang, C., Yeh, H., Liu, M., Hong, C., Tu, P., . . . Lin, C. (2013). Complexity of
548 spontaneous BOLD activity in default mode network is correlated with cognitive function in
549 normal male elderly: A multiscale entropy analysis. *Neurobiology of Aging*, 34(2), 428-438.
- 550Yang, X. (2010). A wheel of time: The circadian clock, nuclear receptors, and physiology. *Genes*
551 *& Development*, 24(8), 741-747. doi:10.1101/gad.1920710 [doi]
- 552
- 553
- 554

555**Figures**

556**Figure 1.** Optimal movement variability is represented in the middle of the figure as an inverted

557U-shape relationship between complexity and predictability (N. Stergiou et al., 2006).

558Practically, at this optimal state of movement variability the biological system is in a healthy

559state and is characterized by the largest possible effective complexity (i.e., the uppermost point

560along the inverted U-shaped function), attaining high values only in the intermediate region

561between excessive order (i.e. maximum predictability) and excessive disorder (i.e. no

562predictability). However, complexity is affected by the dominant dynamics across scales. Thus,


563the theory of Optimal Movement Variability could actually be generalized into a theory of

564Complex Adaptive Behavior where complexity (and higher dexterity) is associated with fractally

565nested scales of activity supported by non-linear couplings of components both within and across

566scales (Harrison & Stergiou, 2015).

VARIABILITY AND AGING



567

568

569

570