

# Motivic Zeta Function

Maximilian Rath

Geboren am 2. Januar 1995 in Reutlingen

2.2.2016

Bachelorarbeit Mathematik

Prof. Dr. Daniel Huybrechts

Martin Ulirsch

MATHEMATISCHES INSTITUT

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER  
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN



# 1 Introduction

In 1949 André Weil, in the paper [Wei49], formulated his famous conjectures which predicted certain deep properties of the Hasse–Weil zeta function for a variety  $X$  over a finite field  $k = \mathbb{F}_q$ . The zeta function  $Z(X, t) \in \mathbb{Q}[[t]]$  is defined as

$$Z(X, t) = \exp \left( \sum_{m=0}^{\infty} \frac{|X(\mathbb{F}_{q^m})|}{m} t^m \right).$$

Dwork first established the rationality of the zeta function, the most basic of the three conjectures. Later, using his new scheme theoretic approach to algebraic geometry, Grothendieck gave another proof for the special case of smooth projective varieties.

In [Kap00] Kapranov generalized the Hasse–Weil zeta function, defining it not only for varieties over finite fields but varieties in general. The construction takes place in the so called *Grothendieck ring of varieties*  $K_0[\mathcal{V}_k]$  over some field  $k$ , which consists of isomorphism classes of varieties subject to some additional relations (for the complete definition see TODO), and is given as

$$Z_{\mu}(X, t) := \sum_{n=0}^{\infty} \mu([\mathrm{Sym}^n(X)]) t^n$$

where  $\mu$  is a ring homomorphism from  $K_0[\mathcal{V}_k]$  to some arbitrary ring (also called a *motivic measure*). It is a natural question to ask if this zeta function is still a rational function (at least if  $\mu$  takes values in a field). Kapranov was able to prove this for smooth projective curves ([Kap00][TODO]) and conjectured that it is true in general. But it turns out that in this more general setting the zeta function is not as well behaved in terms of rationality as in the classical case: Larsen and Lunts show in [LL03] that the zeta function even fails to be rational for surfaces when working over  $\mathbb{C}$ .

**Theorem 1.1** ([LL03, Thm. 1.6]). *Assume that  $k = \mathbb{C}$ . There exists a field  $\mathcal{H}$  and a motivic measure  $\mu_h: K_0[\mathcal{V}_k] \rightarrow \mathcal{H}$  with the following property: if  $X$  is a smooth complex projective surface such that the geometric genus  $P_g(X) = h^{2,0}(X)$  is at least two, then the zeta-function  $\zeta_{\mu_h}(X, t)$  is not a rational function.*

In this thesis we will first introduce the Grothendieck ring in Section 2 and investigate its structure. This section culminates in a structure theorem due to Bittner ([Bit04]) which will allow us to simplify the original proof by Larsen and Lunts substantially. The second section is devoted to the definition of the motivic zeta function, which entails defining and giving examples for the symmetric product. There we will also see the connection between the classical Hasse–Weil zeta function and the motivic one. Construction of the motivic measure  $\mu_h$  under which  $Z_{\mu_h}(X, t)$  becomes irrational is content of Section TODO while irrationality is then finally proven in the last section.

## 2 The Grothendieck Ring of Varieties and Motivic Measures

**Definiton 2.1.** In the following the term *k-variety* always means a separated, reduced scheme of finite type over a field  $k$ . We will write  $\mathcal{V}_k$  for the category of  $k$ -varieties.

**Definiton 2.2.** Let  $k$  be a perfect field. Consider the abelian group of formal linear combinations of isomorphism classes of varieties, subject to relations of the form

$$[X \setminus Y] = [X] - [Y]$$

where  $Y$  is closed in  $X$ . With multiplication given by

$$[X][Y] = [X \times Y].$$

This forms a ring, called the *Grothendieck ring of varieties* and denoted by  $K_0[\mathcal{V}_k]$ .

A *motivic measure* with values in a ring  $A$  is a ring homomorphism  $\mu: K_0[\mathcal{V}_k] \rightarrow A$ . The identity function  $\text{id}: K_0[\mathcal{V}_k] \rightarrow K_0[\mathcal{V}_k]$  is called the *universal motivic measure*.

The reason that we require  $k$  to be perfect is that for nonperfect  $k$  the product of two varieties need not be reduced. Take for example  $k = k_0(t)$  with  $\text{char } k = p$  and  $X = \text{Spec}(k[x]/(x^p - t))$ ,  $Y := \text{Spec}(K) := \text{Spec}(k(t^{\frac{1}{p}}))$ .  $X$  is a variety since  $x^p - t$  is irreducible, hence  $X$  is reduced. But the product can be described as

$$X \times Y = \text{Spec}(K[x]/(x^p - t)) = \text{Spec}(K[X]/((x - t^{\frac{1}{p}})^p).$$

And hence is not reduced. We might adapt the definition to this case by changing the definition of the product to  $[X][Y] := [(X \times Y)_{\text{red}}]$ , taking the induced reduced scheme structure on  $X \times Y$ , but since we are only working in characteristic zero or over finite fields we will not bother with this technical issue any more.

Since the product is commutative up to isomorphism this is a commutative ring with 1 (equal to  $[\text{Spec}(k)]$ ). The cut and paste relation furthermore gives us

$$0 = [\emptyset] - [\emptyset] = [\emptyset \setminus \emptyset] = [\emptyset]$$

**Example 2.3.** For a finite field  $k$ , consider the function <sup>1</sup>

$$\begin{aligned} \psi: \mathcal{V}_k &\rightarrow \mathbb{Z} \\ X &\mapsto |X(k)| \end{aligned}$$

---

<sup>1</sup>not bothering with any set-theoretic issues

1. If  $X$  and  $Y$  are isomorphic  $k$ -varieties, then  $\psi(X) = \psi(Y)$ .
2. If  $Y \subset X$  is closed subvariety, then  $\psi(X \setminus Y) = \psi(X) - \psi(Y)$ .
3. If  $X$  and  $Y$  are two varieties, then  $\psi(X \times Y) = \psi(X)\psi(Y)$ .

Hence  $\psi$  actually factors over  $K_0[\mathcal{V}_k]$  and gives rise to a motivic measure.

**Example 2.4.** Using the decomposition  $\mathbb{P}_k^n = \mathbb{P}_k^{n-1} \amalg \mathbb{A}_k^n$  where  $\mathbb{P}_k^{n-1}$  is a (closed) hyperplane in  $\mathbb{P}_k^n$  we get  $[\mathbb{P}_k^n] = [\mathbb{P}_k^{n-1}] + [\mathbb{A}_k^n]$  in the Grothendieck ring. Inductively this yields the identity

$$[\mathbb{P}_k^n] = \sum_{m=0}^n [\mathbb{A}_k^1]^m$$

We also denote the isomorphism class of the affine line as  $\mathbb{L}$ .

This calculation also gives rise to an example of two irreducible varieties that have the same equivalence class in  $K_0[\mathcal{V}_k]$  but are not isomorphic:  $\mathbb{P}^n \times (\mathbb{A}^1 \setminus \{0\})$  and  $\mathbb{A}^{n+1}$ . These are isomorphic since the global section of the former are  $\text{TODO}$ , but in the Grothendieck ring we get

$$\begin{aligned} [\mathbb{P}^n \times (\mathbb{A}^1 \setminus \{0\})] &= [\mathbb{P}^n] ([\mathbb{A}^1] - 1) \\ &= \sum_{m=0}^n \mathbb{L}^{m+1} - \sum_{m=0}^n \mathbb{L}^m \\ &= \mathbb{L}^{n+1} \end{aligned}$$

**Remark 2.5.** By [Har77, Prop. 10.1 (d)] the product of two smooth varieties over  $k$  is again smooth. Hence the isomorphism classes of smooth irreducible complete varieties form a multiplicative monoid, in the following denoted by  $\mathcal{M}$ .

We now investigate the structure of the Grothendieck ring. First we note that it suffices to take irreducible varieties to generate  $K_0[\mathcal{V}_k]$ . To see this, take a variety  $X$  with irreducible components  $X_1, \dots, X_n$  and set  $U_i := X_i \setminus \bigcup_{j \neq i} X_j$  and  $U := \bigcup_i U_i$ . By construction, the last union is actually a disjoint union, and hence  $[U] = \sum_i [U_i]$ . Together this gives

$$[X] = [X \setminus U] + \sum_i [U_i].$$

Now the  $U_i$  are irreducible as open subsets of irreducible sets, and  $[X \setminus U]$  has dimension strictly smaller than the dimension of  $X$  since we removed a proper open subset from each irreducible component. Hence we can inductively rewrite  $[X \setminus U]$ , losing at least one dimension in each step, with the base case of  $\dim X = 0$  being trivial, as  $X$  is then just a collection of points.

In characteristic zero we can use a weak form of Hironaka's Theorem on the resolution of singularities, namely that for every irreducible, projective variety  $X$  there is a smooth, projective variety that is birational to  $X$  to restrict the set of needed generators even more. This observation culminates in a structure theorem about the Grothendieck ring proven by Bittner in [Bit04]

**Theorem 2.6** (Bittner, [Bit04, Thm. 3.1]). *Let  $k$  be a field of characteristic zero. Then  $K_0[\mathcal{V}_k]$  is generated by smooth, integral, projective varieties. Furthermore, if  $k$  is algebraically closed, it suffices to consider relations of the form*

$$[X] - [f^{-1}(Z)] = [Y] - [Z]$$

where  $X, Y$  are smooth complete varieties and  $f: X \rightarrow Y$  is a morphism which is a blowup with a smooth center  $Z \subset Y$ .

*Proof.* We will only proof the first part of the statement. As remarked before, we can restrict our attention to irreducible varieties. We once again argue by induction on the dimension. So let  $X$  be an irreducible variety of dimension  $n$ , by choosing some nonempty affine subvariety and passing to its projective closure we find  $X'$  projective, irreducible and birational to  $X$ . Resolving singularities we find  $\tilde{X}$  smooth, projective, irreducible birational to  $X$ . Hence we find isomorphic open subset  $U \subset X$ ,  $V \subset \tilde{X}$  and hence

$$[X \setminus U] - [\tilde{X} \setminus V] = [X] - [U] - [\tilde{X}] + [V] = [X] - [\tilde{X}].$$

Thus  $[X]$  can be written as the sum of  $[\tilde{X}]$  and some varieties of lower dimension for which we can invoke the induction hypothesis.  $\square$

We will use the slightly stronger conclusion we just shown, hence we state it as a separate corollary.

**Corollary 2.7.** *For each (irreducible) variety  $X$  there exists a smooth, irreducible, projective variety  $Y$  birational to  $X$  such that  $[X] = [Y] + \sum_i^N m_i [W_i]$  where the  $[W_i]$  are smooth, irreducible, projective varieties of dimension strictly smaller than the dimension of  $X$ .*

### 3 Symmetric Products and the Motivic Zeta Function

We now want to make the concept of the symmetric product precise, i.e. how it can be defined as the quotient of the (regular) product by the action of the symmetric group.

**Definition 3.1.** Let  $X$  be a  $S$ -Scheme,  $G$  a group acting on  $X$  via  $S$ -automorphisms (i.e there is a group homomorphism  $\sigma: G \rightarrow \text{Aut}_S(X)$ ), we

also write  $\sigma_g$  for the image of  $g$  under  $\sigma$ .) Then a variety  $Y$  together with a morphism  $\pi: X \rightarrow Y$  over  $S$  is called the quotient (also denoted by  $X/G$ ) of  $Y$  by  $G$  if  $\pi$  is  $G$ -invariant for all  $g \in G$   $\pi \circ g = \pi$  and is universal in this respect, i.e. for every other  $G$ -invariant morphism  $f: X \rightarrow Z$  there exists a unique  $G$ -invariant morphism  $g: Z \rightarrow Y$  that makes the following diagram commute

$$\begin{array}{ccc} X & \xrightarrow{f} & X/G \\ \downarrow \pi & \nearrow g & \\ Z & & \end{array}$$

We first construct the quotient in the affine case. Assume  $X = \text{Spec}(A)$  with  $G$  acting on  $X$  (from the right) via automorphisms. By the equivalence of categories between affine schemes and commutative rings, we get a left-action of  $G$  on  $A$ , and we simply get the quotient as the affine scheme  $X \rightarrow X/G = \text{Spec}(A^G)$  coming from the subring of  $G$ -invariant elements, together with the natural inclusion  $A^G \rightarrow A$ . By construction this is the categorical quotient defined above.

We now turn to a relevant special case: the quotient of an quasiprojective variety  $X$  by the action of a finite group  $G$ , using the following abstract result from [Gro71]

**Lemma 3.2** ([Gro71][Prop. 1.8]). *Let  $X$  be a scheme with a finite group  $G$  acting on it via automorphisms. If there exists an open cover by  $G$ -invariant affine sets then the quotient  $X/G$  exists. We call such an action admissible*

**Corollary 3.3.** *If  $X$  is quasiprojective, then the quotient by the action of a finite group always exists.*

*Proof.* Let  $X \subset \mathbb{P}_k^n$  be locally closed with a finite group  $G = \{g_1, \dots, g_m\}$  acting on it. Consider the orbit  $\{g_1(x), \dots, g_m(x)\} \subset X$  of a point  $x \in X$ , and denote by  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  the corresponding graded prime ideals corresponding to  $\{x\}$ , and by  $\mathfrak{q}$  the graded prime ideal corresponding to  $\bar{X} \setminus X$  (taken to be the irrelevant ideal if this set is empty). Hence the ideal  $\mathfrak{q}$  is not contained in any of the  $\mathfrak{p}_i$ s hence, by the graded version of the prime avoidance lemma we find an element with positive degree  $y \in \mathfrak{q}$  that is contained in none of the  $\mathfrak{p}_i$  and hence the hypersurface  $V_+(y)$  contains  $\bar{X} \setminus X$  but not the  $g_1(x), \dots, g_m(x)$ , so  $U := D_+(y)$  is an affine set that lies in  $X$  and contains the orbit of  $x$  under  $G$ . Now consider  $U' := \bigcap_{g \in G} g(U)$ . This is clearly  $G$ -invariant, and since  $U$  contained the orbit of  $y$  we still find that  $y$  lies in  $U'$ . Hence we have found a open cover by  $G$ -invariant affine sets, and thus the quotient exists by the above lemma.  $\square$

**Remark 3.4.** If  $X = \text{Proj}(S)$  and the  $G$ -action is a *graded* action (i.e. the induced action on  $S$  respects the grading, we may construct the quotient, like in the affine case, as  $X/G = \text{Proj}(S^G)$

**Definiton 3.5.** Let  $X$  be a quasiprojective variety, then the  $n$ -fold *symmetric product*  $\text{Sym}^n(X)$  is defined as the quotient of  $X^n$  (which is again quasiprojective via the Segre-Embedding) by the natural action of the symmetric group in  $n$  letters.

**Example 3.6.**

**Definiton 3.7.** Let  $\mu: K_0[\mathcal{V}_k] \rightarrow A$  be a motivic measure and  $X$  a variety. Then we define the *motivic zeta function*  $\zeta_\mu(X, t) \in A[[t]]$  as

$$\zeta_\mu(X, t) := \sum_{n=0}^{\infty} \mu([\text{Sym}^n(X)])t^n$$

**Example 3.8.** We can now compute a first zeta function, namely  $\zeta_{id}(\mathbb{P}_k^1, t)$ . Using the identity and notation from example 2.4 we calculate

$$\begin{aligned} \zeta_{id}(\mathbb{P}_k^1, t) &= \sum_{n=0}^{\infty} [\text{Sym}^n(\mathbb{P}_k^1)]t^n = \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \mathbb{L}^k \right) t^n \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^n (\mathbb{L}t)^k t^{n-k} \\ &= \left( \sum_{n=0}^{\infty} t^n \right) \left( \sum_{n=0}^{\infty} (\mathbb{L}t)^n \right) = \frac{1}{(1-t)(1-\mathbb{L}t)}. \end{aligned}$$

Hence  $\zeta_{id}(\mathbb{P}_k^1, t)$  is in fact a rational function.

## 4 Constructing Motivic Measures

We will now construct a measure  $\mu_h$  for which we will show irrationality of the zeta function.

The first important result on the way is a lemma that helps us to construct such measures by extending maps from the monoid of smooth, irreducible and complete varieties. For this we quickly recall the notion of a monoid ring.

**Definiton 4.1.** Let  $G$  be a (multiplicative) monoid. Write  $\mathbb{Z}[G]$  for the free abelian group over  $G$ , i.e. the set of formal sums  $\sum_{g \in G} n_g g$  with the  $m_g$  being integers, only finitely many non zero. Defining multiplication as

$$\left( \sum_{g \in G} n_g g \right) \left( \sum_{g \in G} m_g g \right) = \left( \sum_{g \in G} \left( \sum_{hl=g} n_h m_l \right) g \right)$$

this forms a ring: the *monoid ring* over  $G$ .



**Theorem 4.2.** *Set  $k = \mathbb{C}$ . Let  $G$  be an abelian monoid and  $\mathbb{Z}[G]$  the corresponding monoid ring. As above, denote by  $\mathcal{M}$  the multiplicative monoid of irreducible smooth complete varieties. Let*

$$\psi: \mathcal{M} \rightarrow G$$

*be a homomorphism of monoids such that*

1.  $\psi([X]) = \psi([Y])$  if  $X$  and  $Y$  are birational;
2.  $\psi([\mathbb{P}^n]) = 1$  for all  $n \geq 0$ .

*Then  $\psi$  can be uniquely extended to a ring homomorphism*

$$\phi: K_0[\mathcal{V}_{\mathbb{C}}] \rightarrow \mathbb{Z}[G]$$

*Proof.* By 2.6 we know that the Grothendieck ring (as an abelian group) is generated by  $\mathcal{M}$ , so after extending  $\psi$  to the free abelian group over  $\mathcal{M}$  we have to check that  $\psi$  preserves the relations of the blowup presentation of  $K_0[\mathcal{V}_{\mathbb{C}}]$  and hence factors over it, i.e. given a blow-up  $f: X \rightarrow Y$  with smooth center  $Z \in X$  then  $\psi([X]) - \psi([f^{-1}(Z)]) = \psi([Y]) - \psi([Z])$ . Note that  $[X]$  and  $[Y]$  are birational since  $f$  is a blowup.

Now since  $X$  and  $Z$  are nonsingular the exceptional divisor  $f^{-1}(Z)$  is isomorphic to the projective space bundle  $\mathbb{P}(\mathcal{I}/\mathcal{I}^2)$  where  $\mathcal{I}$  is the ideal sheaf associated to  $Z$  (see for example [Har77, II.8, Thm 8.24]), which is a  $\mathbb{P}^n$ -bundle, i.e. locally trivializes to  $\mathbb{P}_k^n \times U$  for some open set  $U \subset Z$  and hence is birational to  $\mathbb{P}_k^n \times Z$ . We can thus write  $\psi([f^{-1}(Z)])$  as

$$\psi([f^{-1}(Z)]) = \psi([Z \times \mathbb{P}^n]) = \psi([Z][\mathbb{P}^n]) = \psi([Z])\psi([\mathbb{P}^n]) = \psi([Z])$$

Hence we can linearly extend  $\psi$  to define a morphism  $\phi: K_0[\mathcal{V}_{\mathbb{C}}] \rightarrow \mathbb{Z}[G]$   $\square$

Before constructing a suitable monoid homomorphism that will extend to our desired measure, we recall the definition of the Hodge numbers of a variety as well as two basic results (all these are exercises in [Har77, Ch. II]).

**Definiton 4.3.** Let  $X$  be a projective variety. The  $i$ -th *Hodge number*  $h^{i,0}$  is given as the dimension of  $H^0(X, \Omega_X^i)$ , which is a finite dimensional  $k$ -vector space by a theorem of Serre ([Ser55, §3 Prop. 7]). We will also write  $P_g(X)$  for  $h^{d,0}(X)$ , called the geometric genus of a nonsingular variety  $X$ , with  $d$  being the dimension of  $X$ .

**Lemma 4.4** ([Har77, II Ex. 8.8]). *The Hodge numbers  $h^{i,0}$  are a birational invariant of a variety: let  $X, Y$  be birationally equivalent, smooth, projective varieties then  $h^{i,0}(X) = h^{i,0}(Y)$ .*

**Lemma 4.5** (Chow, [Har77, II Ex. 4.10]). *Every proper variety is birational to a projective variety.*

**Definiton 4.6.** Denote by  $C \subset \mathbb{Z}[t]$  the multiplicative monoid of polynomials with positive leading coefficient. For a smooth projective complex variety  $Z$  of dimension  $d$  define

$$\Psi_h(X) := 1 + h^{1,0}(X)t + \cdots + h^{d,0}(X)t^d \in C.$$

(Note that the Hodge numbers might be zero, but  $\Psi_h(X)$  still has a positive leading coefficient, just its degree might be smaller than the dimension of  $Z$ ) By Chow's lemma we can also define  $\Psi_h$  for a smooth complete variety  $Z$  by choosing a smooth projective variety  $X$  which is birational to  $Z$  and setting  $\Psi_h(Z) := \Psi_h(X)$ . This is well defined by Lemma 4.4.

We now check that  $\Psi_h$  satisfies all conditions of Theorem 4.2. Independence of birational equivalence class was the content of Lemma 4.4.

To check multiplicativity, we use the following Lemma.

**Lemma 4.7.** *Let  $X, Y$  be smooth, irreducible, projective  $k$ -varieties. Then the following equality holds:*

$$h^{p,0}(X \times_k Y) = \sum_{i+j=p} h^{i,0}(X)h^{j,0}(Y)$$

*Proof.* TODO □

With this we directly calculate

$$\begin{aligned} \Psi_h(X)\Psi_h(Y) &= \left( \sum_n h^{i,0}(X)t^i \right) \left( \sum_n h^{j,0}(Y)t^j \right) = \sum_n \left( \sum_{i+j=n} h^{i,0}(X)h^{j,0}(Y) \right) t^n \\ &= \sum_n h^{n,0}(X \times_k Y) = \Psi_h(X \times_k Y) \end{aligned}$$

## 5 Irrationality

The constructed motivic measure does not yet yield values in a field, so we would like to pass to the quotient field of  $\mathbb{Z}[\mathcal{M}]$ . That we are able to do so is the content of the next lemma.

**Lemma 5.1.** *Let  $A$  be a factorial ring, and  $S \subset A$  a multiplicative submonoid such that 1 is the only unit in  $S$ . Then the monoid ring  $\mathbb{Z}[S]$  is a polynomial ring (in possibly infinitely many variables), and hence an integral domain.*

*Proof.* Since  $A$  is factorial, every  $s \in S$  has a unique factorization, and since 1 is the only unit in  $S$ ,  $s$  can be uniquely written as product of prime elements, hence if we take  $B$  to be the polynomial ring over the formal variables  $\{x_s | s \in S, s \text{ prime}\}$  we get an isomorphism of rings

$$\begin{aligned} B &\rightarrow \mathbb{Z}[S] \\ x_s &\mapsto s \end{aligned}$$

□

**Definiton 5.2.** Denote by  $\mathcal{H}$  the quotient field of  $\mathbb{Z}[C]$  where, as above,  $C$  denotes the submonoid of  $\mathbb{Z}[t]$  consisting of polynomials with positive leading coefficients. Since  $-1$  is not contained in  $C$  it satisfies the conditions of the previous lemma. We define the motivic measure  $\mu_h: K_0[\mathcal{V}_k] \rightarrow \mathcal{H}$  as the measure obtained extending  $\Psi_h$  as by Theorem 4.2

**Lemma 5.3.** *Let  $Y_1, \dots, Y_s, Z$  be irreducible varieties of dimension  $d$  over a field of characteristic zero such that  $\mu_h([Z]) = \sum_i n_i \mu_h([Y_i])$  for some  $n_i \in \mathbb{Z}$  and  $P_g(Z) \neq 0$  then  $P_g(Z) = P_g(Y_i)$  for some  $i$*

*Proof.* By Corollary 2.7, we find smooth, irreducible projective varieties  $\overline{Z}, \overline{Y}_1, \dots, \overline{Y}_s$  in the same birational class as the original varieties allowing us to rewrite the original equality as

$$\mu_h([\overline{Z}]) = \sum n_i \mu_h([\overline{Y}_i]) + \sum l_i \mu_h([X_i])$$

the  $X_i$  being smooth irreducible varieties of dimension  $< d$ . But now, since  $\mu_h$  was obtained as an extension of  $\Psi_h$  which was defined on smooth, irreducible, projective varieties, this is actually an equation in  $\mathbb{Z}[C]$ , namely

$$\Psi_h(\overline{Z}) = \sum n_i \Psi_h(\overline{Y}_i) + \sum l_i \Psi_h(X_i)$$

Since there are no (additive) relations between elements of  $C$  in the monoid ring, one of the polynomials  $\Psi_h(\dots)$  on the left hand side, must actually be the polynomial on the left hand side, but with the dimension of  $X_i$  being strictly smaller than that of  $Z$  and the assumption that  $P_g(\overline{Z}) = h^{d,0}(\overline{Z}) \neq 0$  it cannot be one of the  $\Psi_h(X_i)$  because all these polynomials have strictly smaller degree. Hence we have  $\Psi_h(\overline{Z}) = \Psi_h(\overline{Y}_i)$  for some  $i$  and in particular, since they are of the same dimension, the genus of  $\overline{Z}$  and  $Y_i$  must agree. □

Now let  $X$  be a smooth projective surface with  $P_g(X) \geq 2$ . We will show that  $\zeta_{\mu_h}(X, t) \in \mathcal{H}[[t]]$  is not rational, thus proving Theorem 1.1. We need one technical lemma about the genus of symmetric powers of  $X$ , which we cannot prove here.

**Lemma 5.4** ([LL03, Lem. 3.8]). *Let  $X$  be a smooth, projective surface over  $\mathbb{C}$ . Then*

$$P_g(\text{Sym}^n(X)) = \binom{P_g(X) + n - 1}{P_g(X) - 1}$$

We will use the following rationality criterion for power series.

**Lemma 5.5.** *Let  $k$  be a field and suppose the power series  $\sum a_i t^i \in k[[t]]$  is a rational function (i.e. an element in  $k(t)$ ) then there exist  $n, n_0 > 0$  such that for each  $m > n_0$  the determinant*

$$\begin{vmatrix} a_m & a_{m+1} & \cdots & a_{m+n} \\ a_{m+1} & a_{m+2} & \cdots & a_{m+n+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m+n} & a_{m+n+1} & \cdots & a_{m+2n} \end{vmatrix}$$

*vanishes.*

*Proof.* Denote the above Matrix by  $A$ . Since  $\sum a_i t^i$  is rational, we may write it as

$$\sum a_i t^i = \frac{p_0 + \cdots + p_e t^e}{q_0 + \cdots + q_d t^d}$$

with not all  $q_i$  being zero. Hence  $p_0 + \cdots + p_e t^e = (\sum a_i t^i) (q_0 + \cdots + q_d t^d) = \sum_{i=0}^{\infty} (\sum_{k+l=i} a_k q_l) t^i$ . This means that for  $m > e$  we get  $0 = \sum_{k+l=m} a_k q_l$ . now note that  $q_l = 0$  for  $l > n$  hence we may rewrite this as

$$0 = \sum_{l=0}^n q_l a_{m-l} = \sum_{l=0}^n q_{n-l} a_{m-n+l}.$$

Thus we have shown that

$$\begin{pmatrix} a_m & a_{m+1} & \cdots & a_{m+n} \\ a_{m+1} & a_{m+2} & \cdots & a_{m+n+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m+n} & a_{m+n+1} & \cdots & a_{m+2n} \end{pmatrix} \begin{pmatrix} q_n \\ \vdots \\ q_1 \\ q_0 \end{pmatrix} = \begin{pmatrix} \sum_{l=0} a_{m+l} q_{n-l} \\ \vdots \\ \sum_{l=0} a_{m+n+l} q_{n-l} \end{pmatrix} = 0$$

for  $m$  big enough. Since one of the  $q_l$  is non zero this means that  $A$  has nonempty kernel and hence its determinant vanishes.  $\square$

*Proof of Theorem 1.1.* Assume that  $\zeta_{\mu_h}(X, t)$  is rational, hence by the above criterion there is a  $n$  such that for  $m$  big enough

$$\begin{aligned} & \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) \mu_h \left( \prod_{i=1}^{n+1} \text{Sym}^{m+i+\sigma(i)-2}(X) \right) = 0 \\ \Leftrightarrow & \mu_h \left( \prod_{i=0}^n \text{Sym}^{m+2i}(X) \right) = - \sum_{\substack{\sigma \in S_{n+1} \\ \sigma \neq id}} \text{sgn}(\sigma) \mu_h \left( \prod_{i=0}^n \text{Sym}^{m+i+\sigma(i+1)-1}(X) \right) \end{aligned}$$

Now we can apply Lemma 5.3 to conclude that there is a permutation  $\sigma$ , not the identity permutation, such that

$$P_g\left(\prod_{i=0}^n \mathrm{Sym}^{m+2i}(X)\right) = P_g\left(\prod_{i=0}^n \mathrm{Sym}^{m+i+\sigma(i+1)-1}(X)\right)$$

By Lemma 5.4 (and using the fact that the genus is multiplicative), we get

$$\prod_{i=0}^n \binom{P_g(X) + m + 2i - 1}{P_g(X) - 1} - \prod_{i=0}^n \binom{P_g(X) + m + i + \sigma(i+1) - 2}{P_g(X) - 1} = 0$$

By assumption  $P_g(X) \geq 2$  hence the left hand side, considered as a polynomial in  $m$ , is not the zero polynomial. So by taking  $m$  large enough we obtain a contradiction.  $\square$

## References

- [Bit04] Franziska Bittner. The universal Euler characteristic for varieties of characteristic zero. *Compos. Math.*, 140(4):1011–1032, 2004.
- [Gro71] Alexander Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*, volume 224 of *Lecture notes in mathematics*. Springer-Verlag, 1971.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Kap00] Mikhail Kapranov. The elliptic curve in the S-duality theory and Eisenstein series for Kac-Moody groups, 2000.
- [LL03] Michael Larsen and Valery A. Lunts. Motivic measures and stable birational geometry. *Mosc. Math. J.*, 3(1):85–95, 259, 2003.
- [Ser55] Jean-Pierre Serre. Faisceaux algébriques cohérents. *Ann. of Math. (2)*, 61:197–278, 1955.
- [Wei49] André Weil. Numbers of solutions of equations in finite fields. *Bull. Amer. Math. Soc.*, 55(5):497–508, 05 1949.