

Contents

1 Categories	2
1-1	2
1-2	5
1-3	7
1-4	9
1-5	10
1-6	12
1-7	13
1-8	15
1-9	17
1-10	20
1-11	22
1-12	26
1-13	29
1-14	30

1 Categories

Problem 1-1: The objects of **Rel** are sets, and an arrow $A \rightarrow B$ is a relation from A to B , that is, a subset $R \subseteq A \times B$. The equality relation $\{\langle a, a \rangle \in A \times A \mid a \in A\}$ is the identity arrow on a set A . Composition in **Rel** is to be given by

$$\{\langle a, c \rangle \in A \times C \mid \exists b(\langle a, b \rangle \in R \& \langle b, c \rangle \in S)\}$$

for $R \subseteq A \times B$ and $S \subseteq B \times C$.

- (a) Show that **Rel** is a category.
- (b) Show also that there is a functor $G : \mathbf{Sets} \rightarrow \mathbf{Rel}$ taking objects to themselves and each function $f : A \rightarrow B$ to its graph,

$$G(f) = \{\langle a, f(a) \rangle \in A \times B \mid a \in A\}$$

- (c) Finally, show that there is a functor $C : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$ taking each relation $R \subseteq A \times B$ to its converse $R^c \subseteq B \times A$, where,

$$\langle a, b \rangle \in R^c \Leftrightarrow \langle b, a \rangle \in R$$

Max:

- (a) By definition, the composition of two relations is another relation, so composition is closed. It suffices to show the identity and associativity rules.

For identity, let $1_A \subseteq A \times A$ and $R \subseteq A \times B$ for some sets A, B . Then $R \circ 1_A$ is,

$$\{\langle a, b \rangle \in A \times B \mid \exists a'(\langle a, a' \rangle \in 1_A \wedge \langle a', b \rangle \in R)\}$$

Since 1_A is the identity relation, $a = a'$, so the set is equivalent to,

$$\{\langle a, b \rangle \in A \times B \mid \langle a, b \rangle \in R\} = R$$

The case for $1_A \circ R$ follows similarly.

For associativity, let $R \subseteq A \times B$, $S \subseteq B \times C$, and $T \subseteq C \times D$ for sets A, B, C, D . Then,

$$\begin{aligned} T \circ (S \circ R) &= \{\langle a, d \rangle \in A \times D \mid \exists c(\langle a, c \rangle \in (S \circ R) \wedge \langle c, d \rangle \in T)\} \\ &= \{\langle a, d \rangle \in A \times D \mid \exists c(\exists b(\langle a, b \rangle \in R \wedge \langle b, c \rangle \in S) \wedge \langle c, d \rangle \in T)\} \\ &= \{\langle a, d \rangle \in A \times D \mid \exists c \exists b(\langle a, b \rangle \in R \wedge \langle b, c \rangle \in S \wedge \langle c, d \rangle \in T)\} \\ &= \{\langle a, d \rangle \in A \times D \mid \exists b \exists c(\langle a, b \rangle \in R \wedge \langle b, c \rangle \in S \wedge \langle c, d \rangle \in T)\} \\ &= \{\langle a, d \rangle \in A \times D \mid \exists b(\langle a, b \rangle \in R \wedge \exists c(\langle b, c \rangle \in S \wedge \langle c, d \rangle \in T))\} \\ &= \{\langle a, d \rangle \in A \times D \mid \exists b(\langle a, b \rangle \in R \wedge \langle b, d \rangle \in T \circ S)\} \\ &= (T \circ S) \circ R \end{aligned}$$

- (b) $G(f)$ is clearly a subset of $G(A) \times G(B) = A \times B$, so it suffices to show identities and composition are preserved.

For identities, let $\iota_A : A \rightarrow A$ be the identity function on set A . Then,

$$G(\iota_A) = \{\langle a, \iota_A(a) \rangle \in A \times A \mid a \in A\} = \{\langle a, a \rangle \in A \times A \mid a \in A\} = 1_A$$

For composition, let $f : A \rightarrow B$ and $g : B \rightarrow C$. Then,

$$\begin{aligned}
G(g) \circ G(f) &= \{\langle a, c \rangle \in A \times C \mid \exists b (\langle a, b \rangle \in G(f) \wedge \langle b, c \rangle \in G(g))\} \\
&= \{\langle a, c \rangle \in A \times C \mid \exists b (b = f(a) \wedge c = g(b))\} \\
&= \{\langle a, c \rangle \in A \times C \mid \exists b (c = g(f(a)))\} \\
&= \{\langle a, c \rangle \in A \times C \mid c = g(f(a))\} \\
&= \{\langle a, c \rangle \in A \times C \mid \exists b (c = (g \circ f)(a))\} = G(g \circ f)
\end{aligned}$$

- (c) Since $R^c \subseteq C(B) \times C(A) = B \times A$, morphisms are compatible with objects, so it suffices to show identities and composition are preserved.

For identities, let $1_A : A \rightarrow A$ be the identity relation on set A . Then,

$$\langle a, a \rangle \in (1_A)^c \iff \langle a, a \rangle \in 1_A$$

$$\text{So } (1_A)^c = 1_A.$$

For composition, let $R \subseteq A \times B$ and $S \subseteq B \times C$. Then,

$$\begin{aligned}
\langle a, c \rangle \in (S \circ R)^c &\iff \langle c, a \rangle \in S \circ R \\
&\iff \exists b (\langle c, b \rangle \in R \wedge \langle b, a \rangle \in S) \\
&\iff \exists b (\langle a, b \rangle \in S^c \wedge \langle b, c \rangle \in R^c) \\
&\iff \langle a, c \rangle \in R^c \circ S^c
\end{aligned}$$

$$\text{So } (S \circ R)^c = R^c \circ S^c.$$

□

Kyle:

- (a) For **Rel** to be a category, it must satisfy:
- Composition. For all morphisms f and g where $\text{cod } f = \text{dom } g$, their composition $g \circ f$ must exist.
 - Associativity. $h \circ (g \circ f) = (h \circ g) \circ f$
 - Identity. For all objects, the identity morphism 1_A must exist and must satisfy

$$\forall f : A \rightarrow B, f \circ 1_A = f = 1_B \circ f$$

Composition in **Rel** is defined

$$S \circ R := \{\langle a, c \rangle \in A \times C \mid \exists b (\langle a, b \rangle \in R \wedge \langle b, c \rangle \in S)\}$$

$$A \xrightarrow{R} B \xrightarrow{S} C$$

Identity is defined

$$1_A := \{\langle a, a \rangle \in A \times A \mid a \in A\}$$

To show associativity, suppose we have relations in the diagram below:

$$A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D$$

$$\begin{aligned}
T \circ (S \circ R) &= T \circ \{\langle a, c \rangle \mid \exists b \langle a, b \rangle \wedge \langle b, c \rangle\} \\
&= \{\langle a, d \rangle \mid \exists b (\langle a, b \rangle \wedge \langle b, c \rangle) \wedge \exists c (\langle b, c \rangle \wedge \langle c, d \rangle)\} \\
&= \{\langle b, d \rangle \mid \exists c \langle b, c \rangle \wedge \langle c, d \rangle\} \circ R \\
&= (T \circ S) \circ R
\end{aligned}$$

This shows associativity. Intuitively, the above proof shows that there must be at least one element $b \in B$ which bridges the gap between A and C , and there must be at least one element $c \in C$ which bridges the gap between B and D . However, these two bridges don't need to meet in the middle.

To show identity we insert the definition of identity into the definition of composition.

$$\begin{aligned}
R \circ 1_A &= \{\langle a, b \rangle \mid \exists a \in A (\langle a, a \rangle \in 1_A \wedge \langle a, b \rangle \in R)\} \\
&= \{\langle a, b \rangle \mid \exists a \in R \langle a, b \rangle\} \quad \text{Because } \forall a, \langle a, a \rangle \in 1_A \\
&= R
\end{aligned}$$

$$\begin{aligned}
1_B \circ R &= \{\langle a, b \rangle \mid \exists b \in B (\langle b, b \rangle \in 1_B \wedge \langle a, b \rangle \in R)\} \\
&= \{\langle a, b \rangle \mid \exists a \in R \langle a, b \rangle\} \quad \text{Because } \forall b, \langle b, b \rangle \in 1_B \\
&= R
\end{aligned}$$

Because composition satisfies identity and associativity, **Rel** is a category.

- (b) To show that there is a functor $G : \mathbf{Sets} \rightarrow \mathbf{Rel}$, just consider that a function is a particular kind of relation, so any morphisms in **Sets** are automatically morphisms in **Rel**; they can be transported over exactly as they are. The same is not true in the other direction $H : \mathbf{Rel} \rightarrow \mathbf{Sets}$, since some relations are not functions.
- (c) The functor $C : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$ takes sets to themselves, identity arrows to themselves, and each relation $C : \langle a, b \rangle \mapsto \langle b, a \rangle$. The two properties of functoriality are preservation of composition and identity, and C satisfies both, it is a functor.

□

Problem 1-2: Consider the following isomorphisms of categories and determine which ones hold.

- (a) $\mathbf{Rel} \cong \mathbf{Rel}^{\text{op}}$
- (b) $\mathbf{Sets} \cong \mathbf{Sets}^{\text{op}}$
- (c) For a fixed set X with a power set $P(X)$, as poset categories $P(X) \cong P(X)^{\text{op}}$ (the arrows in $P(X)$ are subset inclusions $A \subseteq B$ for all $A, B \subseteq X$).

Max:

- (a) Consider the functor from problem 1-1(c). We can define in the same way a converse functor from $\mathbf{Rel} \rightarrow \mathbf{Rel}^{\text{op}}$ taking each relation to its converse, and since converses are involutions, this is a two-sided inverse.
- (b) Suppose F is an isomorphism from $\mathbf{Sets} \rightarrow \mathbf{Sets}^{\text{op}}$. Then for any sets A, B , we would have that $\text{Hom}_{\mathbf{Sets}}(A, B) \cong \text{Hom}_{\mathbf{Sets}^{\text{op}}}(F(A), F(B)) \cong \text{Hom}_{\mathbf{Sets}}(F(B), F(A))$, i.e., $|A^B| = |F(B)^{F(A)}|$.

Letting B be the null set, this would mean (for non-empty A), that $0 = |A^{\{\}}| = |F(\{\})^{F(A)}|$. Since there is only one null set, there is some non-empty A such that $F(A)$ is also non-empty. But there is always at least one set-function from any set to a non-empty set, which is a contradiction.

- (c) We can define a functor $F : P(X) \rightarrow P(X)^{\text{op}}$ by mapping sets to their complement in X and reversing inclusions. This is evidently functorial, and bijective on objects (since compliments are involutions) and on morphisms since $A \subseteq B \iff X - B \subseteq X - A$. It follows that F is an isomorphism of categories.

□

Kyle: For there to be an isomorphism of categories there must exist a two functors F and G between them which compose to form the identity functors for each category. For this to happen, the objects and the morphisms must stand in one-to-one correspondence between the categories.

- (a) $\mathbf{Rel} \cong \mathbf{Rel}^{\text{op}}$. This is an isomorphism, since we can use the functor C from exercise 1. Call its inverse D , which maps sets to themselves and reverses the order of the elements in the relation. That is, $D \circ C = 1_{\mathbf{Rel}}$ and $C \circ D = 1_{\mathbf{Rel}^{\text{op}}}$. Since the two sided inverse for the functor exists, the categories are isomorphic.
- (b) $\mathbf{Sets} \not\cong \mathbf{Sets}^{\text{op}}$. There is an asymmetry in the initial and terminal objects of \mathbf{Sets} we can use to show the nonexistence of an isomorphic functor. Isomorphic functors preserve the cardinality of homsets. As a consequence, they map initial objects to initial objects and terminal objects to terminal objects. Isomorphic functors must also bijectively map objects to objects. But \mathbf{Sets} has one distinct initial object and many distinct terminal objects (i.e. all singleton sets), whereas $\mathbf{Sets}^{\text{op}}$ has one distinct terminal object (i.e. \emptyset^{op}) and many initial objects. So no such bijection can exist between objects, so the functor cannot be constructed, so the two categories are not isomorphic.
- (c) $\mathcal{P}(X) \cong \mathcal{P}(X)^{\text{op}}$. Let F be a functor that maps each subset of X to its complement. The morphism in $\mathcal{P}(X)$ is the subset relation \subseteq , which is the unique morphism between any two objects. Let F map each \subseteq to the unique \supseteq iff it exists in $\mathcal{P}(X)$. For any two subsets

A and B where $A \subseteq B$, $F : A \mapsto A^C$, $F : B \mapsto B^C$, and $F : \subseteq \rightarrow \supseteq$. Every set in the powerset has a unique complement, so the functor's object mapping is bijective. For two arbitrary objects $A, B \in \mathcal{P}(X)$, there is a single unique morphism in the opposite category iff $A \subseteq B$ in $\mathcal{P}(X)$.

- \rightarrow Suppose that $A \subseteq B$. Then $B^C \subseteq A^C$, so $B^{C^{\text{op}}} \supseteq A^{C^{\text{op}}}$, so $A^{\text{op}} \supseteq B^{\text{op}}$.
- \leftarrow Suppose that $A^{\text{op}} \supseteq B^{\text{op}}$. Then $B^{C^{\text{op}}} \supseteq A^{C^{\text{op}}}$, so $B^C \subseteq A^C$, so $A \subseteq B$

□

Problem 1-3:

- (a) Show that in **Sets**, the isomorphisms are exactly the bijections.
- (b) Show that in **Monoids**, the isomorphisms are exactly the bijective homomorphisms.
- (c) Show that in **Posets**, the isomorphisms are *not* the same as bijective homomorphisms.

Max:

- (a) This follows from elementary set theory, that if $f : A \rightarrow B$ and $g : B \rightarrow A$ are such that $g \circ f$ and $f \circ g$ are identities, then f and g are surjective and injective and thus bijective.
- (b) As a **Monoids** morphism is a set function that preserves the monoid structure, any isomorphism is a bijection by (a).

Conversely, any bijection $f : M \rightarrow N$ between monoids M, N satisfies,

$$f(m \cdot m') = f(m) \cdot f(m')$$

For all $m, m' \in M$.

Now if $n, n' \in N$ so that $n = f(m)$ and $n' = f(m')$ (which we can assume by surjectivity of f), we have,

$$f^{-1}(n \cdot n') = f^{-1}(f(m) \cdot f(m')) = f^{-1}(f(m \cdot m')) = m \cdot m' = f^{-1}(n) \cdot f^{-1}(n')$$

- (c) By elementary set theory, any homomorphism which is an isomorphism must have its categorical inverse be the set function inverse. So it suffices to find a bijective homomorphism whose set theoretic inverse is not a homomorphism.

Let $\mathcal{P}(2)$ be the poset of subsets of a 2-element set $2 = \{a, b\}$ under inclusion, and $\mathbf{4}$ be the chain on four elements $1 \leq 2 \leq 3 \leq 4$. Then we can define a function $f : \mathcal{P}(2) \rightarrow \mathbf{4}$ like,


$$\begin{aligned} \{\} &\mapsto 1 \\ \{a\} &\mapsto 2 \\ \{b\} &\mapsto 3 \\ \{a, b\} &\mapsto 4 \end{aligned}$$

We have clearly exhibited an order preserving function which is also bijective. However, the inverse map is not a morphism since in particular $2 \leq 3$ but it is not the case that $\{a\} \subseteq \{b\}$.

□

Kyle:

In a category, isomorphisms between objects are a pair of morphisms between two objects A and B which compose to form the two identity morphisms.



$$1_A = g \circ f$$

$$1_B = f \circ g$$

- (a) In **Sets**, a bijective function has a two sided inverse which composes to form the two identities on sets, so it is an isomorphism.
- (b) Our goal is to show that in **Monoids**, a morphism is iso iff it is bijective on sets.
- → Let $\varphi : A \rightarrow B$ be an iso morphism. That is, it has a two sided inverse ψ such that $\psi \circ \varphi = 1_A$ and $\varphi \circ \psi = 1_B$. To be an inverse on monoids, it must be a bijection on the underlying sets.
 - ← Let $\varphi : A \rightarrow B$ be a bijective monoid homomorphism. Define $\psi : B \rightarrow A$ to be a map which sends $\psi : \varphi(a) \mapsto a$. Since φ is a bijection on sets, this is well defined. To prove that it is also a monoid homomorphism, observe that it sends identities to identities, since $\varphi(e_A) = e_B$, so $\psi : e_B \mapsto e_A$. To show it satisfies the homomorphism property, observe that $\psi(\varphi(a_1) \circ \varphi(a_2)) = \psi(\varphi(a_1 \circ a_2)) = a_1 \circ a_2 = \psi(\varphi(a_1)) \circ \psi(\varphi(a_2))$. So indeed, ψ is a homomorphism, so it exists as a morphism in **Monoids**. To show that ψ is the two-sided inverse of φ , observe that $\psi \circ \varphi(a) = \psi(\varphi(a)) = a$, and $\varphi \circ \psi(\varphi(a)) = \varphi(\psi(\varphi(a))) = \varphi(a)$. Since there exists a two sided inverse, φ is an isomorphism.
- (c) In **Posets**, suppose we have two posets A and B , both containing the same elements $\{c, d\}$, but in A there is no relation between c and d , whereas in B , $c \leq d$. Define a functor F which maps each element to itself. This is clearly bijective on sets. It is a homomorphism, because it preserves the existing relations $c \leq c$ and $d \leq d$. However, F has no inverse, since the relation $c \leq d$ must be mapped to $\text{Hom}_A(c, d)$, which is empty. Since we have a bijective homomorphism with no inverse in **Posets**, the isomorphisms are not the same as the bijective homomorphisms.

□

Problem 1-4: Let X be a topological space and preorder the points by *specialization*: $x \leq y$ iff y is contained in every open set that contains x . Show that this is a preorder, and that it is a poset if X is T_0 (for any two distinct points, there is some open set containing one but not the other). Show that the ordering is trivial if X is T_1 (for any two distinct points, each is contained in an open set not containing the other).

Max: Identity is clear since x is trivially contained in every open set that contains itself.

For transitivity, suppose $x \leq y$ and $y \leq z$. Then if U is open in X , $x \in U \Rightarrow y \in U \Rightarrow z \in U$ so in particular $x \in U \Rightarrow z \in U$ so $x \leq z$. It follows that specialization is at least a preorder.

If X is T_0 , then suppose $x \leq y$ and $y \leq x$. If $x \neq y$, we can choose without loss of generality an open U such that $x \in U$ and $y \notin U$, but the latter is a contradiction since $x \leq y$.

Finally if X is T_1 , and $x \leq y$, $x \neq y$ creates a contradiction, because we can choose some open U, V so that $x \in U$, $y \in V$ and $U \cap V = \emptyset$, so in particular $x \in U$ and $y \notin U$, but by $x \leq y$, $y \in U$. So the order is trivial - the only pairs x, y for which $x \leq y$ are those where $x = y$. \square

Kyle: Skipped \square

Problem 1-5: For any category \mathbf{C} , define a functor $U : \mathbf{C}/C \rightarrow \mathbf{C}$ from the slice category over an object C that “forgets about C .” Find a functor $F : \mathbf{C}/C \rightarrow C^\rightarrow$ to the arrow category such that $\mathbf{dom} \circ F = U$.

Max: For U , define $U(f) = \mathbf{dom} f$ on objects. Then considering $g : f \rightarrow f'$ as a function $\mathbf{dom} f \rightarrow \mathbf{dom} f'$ defines the action on morphisms.

Since we define identities in the slice categories as identities on domains, identities are preserved. Similarly, since we define the composition in \mathbf{C}/C as the composition of the underlying functions on domains, it follows that U preserves composition.

$$\begin{array}{ccc}
 \begin{array}{ccc}
 a & \xrightarrow{f} & c \\
 1_a \downarrow & \nearrow f' & \\
 a & & a
 \end{array} & \mapsto & \begin{array}{ccc}
 a & \xrightarrow{f} & c \\
 g \downarrow & \nearrow f' & \\
 b & & c
 \end{array} \\
 & & \begin{array}{ccc}
 a & \xrightarrow{f} & c \\
 g \downarrow & \nearrow f' & \\
 b & & c
 \end{array} \\
 & & \begin{array}{ccc}
 a & \xrightarrow{f} & c \\
 g \downarrow & \nearrow f' & \\
 b & & c
 \end{array}
 \end{array}$$

Now define F on objects to take each morphism $f : a \rightarrow c$ to itself, and each morphism $g : f \rightarrow f'$ to $\langle g, 1_C \rangle$. Letting $f \xrightarrow{g} f' \xrightarrow{g'} f''$, we have,

$$F(g') \circ F(g) = \langle g', 1_C \rangle \circ \langle g, 1_C \rangle = \langle g' \circ g, 1_C \rangle = F(g' \circ g)$$

Then the composition of functors $\mathbf{dom} \circ F$ has, for objects $f \in \text{ob } \mathbf{C}/C$,

$$(\mathbf{dom} \circ F)(f) = \mathbf{dom} F(f) = \mathbf{dom} f = U(f)$$


And for morphisms $g : f \rightarrow f'$,

$$(\mathbf{dom} \circ F)(g) = \mathbf{dom} \langle g, 1_C \rangle = g = U(g)$$


□

Kyle:

5) The slice category $U: \mathcal{C}/\mathcal{C} \rightarrow \mathcal{C}$ can be defined as a mapping from each object $x \xrightarrow{f} c$ to X and from each morphism $x \xrightarrow{g} x'$ to $X \xrightarrow{g} X'$.



Let $F: \mathcal{C}/\mathcal{C} \rightarrow \mathcal{C}'$ be a functor such that taking the $\text{dom} \circ F$ yields the same as taking U . The arrow category \mathcal{C}'^\rightarrow is a category \mathcal{A} in which all the morphisms in \mathcal{C} become objects, and morphism are pairs of arrows, which commute. In the original category, we see that all commutative triangles terminating in C can be drawn as commutative squares.



Let dom be the functor which maps each $x \xrightarrow{f} c \mapsto X$, the object in the arrow category gets mapped to its domain, and each arrow (a, b) gets mapped to $(a, b) \mapsto a$.

Then this is exactly U . So the functor $F: \mathcal{C}/\mathcal{C} \rightarrow \mathcal{C}'$ maps each object $f \mapsto f$ to itself, and it maps each morphism $(a, 1_c) \mapsto a$.

□

Problem 1-6: Construct the “coslice category” C/\mathbf{C} of a category \mathbf{C} under an object C from the slice category \mathbf{C}/C and the “dual category” operation $-^{\text{op}}$.

Max: Consider the category $(\mathbf{C}^{\text{op}}/C^{\text{op}})^{\text{op}}$. It has as objects maps $f^{\text{op}} : X^{\text{op}} \rightarrow C^{\text{op}}$ in \mathbf{C}^{op} , and as morphisms maps g^{op} satisfying,


$$\begin{array}{ccc} X^{\text{op}} & \xrightarrow{f^{\text{op}}} & C^{\text{op}} \\ h^{\text{op}} \downarrow & \nearrow f'^{\text{op}} & \\ Y^{\text{op}} & & \end{array}$$

Which “unravels” (by reversing arrows) to the following equivalent diagram in \mathbf{C} ,

$$\begin{array}{ccc} & f & \\ X & \xleftarrow{ } & C \\ h \uparrow & \nearrow f' & \\ Y & & \end{array}$$

So we find that the category $(\mathbf{C}^{\text{op}}/C^{\text{op}})^{\text{op}}$ is nothing more than the category of maps f with domain C , and morphisms $h : Y \rightarrow X$ satisfying the above diagrams. This is precisely what we would expect from a “coslice” category C/\mathbf{C} . \square

Kyle:



\square


Problem 1-7: Let $2 = \{a, b\}$ be any set with exactly 2 elements a and b . Define a functor $F : \text{Sets}/2 \rightarrow \text{Sets} \times \text{Sets}$ with $F(f : X \rightarrow 2) = (f^{-1}(a), f^{-1}(b))$. Is this an isomorphism of categories? What about the analogous situation with a one-element set $1 = \{a\}$ instead of 2 ?

Max: Consider the object $(\{1\}, \{1\})$ in $\text{Sets} \times \text{Sets}$, and suppose it was the image of some function $f : X \rightarrow 2$ under F , so $F(f) = (f^{-1}(a), f^{-1}(b)) = (\{1\}, \{1\})$.

$f^{-1}(a) \cap f^{-1}(b) = \emptyset$ by elementary set theory, but $f^{-1}(a) = \{1\}$ and $f^{-1}(b) = \{1\}$, so $f^{-1}(a) \cap f^{-1}(b) = \{1\} \cap \{1\} = \{1\}$. So $\emptyset = \{1\}$, a contradiction. Thus F is not surjective on objects and cannot be an isomorphism of categories.


We don't run into this problem for the one-element set $1 = \{a\}$ - here we can define an inverse functor $G : \text{Sets} \rightarrow \text{Sets}/1$ that maps each set X to the unique function $X \rightarrow 1$ and acts identically on functions. It is straight-forward to show that this is functorial and a two-sided inverse to F defined analogously as with 2 . \square

Kyle:




Define the functor $F : \text{Sets}/2 \rightarrow \text{Sets} \times \text{Sets}$ to map each object $F : f \mapsto (f^{-1}(a), f^{-1}(b))$, and each morphism $F : h \mapsto (h|_{f^{-1}(a)}, h|_{f'^{-1}(a)})$ as shown in the above diagram. To see how this has no inverse, we can construct an object in $\text{Sets} \times \text{Sets}$ which cannot be mapped to by F . Consider two sets with a nonempty intersection, e.g. $A := \{c, d\}$ and $B := \{c, d, e\}$. Clearly these sets exist, and so their product $A \times B$ exists in $\text{Sets} \times \text{Sets}$. Assume that there exists an object f in the slice category which F maps to $A \times B$. f is a function in Sets $f : X \rightarrow \{a, b\}$. Consider the two fibers of f , $f^{-1}(a)$ and $f^{-1}(b)$. The fibers of any function partition the domain, which in this case is X . In other words, $\neg \exists x (x \in f^{-1}(a) \wedge x \in f^{-1}(b))$. But $f^{-1}(a) = A$ and $f^{-1}(b) = B$, which were defined to have a nonempty intersection, so f cannot exist, so F has no inverse, so this is not an isomorphism of categories.

(b)



In slices over $\{\ast\}$, F maps objects (morphisms to $\{\ast\}$) to their fibers, which, because there is only one fiber, is the entire domain. So this functor is the same as the forgetful functor $U: \mathcal{C}/\ast \rightarrow \mathcal{C}$.



Since $\{\ast\}$ has morphisms out of it in Set , in particular $1\ast$, that object will exist in the slice category. This is true for any category. So for any category with a terminal object, taking the slice out of it will produce a category isomorphic to the original category. In particular, $\text{Set}/\ast \cong \text{Set}$. Since $U: \text{Set}/\ast \rightarrow \text{Set}$ is full and faithful,

□

Problem 1-8: Any category \mathbf{C} determines a preorder $P(\mathbf{C})$ by defining a binary relation \leq on the objects by

$$A \leq B \text{ if and only if there is an arrow } A \rightarrow B$$

Show that P determines a functor from categories to preorders, by defining its effect on functors between categories and checking the required conditions. Show that P is a (one-sided) inverse to the evident inclusion functor of preorders into categories.

Max: If F is a functor from $\mathbf{C} \rightarrow \mathbf{D}$, let $P(F)$ coincide with the action of F on objects.

Then if $C \leq C'$ in $P(\mathbf{C})$, there is some morphism $f : A \rightarrow B$ by definition. Then $F(f)$ is a morphism from $F(A) \rightarrow F(B)$ in \mathbf{D} by functoriality, so $F(A) \leq F(B)$ in $P(\mathbf{D})$. It follows that $P(F)$ is order-preserving, so this mapping of morphisms is well-defined.

The identity functor $\mathbf{1}_{\mathbf{C}}$ gets mapped by P to the identity set-function on $P(\mathbf{C}) = \text{ob } \mathbf{C}$, so identities are carried to identities.

If $F : \mathbf{C} \rightarrow \mathbf{D}$ and $G : \mathbf{D} \rightarrow \mathbf{E}$, then $P(G \cdot F)$ goes to the set-function composition of F and G on objects, which is exactly $P(G) \circ P(F)$, so composition is preserved.

Finally to show that P is a one-sided inverse let $I : \mathbf{Pre} \rightarrow \mathbf{Cat}$ be the inclusion functor regarding pre-orders as categories. Consider the category $P(I(X))$: its objects are the objects of $I(X)$, which are the objects of X , and we have,


$$a \underset{P(I(X))}{\leq} b \iff \text{Hom}_{I(X)}(a, b) \neq \emptyset \iff a \underset{X}{\leq} b$$

So that the two pre-orders, X and $P(I(X))$, are equal.

Finally if f is an order-preserving map $X \rightarrow Y$, both I and P preserve f 's action on objects by definitions, so $P(I(f)) = f$. \square

Kyle:

Let P be a functor from the category \mathbf{Cat} to the category \mathbf{Pre} as depicted in the diagram.



Let P take each category \mathcal{C} to its corresponding preorder $P(\mathcal{C})$. Define the action of P on a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ on F 's object maps and morphism maps separately. Let P take each object map $F_a : a \rightarrow Fa$ as the corresponding map in the preorder homomorphism $a \rightarrow Fa$, and let P take each morphism map $F_f : f \rightarrow Ff$ to the corresponding map between relations $a \leq b \rightarrow Fa \leq Fb$.

We can check that P satisfies the 3 properties defining a functor. Observe that P maps $F : \mathcal{C} \rightarrow \mathcal{D}$ to an element in $\text{Hom}_{\text{Pre}}(P(\mathcal{C}), P(\mathcal{D}))$. To see that it preserves identity $P : 1_{\mathcal{C}} \rightarrow 1_{P(\mathcal{C})}$, observe that it takes each identity morphism in the category 1_a to $a \leq a$ and each identity morphism map to the identity relation map. To see that it preserves composition, observe that $P(G \circ F) = P(G) \circ P(F)$, since objects inside categories are mapped to themselves and a relation morphism $a \leq b \rightarrow G \circ Fa \leq G \circ Fa$ iff there is a composition of morphisms in Cat .

To show that P is a one-sided inverse to the evident inclusion functor of preorders into categories, observe that while there is a bijection of objects, homsets of cardinality ≥ 1 get mapped to a homset of cardinality 1, while empty homsets get mapped to empty homsets, so P is necessarily lossy for homsets of cardinality > 1 , whereas the evident inclusion functor is faithful on homsets.

□

Problem 1-9: Describe the free categories on the following graphs by determining their objects, arrows, and composition operations.

(a)

$$a \xrightarrow{e} b$$

(b)

$$\begin{array}{ccc} & e & \\ a & \xrightleftharpoons[]{} & b \\ & f & \end{array}$$

(c)

$$\begin{array}{ccc} & e & \\ a & \xrightarrow{e} & b \\ & g \searrow & \downarrow f \\ & & c \end{array}$$

(d)

$$\begin{array}{ccc} & e & \\ a & \xrightleftharpoons[]{} & b \\ & h \swarrow & \downarrow f \\ & g \searrow & \downarrow \\ & & c \end{array}$$

Max: In each case let denote by G the graph and by $\mathbf{C}(G)$ the free category on G . Since rules for composing paths and the objects of $\mathbf{C}(G)$ are given by definition, we see it sufficing to only characterize the arrows of $\mathbf{C}(G)$.

- (a) Clearly a path cannot have more than one edge, since e is not compatible with itself under path composition. It follows that there are exactly three morphisms on $\mathbf{C}(G)$:
- $\text{Hom}_{\mathbf{C}(G)}(a, a) = \{1_a\}$
 - $\text{Hom}_{\mathbf{C}(G)}(b, b) = \{1_b\}$
 - $\text{Hom}_{\mathbf{C}(G)}(a, b) = \{e\}$

- (b) We claim that every path in $\mathbf{C}(G)$ is alternating between e and h . We prove this by induction on path length n .

The case $n = 0$ is immediate.

Now suppose that the claim holds for paths of length n , and consider a path of length $n + 1$, which by induction will be of the form,

$$p \circ A$$

For some edge p and path A alternating in e and h .

If A is empty, the result is immediate since p must be e or h .

If A starts with e , then the only option for p is h and the result follows. The case A starting with h is similar.

Now that we know what all paths look like in $\mathbf{C}(G)$, we can determine all the arrows like so:

- $\text{Hom}_{\mathbf{C}(G)}(a, a) = \{(he)^n\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(b, b) = \{(eh)^n\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(a, b) = \{(eh)^n e\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(b, a) = \{(he)^n h\}_{n \geq 0}$

- (c) Suppose there is a path of length 3, pqr . $s(p) = t(q)$ cannot be c as that would leave no options for p . This means $s(q) = t(r)$ cannot be b (this would force $t(q) = c$), which means $s(r)$ cannot be a (this would force $t(r) = b$).

Similarly, $t(r)$ cannot be c as that would leave no options for q , which means $s(r)$ cannot be b either.

Finally $s(r)$ cannot be c since there are no edges of source c .

It follows that there are no paths of length 3, and in fact greater than 3 neither since those paths would have length-3 subpaths.

From this we know that all paths are at most length 2, and a simple check shows that only the following are valid:

- $\text{Hom}_{\mathbf{C}(G)}(a, a) = \{1_a\}$
- $\text{Hom}_{\mathbf{C}(G)}(b, b) = \{1_b\}$
- $\text{Hom}_{\mathbf{C}(G)}(a, b) = \{e\}$
- $\text{Hom}_{\mathbf{C}(G)}(b, a) = \{f\}$
- $\text{Hom}_{\mathbf{C}(G)}(a, c) = \{g, fe\}$
- $\text{Hom}_{\mathbf{C}(G)}(c, c) = \{1_c\}$

- (d) We claim that every path in $\mathbf{C}(G)$ is of the form,

$$p \circ A$$

Where p is either an empty path or one of f, g , and A is an alternating sequence of e and h . We prove this by induction on path length n .

The case $n = 0$ is immediate.

Now suppose that the claim holds for paths of length n , and consider a path of length $n + 1$, which by induction will be of the form,

$$p_0 \circ p \circ A$$

For some edge p_0 , p as described above and path A alternating in e and h .

If p is empty, then the result is immediate since if p_0 is f or g it is directly of the needed form, and if its either e or h then the entire path is alternating.

If p is f , then p_0 must be an edge from c . This forces p_0 to be empty and the result follows. The case $p = g$ is similar.

Now that we know what all paths look like in $\mathbf{C}(G)$, we can determine all the arrows like so:


- $\text{Hom}_{\mathbf{C}(G)}(a, a) = \{(he)^n\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(b, b) = \{(eh)^n\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(a, b) = \{(eh)^n e\}_{n \geq 0}$

- $\text{Hom}_{\mathbf{C}(G)}(b, a) = \{(he)^n h\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(b, c) = \{f(eh)^n\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(a, c) = \{f(eh)^n e\}_{n \geq 0} \cup \{g(he)^n\}_{n \geq 0}$
- $\text{Hom}_{\mathbf{C}(G)}(c, c) = \{1_c\}$

And all other hom-sets are empty.

□

Kyle:



□


Problem 1-10: How many free categories on graphs are there which have exactly six arrows? Draw the graphs that generate these categories.

Max: First of all, the underlying graph cannot contain any cycles, since any cycle is a path that can be composed with itself infinitely, resulting in infinitely many arrows.


Then there cannot be more than 6 nodes + edges since each node admits an identity arrow to itself. This gives us a path to a solution by enumerating all (finitely many) non-isomorphic directed acyclic graphs with $|V| + |E| \leq 6$.

Doing this for all possible number of nodes, we get:


- 0) 0 nodes generates the empty category, which has no arrows.
- 1) With 1 node our only option is to have no edges, since we can't have a loop, which generates the singleton category.
- 2) With 2 nodes our only option is to have four edges from one node to the other:



- 3) We get four graphs whose free category has 6 arrows:



- 4) We have the following four graphs on four nodes:



To cut back on checking, we can notice first that there can be at most two edges (by the above logic $|V| + |E| \leq 6$). If there are two edges, there are going to be two disconnected graphs, either of order 3 or 2. If 3, the size 3 subgraph has to end up being as a free category with 5 arrows, all of whom we enumerated while working through case (3). If 2, there is only one possibility enumerated last above. The case of one edge yields only 5 morphisms, and no edges yields only 4.

- 5) There is just one such graph, namely



We can conclude this by noticing there can be at most one edge, and any such choice forces the above graph (up to isomorphism). There cannot be no edge since this yields just five morphisms.


- 6) With 6 nodes we must not select any edges since then we'd have more than 6 arrows.
 This gives us a single option, the discrete category of 6 elements:

★ ★ ★

★ ★ ★

□

Kyle:



□

Problem 1-11: Show that the free monoid functor

$$M : \mathbf{Sets} \rightarrow \mathbf{Mon}$$

exists, in two different ways:

- (a) Assume the particular choice $M(X) = X^*$ and define its effect

$$M(f) : M(A) \rightarrow M(B)$$

on a function $f : A \rightarrow B$ to be

$$M(f)(a_1 \dots a_k) = f(a_1) \dots f(a_k), \quad a_1, \dots a_k \in A$$

- (b) Assume only the UMP of the free monoid and use it to determine M on functions, showing the result to be a functor.

Reflect on how these two approaches are related.

Max:

- (a) Let $f : A \rightarrow B$ and $g : B \rightarrow C$. Then,

$$M(g \circ f)(a_1 \dots a_k) = g(f(a_1)) \dots g(f(a_n)) = M(g)(f(a_1) \dots f(a_n)) = M(g)(M(f)(a_1 \dots a_n))$$

And, for the identity function $\iota_A : A \rightarrow A$,

$$M(\iota_A)(a_1 \dots a_n) = \iota_A(a_1) \dots \iota_A(a_n) = a_1 \dots a_n$$

So M as defined is a functor.

- (b) Let $A, B \in \text{ob}(\mathbf{Sets})$ and $i_A : A \rightarrow |M(A)|$ and $i_B : B \rightarrow |M(B)|$ be the canonical maps from the UMP for A, B respectively. Then for any set function $f : A \rightarrow B$, we can define $M(f) \in \text{Hom}_{\mathbf{Mon}}(M(A), M(B))$ as the unique monoid morphism making the following square commute by UMP:

$$\begin{array}{ccc} |M(f)| & & \\ |M(A)| & \xrightarrow{\quad} & |M(B)| \\ \uparrow i_A & & \uparrow i_B \\ A & \xrightarrow{\quad f \quad} & B \end{array}$$

If ι_A is the identity function $A \rightarrow A$, then the identity monoid morphism $I_A : M(A) \rightarrow M(A)$ coincides with the set identity function $|M(A)| \rightarrow |M(A)|$. Then the diagram below clearly commutes:

$$\begin{array}{ccc} |I_A| = \iota_{M(A)} & & \\ |M(A)| & \xrightarrow{\quad} & |M(A)| \\ \uparrow i_A & & \uparrow i_A \\ A & \xrightarrow{\quad \iota_A \quad} & A \end{array}$$

So $I_A = M(\iota_A)$ by uniqueness of the UMP.

Finally for composition, suppose we have sets A, B, C and $f : A \rightarrow B, g : B \rightarrow C$, then, we have:


$$\begin{array}{ccccc} |M(g)| \circ |M(f)| & = & |M(g) \circ M(f)| \\ \text{---} & & \text{---} \\ |M(A)| & \xrightarrow{|M(f)|} & |M(B)| & \xrightarrow{|M(g)|} & |M(C)| \\ \uparrow i_A & & \uparrow i_B & & \uparrow i_C \\ A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ \text{---} & & \text{---} & & \text{---} \\ & & & & g \circ f \end{array}$$

Since the inner two squares commute, so does the outer square. It follows by uniqueness of UMP that $M(g) \circ M(f) = M(g \circ f)$.

□

Kyle:

- (a) Here we assume the free monoid functor is the Kleene star operation.



Our goal is to show that M is a functor. Functorality requires preservation of identity and composition. M maps 1_A to the identity homomorphism on the free monoid.

Composition follows:

$$\begin{aligned} M(g \circ f)(a_1 \dots a_k) &= (g \circ f)(a_1) \dots (g \circ f)(a_k) \\ &= g(f(a_1)) \dots g(f(a_k)) \\ &= M(g) \circ M(f)(a_1 \dots a_k) \end{aligned}$$


Intuitively, the monoid homomorphism $M(g \circ f)$ takes each letter in the monoid MA and replaces it according to the composition function $g \circ f$, which is the same as replacing them by f first and then by g . So the monoid homomorphism mapped to from the composition of functions is the same as the composition of the two individual monoid homomorphisms.

- (b) Here we don't assume the free monoid functor is the Kleene star operation. Instead we assume the UMP: that given


- A : a generating set.
- $i : A \rightarrow |M(A)|$, An arbitrary function i mapping from the generating set to the (underlying set of the) free monoid
- $f : A \rightarrow |N|$, an arbitrary function f mapping to the underlying set of an arbitrary monoid,

Then there exists a unique monoid homomorphism $\bar{f} : M(A) \rightarrow N$ such that $|\bar{f}| \circ i = f$.

Our first goal is to determine what the functor M will do to an arbitrary function. Let A, B be arbitrary sets, and let $f^* : A \rightarrow B$ be an arbitrary function. Let $M(A)$ and $M(B)$ be the free monoids of those sets, and let i_A and i_B be the canonical injections from A and B into the underlying sets of their respective free monoids. We can define a composite function $f : A \rightarrow |M(B)|$ to be $f := i_B \circ f^*$. By the UMP, we know that, because there is a function $f : A \rightarrow |M(B)|$, there exists a unique homomorphism $\bar{f} : M(A) \rightarrow M(B)$ such that $|\bar{f}| \circ i_A = f$. Thus we have 2 commutative triangles (as shown below), so the entire square in **Sets** commutes.




But this still doesn't answer our question, what does M do to f^* ? In order to be a functor, M must map f^* to a morphism that preserves the domain and codomain under the functor. That is, $M : (f : A \rightarrow B) \mapsto (Mf : M(A) \rightarrow M(B))$ But as we said, above, there is a unique morphism $\bar{f} : M(A) \rightarrow M(B)$, so it must be the case that $M : f^* \mapsto \bar{f}$



To check that M is a functor it suffices to check for preservation of identity and composition. The identity function 1_A gets mapped to the identity monoid homomorphism.

The composition of two arbitrary functions $g^* \circ f^*$ is shown in the commutative diagram:



Here we have defined $g := i_C \circ g^*$. By the UMP of $M(B)$, we know the existence and uniqueness of \bar{g} . Let h^* be defined to be the composite $g^* \circ f^*$, and $h : A \rightarrow |M(C)| := i_C \circ h^*$. By the UMP of $M(A)$ we also know that because there is a function $h : A \rightarrow |M(C)|$, there exists a unique homomorphism $\bar{h} : M(A) \rightarrow M(C)$ such that $|\bar{h}| \circ i_A = h$. But this \bar{h} is exactly where M sends h^* , and as it is the unique morphism from $M(A) \rightarrow M(C)$, it must equal $\bar{g} \circ \bar{f}$. So functorial composition is satisfied.

□

Problem 1-12: Verify the UMP for free categories on graphs, defined as above with arrows being sequences of edges. Specifically, let $\mathbf{C}(G)$ be the free category on the graph G , so defined, and $i : G \rightarrow U(\mathbf{C}(G))$ the graph homomorphism taking vertices and edges to themselves, regarded as objects and arrows in $\mathbf{C}(G)$. Show that for any category \mathbf{D} and graph homomorphism $h : G \rightarrow U(\mathbf{D})$, there is a unique functor

$$\bar{h} : \mathbf{C}(G) \rightarrow \mathbf{D}$$

with

$$U(\bar{h}) \circ i = h,$$

where $U : \mathbf{Cat} \rightarrow \mathbf{Graph}$ is the underlying graph functor.

Max: The condition on \bar{h} mandates that \bar{h} takes each node a , regarded as an element of $\mathbf{C}(G)$, to the object $h(a)$ in \mathbf{D} , and that it takes each edge e , regarded as a morphism of $\mathbf{C}(G)$, to the edge representing the morphism $h(e)$ in $U(\mathbf{D})$.

That this uniquely pins down \bar{h} follows from the fact that it is a graph homomorphism, since we know it on objects and by definition any morphism p in $\mathbf{C}(G)$ can be written as some finite sequence of edges,

$$p = e_1 \circ \dots \circ e_n$$

So that $\bar{h}(p)$ is equal to $\bar{h}(e_1) \circ \dots \circ \bar{h}(e_n)$ by the fact that \bar{h} is a functor.

Now since each non-identity morphism p in $\mathbf{C}(G)$ is uniquely expressible as some sequence of edges, it is no issue to define $\bar{h}(p)$ as $h(e_1) \circ \dots \circ h(e_n)$, where we send the empty sequence to the identity morphism on $h(a)$, thus preserving identities, and we send objects to their image in h .


Then for any non-empty composition-compatible paths $p = e_1 \dots e_n$ and $q = s_1 \dots s_m$, we have,

$$\begin{aligned} \bar{h}(p \circ q) &= \bar{h}(e_1 \dots e_n \circ s_1 \dots s_m) = \bar{h}(e_1 \dots e_n s_1 \dots s_m) \\ &= h(e_1) \circ \dots \circ h(e_n) \circ h(s_1) \circ \dots \circ h(s_m) \\ &= \bar{h}(e_1 \dots e_n) \circ \bar{h}(s_1 \dots s_m) \\ &= \bar{h}(p) \circ \bar{h}(q) \end{aligned}$$


For empty paths then the preservation of composition is immediate. □

Kyle:


Given



we are asked to show:



The functor \mathbf{C} which takes the graph G to its free category $\mathbf{C}(G)$ takes every path in G between nodes A and B to a morphism between objects A and B . This means that every morphism in $\mathbf{C}(G)$ is uniquely determined by the ordered set of edges which make up the corresponding path; if two ordered sets of edges are equal (they have the same edge set in the same order), then they map to the same morphism in $\mathbf{C}(G)$. Because h is a graph homomorphism, it maps vertices to vertices, and edges to edges in such a way that the source and target of each edge get mapped according to the vertex mapping.



Let p be an arbitrary path in the graph G from vertex $A \rightarrow B$. p is also a morphism in $\mathbf{C}(G)$, by the definition of $\mathbf{C}(G)$. p is composed of edges in the graph, which we can write $p = (e_1, \dots, e_n)$. Let p' be the path generated by applying h elementwise to p . That is, $p' := (h(e_1) \dots h(e_n))$. Let us define \bar{h} to be the functor that maps objects according to the graph

homomorphism h and morphisms $\bar{h} : p \mapsto p'$. p is uniquely defined by the edges that make it up, so this functor is the unique functor mapping the free category $\mathbf{C}(G)$ to D .

□

Problem 1-13: Use the Cayley representation to show that every small category is isomorphic to a “concrete” one, that is, one in which the objects are sets and the arrows are functions between them.

Max: Define a functor $\overline{(-)} : \mathbf{C} \rightarrow \overline{\mathbf{C}}$ on objects as sending $C \rightarrow \overline{C}$ and on morphisms $g : C \rightarrow D$ to $\bar{g} : \overline{C} \rightarrow \overline{D}$. It suffices to show that $\overline{(-)}$ is a functor, is bijective on objects, and is bijective on Hom-sets.

For preservation of identity, $\overline{1_A}$ acts by post-composition with the identity, and therefore is the set-functional identity $\overline{A} \rightarrow \overline{A}$; since composition occurs in $\overline{\mathbf{C}}$ as set-function composition, $\overline{1_A}$ is the identity in $\overline{\mathbf{C}}$.

For functorial composition, if $f : A \rightarrow B$ and $g : B \rightarrow C$ in \mathbf{C} , and if $h \in \overline{A}$, then,

$$(\bar{g} \circ \bar{f})(h) = \bar{g}(\bar{f}(h)) = \bar{g}(f \circ h) = g \circ (f \circ h) = (g \circ f) \circ h = (\overline{g \circ f})(h)$$


So that $\bar{g} \circ \bar{f} = \overline{g \circ f}$, i.e. $\overline{(-)}$ preserves composition.

For bijectivity on objects, surjectivity is clear by definition since the objects are precisely the elements \overline{C} for $C \in \text{ob}(\mathbf{C})$. Injectivity is also clear since we can always recover the original object C uniquely from the identity $1_C \in \overline{C}$, which could never be in any other \overline{D} for $D \neq C$.

For bijectivity on morphisms, surjectivity is again clear since every morphism is defined as \bar{g} for some $g \in \text{mor}(\mathbf{C})$. Now if $g : A \rightarrow B$ for objects A, B in \mathbf{C} , \bar{g} acts by post-composition with g from $\overline{A} \rightarrow \overline{B}$. It follows that the original g can be recovered uniquely as $\bar{g}(1_A)$, so we also have injectivity. \square

Kyle:

Let $F : \mathbf{C} \rightarrow \overline{\mathbf{C}}$ be a functor from a small factor to its Cayley representation, in which F maps objects $F : C \mapsto \{\text{morphisms ending at } C\}$ and maps morphisms $F : f \mapsto \{\text{functions generated by postcomposition with } f\}$, as shown in the diagram below.



To show $\mathbf{C} \simeq \overline{\mathbf{C}}$, it suffices to show that F is bijective in objects and that the functors are full and faithful. F is obviously bijective in objects, since each codomain object is defined in terms of the ending of morphisms, which is distinct. For every morphism in the homset $f \in \text{Hom}_{\mathbf{C}}(A, B)$, there is one morphism $\bar{f} \in \text{Hom}_{\overline{\mathbf{C}}}(A, B)$ by construction, so F is full. To show that F is faithful, suppose that $g, g' \in \text{Hom}_{\mathbf{C}}(B, C)$ such that $g \neq g'$. g acts on each element in \bar{B} by precomposition, and likewise for g' . Because $g \neq g'$, for any arbitrary $f \in \bar{B}$, $g \circ f \neq g' \circ f$. Because their actions on arbitrary elements are distinct, as functions they are not equal $\bar{g} \neq \bar{g}'$, so F is faithful, and therefore \mathbf{C} and $\overline{\mathbf{C}}$ are isomorphic. \square

Problem 1-14: The notion of a category can also be defined with just one sort (arrows) . rather than two (arrows and objects); the domains and codomains are taken to be certain arrows that act as units under composition, which is partially defined. Read about this definition in section I.1 of Mac Lane's *Categories for the Working Mathematician*, and do the exercise mentioned there, showing that it is equivalent to the usual definition.

Max:

- (\Rightarrow) Suppose we have an axiomatization of \mathbf{C} as objects and arrows. Then consider just the collection of arrows, with composition defined as in \mathbf{C} . We show each axiom is satisfied.
 - (i) If $(k \circ g)$ and f are composable then it is not hard to see that g is composable with f and k is composable with $g \circ f$. Then by associativity $(k \circ g) \circ f = k \circ (g \circ f)$.
 - (ii) If $k \circ g$ and $g \circ f$ are valid compositions then by considering the constraints on domains/codomains, we find that $k \circ g \circ f$ is valid.
 - (iii) For any arrow $g : A \rightarrow B$ we have $g \circ 1_A = g$ and $1_B \circ g = g$, so there exist such arrows.
- (\Leftarrow) With the collection of arrows as A , and compositability of g, f as gf when allowed, we will define a category \mathbf{C} as follows:
 - For objects, the collection of identity arrows u so that $uf = f$ and $gu = u$ when such compositions are defined.
 - For morphisms from $u \rightarrow u'$, the collection of arrows f so that uf is defined and fu' is also defined (and necessarily both equal to f).
 - For composition, $g \circ f = gf$

First of all, composition is well-defined since if $uf, fu', u'g$, and gu'' are defined for identity arrows $u, u', u'',$ then by axiom (ii) $fu'g$ is defined and equal to $(fu')g = fg$ since $fu' = f$.

Now by axiom (iii), any identity u has uu' defined for some other identity u' . But then by the definition of identities,

$$u = uu' = u'$$

It follows that $uu = u$ so that u is a morphism in \mathbf{C} from $u \rightarrow u$ with $u \circ f = f$ and $g \circ u = g$ for any g, f where such equations are defined. It follows that each object u has a two-sided identity in \mathbf{C} .

Associativity follows directly from axiom (i). □

Kyle: Skipped □

