

Bounded step walks on irreducibles in quadratic orders

Yongxian Zhang

School of Computer Science and Engineering
South China University of Technology

mxymmxym1@gmail.com

ORCID: 0009-0000-3864-3536

8 October 2026

Abstract

For every order in a quadratic number field, every fixed full planar lattice embedding, and every fixed step bound, we give a proof that the bounded-step graph on irreducible elements has uniformly bounded component cardinalities. The argument transfers the geometric and entropy method of OpenAI's *Bounded-Step Walks on Gaussian Primes* to an explicit five-hypothesis lattice interface. Ray class fields supply principal split primes in nonmaximal orders. A bounded-norm isolation lemma and a graph restoration lemma handle the infinitely many associates that occur in real quadratic fields. The conclusion includes $\mathbb{Z}[\sqrt{2}]$ in coefficient or Minkowski coordinates. The analytic scale choices are not quantified.

Attribution and verification status. The geometric sampling, signed-prime separation, entropy enrichment, coverage transfer, multiscale schedule and information telescope adapt OpenAI's Gaussian-prime manuscript [1], especially Sections 3–8. They are not presented as independently invented methods. The extension is the arithmetic interface for every quadratic order and restoration of the bounded-norm exceptional elements. The principal all-order conclusion has now been proved in Lean without unproved arithmetic bridge assumptions. The readable proof below uses standard class field theory and Chebotarev. The formal proof instead closes a weaker Dirichlet-supply route sufficient for the same principal theorem; its strong natural-density/PNT target is not asserted. These are two sufficient routes to the same conclusion.

1 Statement and scope

Let $K/\mathbb{Q}$ be a quadratic number field and let $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K$ be an order, with conductor $f \geq 1$. Fix a real-linear isomorphism from its two-dimensional coefficient space to $\mathbb{R}^2$; write ι for its restriction to $\mathcal{O}$ and $\Lambda = \iota(\mathcal{O})$. Thus Λ is a full lattice. An irreducible is a nonzero nonunit that has no factorization into two nonunits of $\mathcal{O}$. No assumption of unique factorization is made.

For $D \geq 0$, let $G_D(\mathcal{O}, \iota)$ have vertex set $\text{Irr}(\mathcal{O})$, joining distinct x, y when $\|\iota(x) - \iota(y)\| \leq D$.

Theorem 1.1 (Quadratic order moat theorem). *For every quadratic order $\mathcal{O}$, every embedding ι as above, and every finite $D \geq 0$, there is a finite constant $B_{\mathcal{O}, \iota, D}$ such that every connected component of $G_D(\mathcal{O}, \iota)$ has at most $B_{\mathcal{O}, \iota, D}$ vertices. Consequently, every injective walk on irreducibles with steps at most D has at most $B_{\mathcal{O}, \iota, D}$ terms.*

All fixed norms on the two-dimensional real coefficient space are equivalent, so the same conclusion holds for any such norm. In a real quadratic field one may use the Minkowski embedding $x \mapsto (\sigma_1(x), \sigma_2(x))$ or an integral-basis coefficient embedding. A *single* real embedding $K \hookrightarrow \mathbb{R}$ does not give a discrete planar lattice and is outside the statement. The theorem has no asserted uniformity as K , f , ι , or D vary, and the analytic argument does not quantify its scale thresholds.

Proof structure. Sections 2–8 prove the abstract finite-sieve statement, including its entropy engine. Section 9 constructs the requisite residue maps. Section 10 restores the exceptional irreducibles and completes the proof. The construction is source-dependent in the precise sense stated above; the Gaussian theorem alone would not imply Theorem 1.1.

2 An arithmetic interface for planar sieving

Let $\Lambda \subset \mathbb{R}^2$ be a full lattice of covolume c_Λ . Let $\mathcal{P}$ be a fixed set of rational primes. For each $p \in \mathcal{P}$, suppose that two surjective additive maps

$$\phi_{p,+}, \phi_{p,-} : \Lambda \longrightarrow \mathbb{F}_p$$

are given. A nonzero lattice vector is *primitive* if it is not av for another lattice vector v and an integer $|a| > 1$. Impose the following fixed-data hypotheses.

(A1) For distinct $p_1, \dots, p_r \in \mathcal{P}$ and arbitrary signs σ_i , the joint map $\Lambda \rightarrow \prod_i \mathbb{F}_{p_i}$ is surjective. Its kernel therefore has index $\prod_i p_i$.

(A2) $\ker \phi_{p,+} \cap \ker \phi_{p,-} = p\Lambda$.

(A3) For a fixed $c > 0$, every nonzero vector in either kernel at p has length at least $c\sqrt{p}$.

(A4) For a fixed $C \geq 1$, every primitive $v \in \Lambda$ and finite $S \subset \mathcal{P}$ satisfy

$$\prod_{p \in S: \phi_{p,+}(v)=0 \text{ or } \phi_{p,-}(v)=0} p \leq C\|v\|^2.$$

(A5) For a fixed $\rho > 0$,

$$\#\{p \in \mathcal{P} : T \leq p \leq 2T\} = (\rho + o(1)) \frac{T}{\log T} \quad (T \longrightarrow \infty).$$

For finite $S \subset \mathcal{P}$, define the periodic avoiding set

$$A(S) = \{x \in \Lambda : \phi_{p,+}(x) \neq 0, \phi_{p,-}(x) \neq 0 \text{ for every } p \in S\}.$$

Theorem 2.1 (Finite planar sieve). *Under (A1)–(A5), for each finite $D \geq 0$ there is a finite $S_D \subset \mathcal{P}$ such that $A(S_D)$ contains no infinite injective D -step walk. Writing $Q = \prod_{p \in S_D} p$, every component of its D -step graph has at most Q^2 vertices.*

A fixed scalar rescaling lets us assume that the shortest nonzero vector of Λ has length at least one. It only changes the fixed constants. The assertion is immediate if $D < 1$, so assume $D \geq 1$. Throughout the proof, constants in $O(\cdot)$ may depend on the fixed interface, lattice and D , but not on the walk, its exact starting time, or the asymptotic parameters unless specified.

The component conclusion follows from the no-walk conclusion. The graph is locally finite; an infinite component has an infinite simple path by the finitely branching path-tree argument. Each map kills $p\Lambda$, so translations by $Q\Lambda$ preserve $A(S_D)$. If two vertices of a finite component differ by nonzero $v \in Q\Lambda$, translating that component by v gives an intersecting component, hence the same component. A nonempty finite set cannot be invariant under a nonzero translation. The component thus injects into $\Lambda/Q\Lambda$, which has Q^2 elements.

3 Geometric separation and entropy conventions

We use natural logarithms. For finite variables, H denotes Shannon entropy and $I(X; Y | Z) = H(X | Z) - H(X | Y, Z)$. Write $h(u) = -u \log u - (1 - u) \log(1 - u)$ and use total variation with its usual factor $1/2$. The following elementary facts will be used repeatedly.

Lemma 3.1 (Finite information estimates). *For finite random times T_0, T_1 on a deterministic D -step lattice walk, if $|T_1 - T_0| \leq n$, then*

$$H(z_{T_1} - z_{T_0}) \leq 2 \log(n + 1) + O(1).$$

For any vector F of additive residue maps,

$$H(F(z_{T_1})) \geq H(F(z_{T_0})) - H(z_{T_1} - z_{T_0}).$$

If two laws on a product alphabet of size at most a are at total variation distance $\varepsilon \leq 1/2$, their conditional entropies differ by at most $2h(\varepsilon) + 2\varepsilon \log a$. Finally, if X has at most a values and a data-dependent list $L(Y)$ obeys $\mathbb{P}\{X \in L(Y), |L(Y)| \leq b\} = u$, then

$$\log a - H(X | Y) \geq u \log(a/b) - h(u).$$

Proof. A displacement lies in a disk of radius Dn , containing $O((n + 1)^2)$ lattice points. The second inequality follows because the final residue and displacement determine the initial residue. For continuity, couple the two laws with mismatch probability ε and apply the entropy bound $h(\varepsilon) + \varepsilon \log a$ to the joint law and to the conditioning marginal. For the list bound, introduce the indicator of the specified success event and use $H(X | Y) \leq h(u) + u \log b + (1 - u) \log a$. $\square$

We also use Pinsker's inequality $\log a - H(X) \geq 2 \text{TV}(\mathcal{L}(X), \text{Unif}_a)^2$, Hoeffding's bounded-sum inequality, and the Chernoff bound for independent Bernoulli trials. These statements are applied only to finite variables. The deterministic walk itself is never randomized: all randomness below comes from finitely supported times, endpoint choices, and auxiliary residue selections.

Lemma 3.2 (Many differences). *Let E be the vertices of a self-avoiding D -step segment of $n \geq 1$ steps. Let R be its diameter, choose a diameter line, and let W be the larger of 1 and the greatest distance from E to that line. Set $A = RW$. Then*

$$n \ll_{\Lambda} A \leq D^2 n^2, \quad |E - E| \gg_{\Lambda, D} A.$$

Proof. The projection of E to a diameter line lies between its endpoints; otherwise one endpoint would be farther than the diameter. Thus E lies in a rectangle of length R and width $2W$, with $R \geq W \geq 1$. Packing fixed disjoint disks around lattice points proves $n + 1 \ll_{\Lambda} RW$; also $R \leq Dn$ and $W \leq R$.

If $W = 1$, then $|E - E| \geq n + 1 \geq R/D$. Otherwise choose diameter endpoints p_0, p_1 and a point p_2 at transverse distance W . Put $a = p_1 - p_0$, $b = p_2 - p_0$. The torus $\mathbb{R}^2/(\mathbb{Z}a + \mathbb{Z}b)$ has area A .

Following the linearly interpolated walk in either time direction gives paths α, γ from p_0 to p_1, p_2 . The map

$$(s, t) \mapsto \alpha(s) - \gamma(t) \pmod{\mathbb{Z}a + \mathbb{Z}b}$$

descends from the parameter square to its torus. An endpoint-preserving homotopy to straight segments shows it has degree -1 , hence is onto: a map missing a point factors through a punctured torus, whose second homology is zero. Each interpolated point is within D of a vertex, so radius- $2D$ disks about projected differences cover the torus. Their total area is at most $4\pi D^2|E - E|$. $\square$

Lemma 3.3 (A randomly signed product separates a thin rectangle). *There are $a, b, A_0 > 0$, depending only on c_Λ, C , with the following property. Take distinct $p_1, \dots, p_r \in \mathcal{P} \cap [T, 2T]$, $T \geq 5$, and set $P_0 = \prod_i p_i$. Let $\mathcal{R}$ be an origin-centered rectangle, in any orientation, with half-lengths $R_1 \geq W_1 \geq 1$. If*

$$R_1 W_1 \leq P_0^{1-d}, \quad 0 < d < \frac{1}{4}, \quad d^3 r \geq A_0,$$

then independent fair signs satisfy

$$\mathbb{P} \left\{ (\mathcal{R} \setminus \{0\}) \cap \bigcap_i \ker \phi_{p_i, \sigma_i} \neq \emptyset \right\} \leq a e^{-b d^3 r}.$$

Proof. Call a sign vector bad when it has a nonzero witness. If two bad vectors differ in $h \leq dr/4$ signs, their witnesses w, w' lie in the common-kernel lattice of the unchanged coordinates. With P_{com} the exact product of those primes, (A1) gives

$$\det(w, w') \in c_\Lambda P_{\text{com}} \mathbb{Z}, \quad P_{\text{com}} \geq \frac{P_0}{(2T)^h}.$$

The rectangle gives $|\det(w, w')| \leq 2R_1 W_1 \leq 2P_0^{1-d}$. The logarithm of the ratio of the nonzero determinant lower bound to this upper bound is at least

$$dr \log T - \frac{dr}{4} \log(2T) + \log(c_\Lambda/2) > 0$$

once A_0 is sufficiently large. Thus the witnesses are collinear. This also proves that each bad vector has a unique witness line.

Fix a line with primitive lattice direction v . By (A2), at most one of the two maps at a prime kills v . If a selected map does not kill v , it kills mv , $m \in \mathbb{Z}$, precisely when $p \mid m$. If $P_\sigma(v)$ is the product of selected primes whose maps kill v , a nonzero common-kernel point on this line has length at least $\|v\| P_0 / P_\sigma(v)$. Since every point of $\mathcal{R}$ has length at most $\sqrt{2} R_1 \leq \sqrt{2} P_0^{1-d}$, a witness forces

$$\log P_\sigma(v) \geq \log \|v\| + d \log P_0 - \frac{1}{2} \log 2.$$

On the other hand, independence, fair signs and (A4) imply

$$\mathbb{E} \log P_\sigma(v) \leq \frac{1}{2} \log(C \|v\|^2) = \log \|v\| + \frac{1}{2} \log C.$$

The required upward deviation is at least $\frac{d}{2} r \log T$ after increasing A_0 . Each summand has range at most $\log(2T)$; Hoeffding gives a fixed-line probability at most $e^{-b_0 d^2 r}$.

It remains to combine lines without counting them. Distinct witness-line classes in the r -dimensional Boolean cube have Hamming separation greater than $dr/4$. Their neighborhoods of radius $h_0 = \lfloor dr/10 \rfloor$ are disjoint. The cube edge-isoperimetric bound yields

$$|\partial_v F| \geq \frac{|F|}{r} \log_2 \frac{2^r}{|F|}.$$

Here the edge bound $|\partial_e F| \geq |F| \log_2(2^r/|F|)$ follows by induction after splitting the cube into two faces, and division by the maximum degree r gives the displayed vertex bound. While an enlarging class has density at most $e^{-b_0 d^2 r/2}$, each enlargement increases its size by a factor at least $1 + b_1 d^2$. After h_0 enlargements this is at least $e^{b_2 d^3 r}$. If the threshold was crossed earlier, its ratio to the initial density, at most $e^{-b_0 d^2 r}$, is already large enough. Summing the disjoint enlarged classes proves the assertion. Fixed covolume and norm constants have only altered the threshold A_0 . $\square$

4 Entropy enrichment

Fix a batch of k primes in $[T, 2T]$ and put $\bar{\ell} = k^{-1} \sum_p \log p$. A signed s -selection chooses s distinct batch primes uniformly and gives each an independent fair sign. Write F_S for the resulting residue vector and, for a finitely supported Λ -valued variable Z , set

$$f_Z(s) = \mathbb{E}_S H(F_S(Z)).$$

Each entropy is computed with S fixed, before averaging; it is not the entropy of a mixture including S . Exchangeability and decreasing conditional entropy imply discrete concavity of f_Z , as well as $f_Z(0) = 0$ and $f_Z(s) \leq s\bar{\ell}$.

Fix an infinite deterministic self-avoiding D -step walk $(z_t)_{t \geq 0}$, and a finite random time τ with $Z = z_\tau$. Given $U > 0$, $0 < g < 10^{-2}$, let $n = \lfloor e^{gU} \rfloor$. At exact time $\tau = t$, sample h uniformly from the *distinct differences* of $E_t = \{z_t, \dots, z_{t+n}\}$. For each difference fix an ordered endpoint pair (X, Y) with $X - Y = h$, then let V be a fair choice between X and Y . This is a forward time transition by at most n steps; its law does not involve residue selections.

Lemma 4.1 (Enrichment). *Set $q = \lfloor U/\bar{\ell} \rfloor$ and $b = \lfloor g^2 U/\bar{\ell} \rfloor$. In any regime with $T \rightarrow \infty$, $q \leq k$, and*

$$\frac{g^j U}{\log T} \longrightarrow \infty \quad \text{for every fixed } j > 0,$$

one has

$$\frac{f_V(b)}{b\bar{\ell}} \geq \frac{1}{2} \left(1 + \frac{f_Z(q)}{q\bar{\ell}} \right) - O(g). \quad (4.1)$$

The error constants and sufficiently-large-parameter thresholds are uniform over the walk, the batch, every exact start, and every finitely supported starting-time law. In particular, $1 \leq b < q$ eventually.

Proof. Choose a signed q -selection Q and a uniform b -prefix S , and set $B_X = F_S(X)$, $B_Y = F_S(Y)$. At exact time t , Lemma 3.2 gives

$$gU - O(1) \leq \log A \leq 2gU + O(1).$$

Let $r = \lceil (1 + 10g) \log A/\bar{\ell} \rceil$. Uniformly in t , the admissible regime gives

$$\frac{1}{2}gq \leq r \leq 3gq \leq q - b$$

for large parameters. Select r further coordinates J from $Q \setminus S$. The sum of their prime logarithms has mean $r\bar{\ell}$ and variance $O(r)$: all logarithms lie in an interval of length $\log 2$, and sampling is without replacement. Hence its probability of being below $(1 + 5g) \log A$ is $O((g^2 r \bar{\ell}^2)^{-1})$.

Differences of differences lie in an origin-centered rectangle whose half-length product is at most $16A$. Lemma 3.3, with d a fixed positive multiple of g , therefore makes F_J injective on $E_t - E_t$, apart from probability

$$\varepsilon = O\left(\frac{1}{g^2 r \bar{\ell}^2} + e^{-b_3 g^3 r}\right) = o(g).$$

The factor 16 is absorbed because $g \log A \rightarrow \infty$; also $g^3 r \rightarrow \infty$ with all needed margins. Since h is uniform on distinct differences, conditioning on B_X, B_Y costs at most $2b \log(2T)$, and

$$\mathbb{E}H(F_J(h) \mid B_X, B_Y, \tau = t) \geq (1 - \varepsilon)(\log A - O(1)) - 2b \log(2T).$$

Conditional subadditivity, divided by r , yields for a uniform remaining coordinate j

$$\mathbb{E}H(\phi_j(h) \mid B_X, B_Y, \tau = t) \geq (1 - O(g))\bar{\ell}. \quad (4.2)$$

Although r depends on t , the one-coordinate marginal is the same uniform marginal at each fixed time. We may therefore average over t and remove conditioning on τ .

For fixed indices, the chain rule and $h = X - Y$ give

$$\begin{aligned} H(\phi_j(h) \mid B_X, B_Y) &\leq H(\phi_j(X) \mid B_X) + H(\phi_j(Y) \mid B_Y) \\ &\quad - H(\phi_j(X) \mid B_X, B_Y, h). \end{aligned}$$

Conditioning on the full displacement h , rather than only $\phi_j(h)$, decreases the subtracted entropy and keeps the inequality in this direction. Subadditivity over the $q - b$ remaining coordinates bounds their average subtracted term below by

$$\frac{f_Z(q) - 2b \log(2T) - H(h) - H(X - Z)}{q - b}.$$

Indeed the full X residue determines the full Z residue when $X - Z$ is given; conditioning additionally on B_X, B_Y, h costs at most their joint entropy. Both displacements are supported in a disk of radius Dn , so

$$H(h) + H(X - Z) \leq 4 \log(n + 1) + O(1) = O(gU).$$

The last lower bound is consequently $f_Z(q)/q - O(g)\bar{\ell}$. The first two terms in the chain-rule inequality average to forward differences of f_X and f_Y at b , each at most $f_X(b)/b$ or $f_Y(b)/b$ by concavity. Finally entropy is concave under the fair endpoint mixture, so $f_V(b) \geq (f_X(b) + f_Y(b))/2$. Combining with (4.2) proves (4.1). $\square$

5 Coverage from shared displacement data

A law on a group of order p has (τ, β) -coverage if fewer than $p^{1-\beta}$ residues have mass less than τ/p . In the following lemma, a coordinate means one of the $2k$ signed batch coordinates, with uniform averaging over them. This step is the abstract probabilistic mechanism of [1, Theorem 5.2].

Lemma 5.1 (Coverage transfer). *Let $T \geq 2$ be real. Let k, s, n be integers with $1 \leq s \leq k$ and $n \geq 1$, and let $0 < \eta \leq 1$, $0 \leq \eta' \leq 1$, and $\varepsilon, \kappa \geq 0$. Let A be finite, $Y = y(A) \in \Lambda$, and, conditional on A , let $h_1, \dots, h_n$ be independent samples of a finitely supported displacement law depending on A . Put $Z = Y + h_1$ and $\mathcal{H} = (h_1, \dots, h_n)$. All auxiliary signed selections are independent of this experiment.*

At each exact $A = a$, suppose all but a fraction η' of signed coordinates have $(\tau + 4\lambda, \beta')$ -coverage for Z . Suppose further that

$$\mathbb{E}_S H(F_S(Y)) \geq (1 - \varepsilon)s\bar{\ell}, \quad H(\mathcal{H}) \leq \kappa s\bar{\ell},$$

and that the following inequalities hold for every batch prime p :

$$0 < \tau \leq \frac{3}{8}, \quad 0 < \lambda \leq \frac{1}{64}, \quad 0 < \beta' \leq \beta < 1, \quad (5.1)$$

$$\eta' \leq \frac{\eta\lambda}{100}, \quad \varepsilon + \kappa \leq \frac{\eta\lambda\beta'}{16}, \quad (5.2)$$

$$p \exp\left(-\frac{1}{2}\lambda^2 np^{-\beta}\right) \leq \lambda/100, \quad (5.3)$$

$$\lambda^{-1} < p^{\beta'/2}, \quad \log(2e/\lambda) \leq \frac{1}{4}\beta' \log T. \quad (5.4)$$

Then all but a fraction η of coordinates have (τ, β) -coverage for the unconditional law of Z .

Proof. Conditioning on $\mathcal{H}$ costs its entropy only once for the selected vector. Subadditivity then gives the shared-data estimate

$$\text{av}_c[\log p_c - H(\phi_c(Y) \mid \mathcal{H})] \leq (\varepsilon + \kappa)\bar{\ell}. \quad (5.5)$$

Suppose instead that a fraction $\theta > \eta$ is deficient. For each deficient coordinate c , choose $u_p = \lceil p^{1-\beta} \rceil$ low-mass residues K_c and define

$$L_c(\mathcal{H}) = \left\{ x : \sum_{\ell=1}^n \mathbf{1}_{x+\phi_c(h_\ell) \in K_c} \leq n(\tau + \lambda)u_p/p \right\}.$$

The true residue $\phi_c(Y)$ is on this list with probability at least $1 - \tau/(\tau + \lambda) \geq \lambda$, by Markov's inequality and the common one-trial marginal.

At exact $A = a$ where conditional coverage holds, let $E_{a,c}$ be the exceptional displacement residues; $|E_{a,c}| < p^{1-\beta'}$. The counting identity

$$\sum_x |(K_c - x) \cap E_{a,c}| = u_p |E_{a,c}|$$

shows that, except for fewer than $\lambda^{-1}p^{1-\beta'}$ values of x , at most λu_p points of $K_c - x$ are exceptional. For every other x , one trial has hit probability at least

$$(\tau + 4\lambda)(1 - \lambda)u_p/p \geq (\tau + 2\lambda)u_p/p.$$

Conditional independence and the Chernoff lower-tail bound put the pass probability below $\exp(-\lambda^2 np^{-\beta}/2)$. Taking a union bound over candidates and using (5.4), the list has at most $p^{1-\beta'/2}$ entries with probability at least $1 - \lambda/100$.

Average first over A , then over deficient coordinates. The fraction of pairs lacking conditional coverage is at most $\eta'/\theta < \lambda/100$. The average probability a_c of simultaneous small-list success is therefore $a = \text{av}_c a_c \geq \lambda/2$. Lemma 3.1 gives

$$\log p_c - H(\phi_c(Y) \mid \mathcal{H}) \geq a_c \frac{\beta'}{2} \log p_c - h(a_c).$$

Concavity gives $\text{av } h(a_c) \leq h(a) \leq a \log(2e/\lambda)$, using $a \geq \lambda/2$. Thus the average deficit over deficient coordinates is at least $\lambda\beta' \log T/8$. Since $\bar{\ell} \leq 2 \log T$ and $\theta > \eta$, this contradicts (5.5) and (5.2). $\square$

6 A finite schedule with uniform margins

We now construct one common sampling law that works simultaneously for finitely many widely separated prime batches. All constants are chosen before the final integer parameter M tends to infinity. This is the multiscale design of [1, Section 6], with the density and lattice constants allowed by (A1)–(A5).

Let $K_D = \#\{v \in \Lambda : \|v\| \leq D\}$, including zero. Set

$$\beta_0 = 1/200, \quad e_0 = \beta_0/50.$$

Choose constants in the following order.

1. By (A5), choose $a_0 > 0$ such that $a_0 T / \bar{\ell} \leq k_T$ for all large T , where $k_T = \#(\mathcal{P} \cap [T, 2T])$; any $a_0 < \rho/2$ works.
2. Choose an integer ℓ_0 with $2^{-\ell_0} < e_0/3$. Choose $g_0 \in (0, 10^{-2})$ small enough that the enrichment and displacement errors in the top-band argument below total less than $2e_0/3$. Those constants depend only on already fixed data.
3. Choose $c_0 > 0$ with $c_0 < a_0 g_0^{2(\ell_0+1)}/2$.
4. Choose a fixed $B > 2$ so large that, for selected geometric batches,

$$\sum_{T' < T} k_{T'} \log(2T') \leq e_0 c_0 T. \quad (6.1)$$

Indeed (A5) gives $k_{T'} \log(2T') \leq C_0 T'$ eventually; summing $T' = B^j < T$ gives at most $C_0 T / (B - 1)$.

Let $M \rightarrow \infty$ through integers, $X_m = 100^m$, and use all batches at $T = B^j$ for which

$$X_m \leq \log T \leq 1.05 X_m \quad \text{for some } \lceil M/2 \rceil \leq m \leq M.$$

These batches are disjoint since $B > 2$. Put $T_+ = \exp(1.05 X_M)$. From any exact starting time, perform the enrichment transitions of Section 4 in decreasing U , using these three bands in each window m :

$$\begin{aligned} 0.8 X_m \leq \log U \leq 1.3 X_m : \quad & g = g_0, \\ 0.20 X_m \leq \log U \leq 0.30 X_m : \quad & g = g_1 = e^{-M^3}, \\ 0.05 X_m \leq \log U \leq 0.10 X_m : \quad & g = g_2 = e^{-100 M^{20}}. \end{aligned}$$

Within a band, successive U have ratio g^2 ; take the geometric progression down from its upper endpoint while inside the band. Each transition uses fresh randomness conditional on the exact current time. Finally add an independent uniform offset in $\{0, \dots, N - 1\}$, with $N = \lceil T_+^{10} \rceil$, and call the terminal time t_* . This entire schedule is independent of all residue selections. Although the time support is very large, it is finite.

6.1 Displacement support and smoothing

At a checkpoint just before scale V in a band with parameter g , every remaining forward time displacement is at most

$$S_V = N - 1 + \sum_{\text{remaining } U} \lfloor \exp(g_U U) \rfloor.$$

Uniformly over all checkpoints, for large M ,

$$\log(1 + S_V) \leq 2gV. \quad (6.2)$$

To see this, later scales in the same band have $g_U U \leq gV$. Gaps between successive bands are fixed positive multiples of X_m ; even across windows the smallest gap is $0.05X_m - 1.3X_{m-1} = 0.037X_m$. They dominate all parameter ratios, since $100M^{20} = o(X_{\lceil M/2 \rceil})$. There are $\exp(O(M))$ operations. Moreover

$$\log(gV) \geq 0.05X_{\lceil M/2 \rceil} - 100M^{20}, \quad \log N = O(X_M) = o(gV).$$

Taking the logarithm of the sum proves (6.2). Consequently every future physical displacement has entropy $O(gV)$, from one deterministic disk support valid over all exact starts. This unconditional support assertion is stronger than a conditional entropy bound.

The final uniform offset also gives, conditional on any earlier exact checkpoint,

$$\text{TV}(\mathcal{L}(t_*), \mathcal{L}(t_* + a)) \leq T_+^{-9}, \quad 0 \leq a \leq T_+. \quad (6.3)$$

It follows by coupling a uniform interval with its shift, then mixing the earlier time.

6.2 Top-band entropy that persists

For a batch T , begin at the first top scale $U \leq a_0 T$, which is greater than $g_0^2 a_0 T$. After ℓ_0 transitions let V be the next scale and record $q_T = \lfloor V/\bar{\ell} \rfloor$. Adjacent input and output coordinate counts match exactly because the scale ratio is g_0^2 . If d_j is the normalized entropy deficit, Lemma 4.1 gives

$$d_{j+1} \leq \frac{1}{2}d_j + C_1 g_0.$$

Preservation to any later checkpoint loses only $O(g_0 V)$ entropy by (6.2) and Lemma 3.1. The choices above imply

$$f_Z(q_T) \geq (1 - e_0)q_T \bar{\ell}, \quad q_T \bar{\ell} \geq c_0 T \quad (6.4)$$

at every later checkpoint. These scales lie inside the top band for large M . This is where the supply density influences a_0, c_0 , and ultimately the spacing B ; no specified numerical density is needed.

6.3 Middle checkpoints and bottom accuracy

For each batch, put

$$\beta_i = 2^{-i}\beta_0, \quad r_0 = \frac{1}{4}, \quad r_{i+1} = r_i - 2\beta_i.$$

Stop at the least J with $\beta_J \log T \leq M^{20}$. Then $J = O(M)$,

$$M^{20}/2 < \beta_J \log T \leq M^{20}, \quad 0.23 < r_i \leq 0.25.$$

Let A_i be the random checkpoint time just before the first scheduled $U \leq T^{r_i}$. It lies in the middle band at a scale in $(g_1^2 T^{r_i}, T^{r_i}]$. From any exact start at A_i , all displacements to later checkpoints lie in a common finite disk $\mathcal{D}_i$ with

$$\log|\mathcal{D}_i| \leq C_2 T^{r_i}. \quad (6.5)$$

For $i < J$, the first $\lceil M^3 \rceil$ middle transitions after A_i fit before A_{i+1} : they consume $O(M^6)$ logarithmic scale, whereas the available gap is $2\beta_i \log T > 2M^{20}$. At the recorded coordinate count s_i , enrichment and subsequent preservation give, from every exact A_i start,

$$f_{z_{A_{i+1}}}(s_i) \geq (1 - e^{-c_1 M^3}) s_i \bar{\ell}, \quad (6.6)$$

$$s_i \bar{\ell} \geq \exp(r_i \log T - O(M^6)). \quad (6.7)$$

Here one may fix $c_1 > 0$ sufficiently small. The bottom band, starting from exact A_J , has room for $\lceil 200M^{20} \rceil$ transitions, whose total logarithmic scale use is $O(M^{40}) = o(X_m)$. With $g_2 = e^{-100M^{20}}$, the resulting normalized deficit, preserved to t_* , is at most $e^{-90M^{20}}$. Concavity therefore gives

$$f_{z_{t_*}}(1) \geq (1 - e^{-90M^{20}}) \bar{\ell}. \quad (6.8)$$

All invocations of enrichment are admissible uniformly: the smallest U has logarithm a positive multiple of $X_{\lceil M/2 \rceil}$, which dominates every fixed power loss from g_2 and $\log T$.

6.4 Backward coverage induction

Set

$$\tau_i = \frac{3}{8} - 2^{-i-3}, \quad \lambda_i = 2^{-i-6}, \quad \eta_0 = 1/100, \quad \eta_{i+1} = \eta_i \lambda_i / 100.$$

Then $\tau_{i+1} = \tau_i + 4\lambda_i$, $\log(1/\lambda_i) = O(M)$, and $\eta_i \lambda_i \beta_{i+1} \geq e^{-O(M^2)}$.

At an exact A_J start, failure of (τ_J, β_J) -coverage entails total variation from uniform at least $(1 - \tau_J)p^{-\beta_J} \geq c_2 e^{-M^{20}}$. Pinsker and (6.8) show that the fraction of failing signed coordinates is at most

$$C_3 \bar{\ell} e^{-88M^{20}} < \eta_J.$$

For the induction step fix exact A_i , run to $Y = z_{A_{i+1}}$, and repeat the continuation to t_* independently *conditional on the exact next checkpoint time*, using $n_i = \lceil T^{6\beta_i/5} \rceil$ trials. Their shared displacement vector has unconditional entropy at most $C_2 n_i T^{r_{i+1}}$ by (6.5). Relative to (6.7), its cost obeys

$$\kappa_i \leq C_4 \exp\left(-\frac{4}{5}\beta_i \log T + O(M^6)\right) \leq e^{-M^3}.$$

Together with (6.6), this is below $\eta_i \lambda_i \beta_{i+1} / 16$. Also

$$n_i p^{-\beta_i} \geq \frac{1}{2} \exp(M^{20}/5).$$

Thus $\lambda_i^2 n_i p^{-\beta_i} / 2$ dominates $\log p + \log(100/\lambda_i)$, because $\log p = O(100M)$, while the former grows as $\exp(M^{20}/5 - O(M))$. The logarithmic conditions (5.4) follow from $\beta_{i+1} \log T > M^{20}/2$ and $\log(1/\lambda_i) = O(M)$. Lemma 5.1 applies with $(\beta, \beta') = (\beta_i, \beta_{i+1})$ and proves the induction.

We have established the following uniform statement: from every exact A_0 start, at least 99% of signed batch coordinates have terminal mass at least $1/(4p)$ outside fewer than $p^{199/200}$ exceptional residues. Exact-start uniformity permits later posterior mixtures without assuming independence of a checkpoint and its residues.

7 The information cost of avoiding zero

Suppose that the hypothetical infinite walk avoids both zero classes for every selected prime. For each batch choose q_T distinct primes with independent fair signs, independently across batches; let $\mathcal{C}$ denote all choices. These choices are independent of the schedule. Set

$$L_T = 2^{\lfloor \log_2(T^{2/5}) \rfloor}.$$

Let X'_T be the selected T -residues at t_* , let O'_T contain selected smaller-batch residues there, and let $W_L(t)$ be the word of the next L increments after time t .

Proposition 7.1 (A charge for each batch). *There is a constant $c_* > 0$, depending only on the fixed data, such that for every selected batch and all sufficiently large M ,*

$$\frac{\mathbb{E}_{\mathcal{C}} I(X'_T; W_{L_T}(t_*) \mid O'_T)}{L_T} \geq \frac{c_*}{\log T}.$$

Proof. Fix T and abbreviate $L = L_T$. At $A_0(T)$ let X, O be the corresponding current- and smaller-batch residue vectors. Let $s = z_{t_*} - z_{A_0}$ and $W = W_L(t_*)$. Equations (6.4) and (6.1) give

$$\mathbb{E}_{\mathcal{C}} H(X \mid O) \geq (1 - 2e_0)q_T \bar{\ell}, \quad H(s) \leq C_5 T^{1/4}. \quad (7.1)$$

Conditionally on $\mathcal{C}, O, X$, draw

$$m = \left\lceil K_0 \frac{T}{L} \log T \right\rceil$$

independent packages (s_a, W_a) from the original conditional law. For each selected coordinate, test every candidate checkpoint residue x against every package, retaining it exactly when every tested position avoids zero. The true coordinate of X always survives, since the actual walk avoids zero. One may use the L positions at offsets $0, \dots, L-1$ in each package.

The package law conditional on $(O, X, \mathcal{C})$ is a posterior mixture of fresh continuation laws from exact A_0 times. At each such time at most η_0 of all signed coordinates lack the coverage proved above. By the tower property, the expected number of selected coordinates whose posterior probability of good coverage is below $1/2$ is at most $2\eta_0 q_T$. This statement averages over both residue choices and the posterior; it does not assume that the checkpoint and $\mathcal{C}$ remain independent after conditioning on O, X . Call the other coordinates favorable.

For a favorable coordinate, at least half of the posterior weight is on good exact starts. Each good start excludes fewer than $p^{1-\beta_0}$ displacement residues. Averaging these exceptional sets shows that all but at most $4p^{1-\beta_0}$ candidate residues have terminal displacement hit probability at least $\tau_0/(4p)$, with $\tau_0 = 1/4$. To see the factor 4, the total posterior exceptional mass, summed over candidates, is at most $p^{1-\beta_0}$; candidates with exceptional mass above $1/4$ number at most four times this quantity. Each remaining candidate has good, nonexceptional posterior weight at least $1/4$.

Conditional smoothing (6.3) persists under this posterior mixture. For large T each of the L offsets therefore has hit probability at least $\tau_0/(8p)$. Within one package these hit events are disjoint: two hits would give a nonzero vector in the same kernel with length at least $c\sqrt{p}$ by (A3), while its length is at most

$$DL \leq DT^{2/5} < c\sqrt{T} \leq c\sqrt{p}.$$

The vector is nonzero because the walk is self-avoiding. Hence one package rejects every such candidate with probability at least $c_5 L/p$. Conditional independence of the repeated packages gives expected survivor count at a favorable coordinate at most

$$4p^{1-\beta_0} + p \exp(-c_5 Lm/p) \leq 5p^{1-\beta_0}$$

when the fixed K_0 is large enough.

The true residue is supported on its survivor list. Subadditivity, Jensen's inequality for the logarithm, and the full alphabet bound on unfavorable coordinates yield

$$\mathbb{E}_{\mathcal{C}} H(X \mid O, \text{all packages}) \leq (1 - \beta_0/2)q_T \bar{\ell}$$

for large T . The loss from an expected $2\eta_0 q_T$ unfavorable coordinates is harmless here: their lost saving is at most $2\eta_0 \beta_0 q_T \log(2T)$, and the additive $q_T \log 5$ is negligible. From (7.1), the packages thus carry at least $(\beta_0/3)q_T \bar{\ell}$ information about X given O . Their conditional independence given $X, O, \mathcal{C}$ bounds this by $m \mathbb{E}_{\mathcal{C}} I(X; s, W \mid O)$.

Given s , additive translation bijectively changes (X, O) into (X'_T, O'_T) . The chain rule gives

$$\begin{aligned} I(X; s, W \mid O) &\leq H(s) + I(X; W \mid O, s) \\ &= H(s) + I(X'_T; W \mid O'_T, s) \\ &\leq I(X'_T; W \mid O'_T) + 2H(s). \end{aligned}$$

No independence of s is required. Consequently

$$\mathbb{E}_{\mathcal{C}} I(X'_T; W \mid O'_T) \geq \frac{\beta_0 q_T \bar{\ell}}{3m} - 2H(s) \geq c_6 \frac{L}{\log T} - C_6 T^{1/4}.$$

Since $L > T^{2/5}/2$, the final error is $o(L/\log T)$, proving the claim. $\square$

8 A common-law telescope

Fix $\mathcal{C}$ and order the selected batches $T_1 < \dots < T_n$, writing $L_j = L_{T_j}$. Let $Q_j(t)$ be the selected residues through batch j at z_t , with Q_0 empty. Define

$$a_j = \frac{H(W_{L_j}(t_*) \mid Q_{j-1}(t_*))}{L_j}, \quad b_j = \frac{H(W_{L_j}(t_*) \mid Q_j(t_*))}{L_j}.$$

The normalized information charge is $a_j - b_j$. The nondecreasing dyadic L_j divide subsequent lengths. Partition $W_{L_{j+1}}(t_*)$ into L_j -blocks. Its initial Q_j and preceding increments determine each block's starting residue vector. The entropy chain rule, followed by removal of additional conditioning, bounds the entropy of the long word by the sum of block entropies conditioned only on their respective starting residues.

Every block-start offset is at most $T_+^{2/5} \leq T_+$. By (6.3), the joint law of its word and starting residue vector is within T_+^{-9} of the corresponding joint law at t_* . The logarithm of the full residue alphabet is at most

$$\sum_T k_T \log(2T) = O(T_+),$$

by (A5) and geometric spacing, and the word alphabet has logarithm at most $L_j \log K_D = O_D(T_+)$. Lemma 3.1 therefore bounds the conditional entropy error of each block by $C_7 T_+^{-7}$. Importantly, continuity is applied to the *joint* laws, not to separately conditioned laws. Dividing the block sum by L_{j+1} yields

$$a_{j+1} \leq b_j + C_7 T_+^{-7}.$$

There are $O(\log T_+)$ batches. Since $a_1 \leq \log K_D$ and $b_n \geq 0$, telescoping gives, uniformly in $\mathcal{C}$ and the walk,

$$\sum_T \frac{I(X'_T; W_{L_T}(t_*) \mid O'_T)}{L_T} \leq \log K_D + o(1).$$

Averaging and applying Proposition 7.1 forces

$$c_* \sum_T \frac{1}{\log T} \leq \log K_D + o(1). \tag{8.1}$$

For each m , the interval $[X_m, 1.05X_m]$ contains at least $c_7 X_m / \log B$ multiples of $\log B$ for large M . Its contribution to the sum is at least $c_7 / (1.05 \log B)$. There are order M windows, contradicting (8.1) as $M \rightarrow \infty$.

Every threshold used above is uniform in the hypothetical walk and its starting times. Fix one sufficiently large M and let S_D be the union of its finitely many prime batches. No infinite self-avoiding walk can avoid the selected zero classes. The periodic argument in Section 2 supplies the bound Q^2 . This completes the proof of Theorem 2.1.

9 Verification for every quadratic order

The arithmetic construction uses only ideals of the maximal order for class field theory; the final residue kernels are genuinely principal ideals of the possibly nonmaximal order $\mathcal{O}$.

Proposition 9.1 (Principal split prime supply). *Let $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K$. There is a set $\mathcal{P}$ of rational primes of positive dyadic density such that, for every $p \in \mathcal{P}$, there are conjugate elements $\alpha_p, \alpha'_p \in \mathcal{O}$ with $|\mathrm{N}_{K/\mathbb{Q}}(\alpha_p)| = p$. Their principal ideals are distinct and satisfy*

$$\mathcal{O}/\alpha_p \mathcal{O} \simeq \mathbb{F}_p, \quad \mathcal{O}/\alpha'_p \mathcal{O} \simeq \mathbb{F}_p, \quad \alpha_p \mathcal{O} \cap \alpha'_p \mathcal{O} = p\mathcal{O}.$$

Proof. Let R_f/K be the ray class field for the finite modulus $f\mathcal{O}_K$, with no infinite places included. Let $L/\mathbb{Q}$ be its finite normal closure. Choose rational primes splitting completely in L , discarding those dividing $\mathrm{disc}(\mathcal{O})$. Chebotarev, in natural-density form, and the prime number theorem give

$$\#(\mathcal{P} \cap [T, 2T]) \sim \frac{1}{[L : \mathbb{Q}]} \frac{T}{\log T}. \quad (9.1)$$

The class field theory inputs used here are ray class reciprocity and existence, and Chebotarev; see [2, Chapter V, Theorems 3.5, 3.6, 3.23 and Remark 3.26(d)]. Only this fixed-extension asymptotic is needed.

For such a prime, $p\mathcal{O}_K = \mathfrak{p}\mathfrak{p}'$ with distinct degree-one primes, and $\mathfrak{p}$ splits in R_f . Triviality of its ray class gives

$$\mathfrak{p} = \alpha \mathcal{O}_K, \quad \alpha \equiv 1 \pmod{f\mathcal{O}_K}.$$

Initially the congruence is local at primes dividing f . Since $\alpha \mathcal{O}_K$ is an integral ideal, $\alpha \in \mathcal{O}_K$; these local congruences then say globally that $\alpha - 1 \in f\mathcal{O}_K$. Thus $\alpha \in 1 + f\mathcal{O}_K \subset \mathcal{O}$. Its conjugate α' also lies in $\mathcal{O}$, generates $\mathfrak{p}'$, and has the same absolute norm p . No positivity at real embeddings is imposed or needed.

Multiplication by α on the rank-two free abelian group $\mathcal{O}$ has determinant $\mathrm{N}(\alpha)$, so $[\mathcal{O} : \alpha\mathcal{O}] = p$. The quotient, a unital ring of prime order, is $\mathbb{F}_p$. The same holds for α' . Because $p \nmid f$, the natural map

$$\mathcal{O}/p\mathcal{O} \longrightarrow \mathcal{O}_K/p\mathcal{O}_K \simeq \mathbb{F}_p \times \mathbb{F}_p$$

is an isomorphism. For completeness, the index $[\mathcal{O}_K : \mathcal{O}] = f$ is prime to p , so its finite cokernel has invertible multiplication by p ; reducing modulo p gives both injectivity and surjectivity. The image of α vanishes in the $\mathfrak{p}$ factor and not the $\mathfrak{p}'$ factor, because $(\alpha) = \mathfrak{p}$. Hence $\alpha\mathcal{O}$ is contained in the first factor kernel, and equality follows from their common index p . Similarly $\alpha'\mathcal{O}$ is the other kernel. Their intersection is $p\mathcal{O}$. $\square$

For the maximal order ($f = 1$), one may take the ordinary Hilbert class field H_K . It is normal over $\mathbb{Q}$: conjugation preserves the defining maximal everywhere-unramified abelian extension,

including the condition at infinite places. If h_K is the ordinary ideal class number, its degree over $\mathbb{Q}$ is $2h_K$ and the density is $1/(2h_K)$. For nonmaximal orders, the possibly smaller density in (9.1) suffices; no formula in terms of an order class number is asserted.

Proposition 9.2 (The interface holds). *Using the two quotient maps in Proposition 9.1, every fixed planar embedding of $\mathcal{O}$ satisfies (A1)–(A5).*

Proof. Identify $\mathcal{O}$ with Λ . Distinct selected index- p ring ideals at different rational primes are comaximal, since they contain their respective rational primes. The Chinese remainder theorem gives (A1), and Proposition 9.1 gives (A2).

The field norm is a homogeneous real quadratic polynomial on the coefficient space. For a fixed embedding there is $C \geq 1$ such that

$$|N(v)| \leq C\|v\|^2 \quad (v \in \Lambda). \quad (9.2)$$

If a nonzero v is in either selected principal ideal, $p \mid |N(v)|$; the norm is a nonzero integer. Thus $\|v\| \geq C^{-1/2}\sqrt{p}$, proving (A3). For a primitive v , both maps at p cannot kill it, by (A2). Every eligible prime divides the nonzero integer $|N(v)|$, so the product of all distinct eligible primes divides that integer and is at most $C\|v\|^2$. This is (A4). Equation (9.1) is (A5). $\square$

In particular, no ideal factorization theorem for $\mathcal{O}$ itself has been invoked. Using all split prime ideals without ensuring principality in $\mathcal{O}$ would not justify the subsequent irreducible-element argument. Also, the field norm need not be positive definite: only its absolute value and its nonvanishing on nonzero field elements were used.

10 Bounded norm exceptions and component restoration

The finite sieve removes every irreducible except associates of finitely many chosen elements. In a real quadratic order the number of these associates can be infinite, so replacing their set by a finite set is not permissible. The following lemmas address this issue directly.

Lemma 10.1 (Isolation for anisotropic binary forms). *Let q be a nondegenerate rational quadratic form on a rank-two lattice, anisotropic over $\mathbb{Q}$. For $M, R > 0$, put $E_M = \{x \in \Lambda : 0 < |q(x)| \leq M\}$. There are only finitely many ordered pairs of distinct $x, y \in E_M$ with $\|x - y\| \leq R$.*

Proof. There are finitely many nonzero differences $v = y - x \in \Lambda$ of length at most R . After clearing one denominator of q , there are finitely many possible values $a = q(x)$ and $b = q(x + v)$. For fixed (v, a, b) , write the polarization as $B_q(x, v) = q(x + v) - q(x) - q(v)$. The equations are

$$q(x) = a, \quad B_q(x, v) = b - a - q(v).$$

Nondegeneracy makes the second equation an affine rational line. Its direction has a nonzero rational spanning vector w , and anisotropy gives $q(w) \neq 0$. Restricting $q(x) = a$ to the line therefore gives a genuine quadratic polynomial equation, with at most two real solutions. There are finitely many triples (v, a, b) . $\square$

For a quadratic field the norm is anisotropic over $\mathbb{Q}$, since $N(x) = 0$ only when $x = 0$. A sharper statement is available using its multiplicative structure.

Lemma 10.2 (An explicit radius for close bounded-norm pairs). *Use the real Minkowski norm $\|x\|_M = (|\sigma_1 x|^2 + |\sigma_2 x|^2)^{1/2}$ in a real quadratic field, and the complex absolute value in an imaginary quadratic field. If distinct $x, y \in \mathcal{O}$ satisfy*

$$0 < |N(x)|, |N(y)| \leq M, \quad \|x - y\|_M \leq R,$$

then

$$\|x\|_M, \|y\|_M \leq (2M + 2)R.$$

For any fixed planar embedding, the same conclusion holds with right-hand side $\kappa(2M + 2)R$, where κ is the product of the operator norms of the linear change from Minkowski coordinates and its inverse.

Proof. Let $v = y - x \neq 0$ and $t = x/v \in K$. Since $N(v)$ is a nonzero integer,

$$|N(t)| \leq M, \quad \text{Tr}(t) = \frac{N(y) - N(x) - N(v)}{N(v)}, \quad |\text{Tr}(t)| \leq 2M + 1.$$

The conjugates of t are roots of $u^2 - \text{Tr}(t)u + N(t)$. The elementary root bound for a monic polynomial gives $\max_\sigma |\sigma(t)| \leq 1 + \max(|\text{Tr}(t)|, |N(t)|) \leq 2M + 2$. Multiplication by t has operator norm at most this quantity in the stated Minkowski norm, hence $\|x\|_M = \|vt\|_M \leq (2M + 2)R$. Interchanging x, y proves the same bound for y . Equivalence of the two coordinate norms gives the last assertion. $\square$

Lemma 10.3 (Restoring exceptional vertices). *Let a graph G on a subset V of a lattice have edges of length at most D and maximum degree at most Δ . Let $E \subset V$. Suppose every component of $G \setminus E$ has at most B vertices. On E , join two distinct points whenever their distance is at most $D(B + 1)$, and suppose this enlarged exceptional graph has component size at most M_E . Then every component of G has at most*

$$\max\{B, M_E(1 + \Delta B)\}$$

vertices.

Proof. Between successive exceptional vertices on a simple path there is either one edge or a passage through a single component of $G \setminus E$, using at most B intermediate vertices. Their distance is at most $D(B + 1)$. Thus all exceptional vertices of a G -component belong to one enlarged exceptional component and number at most M_E . At most ΔM_E nonexceptional components can meet them, each contributing at most B vertices. A component with no exceptional vertex already has size at most B . $\square$

Proof of Theorem 1.1. Apply Proposition 9.2 and Theorem 2.1, obtaining S_D and $Q = \prod_{p \in S_D} p$. Put $B = Q^2$. If S_D is empty, every irreducible is in the avoiding set and the conclusion is immediate. Otherwise let E consist of irreducibles divisible in $\mathcal{O}$ by one of $\alpha_p, \alpha'_p, p \in S_D$. Each chosen element has absolute norm p , so is a nonunit. If it divides an irreducible, the other factor must be a unit. Thus every element of E is an associate of one of these finitely many generators.

A unit of $\mathcal{O}$ has norm ± 1 , since multiplication by it is an invertible integral matrix. Consequently, with $M = \max_{p \in S_D} p$,

$$E \subset \{x \in \mathcal{O} : 0 < |N(x)| \leq M\}.$$

Every irreducible outside E is in $A(S_D)$, so each component after deleting E has at most B vertices. The graph degree is at most $\Delta = \#\{v \in \Lambda : 0 < \|v\| \leq D\} < \infty$.

By Lemma 10.1, the exceptional graph with edge radius $D(B + 1)$ has only finitely many edges, hence a uniform component bound M_E . More concretely, Lemma 10.2 puts both endpoints of every such edge in the finite lattice ball of radius

$$R_* = \kappa(2M + 2)D(Q^2 + 1).$$

One may take M_E to be the larger of 1 and the number of lattice points in that ball. All exceptional vertices outside it are isolated. Lemma 10.3 gives the finite bound

$$B_{O, \iota, D} = \max\{Q^2, M_E(1 + \Delta Q^2)\}.$$

This expression establishes uniformity over all starting irreducibles, but the analytic argument has not assigned numerical values to its scale choices. $\square$

Corollary 10.4. *For each fixed D , the coefficient-plane graph on irreducibles $a + b\sqrt{2} \in \mathbb{Z}[\sqrt{2}]$, with distance $\sqrt{(a - c)^2 + (b - d)^2}$ between $a + b\sqrt{2}$ and $c + d\sqrt{2}$, has uniformly bounded component sizes. The same is true in its two-real-embedding Minkowski metric.*

11 Context and limits of the present draft

The Gaussian case is the result and proof source of OpenAI [1]. Prasad [3] studies computations and walks in class-number-one imaginary quadratic fields. Li, Miller, Popescu, Sarnecki and Wattanawanichkul [4] formulate the corresponding bounded-walk question in $\mathbb{Z}[\sqrt{2}]$ and prove a restricted asymptote-strip result. Prime-free discs, such as those studied by Khale [5], give a different kind of obstruction: a planar path can go around a hole, so holes alone do not establish this component conclusion.

The all-order statement here is proposed as a source-dependent extension. A limited literature check found these predecessors but does not establish priority. A broader literature review remains appropriate; formal verification of the principal conclusion is complete, and external human peer review is not claimed. No claim is made about higher-degree number fields, a single real embedding, a practical closed-form moat size, or bounds uniform over orders. The separate algorithmic certificate supplements are outside this preprint and its formal coverage.

Source snapshot. The inherited proof was read at repository commit

adc7f1241b42e322a6451854ab7e4b4c146bf78a.

The source PDF has SHA-256

a032c3f94f247e90de8077457439a0aad38325ae6f2557c2d08bfb63bfae78d4.

Its source family is 028. The accompanying Gaussian formal statement concerns uniform component bounds and bounded injective walks on Gaussian irreducibles; the analogous all-quadratic-order principal theorem now has the complete public formalization specified below.

12 Formal verification and reproduction

The principal theorem has a complete Lean proof. The precise scope is: All quadratic orders, full planar real-linear embeddings, every fixed finite step bound; uniformly bounded component cardinalities and no injective infinite bounded-step walk. No effective numerical bound or strong

prime-ideal PNT is claimed as a formalized result. The public proof source snapshot is fixed at commit 02af7db99db789fecb0d5f5f8f37b1a4e40762a0 in <https://github.com/mxym/math>. The proof packages, with their own dependency locks, audit records and reproduction instructions, are:

- `formalizations/quadratic-orders-main`.

The main formal entry points are:

- `Entry002.arithmeticSupply_mainTarget_proved`.

Lean is fixed to version 4.34.1, compiler commit 5045d0056413266e57c625dcd7c365b10e377c52; Mathlib is fixed to d13f23b723b8a846827a245b89c10fc7d3f11612. The audited mathematical roots depend only on the standard axioms `propext`, `Classical.choice`, and `Quot.sound`. The archived audits state their actual compilation and replay scopes; publication of this typeset version does not imply a new execution of every previous empty-kernel replay. The companion README maps each printed principal result to its formal theorem and explains how to repeat verification.

Research and preparation disclosures

This research, formalization and manuscript preparation used AI assistance. The author is responsible for the manuscript. The principal claims are supported by the readable proofs and the public Lean artifacts specified above. Formal verification is not external human peer review; no such review is claimed. No external funding was received. No worldwide priority or journal acceptance is asserted.

References

- [1] OpenAI, *Bounded-Step Walks on Gaussian Primes*, 26 September 2026. Pinned manuscript in the OpenAI math repository. See Sections 3–8 for the inherited geometric and entropy method, and the family 028 scope note for the source’s formal artifacts.
- [2] J. S. Milne, *Class Field Theory*, version 4.03, 2020. Author’s course notes. Chapter V, Theorems 3.5 and 3.6 (reciprocity and existence), Theorem 3.23 and Remark 3.26(d) (Chebotarev and natural density); Introduction, Theorem 0.4 (Hilbert class field).
- [3] S. Prasad, *Walks on Primes in Imaginary Quadratic Fields*, 2014. arXiv:1412.2310.
- [4] B. Li, S. J. Miller, T. Popescu, D. Sarnecki and N. Wattanawanichkul, *Modeling Random Walks to Infinity on Primes in $\mathbb{Z}[\sqrt{2}]$* , 2020, revised 2022. arXiv:2011.07386.
- [5] T. Khale, *Prime-free discs in imaginary quadratic fields*, 2025. arXiv:2510.14006.