

PMATH 351: Real Analysis

Instructor Stephen New
 \LaTeX er Iris Jiang

Spring 2020

Contents

1	Cardinality	2
2	Metric Spaces	12

1. Cardinality

Definition 1.1 — domain, range, image, inverse image.

Let X and Y be sets and let $f : X \rightarrow Y$. Recall the **domain** of f and the **range** of f are the sets

$$\text{Domain}(f) = X, \text{Range}(f) = f(X) = \{f(x) | x \in X\}$$

for $A \subseteq X$, the **image** of A under f is the set

$$f(A) = \{f(x) | x \in A\}$$

For $B \subseteq Y$, the **inverse image** of B under f is the set

$$f^{-1}(B) = \{x \in X | f(x) \in B\}$$

Definition 1.2 — Composite.

Let X, Y and Z be sets, let $f : X \rightarrow Y$ and let $g : Y \rightarrow Z$. We define the **composite** function $(g \circ f)(x) = g(f(x))$ for all $x \in X$

Definition 1.3 — injective, surjective, bijective.

We say that f is **injective** (or **one-to-one**) when for every $y \in Y$ there exists **at most** one $x \in X$ such that $f(x) = y$. Equivalently, f is injective when for all $x_1, x_2 \in X$, if $f(x_1) = f(x_2)$ then $x_1 = x_2$.

We say that f is **surjective** (or **onto**) when for every $y \in Y$ there exists **at least** one $x \in X$ such that $f(x) = y$. Equivalently, f is surjective when $\text{Range}(f) = Y$

We say that f is **bijective** (or **invertible**) when f is both injective and surjective, that is when for every $y \in Y$ there exists exactly one $x \in X$ such that $f(x) = y$. When f is both injective and surjective, that is when for every $y \in Y$ there exists exactly one $x \in X$ such that $f^{-1} : Y \rightarrow X$ such that for all $y \in Y$, $f^{-1}(y)$ is equal to the unique element $x \in X$ such that $f(x) = y$. Note that when f is bijective so is f^{-1} , and in this case we have $(f^{-1})^{-1} = f$

Theorem 1.1 Let $f : X \rightarrow Y$ and let $g : Y \rightarrow Z$. Then

- (1) If f and g are both injective then so is $g \circ f$
- (2) If f and g are both surjective then so is $g \circ f$
- (3) If f and g are both invertible then so is $g \circ f$, and in this case $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$

Proof.

- (1) Suppose that f and g are both injective. Let $x_1, x_2 \in X$. If $g(f(x_1)) = g(f(x_2))$ then since g is injective we have $f(x_1) = f(x_2)$, and then since f is injective we have $x_1 = x_2$. Thus $g \circ f$ is injective.
- (2) Suppose that f and g are both surjective. Given $z \in Z$, since g is surjective we can choose $y \in Y$ so that $g(y) = z$, then since f is surjective we can choose $x \in X$ so that $f(x) = y$, and then we have $g(f(x)) = g(y) = z$. Thus $g \circ f$ is surjective.
- (3) Follows (1) and (2). ■

Definition 1.4 — identity function.

For a set X , we define the **identity function** on X to be the function $I_X : X \rightarrow X$ given by $I_X(x) = x$ for all $x \in X$. Note that for $f : X \rightarrow Y$ we have $f \circ I_X = f$ and $I_Y \circ f = f$.

Definition 1.5 — inverse.

Let X and Y be sets and let $f : X \rightarrow Y$. A **left inverse** of f is a function $g : Y \rightarrow X$ given by $g \circ f = I_X$. Equivalently, a function $g : Y \rightarrow X$ is a left inverse of f when $g(f(x)) = x$ for all $x \in X$.

A **right inverse** of f is a function $h : Y \rightarrow X$ such that $f \circ h = I_Y$. Equivalently, a function $h : Y \rightarrow X$ is a right inverse of f when $f(h(y)) = y$ for all $y \in Y$.

Theorem 1.2 Let X and Y be nonempty sets and let $f : X \rightarrow Y$. Then

- (1) f is injective $\iff f$ has a left inverse.
- (2) f is surjective $\iff f$ has a right inverse.
- (3) f is bijective $\iff f$ has a left inverse g and a right inverse h , and in this case we have $g = h = f^{-1}$.

Proof.

- (1) Suppose first that f is injective. Since $X \neq \emptyset$ we can choose $a \in X$ and then define $g : Y \rightarrow X$ as follows: if $y \in \text{Range}(f)$ then (using the fact the f is injective) we define $g(y)$ to be the unique element $x_y \in X$ with $f(x_y) = y$, and if $y \notin \text{Range}(f)$, then we define $g(y) = a$. Then for every $x \in X$ we have $y = f(x) \in \text{Range}(f)$, so $g(y) = x_y = x$, that is $g(f(x)) = x$. Conversely, if f has a left inverse, say g , then f is injective since for all $x_1, x_2 \in X$, if $f(x_1) = f(x_2)$ then $x = g(f(x_1)) = g(f(x_2)) = x_2$.
- (2) Suppose first that f is onto. For each $y \in Y$, choose $x_y \in X$ with $f(x_y) = y$, then define $g : X \rightarrow Y$ by $g(y) = x_y$ (We need the Axiom of Choice for this). Then g is a right inverse of f since for every $y \in Y$ we have $f(g(y)) = f(x_y) = y$. Conversely, if f has a right inverse, say g , then f is onto since given any $y \in Y$ we can choose $x = g(y)$ and then we have $f(x) = f(g(y)) = y$.
- (3) Suppose first that f is bijective. The inverse function $f^{-1} : Y \rightarrow X$ is a left inverse for f because given $x \in X$ we can let $y = f(x)$ and then $f^{-1}(y) = x$ so that $f^{-1}(f(x)) = f^{-1}(y) = x$. Similarly, f^{-1} is a right inverse for f because given $y \in Y$ we can let x be the unique element in X with $y = f(x)$ and then we have $x = f^{-1}(y)$ so that $f(f^{-1}(y)) = f(x) = y$. Conversely, suppose that g is a left inverse for f and h

is a right inverse for f . Since f has a left inverse, it is injective by (1). Since f has a right inverse, it is surjective by (2). Since f is injective and surjective, it is bijective. As shown above, the inverse function f^{-1} is both a left inverse and a right inverse. Finally, note that $g = f^{-1} = h$ because for all $y \in Y$ we have

$$g(y) = g(f(f^{-1}(y))) = f^{-1}(y) = f^{-1}(f(h(y))) = h(y)$$

■

Corollary 1.3

Let X and Y be sets. Then there exists an injective map $f : X \rightarrow Y$ if and only if there exists a surjective map $g : Y \rightarrow X$.

Proof. Suppose $f : X \rightarrow Y$ is an injective map. Then f has a left inverse. Let g be a left inverse of f . Since $g \circ f = I_X$, we see that f is a right inverse of g . Since g has a right inverse, g is surjective. Thus, there is a surjective map $g : Y \rightarrow X$. Similarly, if $g : Y \rightarrow X$ is surjective, then it has a right inverse $f : X \rightarrow Y$ which is injective. ■

Definition 1.6 — same cardinality, less than or equal to, less than.

Let A and B be sets. We say that A and B have the **same cardinality**, and write $|A| = |B|$, when there exists a bijective map: $f : A \rightarrow B$ (or equivalently when there exists a bijective map $g : B \rightarrow A$).

We say that the cardinality of A is **less than or equal to** the cardinality of B , and write $|A| \leq |B|$, when there exists an injective map $f : A \rightarrow B$ (or equivalently a surjective map $g : B \rightarrow A$).

We say that the cardinality of A is **less than** the cardinality of B , and write $|A| < |B|$, when $|A| \leq |B|$ and $|A| \neq |B|$, (that is when there exists an injective map $f : A \rightarrow B$ but there does not exist a bijective map $g : A \rightarrow B$).

We also write $|A| \geq |B|$ when $|B| \leq |A|$; and $|A| > |B|$ when $|B| < |A|$.

■ **Example 1.1** Let $\mathbb{N} = \{n \in \mathbb{Z} | n \geq 0\} = \{0, 1, 2, \dots\}$.

- (1) The map $f : \mathbb{N} \rightarrow 2\mathbb{N}$ given by $f(k) = 2k$ is bijective, so $|2\mathbb{N}| = |\mathbb{N}|$.
- (2) The map $g : \mathbb{N} \rightarrow \mathbb{Z}$ given by $g(2k) = k$ and $g(2k+1) = -k-1$ for $k \in \mathbb{N}$ is bijective, so we have $|\mathbb{Z}| = |\mathbb{N}|$.
- (3) The map $h : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ given by $h(k, l) = 2^k(2l+1) - 1$ is bijective, so we have $|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$.

Theorem 1.4 For all sets A , B and C

- (1) $|A| = |A|$
- (2) If $|A| = |B|$ then $|B| = |A|$
- (3) If $|A| = |B|$ and $|B| = |C|$, then $|A| = |C|$
- (4) $|A| \leq |B| \iff (|A| = |B| \text{ or } |A| < |B|)$
- (5) If $|A| \leq |B|$ and $|B| \leq |C|$, then $|A| \leq |C|$

Proof.

- (1) holds because the identity function $I_A : A \rightarrow A$ is bijective.
- (2) holds because if $f : A \rightarrow B$ is bijective then so is $f^{-1} : B \rightarrow A$.
- (3) holds because if $f : A \rightarrow B$ and $g : B \rightarrow C$ are bijective then so is the composite $g \circ f : A \rightarrow C$

■

Definition 1.7 — finite, infinite, countable.

Let A be a set. For each $n \in \mathbb{N}$, let $S_n = \{0, 1, 2, \dots, n-1\}$. For $n \in \mathbb{N}$, we say that the cardinality of A is equal to n , or that A has n **elements**, and we write $|A| = n$, when $|A| = |S_n|$.

We say that A is **finite** when $|A| = n$ for some $n \in \mathbb{N}$. We say A is **infinite** when A is not finite. We say that A is **countable** when $|A| = |\mathbb{N}|$.

Note 1.1 When a set A is finite with $|A| = n$, and when $f : A \rightarrow S_n$ is a bijection, if we let $a_k = f^{-1}(k)$ for each $k \in S_n$ then we have $A = \{a_0, a_1, \dots, a_{n-1}\}$ with the elements a_k distinct. Conversely, if $A = \{a_0, a_1, \dots, a_{n-1}\}$ with the elements a_k all distinct, then we define a bijection $f : A \rightarrow S_n$ by $f(a_k) = k$. Thus we see that A is finite with $|A| = n$ if and only if A is of the form $A = \{a_0, a_1, \dots, a_{n-1}\}$ with the elements a_k all distinct. Similarly, a set A is countable if and only if A is of the form $A = \{a_0, a_1, a_2, \dots\}$ with the elements a_k all distinct.

Note 1.2 For $n \in \mathbb{N}$, if A is a finite set with $|A| = n + 1$ and $a \in A \setminus \{a\} = n$. Indeed, if $A = \{a_0, a_1, \dots, a_n\}$ with the elements a_i distinct, and if $a = a_k$ so that we have $A \setminus \{a\} = \{a_0, a_1, \dots, a_{k-1}, a_{k+1}, \dots, a_n\}$, then we can define a bijection $f : S_n \rightarrow A \setminus \{a\}$ by $f(i) = a_i$ for $0 \leq i < k$ and $f(i) = a_{i+1}$ for $k \leq i < n$.

Theorem 1.5 Let A be a set. Then the following are equivalent:

- (1) A is infinite
- (2) A contains a countable subset
- (3) $|\mathbb{N}| \leq |A|$
- (4) There exists a map $f : A \rightarrow A$ which is injective but not surjective

Proof.

- (1) \implies (2) Suppose A is infinite. Since $A \neq \emptyset$ we can choose an element $a_0 \in A$. Since $A \neq \{a_0\}$ we can choose an element $a_1 \in A \setminus \{a_0\}$. Since $A \neq \{a_0, a_1\}$ we can choose $a_2 \in A \setminus \{a_0, a_1\}$. Continue this procedure: having chosen distinct elements $a_0, a_1, \dots, a_{n-1} \in A$, since $A \neq \{a_0, a_1, \dots, a_{n-1}\}$ we can choose $a_n \in A \setminus \{a_0, a_1, \dots, a_{n-1}\}$. In this way we obtain $\{a_0, a_1, a_2, \dots\} \subseteq A$.
- (2) \iff (3) Suppose that A contains a countable subset, say $\{a_0, a_1, a_2, \dots\} \subseteq A$ with the element a_i distinct. Since a_i are distinct, the map $f : \mathbb{N} \rightarrow A$ given by $f(k) = a_k$ is injective, and so we have $|\mathbb{N}| \leq |A|$. Conversely as a map from $\mathbb{N} \rightarrow f(\mathbb{N})$ where f is bijective, so we have $|\mathbb{N}| = |f(\mathbb{N})|$ hence $f(\mathbb{N})$ is a countable subset of A .
- (2) \implies (4) Suppose that A has a countable subset, say $\{a_0, a_1, a_2, \dots\} \subseteq A$ with the element a_i distinct. Define $f : A \rightarrow A$ by $f(a_k) = a_{k+1}$ for all $k \in \mathbb{N}$ and by $f(b) = b$ for all $b \in A \setminus \{a_0, a_1, a_2, \dots\}$. Then f is injective but not surjective (the element a_0 is not in the range of f).
- (4) \implies (1) To prove this we shall prove that if A is finite then every injective map $f : A \rightarrow A$ is surjective. We prove this by induction on the cardinality of A .
The only set A with $|A| = 0$ is the set $A \neq \emptyset$, and then the only function $f : A \rightarrow A$ is the empty function, which is surjective.
Since that base case may appear too trivial, let us consider the next case. Let $n = 1$ and let A be a set with $|A| = 1$, say $A = \{a\}$. The only function $f : A \rightarrow A$ is the function given by $f(a) = a$, which is surjective.
Let $n \geq 1$ and suppose, inductively, that for every set A with $|A| = n$, every injective

map $f : A \rightarrow A$ is surjective. Let B be a set with $|B| = n + 1$ and let $g : B \rightarrow B$ be injective.

Suppose, for a contradiction, that g is not surjective. Choose an element $b \in B$ which is not in the range of g so that we have $g : B \rightarrow B \setminus \{b\}$. Let $A = B \setminus \{b\}$ and let $f : A \rightarrow A$ be given by $f(x) = g(x)$ for all $x \in A$. Since $g : B \rightarrow A$ is injective and $f(x) = g(x)$ for all $x \in A$, f is also injective. Again since g is injective, there is no element $x \in B \setminus \{b\}$ with $g(x) = g(b)$, so there is no element $x \in A$ with $f(x) = g(b)$, and so f is not surjective. Since $|A| = n$, this contradicts the induction hypothesis. Thus g must be surjective.

By the Principle of Induction, for every $n \in \mathbb{N}$ and for every set A with $|A| = n$, every injective function $f : A \rightarrow A$ is surjective. ■

Corollary 1.6

Let A and B be sets.

- (1) If A is countable then A is infinite
- (2) When $|A| \leq |B|$, if B is finite so is A (equivalently if A is infinite then so is B)
- (3) If $|A| = n$ and $|B| = m$ then $|A| = |B|$ if and only if $n = m$
- (4) If $|A| = n$ and $|B| = m$ then $|A| \leq |B|$ if and only if $n \leq m$
- (5) When one of the two sets A and B is finite, if $|A| \leq |B|$ and $|B| \leq |A|$ then $|A| = |B|$

Proof.

- (1) If A is countable then A contains a countable subset (itself), so A is infinite by Theorem 1.5.
- (2) Suppose that $|A| \leq |B|$ and that $|A|$ is infinite. Since A is infinite, we have $|\mathbb{N}| \leq |A|$ (by Theorem 1.5). Since $|\mathbb{N}| \leq |A|$ and $|A| \leq |B|$ we have $|\mathbb{N}| \leq |B|$ (by Theorem 1.4). Since $|\mathbb{N}| \leq |B|$, B is infinite (by Theorem 1.5).
- (3) Suppose that $|A| = n$ and $|B| = m$. If $n = m$ then we have $S_n = S_m$ and so $|A| = |S_n| = |S_m| = |B|$. Conversely, suppose that $|A| = |B|$. Suppose, for a contradiction, that $n \neq m$, say $n > m$, and note that $S_m \subsetneq S_n$. Since $|A| = |B|$ we have $|S_n| = |A| = |B| = |S_m|$ so we must have $n = m$.
- (4) Suppose $|A| = n$ and $|B| = m$. If $n \leq m$ then $S_n \subseteq S_m$ so the inclusion map $I : S_n \rightarrow S_m$ is injective and we have $|A| = |S_n| \leq |S_m| = |B|$. Conversely, suppose that $|A| \leq |B|$ and suppose, for a contradiction, that $n > m$. Since $|A| \leq |B|$ we have $|S_n| = |A| \leq |B| = |S_m|$ so we can choose an injective map $f : S_n \rightarrow S_m$. Since $n > m$ we have $S_m \subsetneq S_n$ so we can consider f as a map $f : S_n \rightarrow S_m$, and this map is injective but not surjective. This contradicts Theorem 1.5, and so $n \leq m$.
- (5) Suppose that one of the two sets A and B is finite, and that $|A| \leq |B|$ and $|B| \leq |A|$. If A is finite then, since $|B| \leq |A|$, (2) implies that B is finite. If B is finite then, since $|A| \leq |B|$, (2) implies that A is finite. Thus, in either case, we see that A and B are both finite. Since A and B are both finite with $|A| \leq |B|$ and $|B| \leq |A|$, we must have $|A| = |B|$ by (3) and (4). ■

Theorem 1.7 Let A be a set. Then $|A| \leq |\mathbb{N}| \iff A$ is finite or countable.

Proof. First we claim that every subset of \mathbb{N} is either finite or countable. Let $A \subseteq \mathbb{N}$ and suppose that A is not finite.

Since $A \neq \emptyset$, we can set $a_0 = \min\{A\}$ (using the Well-Ordering Property of \mathbb{N}). Note that

$\{0, 10, \dots, a_0\} \cap A = \{a_0\}$.

Since $A \neq \{a_0\}$ (so the set $A \setminus \{a_0\}$ is nonempty), we can set $a_1 = \min\{A \setminus \{a_0\}\}$. Then we have $a_0 < a_1$ and $\{0, 1, \dots, a_1\} \cap A = \{a_0, a_1\}$.

Since $A \neq \{a_0, a_1\}$ we can set $a_2 = \min\{A \setminus \{a_0, a_1\}\}$. Then we have $a_0 < a_1 < a_2$ and $\{0, 1, 2, \dots, a_2\} \cap A = \{a_0, a_1, a_2\}$.

We continue the procedure: having chosen $a_0, a_1, \dots, a_{n-1} \in A$ with $a_0 < a_1 < \dots < a_{n-1}$ such that $\{0, 1, \dots, a_{n-1}\} \cap A = \{a_0, a_1, \dots, a_{n-1}\}$. Since $A \neq \{a_0, a_1, \dots, a_{n-1}\}$, we can set $a_n = \min\{A \setminus \{a_0, a_1, \dots, a_{n-1}\}\}$ and then we have $a_0 < a_1 < \dots < a_{n-1} < a_n$ and $\{0, 1, \dots, a_n\} \cap A = \{a_0, a_1, \dots, a_n\}$.

In this way, we obtain a countable set $\{a_0, a_1, a_2, \dots\} \subseteq A$ with $a_0 < a_1 < a_2 < \dots$ with the property that for all $m \in \mathbb{N}$, $\{0, 1, 2, \dots, a_m\} \cap A = \{a_0, a_1, \dots, a_m\}$.

Since $0 \leq a_0 < a_1 < a_2 < \dots$, it follows (by induction) that $a_k \geq k$ for all $k \in \mathbb{N}$. It follows in turn that $A \subseteq \{a_0, a_1, a_2, \dots\}$ because given $m \in A$, since $m \leq a_m$ we have

$$m \in \{0, 1, 2, \dots, m\} \cap A \subseteq \{0, 1, 2, \dots, a_m\} \cap A = \{a_0, a_1, \dots, a_m\}.$$

Thus $A = \{a_0, a_1, a_2, \dots\}$ and the elements a_i are distinct, so A is countable. This proves our claim that every subset of \mathbb{N} is either finite or countable.

Suppose that $|A| \leq |\mathbb{N}|$ and choose an injective map $f : A \rightarrow \mathbb{N}$. Since f is injective, when we consider it as a map $f : A \rightarrow f(A)$, it is bijective, and so $|A| = |f(A)|$. Since $f(A) \subseteq \mathbb{N}$, the previous paragraph shows that $f(A)$ is either finite or countable. If $f(A)$ is finite with $|f(A)| = n$ then $|A| = |f(A)| = |S_n|$, and if $f(A)$ is countable then we have $|A| = |f(A)| = |\mathbb{N}|$. Thus A is finite or countable. ■

Theorem 1.8 Let A be a set. Then

- (1) $|A| < |\mathbb{N}| \iff A$ is finite
- (2) $|\mathbb{N}| < |A| \iff A$ is neither finite nor countable
- (3) if $|A| \leq |\mathbb{N}|$ and $|\mathbb{N}| \leq |A|$ then $|A| = |\mathbb{N}|$

Proof.

(1) By Theorem 1.5

$$\begin{aligned} |A| < |\mathbb{N}| &\iff (|A| \leq |\mathbb{N}| \text{ and } |A| \neq |\mathbb{N}|) \\ &\iff (A \text{ is finite or countable and } A \text{ is not countable}) \\ &\iff A \text{ is finite} \end{aligned}$$

(2) By Theorem 1.7

$$\begin{aligned} |\mathbb{N}| < |A| &\iff (|\mathbb{N}| \leq |A| \text{ and } |\mathbb{N}| \neq |A|) \\ &\iff (A \text{ is not finite and } A \text{ is not countable}) \end{aligned}$$

(3) Suppose that $|A| \leq |\mathbb{N}|$ and $|\mathbb{N}| \leq |A|$. Since $|A| \leq |\mathbb{N}|$, we know that A is finite or countable by Theorem 1.7. Since $|\mathbb{N}| \leq |A|$, we know that A is infinite by Theorem 1.5. Since A is finite or countable and A is not finite, it follows that A is countable. Thus $|A| = |\mathbb{N}|$. ■

Definition 1.8 — at most countable, uncountable.

Let A be a set. When A is countable we write $|A| = \aleph_0$. When A is finite we write $|A| < \aleph_0$. When A is infinite we write $|A| \geq \aleph_0$. When A is either finite or countable we write $|A| \leq \aleph_0$ and we say that A is **at most countable**. When A is neither finite nor

countable we write $|A| > \aleph_0$ and we say that A is **uncountable**.

Theorem 1.9

- (1) If A and B are countable sets, then so is $A \times B$
- (2) If A and B are countable sets, then so is $A \cup B$
- (3) If A_0, A_1, A_2, \dots are countable sets, then so is $\bigcap_{k=0}^{\infty} A_k$
- (4) \mathbb{Q} is countable

Proof.

- (1) Let $A = \{a_0, a_1, a_2, \dots\}$ with the a_i distinct and let $B = \{b_0, b_1, b_2, \dots\}$ with b_i distinct. Since every positive integer can be written uniquely in the form $2^k(2l+1)$ with $k, l \in \mathbb{N}$, the map $f : A \times B \rightarrow \mathbb{N}$ given by $f(a_k, b_l) = 2^k(2l+1) - 1$ is bijective, and so $|A \times B| = |\mathbb{N}|$
- (2) Similar to (1), since the map $g : \mathbb{N} \rightarrow A \cup B$ given by $g(k) = a_k$ is injective, we have $|\mathbb{N}| \leq |A \cup B|$. Since the map $h : \mathbb{N} \rightarrow A \cup B$ given by $h(2k) = a_k$ and $h(2k+1) = b_k$ is surjective, we have $|A \cup B| \leq |\mathbb{N}|$. Since $|\mathbb{N}| \leq |A \cup B|$ and $|A \cup B| \leq |\mathbb{N}|$, we have $|A \cup B| = |\mathbb{N}|$ by Theorem 1.8
- (3) For each $k \in \mathbb{N}$, let $A_k = \{a_{k0}, a_{k1}, a_{k2}, \dots\}$ with the a_{ki} distinct. Since the map $f : \mathbb{N} \rightarrow \bigcap_{k=0}^{\infty} A_k$ given by $f(k) = a_{0,k}$ is injective, $|\mathbb{N}| \leq \left| \bigcap_{k=0}^{\infty} A_k \right|$. Since $\mathbb{N} \times \mathbb{N}$ is countable by (1), and since the map $g : \mathbb{N} \times \mathbb{N} \rightarrow \bigcap_{k=0}^{\infty} A_k$ given by $g(k, l) = a_{k,l}$ is surjective, we have $\left| \bigcap_{k=0}^{\infty} A_k \right| \leq |\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$. By Theorem 1.8, we have $\left| \bigcap_{k=0}^{\infty} A_k \right| = |\mathbb{N}|$.
- (4) Since the map $f : \mathbb{N} \rightarrow \mathbb{Q}$ given by $f(k) = k$ is injective, we have $|\mathbb{N}| \leq |\mathbb{Q}|$. Since the map $g : \mathbb{Q} \rightarrow \mathbb{Z} \times \mathbb{Z}$ given by $g(\frac{a}{b}) = (a, b)$ for all $a, b \in \mathbb{Z}$ with $b > 0$ and $\gcd(a, b) = 1$, is injective, and since $\mathbb{Z} \times \mathbb{Z}$ is countable, we have $|\mathbb{Q}| \leq |\mathbb{Z} \times \mathbb{Z}| = |\mathbb{N}|$. Since $|\mathbb{N}| \leq |\mathbb{Q}|$ and $|\mathbb{Q}| \leq |\mathbb{N}|$, we have $|\mathbb{Q}| = |\mathbb{N}|$

■

Exercise 1.1 Let A be a countable set. Show that the set of finite sequences with terms in A is countable. Show that the set of all finite subsets of A is countable.

Definition 1.9 — power set.

For a set A , let $\mathcal{P}(A)$ denote the **power set** of A , that is the set of all subsets of A , and let 2^A denote the set of all functions from A to $S_2 = \{0, 1\}$

Theorem 1.10

- (1) For every set A , $\mathcal{P}(A) = |2^A|$
- (2) For every set A , $|A| < \mathcal{P}(A)$
- (3) \mathbb{R} is uncountable

Proof.

- (1) Let A be any set. Define a map $g : \mathcal{P}(A) \rightarrow 2^A$ as follows: given $S \in \mathcal{P}(A)$, that is given $S \subseteq A$, we define $g(S) \in 2^A$ to be the map $g(S) : A \rightarrow \{0, 1\}$ given by

$$g(S)(a) = \begin{cases} 1 & \text{if } a \in S, \\ 0 & \text{if } a \notin S. \end{cases}$$

Define map $h : 2^A \rightarrow \mathcal{P}(A)$ as follows: given $f \in 2^A$, that is given a map: $f : A \rightarrow \{0, 1\}$, we define $h(f) \in \mathcal{P}(A)$ to be the subset

$$h(f) = \{a \in A \mid f(a) = 1\} \subseteq A$$

This maps g and h are the inverses of each other because for every $S \subseteq A$ and every $f : A \rightarrow \{0, 1\}$ we have

$$\begin{aligned} f = g(S) &\iff \forall a \in A, f(a) = g(S)(a) \iff \forall a \in A, f(a) = \begin{cases} 1 & \text{if } a \in S, \\ 0 & \text{if } a \notin S. \end{cases} \\ &\iff \forall a \in A, (f(a) = 1 \iff a \in S) \iff \{a \in A \mid f(a) = 1\} = S \\ &\iff h(f) = S \end{aligned}$$

- (2) Let A be any set. Since the map $f : A \rightarrow \mathcal{P}(A)$ given by $f(a) = \{a\}$ is injective, we have $|A| \leq |\mathcal{P}(A)|$. We need to show that $|A| \neq |\mathcal{P}(A)|$. Let $g : A \rightarrow \mathcal{P}(A)$ be any map. Let $S = \{a \in A \mid a \notin g(a)\}$. Note that S cannot be in the range of g because we could choose $a \in A$ so that $g(a) = S$ then, by the definition of S , we would have

$$a \in S \iff a \notin g(a) \iff a \notin S$$

which is impossible. Since S is not in the range of g , the map g is not surjective. Since g was an arbitrary map from A to $\mathcal{P}(A)$, it follows that there is no surjective map from A to $\mathcal{P}(A)$. Thus there is no bijective map from A to $\mathcal{P}(A)$ and so we have $|A| \neq |\mathcal{P}(A)|$.

- (3) We prove \mathbb{R} is uncountable using the fact that every real number has a unique decimal expansion which does not end with an infinite string of 9's. Define a map $g : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ as follows: given $f \in 2^{\mathbb{N}}$, that is given a map $f : \mathbb{N} \rightarrow \{0, 1\}$, we define $g(f)$ to be the real number of $g(f) \in [0, 1)$ with the decimal expansion $g(f) = 0.f(1)f(2)f(3)\dots$, that is $g(f) = \sum_{k=0}^{\infty} f(k)10^{-k-1}$. By the uniqueness of decimal expansions, the map g is injective, so we have $|2^{\mathbb{N}}| \leq |\mathbb{R}|$. Thus $|\mathbb{N}| < |\mathcal{P}(\mathbb{N})| = |2^{\mathbb{N}}| \leq |\mathbb{R}|$, and so \mathbb{R} is uncountable by Theorem 1.8. ■

Theorem 1.11 — Cantor-Schroeder-Bernstein.

Let A and B be sets. Suppose that $|A| \leq |B|$ and $|B| \leq |A|$. Then $|A| = |B|$

Proof. We sketch a proof. Choose injective functions $f : A \rightarrow B$ and $g : B \rightarrow A$. Since the functions $f : A \rightarrow f(A)$, $g : B \rightarrow g(B)$ and $f : g(B) \rightarrow f(g(B))$ are bijective, we have $|A| = |f(A)|$ and $|B| = |g(B)| = |f(g(B))|$. Also note that $f(g(B)) \subseteq f(A) \subseteq B$. Let $X = f(g(B))$, $Y = f(A)$ and $Z = B$. Then we have $X \subseteq Y \subseteq Z$ and we have $|x| = |z|$ and we need to show that $|Y| = |Z|$. The composite $h = f \circ g : Z \rightarrow X$ is a bijective. Define sets Z_n and Y_n for $n \in \mathbb{N}$ recursively by

$$Z_0 = Z, Z_n = h(Z_{n-1}) \text{ and } Y_0 = Y, Y_n = h(Y_{n-1})$$

Since $Y_0 = Y$, $Z_0 = Z$, $Z_1 = h(Z_0) = h(Z) = X$ and $X \subseteq Y \subseteq Z$, we have

$$Z_1 \subseteq Y \subseteq Z_0$$

Also note that for $1 \leq n \in \mathbb{N}$,

$$Z_n \subseteq Y_{n-1} \subseteq Z_{n-1} \implies h(Z_n) \subseteq h(Y_{n-1}) \subseteq h(Z_{n-1}) \implies Z_{n+1} \subseteq Y_n \subseteq Z_n$$

By the Induction Principle, it follows that $Z_n \subseteq Y_{n-1} \subseteq Z_{n-1}$ for all $n \geq 1$, so we have

$$Z_0 \supseteq Y_0 \supseteq Z_1 \supseteq Y_1 \supseteq Z_2 \supseteq Y_2 \supseteq \cdots$$

Let $U_n = \frac{Z_n}{Y_n}$, $U = \bigcup_{n=0}^{\infty} U_n$ and $V = \frac{Z}{U}$. Define $H : Z \rightarrow Y$ by

$$H(x) = \begin{cases} h(x) & \text{if } x \in U \\ x & \text{if } x \in V \end{cases}$$

Verify that H is bijective. ■

Exercise 1.2 Show that $|\mathbb{R}| = |2^{\mathbb{N}}|$

Solution. $g : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ as follows: for $f \in 2^{\mathbb{N}}$ we let $g(f)$ be the real number $g(f) \in [0, 1]$ with decimal expansion $g(f) = 0.f(1)f(2)\cdots$. Then g is injective so $|2^{\mathbb{N}}| \leq |\mathbb{R}|$. Define $h : 2^{\mathbb{N}} \rightarrow [0, 1]$ as follows: for $f \in 2^{\mathbb{N}}$ let $h(f)$ be the real number $h(f) \in [0, 1]$ with binary expansion $h(f) = 0.f(0)f(1)f(2)\cdots$. Then h is surjective so we have $|[0, 1]| \leq |2^{\mathbb{N}}|$. The map $k : \mathbb{R} \rightarrow [0, 1]$ given by $k(x) = \frac{1}{2} + \frac{1}{\pi} \tan^{-1}(x)$ is injective, so we have $|\mathbb{R}| \leq |[0, 1]|$. Since $|\mathbb{R}| \leq |[0, 1]| \leq |2^{\mathbb{N}}|$ and $|2^{\mathbb{N}}| \leq |\mathbb{R}|$, we have $|\mathbb{R}| = |2^{\mathbb{N}}|$ by the Cantor-Schroeder-Bernstein Theorem (1.11) ■

Notation 1.1 For sets A and B , we write A^B to denote the set of functions $f : B \rightarrow A$

Theorem 1.12 Let A and B be finite sets and let $\mathcal{P}(A)$ is the power set of A (that is the set of all subsets of A). Then

- (1) if A and N are disjoint then $|A \cup B| = |A| + |B|$
- (2) $|A \times B| = |A| \cdot |B|$
- (3) $|A^B| = |A|^{|B|}$
- (4) $|\mathcal{P}| = 2^{|A|}$

Proof. The proof is left as an exercise ■

Theorem 1.13 Let A, B, C and D be sets with $|A| = |C|$ and $|B| = |D|$. Then

- (1) if $A \cap B = \emptyset$ and $C \cap D = \emptyset$ then $|A \cup B| = |C \cup D|$
- (2) $|A \times B| = |C \times D|$
- (3) $|A^B| = |C^D|$

Proof. The proof is left as an exercise ■



It is possible to define certain specific sets called **cardinals** such that for every set A there exists a unique cardinal κ with $|A| = |\kappa|$. We can then define the **cardinality** of a set A to be equal to the unique cardinal κ such that $|A| = |\kappa|$ and, in this case, we define the **cardinality** of the set A to be $|A| = \kappa$. In foundational set theory, the natural numbers are defined, formally, to be equal to the sets $0 = \emptyset$, $1 = \{0\} = \{\emptyset\}$, $2 = \{0, 1\} = \{\emptyset, \{\emptyset\}\}$ and, in general, $n + 1 = n \cup \{n\}$ so that the natural number n is equal to the set that we previously denoted by S_n , that is $n = S_n = \{0, 1, \dots, n-1\}$. The finite cardinals are equal to the natural numbers and the countable cardinal \aleph_0 is equal to the set of natural numbers. The previous theorem allows us to define **arithmetic operations** on cardinals which extend the usual arithmetic operations on the natural numbers. Given cardinals κ and λ we define $\kappa + \lambda$, $\kappa \cdot \lambda$ and κ^λ to be the cardinals such that

$$\kappa + \lambda = |(\kappa \times \{0\}) \cup (\lambda \times \{1\})|$$

- $\kappa \cdot \lambda = |\kappa \times \lambda|$
- $\kappa^\lambda = |\kappa^\lambda|$

Theorem 1.14 Let κ, λ and μ be cardinals. Then

- (1) $\kappa + \lambda = \lambda + \kappa$
- (2) $(\kappa + \lambda) + \mu = \kappa + (\lambda + \mu)$
- (3) $\kappa + 0 = \kappa$
- (4) $\lambda \leq \mu \implies \kappa + \lambda \leq \kappa + \mu$
- (5) $\kappa \cdot \lambda = \lambda \cdot \kappa$
- (6) $(\kappa \cdot \lambda) \cdot \mu = \kappa \cdot (\lambda \cdot \mu)$
- (7) $\kappa \cdot 1 = \kappa$
- (8) $\kappa \cdot (\lambda + \mu) = (\kappa \cdot \lambda) + (\kappa \cdot \mu)$
- (9) $\lambda \leq \mu \implies \kappa \cdot \lambda \leq \kappa \cdot \mu$
- (10) $\kappa^{\lambda+\mu} = \kappa^\lambda \cdot \kappa^\mu$
- (11) $(\kappa^\lambda)^\mu = \kappa^{\lambda \cdot \mu}$
- (12) $(\kappa \cdot \lambda)^\mu = \kappa^\mu \cdot \lambda^\mu$
- (13) $\lambda \leq \mu \implies \kappa^\lambda \leq \kappa^\mu$
- (14) $\kappa \leq \lambda \implies \kappa^\mu \leq \lambda^\mu$

Proof. We sketch a proof for (9) and (11) and leave the rest as an exercise.

- (9) Let A, B and C be sets with $|A| = \kappa, |B| = \lambda$ and $|C| = \mu$ and suppose that $|B| \leq |C|$.

We need to show that $|A \times B| \leq |A \times C|$. Let $f : B \rightarrow C$ be an injective map. Define $F : A \times B \rightarrow A \times C$ by $F(a, b) = (a, f(b))$ then verify that F is injective.

- (11) Let A, B and C be sets with $|A| = \kappa, |B| = \lambda$ and $|C| = \mu$. We need to show $|(A^B)^C| = |A^{B \times C}|$. Define $F : (A^B)^C \rightarrow A^{B \times C}$ by $F(f)(b, c) = f(c)(b)$. Verify that F is bijective with inverse $G : A^{B \times C} \rightarrow (A^B)^C$ given by $G(g)(c)(b) = g(b, c)$

■

Exercise 1.3 Show that $\left| \bigcup_{n=0}^{\infty} \mathbb{R}^n \right| = 2^{\aleph_0}$

Exercise 1.4 Find $|\mathbb{R}^{[0,1]}|$

2. Metric Spaces

Definition 2.1 — inner product, orthogonal, homomorphism, isomorphism.

Let $F = \mathbb{R}$ or \mathbb{C} . Let U be a vector space over F . An **inner product** on U (over F) is function $\langle \cdot, \cdot \rangle : U \times U \rightarrow F$ (meaning that if $u, v \in U$ then $\langle u, v \rangle \in F$) such that for all $u, v, w \in U$ and all $t \in F$ we have

(1) (Sesquilinearity)

$$\begin{aligned}\langle u + v, w \rangle &= \langle u, w \rangle + \langle v, w \rangle, \langle tu, v \rangle = t \langle u, v \rangle \\ \langle u, v + w \rangle &= \langle u, v \rangle + \langle u, w \rangle, \langle u, tv \rangle = \bar{t} \langle u, v \rangle\end{aligned}$$

(2) (Conjugate Symmetry)

$$\langle u, v \rangle = \overline{\langle v, u \rangle}$$

(3) (Positiveness Definition)

$$\langle u, u \rangle \geq 0 \text{ with } \langle u, u \rangle = 0 \iff u = 0$$

For $u, v \in U$, $\langle u, v \rangle$ is called the **inner product** of u with v . We say that u and v are **orthogonal** when $\langle u, v \rangle = 0$. An **inner product space** (over F) is a vector space over F equipped with an inner product. Given two inner product spaces U and V over F , a linear map $L : U \rightarrow V$ is called a **homomorphism** of inner product spaces (or we say that L preserves inner product) when $\langle L(x), L(y) \rangle = \langle x, y \rangle$ for all $x, y \in U$. A bijection homomorphism is called an **isomorphism**.

Definition 2.2 — norm (length).

Let U be an inner product space over $F = \mathbb{R}$ or \mathbb{C} . For $u \in U$, we define the **norm** (or **length**) of u to be

$$\|u\| = \sqrt{\langle u, u \rangle}$$

Theorem 2.1 Let U be an inner product space over $F = \mathbb{R}$ or \mathbb{C} . For $u, v \in U$ and $t \in F$ we have

(1) (Scaling) $\|tu\| = |t|\|u\|$

(2) (Positive Definiteness) $\|u\| \geq 0$ with $\|u\| = 0 \iff u = 0$

(3) $\|u + v\|^2 = \|u\|^2 + 2\operatorname{Re} \langle u, v \rangle + \|v\|^2$

- (4) (Polarization Identity) if $F = \mathbb{R}$ then $\langle u, v \rangle = \frac{1}{4}(\|u + v\|^2 - \|u - v\|^2)$ and if $F = \mathbb{C}$ then $\langle u, v \rangle = \frac{1}{4}(\|u + v\|^2 + i\|u + iv\|^2 - \|u - v\|^2 - i\|u - iv\|^2)$
- (5) (The Cauchy-Schwarz Inequality) $|\langle u, v \rangle| \leq \|u\|\|v\|$ with $|\langle u, v \rangle| = \|u\|\|v\|$ if and only if $\{u, v\}$ is linearly dependent
- (6) (The Triangle Inequality) $|\|u\| - \|v\|| \leq \|u\| + \|v\|$

Proof. The first 4 parts are easy to prove.

- (5) Suppose that $\{u, v\}$ is linearly dependent. Then one of u and v is a multiple of the other, say $v = tu$ with $t \in F$. Then we have $|\langle u, v \rangle| = |\langle u, tu \rangle| = |\bar{t}\langle u, u \rangle| = |t|\|u\|^2 = \|u\|\|tu\| = \|u\|\|v\|$. Next suppose that $\{u, v\}$ is linearly independent. Then $1 \cdot v + t \cdot u \neq 0$ for all $t \in F$, so in particular $v - \frac{\langle v, u \rangle}{\|u\|^2}u \neq 0$. Thus we have

$$\begin{aligned} 0 &< \|v - \frac{\langle v, u \rangle}{\|u\|^2}u\|^2 = \left\langle v - \frac{\langle v, u \rangle}{\|u\|^2}u, v - \frac{\langle v, u \rangle}{\|u\|^2}u \right\rangle \\ &= \langle v, v \rangle - \frac{\overline{\langle v, u \rangle}}{\|u\|^2} \langle v, u \rangle - \frac{\langle v, u \rangle}{\|u\|^2} \langle u, v \rangle + \frac{\langle v, u \rangle}{\|u\|^2} \frac{\overline{\langle v, u \rangle}}{\|u\|^2} \langle u, u \rangle \\ &= \|v\|^2 - \frac{|\langle v, u \rangle|^2}{\|u\|^2} \end{aligned}$$

So that $\frac{|\langle v, u \rangle|^2}{\|u\|^2} < \|v\|^2$ and hence $|\langle u, v \rangle| \leq \|u\|\|v\|$

- (6) Using (3) and (5), and the inequality $|\operatorname{Re}(z)| \leq |z|$ for $z \in \mathbb{C}$ (which follows Pythagoras' Theorem in \mathbb{R}^2), we have

$$\begin{aligned} \|u + v\|^2 &= \|u\|^2 + \operatorname{Re} \langle u, v \rangle + \|v\|^2 \leq \|u\|^2 + 2|\langle u, v \rangle| + \|v\|^2 \\ &\leq \|u\|^2 + 2\|u\|\|v\| + \|v\|^2 = (\|u\| + \|v\|)^2 \end{aligned}$$

Taking the square root on both sides gives $\|u + v\| \leq \|u\| + \|v\|$. Finally note that $\|u\| = \|(u + v) - v\| \leq \|u + v\| + \|-v\| = \|u + v\| + \|v\|$ so that we have $\|u\| - \|v\| \leq \|u + v\|$, and similarly $\|v\| - \|u\| \leq \|u + v\|$, hence $|\|u\| - \|v\|| \leq \|u + v\|$ ■

Definition 2.3 — norm, unit vector, normed linear space, homomorphism, isomorphism.

Let $F = \mathbb{R}$ or \mathbb{C} . Let U be a vector space over F . A **norm** on U is a function $\| \cdot \| : U \rightarrow \mathbb{R}$ (meaning that if $u \in U$ then $\|u\| \in \mathbb{R}$) such that for all $u, v \in U$ and all $t \in F$ we have

- (1) (Scaling) $\|tu\| = |t|\|u\|$
- (2) (Positive Definiteness) $\|u\| \geq 0$ with $\|u\| = 0 \iff u = 0$
- (3) (Triangle Inequality) $\|u + v\| \leq \|u\| + \|v\|$

For $u \in U$ the real number $\|u\|$ is called the **norm** (or **length**) of u , and we say that u is a **unit vector** when $\|u\| = 1$. A **normed linear space** (over F) is a vector space equipped with a norm. Given two normed linear spaces U and V over F , a linear map $L : U \rightarrow V$ is called a homomorphism of normed linear spaces (or we say that L preserves norm) when $\|L(x)\| = \|x\|$ for all $x \in U$. A bijection homomorphism is called an **isomorphism**.

Definition 2.4 — distance.

Let $F = \mathbb{R}$ or \mathbb{C} and let U be a normed linear space over F . For $u, v \in U$, we define the **distance** between u and v to be

$$d(u, v) = \|v - u\|$$

Theorem 2.2 Let U be a normed linear space over $F = \mathbb{R}$ or \mathbb{C} . For all $u, v, w \in U$

- (1) (Symmetry) $d(u, v) = d(v, u)$
- (2) (Positive Definiteness) $d(u, v) \geq 0$ with $d(u, v) = 0 \iff u = v$
- (3) (Triangle Inequality) $d(u, w) \leq d(u, v) + d(v, w)$

Proof. The proof is left as exercise ■

Definition 2.5 — metric, distance, metric space, homomorphism, isomorphism.

Let X be a non-empty set. A **metric** on X is a map $d : X \times X \rightarrow \mathbb{R}$ such that for all $a, b, c \in X$ we have

- (1) (Symmetry) $d(a, b) = d(b, a)$
- (2) (Positive Definiteness) $d(a, b) \geq 0$ with $d(a, b) = 0 \iff a = b$
- (3) (Triangle Inequality) $d(a, c) \leq d(a, b) + d(b, c)$

For $a, b \in X$, $d(a, b)$ is called the **distance** between a and b . A **metric space** is a set X which is equipped with a metric d , and we sometimes denote the metric space by X and sometimes by the pair (X, d) . Given two metric spaces (X, d_X) and (Y, d_Y) , a map $f : X \rightarrow Y$ is called a **homomorphism of metric spaces** (or we say that f is **distance preserving**) when $d_Y(f(a), f(b)) = d_X(a, b)$ for all $a, b \in X$. A bijective homomorphism is called an **isomorphism** or an **isometry**.

Note 2.1 Every inner product space is also a normed linear space, using the induced norm given by $\|u\| = \sqrt{\langle u, u \rangle}$. Every normed linear space is also a metric space, using the induced metric given by $d(u, v) = \|v - u\|$. If U is an inner product space over $F = \mathbb{R}$ or \mathbb{C} then every subspace of U is also an inner product space using (the restriction of) the same inner product used in U . If U is a normed linear space over $F = \mathbb{R}$ or \mathbb{C} then every subspace of U is also a normed linear space using the same norm. If X is a metric space then so is every subset of X using the same metric.

■ **Example 2.1** Let $F = \mathbb{R}$ or \mathbb{C} . The **standard inner product** on F^n is given by

$$\langle u, v \rangle = v * u = \sum_{i=1}^n u_i \overline{v_i}$$

The standard inner product induces the **standard norm** on F^n , which is also called the **2-norm** on F^n , given by

$$\|u\|_2 = \|u\| = \sqrt{\langle u, u \rangle} = \left(\sum_{i=1}^n |u_i|^2 \right)^{\frac{1}{2}}$$

The standard norm on F^n induces the **standard metric** on F^n , given by

$$d_2(u, v) = d(u, v) = \|v - u\| = \left(\sum_{i=1}^n |v_i - u_i|^2 \right)^{\frac{1}{2}}$$

The **1-norm** on F^n is given by

$$\|u\|_1 = \sum_{i=1}^n |u_i|$$

and it induces the **1-metric** on F^n given by $d_1(u, v) = \|v - u\|_1$. The **supremum norm** also called **infinity norm**, on F^n is given by

$$\|u\|_\infty = \max\{|u_1|, |u_2|, \dots, |u_n|\}$$

and it induces the **supremum metric** on \mathbf{F}^n given by $d_\infty(u, v) = \|v - u\|_\infty$

■ **Example 2.2** For $\mathbf{F} = \mathbb{R}$ or \mathbb{C} . We write

$$\begin{aligned}\mathbf{F}^\omega &= \{u = (u_1, u_2, u_3, \dots) \mid \text{each } u_i \in \mathbf{F}\} \\ \mathbf{F}^\infty &= \{u \in \mathbf{F}^\omega \mid \text{there exists } n \in \mathbb{Z}^+ \text{ such that } u_k = 0 \text{ for all } k \geq n\}\end{aligned}$$

Recall that \mathbf{F}^∞ is a countable-dimensional vector space with standard basis $\{e_1, e_2, e_3, \dots\}$ where $e_1 = (1, 0, 0, \dots)$, $e_2 = (0, 1, 0, \dots)$ and so on. The **standard inner product** on \mathbf{F}^∞ is given by

$$\langle u, v \rangle = \sum_{i=1}^{\infty} u_i \overline{v_i}$$

and it induces the **standard norm**, also called the **2-norm**, on \mathbf{F}^∞ given by

$$\|u\|_2 = \sqrt{\langle u, u \rangle} = \left(\sum_{i=1}^{\infty} |u_i|^2 \right)^{\frac{1}{2}}$$

The **1-norm** on \mathbf{F}^∞ is given by

$$\|u\|_1 = \sum_{i=1}^{\infty} |u_i|$$

and it induces the **1-metric** on \mathbf{F}^∞ given by $d_1(u, v) = \|v - u\|_1$. The **supremum norm** also called the **infinity norm**, on \mathbf{F}^n is given by

$$\|u\|_\infty = \max\{|u_1|, |u_2|, \dots, |u_n|\}$$

and it induces the **supremum metric** on \mathbf{F}^n given by $d_\infty(u, v) = \|v - u\|_\infty$

■ **Example 2.3** For $\mathbf{F} = \mathbb{R}$ or \mathbb{C} , the standard inner product, the 1-norm, the 2-norm and the ∞ -norm, which are well defined on the vector space \mathbf{F}^∞ , do not extend naturally to give a well defined inner product or well-defined norms on the vector space \mathbf{F}^ω (because the relevant sums do not necessarily converge). But we can, and do, extend there definitions to various subspaces of \mathbf{F}^ω . We define

$$\begin{aligned}\ell_1(\mathbf{F}) &= \{u \in \mathbf{F}^\omega \mid \sum_{i=1}^{\infty} |u_i| < \infty\} \\ \ell_2(\mathbf{F}) &= \{u \in \mathbf{F}^\omega \mid \sum_{i=1}^{\infty} |u_i|^2 < \infty\} \\ \ell_\infty(\mathbf{F}) &= \{u \in \mathbf{F}^\omega \mid \sup\{|u_1|, |u_2|, \dots\} < \infty\}\end{aligned}$$

Verify that $\ell_1(\mathbf{F})$ is a normed linear space using **1-norm** given by $\|u\|_1 = \sum_{i=1}^{\infty} |u_i|$, hence $\ell_1(\mathbf{F})$ is also a metric space using the **1-metric** $d_1(u, v) = \|v - u\|_1$. Verify that $\ell_\infty(\mathbf{F})$ is a normed linear space using the **supremum norm**, also called the **infinity norm**, given by $\|u\|_\infty = \sup\{|u_1|, |u_2|, \dots\}$, hence $\ell_\infty(\mathbf{F})$ is also a metric space using the **supremum metric** $d_\infty = \|v - u\|_\infty$. Verify that $\ell_2(\mathbf{F})$ is an inner product space using the **standard inner product** given by $\langle u, v \rangle = \sum_{i=1}^{\infty} u_i \overline{v_i}$. The standard inner product on $\ell_2(\mathbf{F})$ induces

the **standard norm**, also called the **2-norm**, on $\ell_2(\mathbf{F})$ given by $\|u\|_2 = \left(\sum_{i=1}^{\infty} |u_i|^2 \right)^{\frac{1}{2}}$ and

the **standard metric**, or the **2-metric**, $d_2(u, v) = \|v - u\|_2$.

Since we shall usually work with the field $\mathbf{F} = \mathbb{R}$, for $p = 1, 2$ or ∞ we shall write

$$\ell_p = \ell_p(\mathbb{R})$$

■ **Example 2.4** For $\mathbf{F} = \mathbb{R}$ or \mathbb{C} and for $a, b \in \mathbb{R}$ with $a \leq b$, we write

$$\begin{aligned}\mathcal{F}([a, b], \mathbf{F}) &= \mathbf{F}^{[a, b]} = \{f : [a, b] \rightarrow \mathbf{F}\} \\ \mathcal{B}([a, b], \mathbf{F}) &= \{f : [a, b] \rightarrow \mathbf{F} \mid f \text{ is bounded}\} \\ \mathcal{C}([a, b], \mathbf{F}) &= \{f : [a, b] \rightarrow \mathbf{F} \mid f \text{ is continuous}\}\end{aligned}$$

Recall that for $f : [a, b] \rightarrow \mathbb{C}$ given by $f = u + iv$ where $u, v : [a, b] \rightarrow \mathbb{R}$, the function f is continuous if and only if both u and v are continuous and, in this case, $\int_a^b f = \int_a^b u + i \int_a^b v$. In the space $\mathcal{C}([a, b], \mathbf{F})$ we have the **1-norm**, the **2-norm**, and the **supremum norm**

$$\begin{aligned}\|f\|_1 &= \int_a^b |f| \\ \|f\|_2 &= \left(\int_a^b |f|^2 \right)^{\frac{1}{2}} \\ \|f\|_\infty &= \sup_{a \leq x \leq b} |f(x)|\end{aligned}$$

The supremum norm also gives a well-defined norm on the space $\mathcal{B}([a, b], \mathbf{F})$. The 2-norm on $\mathcal{C}([a, b], \mathbf{F})$ is induced by the inner product $\mathcal{C}([a, b], \mathbf{F})$ given by

$$\langle f, g \rangle = \int_a^b f \bar{g}$$

Since we shall usually work with the field $\mathbf{F} = \mathbb{R}$ we shall write

$$\mathcal{F}[a, b] = \mathcal{F}([a, b], \mathbb{R}), \quad \mathcal{B}[a, b] = \mathcal{B}([a, b], \mathbb{R}) \text{ and } \mathcal{C} = \mathcal{C}([a, b], \mathbb{R})$$

Ⓡ For $\mathbf{F} = \mathbb{R}$ or \mathbb{C} and for $1 \leq p < \infty$, one can show that we can define a norm on \mathbf{F}^n by

$$\|u\|_p = \left(\sum_{i=1}^n |u_i|^p \right)^{\frac{1}{p}}$$

and we can define a norm on \mathbf{F}^ω or on the space $\ell_\infty(\mathbf{F}) = \{u \in \mathbf{F}^\omega \mid \sum_{i=1}^\infty |u_i|^p < \infty\}$ by

$$\|u\|_p = \left(\sum_{i=1}^\infty |u_i|^p \right)^{\frac{1}{p}}$$

Also, we can define a norm on the space $\mathcal{C}([a, b], \mathbf{F})$ by

$$\|f\|_p = \left(\int_a^b |f|^p \right)^{\frac{1}{p}}$$

■ **Example 2.5** For any set $X \neq \emptyset$, the **discrete metric** on X is given by $d(x, y) = 1$ for all $x, y \in X$ with $x \neq y$ and $d(x, x) = 0$ for all $x \in X$.

Definition 2.6 — open ball, closed ball, punctured ball, bounded.

Let X be a metric space. For $a \in X$ and $0 < r \in \mathbb{R}$, the **open ball**, the **closed ball** and the (open) **punctured ball** in X centered at a of radius r are defined to be the sets

$$\begin{aligned}B(a, r) &= B_X(a, r) = \{x \in X \mid d(x, a) < r\} \\ \bar{B}(a, r) &= \bar{B}_X(a, r) = \{x \in X \mid d(x, a) \leq r\} \\ B^*(a, r) &= B_X^*(a, r) = \{x \in X \mid 0 < d(x, a) < r\}\end{aligned}$$

When the metric on X denoted by d_p with $1 \leq p \leq \infty$, we often write $B(a, r)$, $\overline{B}(a, r)$ and $B^*(a, r)$ as $B_p(a, r)$, $\overline{B}_p(a, r)$ and $B_p^*(a, r)$. For $A \subseteq X$, we say that A is **bounded** when $A \subseteq B(a, r)$ for some $a \in X$ and some $0 < r \in \mathbb{R}$.

Exercise 2.1 Draw a picture of the open balls $B_1(0, 1)$, $B_2(0, 1)$ and $B_\infty(0, 1)$ in \mathbb{R}^2 (using the metrics d_1 , d_2 and d_∞).

Definition 2.7 — open, closed.

Let X be a metric space. For $A \subseteq X$, we say that A is **open** (in X) when for every $a \in A$ there exists $r > 0$ such that $B(a, r) \subseteq A$, and we say that A is **closed** (in X) when its complement $A^c = X \setminus A$ is open in X .

■ **Example 2.6** Let X be a metric space. Show that for $a \in X$ and $0 < r \in \mathbb{R}$, the set $B(a, r)$ is open and the set $\overline{B}(a, r)$ is closed.

Proof. Let $a \in X$ and Let $r > 0$. We claim that $B(a, r)$ is open. We need to show that for all $b \in B(a, r)$ there exists $s > 0$ such that $B(b, s) \subseteq B(a, r)$. Let $b \in B(a, r)$ and note that $d(a, b) < r$. Let $s = r - d(a, b)$ and note that $s > 0$. Let $x \in B(b, s)$, so we have $d(x, b) < s$. Then, by the Triangle Inequality, we have

$$d(x, a) \leq d(x, b) + d(b, a) < s + d(a, b) = r$$

and so $x \in B(a, r)$. This shows that $B(b, s) \subseteq B(a, r)$ and hence $B(a, r)$ is open.

Next we claim that $\overline{B}(a, r)$ is closed, that is $\overline{B}(a, r)^c$ is open. Let $b \in \overline{B}(a, r)^c$, that is let $b \in X$ with $b \notin \overline{B}(a, r)$. Since $b \notin \overline{B}(a, r)$ we have $d(a, b) > r$. Let $s = d(a, b) - r > 0$. Let $x \in B(b, s)$ and not that $d(x, b) < s$. Then, by the Triangle Inequality, we have

$$d(a, b) \leq d(a, x) + d(x, b) < d(a, x) + s$$

and so $d(x, a) > d(a, b) - s = r$. Since $d(x, a) > r$ we have $b \notin \overline{B}(a, r)$ and so $x \in \overline{B}(a, r)^c$. This shows that $B(b, s) \subseteq \overline{B}(a, r)^c$ and it follows that $\overline{B}(a, r)^c$ is open and hence that $\overline{B}(a, r)$ is closed. ■

Theorem 2.3 — Basic Properties of Open Sets.

Let X be a metric space

- (1) The sets \emptyset and X are open in X
- (2) If S is a set of open sets in X then the union $\bigcup S = \bigcup_{U \in S} U$ is open in X
- (3) If S is a finite set of open sets in X then the intersection $\bigcap S = \bigcap_{U \in S} U$ is open in X

Proof. (1) The empty set is open because any statement of the form “for all $x \in \emptyset, F$ ” (where F is any statement) is considered to be true (by convention). The set X is open because given $a \in X$ we can choose any value of $r > 0$ and then $B(a, r) \subseteq X$ by the definition of $B(a, r)$.

(2) Let S be any set of open sets in X . Let $a \in \bigcup S = \bigcup_{U \in S} U$. Choose an open set $U \in S$ such that $a \in U$. Since U is open we can choose $r > 0$ such that $B(a, r) \subseteq U$. Since $U \in S$ we have $U \subseteq \bigcup S$. Since $B(a, r) \subseteq U$ and $U \subseteq \bigcup S$ we have $B(a, r) \subseteq \bigcup S$. Thus $\bigcup S$ is open.

(3) Let S be a finite set of open sets in X . If $S \neq \emptyset$ then we use the convention that $\bigcap S = X$, which is open. Suppose that $S \neq \emptyset$. say $S = \{U_1, U_2, \dots, U_m\}$ where

each U_k is an open set. Let $a \in \bigcap S = \bigcap_{k=1}^m U_k$. For each index k , since $a \in U_k$ we can choose $r_k > 0$ so that $B(a, r_k) \subseteq U_k$. Let $r = \min\{r_1, r_2, \dots, r_m\}$. Then for each index k we have $B(a, r) \subseteq B(a, r_k) \subseteq U_k$. Since $B(a, r) \subseteq U_k$ for every index k , it follows that $B(a, r) \subseteq \bigcap_{k=1}^m U_k = \bigcap S$

■

Theorem 2.4 — Basic Properties of Closed Sets.

Let X be a metric space

- (1) The sets \emptyset and X are closed in X
- (2) If S is a set of closed sets in X then the union $\bigcap S = \bigcap_{U \in S} U$ is open in X
- (3) If S is a finite set of closed sets in X then the intersection $\bigcup S = \bigcup_{U \in S} U$ is open in X

Proof. The proof is left as exercise

■

Definition 2.8 — topology, topology space, metric topology, finer, coarser.

A **topology** on a set X is a set T of subsets of X such that

- (1) $\emptyset \in T$ and $X \in T$
- (2) For every set $S \subseteq T$ we have $\bigcup S \in T$
- (3) For every finite subset $S \subseteq T$ we have $\bigcap S \in T$

A **topology space** is a set X with a topology T . When X is a metric space, the set of all open sets in X is a topology on X , which we call the **metric topology** (or the topology **induced** by the metric). When X is any topological space, the sets in the topology T are called the **open sets** in X and their complements are called the **closed sets** in X . When S and T are both topologies on a set X with $S \subseteq T$, we say that the topology T is **finer** than the topology S , and the topology S is **coarser** than the topology T .

■ **Example 2.7** Show that in \mathbb{R}^n , the metrics d_1, d_2 and d_∞ all induce the same topology

Proof. For $a, x \in \mathbb{R}^n$ we have

$$\max_{1 \leq i \leq n} |x_i - a_i| \leq \left(\sum_{i=1}^n |x_i - a_i|^2 \right)^{\frac{1}{2}} \leq \sum_{i=1}^n |x_i - a_i| \leq n \max_{1 \leq i \leq n} |x_i - a_i|$$

and so

$$d_\infty(a, x) \leq d_2(a, x) \leq d_1(a, x) \leq n d_\infty(a, x).$$

It follows that for all $a \in \mathbb{R}^n$ and $r > 0$ we have

$$B_\infty(a, r) \supseteq B_2(a, r) \supseteq B_1(a, r) \supseteq B_\infty(a, \frac{r}{n}).$$

Thus for $U \subseteq \mathbb{R}^n$, if U is open in \mathbb{R}^n using d_∞ then it is open using d_2 , and if U is open using d_2 then it is open using d_1 , and if U is open using d_1 then it is open using d_∞ . ■

■ **Example 2.8** Show that on the space $\mathcal{C}[a, b]$, the topology induced by the metric d_∞ is strictly finer than the topology induced by the metric d_1

Proof. For $f, g \in \mathcal{C}[a, b]$ we have

$$d_1(f, g) = \int_a^b |f - g| \leq \int_a^b \max_{a \leq x \leq b} |f(x) - g(x)| = (b - a) d_\infty(f, g)$$

It follows that for $f \in \mathcal{C}[a, b]$ and $r > 0$ we have

$$B_\infty(f, r) \subseteq B_1(f, (b - a)r)$$

Thus for $U \subseteq \mathcal{C}[a, b]$, if U is open using d_1 then U is also open using d_∞ , and so the topology induced by the metric d_∞ is finer (or equal to) the topology induced by d_1 .

On the other hand, we claim that for $f \in \mathcal{C}[a, b]$ and $r > 0$, the set $B_\infty(f, r)$ is not open in the topology induced by d_1 . Fix $g \in B_\infty(f, r)$ and let $s > 0$. Choose a bump function $h \in \mathcal{C}[a, b]$ with $h \geq 0$, $\int_a^b h < h$ and $\max_{a \leq x \leq b} h(x) > 2r$. Then we have $g + h \in B_1(g, s)$ but $g + h \notin B_\infty(f, r)$. It follows that $B_\infty(f, r)$ is not open in the topology induced by d_1 , as claimed. ■

■ **Example 2.9** For any set X , the **trivial topology** on X is the topology in which the only open sets in X are the sets \emptyset and X , and the **discrete topology** on X is the topology in which every subset of X is open. Note that the discrete metric on a nonempty set X induces the discrete topology on x .

Definition 2.9 — interior, closure, dense.

Let X be a metric space (or a topological space) and let $A \subseteq X$. The **interior** and the **closure** of A (in X) are the sets

$$A^\circ = \bigcup \{U \subseteq X \mid U \text{ is open, and } U \subseteq A\}$$

$$\overline{A} = \bigcap \{K \subseteq X \mid K \text{ is closed, and } A \subseteq K\}$$

We say that A is **dense** in X when $\overline{A} = X$.

Theorem 2.5 Let X be a metric space (or a topological space) and let $A \subseteq U$

- (1) The interior of A is the largest open set which is contained in A . In other words, $A^\circ \subseteq A$ and A° is open, and for every open set U with $U \subseteq A$ we have $U \subseteq A^\circ$
- (2) The closure of A is the smallest closed set which contains A . In other words, $A \subseteq \overline{A}$ and \overline{A} is closed, and for every closed set K with $A \subseteq K$ we have $\overline{A} \subseteq K$

Proof.

- (1) Let $L = \{U \subseteq X \mid U \text{ is open, and } U \subseteq A\}$. Note that A° is open (by Part 2 of Theorem 2.3 or by Part 2 of Definition 2.8) because A° is equal to the union of L , which is a set of open sets. Also note that $A^\circ \subseteq A$ because A° is equal to the union of L , which is a set of subsets of A . Finally note that for any open set U with $U \subseteq A$ we have $U \in L$ so that $U \subseteq \bigcup L = A^\circ$.
- (2) The proof is similar to (1)

■

Corollary 2.6

Let X be a metric space (or a topological space) and let $A \subseteq X$

- (1) $(A^\circ)^\circ = A^\circ$ and $\overline{\overline{A}} = \overline{A}$
- (2) A is open $\iff A = A^\circ$
- (3) A is closed $\iff A = \overline{A}$

Proof. The proof is left as exercise

■

Definition 2.10 — interior point, limit point, isolated point, boundary point, boundary.

Let X be a metric space and let $A \subseteq X$.

An **interior point** of A is a point $a \in A$ such that for some $r > 0$ we have $B(a, r) \subseteq A$.

A **limit point** of A is a point $a \in X$ such that for every $r > 0$ we have $B^*(a, r) \cap A \neq \emptyset$.

An **isolated point** of A is a point $a \in A$ which is not a limit point of A .

A **boundary point** of A is a point $a \in X$ such that for every $r > 0$ we have $B(a, r) \cap A \neq \emptyset$ and $B(a, r) \cap A^c \neq \emptyset$.

The set of all limit points of A is denoted by A' . The **boundary** of A , is the set of all boundary points of A .

Theorem 2.7 — Properties of Interior, Limit and Boundary Points.

Let X be a metric space and let $A \subseteq X$

- (1) A° is equal to the set of all interior points of A
- (2) A is closed $\iff A' \subseteq A$
- (3) $\overline{A} = A \cup A'$
- (4) $\partial A = \overline{A} \setminus A^\circ$

Proof. We leave the proofs of (1) and (4) as exercise.

- (2) Note that when $a \notin A$ we have $B(a, r) \cap A = \emptyset$ and so

$$\begin{aligned}
 A \text{ is closed} &\iff A^c \text{ is open} \\
 &\iff \forall a \in A^c, \exists r > 0, B(a, r) \subseteq A^c \\
 &\iff \forall a \in \mathbb{R}^n, (a \notin A \implies \exists r > 0, B(a, r) \subseteq A^c) \\
 &\iff \forall a \in \mathbb{R}^n, (a \notin A \implies \exists r > 0, B(a, r) \cap A = \emptyset) \\
 &\iff \forall a \in \mathbb{R}^n, (a \notin A \implies \exists r > 0, B^*(a, r) \cap A = \emptyset) \\
 &\iff \forall a \in \mathbb{R}^n, (\forall r > 0, B^*(a, r) \cap A \neq \emptyset \implies a \in A) \\
 &\iff \forall a \in \mathbb{R}^n, (a \in A' \implies a \in A) \\
 &\iff A' \subseteq A.
 \end{aligned}$$

- (3) We shall prove that $A \cup A'$ is the smallest closed set which contains A . It is clear that $A \cup A'$ is closed, that is $(A \cup A')^c$ is open. Let $a \in (A \cup A')^c$ with, that is let $a \in X$ with $a \notin A$ and $a \notin A'$. Since $a \notin A'$ we can choose $r > 0$ so that $B(a, r) \cap A = \emptyset$. We claim that because $B(a, r) \cap A = \emptyset$ it follows that $B(a, r) \cap A' = \emptyset$. Since $b \in B(a, r)$ and $B(a, r)$ is open, we can choose $s > 0$ so that $B(b, s) \subseteq B(a, r)$. Since $b \in A'$ it follows that $B(b, s) \cap A \neq \emptyset$. Choose $x \in B(b, s) \cap A$. Then we have $x \in B(b, s) \subseteq B(a, r)$ and $x \in A$ and so $x \in B(a, r) \cap A$, which contradicts the fact that $B(a, r) \cap A = \emptyset$. Thus $B(a, r) \cap A' = \emptyset$ as claimed. Since $B(a, r) \cap A = \emptyset$ and $B(a, r) \cap A' = \emptyset$, it follows that $B(a, r) \cap (A \cup A') = \emptyset$, hence $B(a, r) \subseteq (A \cup A')^c$. Thus proves that $(A \cup A')^c$ is open, and hence $A \cup A'$ is closed.

It remains to show that for every closed set K in X with $A \subseteq K$ we have $A \cup A' \subseteq K$. Let K be a closed set in X with $A \subseteq K$. Note that since $A \subseteq K$ it follows that $A' \subseteq K'$ because if $a \in A'$ then for all $r > 0$ we have $B(a, r) \cap A \neq \emptyset$ hence $B(a, r) \cap K \neq \emptyset$ and so $a \in K'$. Since K is closed we have $K' \subseteq K$ by (2). Since $A' \subseteq K'$ and $K' \subseteq K$, we have $A' \subseteq K$. Since $A \subseteq K$ and $A' \subseteq K$ we have $A \cup A' \subseteq K$ as required. ■



Let X be a topological space and let $A \subseteq X$, An **interior point** of A is a point $a \in A^\circ$. A **limit point** of A is a point $a \in X$ such that for every open set U in X with $a \in U$ there

exists a point $b \in U \cap A$ with $b \neq a$. The **boundary** of A in X is the set $\partial A = \bar{A} \setminus A^\circ$, and a **boundary point** of A is a point $a \in \partial A$.

Note 2.2 Let X be a metric space and let $P \subseteq X$. Note that P is also a metric space using (the restriction of) the metric used in X . For $a \in P$ and $0 < r \in \mathbb{R}$, note that the open and closed balls in P , centered at a and of radius r , are related to the open and closed balls in X by

$$B_P(a, r) = \{x \in P \mid d(x, a) < r\} = B_X(a, r) \cap P$$

$$\bar{B}_P(a, r) = \{x \in P \mid d(x, a) \leq r\} = \bar{B}_X(a, r) \cap P$$

Theorem 2.8 Let X be a metric space and let $A \subseteq P \subseteq X$

- (1) A is open in $P \iff$ there exists an open set U in X such that $A = U \cap P$
- (2) A is closed in $P \iff$ there exists a closed set K in X such that $A = K \cap P$

Proof.

- (1) Suppose that A is open in P . For each $a \in A$, choose $r_a > 0$ so that $B_P(a, r_a) \subseteq A$, that is $B_X(a, r_a) \cap P \subseteq A$, and let $U = \bigcup_{a \in A} B_X(a, r_a)$. Since U is equal to the union of a set of open sets in X , it follows that U is open in X . Note that $A \subseteq U \cap P$, and, since $B_X(a, r_a) \cap P \subseteq A$ for every $a \in A$, we also have $U \cap P = (\bigcup_{a \in U} B_X(a, r_a)) \cap P = \bigcup_{a \in A} (B_X(a, r_a) \cap P) \subseteq A$. Thus $A = U \cap P$ as required. Conversely, suppose that $A = U \cap P$ with U open in X . Let $a \in A$. Since we have $a \in A = U \cap P$, we also have $a \in U$. Since $a \in U$ and U is open in X we can choose $r > 0$ so that $B_X(a, r) \subseteq U$. Since $B_X(a, r) \subseteq U$ and $U \cap P = A$ we have $B_P(a, r) = B_X(a, r) \cap P \subseteq U \cap P = A$. Thus A is open as required.
- (2) Suppose that A is closed in P . Let B be the complement of A in P , that is $B = P \setminus A$. Then B is open in P . Choose an open set U in X such that $B = U \cap P$. Let K be the complement of U in X , that is $K = X \setminus U$. Then $A = K \cap P$ since for $x \in X$ we have

$$\begin{aligned} x \in A &\iff (x \in P \text{ and } x \notin U \cap P) \iff (x \in P \text{ and } x \notin U) \\ &\iff (x \in P \text{ and } x \in K) \iff (x \in K \cap P) \end{aligned}$$

Conversely, suppose that K is a closed set in P with $A = K \cap P$. Let B be the complement of A in P , that is $B = P \setminus A$, and let U be the complement of K in P , that is $U = P \setminus K$, and note that U is open in P . Then we have $B = U \cap P$ since for $x \in P$ we have

$$\begin{aligned} x \in B &\iff (x \in P \text{ and } x \notin A) \iff (x \in P \text{ and } x \notin K \cap P) \\ &\iff (x \in P \text{ and } x \notin K) \iff (x \in P \text{ and } x \in U) \iff (x \in U \cap P) \end{aligned}$$

Since U is open in P and $B = U \cap P$ we know that B is open in P . Since B is open in P , its complement $A = P \setminus B$ is closed in P . ■



Let X be a topological space and let $P \subseteq X$. Verify, as an exercise, that we can use the topology on X to define a topology on P as follows. Given a set $A \subseteq P$, we define A to be **open** in P when $A = U \cap P$ for some open set U in X . The resulting topology on P is called the **subspace topology**.