

ALGEBRA I

EDWARD O'CALLAGHAN

CONTENTS

1. Prelude	2
2. Introduction	2
3. Groups	2
3.1. Group Homomorphisms	3
3.2. Properties of homomorphisms	4
3.3. Cosets	4
3.4. Factor (or Quotient) groups	4
3.5. Non-commutative Groups	5
3.6. Group actions	5
3.7. Permutations	5
3.8. Permutation parity	6
3.9. Fields	6
4. Exact sequence	7
5. First Isomorphism Theorem	7
6. Determinants	8
7. Vector Spaces	10
7.1. Subspaces	11
8. Direct Sum	14
9. Invariant Subspaces	15
10. Algebras	17
11. Vector space Homomorphisms	18
11.1. Properties of linear maps	19
12. Dual Spaces	19
13. Normed vector space	20
13.1. Linear Operators	20
14. Inner Product Spaces	20

1. PRELUDE

TODO: Fix notation here...

2. INTRODUCTION

In this course we build up the rudiments of some important notions of algebraic structures. That is, an algebraic structure of an arbitrary set, or carrier set, coupled with various finitary operations defined on it. ..

3. GROUPS

Definition 3.1 (Binary operation). A **binary operation** on a set \mathcal{X} is a map $\circ : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{X}$. **N.B.** that the binary operation is *closed*.

Definition 3.2 (Magma). A **magma** is a set \mathcal{M} equipped with a binary operation \circ . We denote the magma as the tuple pair (\mathcal{M}, \circ) .

Definition 3.3 (Semi-group). A **semi-group** is a set \mathcal{G} equipped with binary operation that is *associative*. Hence, a semi-group is a magma where the operation is *associative*; That is, given any $x, y, z \in \mathcal{G}$ then $x \circ (y \circ z) = (x \circ y) \circ z \in \mathcal{G}$. We denote the semi-group as the tuple pair (\mathcal{G}, \circ) , not to be confused with a magma from context.

Definition 3.4 (Monoid). A **semi-group with identity** or, **monoid** for short, is a semi-group (\mathcal{G}, \circ) with a unique identity element $e \in \mathcal{G}$ such that $x \circ e = x = e \circ x \forall x \in \mathcal{G}$

Proof: uniqueness of identity. Assume some other identity e' exists in \mathcal{G} then, $e' = e' \circ e = e \circ e' = e$. \square

Example 3.5. Given $\mathcal{G} = \mathbb{N}$ with the binary law of composition \circ to be defined as arithmetic addition $+$. Then, $(\mathbb{N}, +)$ forms a semi-group with identity 0. Verify the axioms.

Definition 3.6 (Group). A **group** is a monoid where every element has an inverse. A **abelian group** is a group that is commutative.

Example 3.7. Given $\mathcal{G} = \mathbb{Z}$ with the binary law of composition \circ to be defined as arithmetic addition $+$. Then, $(\mathbb{Z}, +)$ forms a semi-group with identity 0. Verify the axioms.

Question 3.8. Why does the set of naturals \mathbb{N} not form a group under addition, however does form a monoid?

Definition 3.9 (Subgroup). A group \mathcal{H} is a **subgroup** of a group \mathcal{G} if the restriction of the binary operation $\circ : \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ is a group operation on \mathcal{H} . In particular, A non-empty subset \mathcal{H} of a group \mathcal{G} is a subgroup of \mathcal{G} if and only if $h_1 \circ h_2 \in \mathcal{H}$ whenever $h_1, h_2 \in \mathcal{H}$, and $h^{-1} \in \mathcal{H}$ whenever $h \in \mathcal{H}$. We denote the subgroup by $\mathcal{H} \leq \mathcal{G}$.

Definition 3.10 (Smallest subgroup). If \mathcal{A} is a subset of a group \mathcal{G} , there is a *smallest* subgroup $\text{Gp}(\mathcal{A})$ of \mathcal{G} which contains \mathcal{A} , the subgroup *generated* by \mathcal{A} .

Example 3.11. Suppose $\mathcal{A} = \{g\}$ then $\text{Gp}(\mathcal{A}) = \text{Gp}(g)$ and so $\text{Gp}(g) = \{g^n : n \in \mathbb{Z}\}$, where $g^0 = e$, g^n is the product of n copies of g where $n > 0$, and g^n is the product of $|n|$ copies of g^{-1} when $n < 0$.

Definition 3.12 (Cyclic group). A group \mathcal{G} is *cyclic* if $\mathcal{G} = \text{Gp}(g)$ for some $g \in \mathcal{G}$. Such a element is called a *generator* of the group.

Definition 3.13 (Group order). If a group \mathcal{G} has finitely many elements, then the *order* $o(\mathcal{G})$ is the number of elements of \mathcal{G} .

Definition 3.14 (Normal subgroup). A subgroup \mathcal{H} of a group \mathcal{G} is a **normal**, or *self-conjugate*, if $g^{-1}hg \in \mathcal{H} \forall g \in \mathcal{G}$ and $h \in \mathcal{H}$. We denote the normal $\mathcal{H} \trianglelefteq \mathcal{G}$.

Definition 3.15 (Simple group). A group \mathcal{G} is **simple** if it has no normal subgroups other than $\{e\}$ and \mathcal{G} .

3.1. Group Homomorphisms. Homomorphisms are structure preserving mappings. In group homomorphisms we preserve the structure of the binary operation \circ as follows;

Definition 3.16 (Homomorphism). Let \mathcal{G} and \mathcal{H} be two groups. Then a mapping

$$\varphi : \mathcal{G} \rightarrow \mathcal{H}$$

is called a *homomorphism* if

$$\varphi(x \circ y) = \varphi(x) \circ \varphi(y) : x, y \in \mathcal{G}$$

It follows that, for some $g \in \mathcal{G}$ we have,

$$\begin{aligned} \varphi(e_g) &= \varphi(g \circ g^{-1}) \\ &= \varphi(g) \circ \varphi(g^{-1}) \\ &= \varphi(g) \circ (\varphi(g))^{-1} \\ &= e_h \in \mathcal{H}. \end{aligned}$$

That is the identity e has been preserved. Hence, it does not matter if we compose in \mathcal{G} and map to \mathcal{H} or take two elements in \mathcal{G} then compose the mapped elements in \mathcal{H} , since the group structure has been preserved.

How much information about the elements inside the structure is, however, another quality to consider. Hence we fix some terminology here.

- A homomorphism that is injective is called monomorphic.
- A homomorphism that is surjective is called epimorphic.

- A homomorphism that is bijective is called isomorphic.

Thus we have the following definitions by considering a group homomorphism $\varphi : \mathcal{G} \rightarrow \mathcal{H}$.

Definition 3.17 (Monomorphic). φ is **monomorphic** if for $\varphi(x) = \varphi(y) \implies x = y \forall x, y \in \mathcal{G}$.

Definition 3.18 (Epimorphic). φ is **epimorphic** if $\forall h \in \mathcal{H} \exists g \in \mathcal{G}$ so that $\varphi(g) = h$.

Definition 3.19 (Isomorphic). φ is **isomorphic** if φ is **both** mono- and epic- morphic.

Some special cases are sometimes of particular interest and we shall outline them now.

Definition 3.20 (Endomorphic). A monomorphism $\mathcal{G} \rightarrow \mathcal{G}$ for a group \mathcal{G} is called an *endomorphism* of \mathcal{G} .

Definition 3.21 (Automorphic). A isomorphism $\mathcal{G} \rightarrow \mathcal{G}$ for a group \mathcal{G} is called an *automorphism* of \mathcal{G} .

Remark. The set $Aut(\mathcal{G})$ of automorphisms of \mathcal{G} forms a group, when composition of mappings is taken as the group law of composition.

3.2. Properties of homomorphisms.

Definition 3.22 (kernel). If $\varphi : \mathcal{G} \rightarrow \mathcal{H}$ is a group homomorphism, then the *kernel* is the set $\ker(\varphi) = \{g \in \mathcal{G} : \varphi(g) = e_{\mathcal{H}}\}$.

If $\varphi : \mathcal{G} \rightarrow \mathcal{H}$ is a group homomorphism, then observe that $\ker(\varphi)$ is a normal subgroup of \mathcal{G} .

3.3. Cosets. Let \mathcal{G} be a group and \mathcal{H} be a subgroup of \mathcal{G} with $g \in \mathcal{G} : g \notin \mathcal{H}$, then

Definition 3.23 (Left Coset). $g\mathcal{H} = \{gh : h \in \mathcal{H}\}$ is a **left coset of \mathcal{H}** in \mathcal{G} .

Definition 3.24 (Right Coset). $\mathcal{H}g = \{hg : h \in \mathcal{H}\}$ is a **right coset of \mathcal{H}** in \mathcal{G} .

Definition 3.25 (Normal Subgroup). If $g\mathcal{H} = \mathcal{H}g$ then \mathcal{H} is a **normal** subgroup of \mathcal{G} , denoted by $\mathcal{H} \trianglelefteq \mathcal{G}$.

3.4. Factor (or Quotient) groups. Let \mathcal{G} be a commutative group and consider a subgroup \mathcal{H} . Then \mathcal{H} determines an equivalence relation in \mathcal{G} given by

$$x \sim x' \text{ iff } x - x' \in \mathcal{H}.$$

..

3.5. Non-commutative Groups. A common class of non-commutative groups are transformation groups. Note:

Definition 3.26 (Transformation). A bijective map $\varphi : X \rightarrow X$ is called a **transformation** of X .

Note. The most trivial case is the *identity map* id_X by $id_X(x) = x, \forall x \in X$.

Hence, there exists a inverse φ^{-1} of φ such that $\varphi^{-1} \circ \varphi = id_X = \varphi \circ \varphi^{-1}$. Now, take two transformations of X , φ and ψ , and let the product $\varphi \circ \psi$ be well defined. Then the set of all transformations of X form the group **Transf**(X). Since, given $\varphi, \psi, \phi \in Transf(X)$ then we have associativity, $\varphi \circ (\psi \circ \phi) = (\varphi \circ \psi) \circ \phi$. We have identity $e = id_X \in Transf(X)$ and so, inverses $\forall \varphi \in Transf(X) \exists! \varphi^{-1} : \varphi \circ \varphi^{-1} = e$. Closure follows from the composition of two transformations φ and ψ , since $(\varphi \circ \psi)^{-1} = \psi^{-1} \circ \varphi^{-1}$.

A transformation group is a type of group action which describes symmetries of objects. More abstractly, since a group \mathcal{G} is a category with a single object in which every morphism is bijective. Then, a group action is a *forgetful functor* \mathcal{F} from the group \mathcal{G} in the category **Grp** to the set category **Set** that is, $\mathcal{F} : \mathcal{G} \rightarrow \mathbf{Set}$.

3.6. Group actions. For any mathematical object (e.g. sets, groups, vector spaces) X a isomorphism of X is a symmetry of X . The set of all isomorphisms of X , or symmetries of X , form a group called the symmetry group of X , denoted $Sym(X)$. More formally;

Definition 3.27 (Group action). An *action* of a group \mathcal{G} on a mathematical object X is a mapping $\mathcal{G} \times X \rightarrow X$, defined by $(g, x) \mapsto gx$ satisfying:

- $e.x = x \forall x \in X$ and
- $(gh).x = g.(g.x) \forall g, h \in \mathcal{G}, x \in X$.

That is, we have the (*left*) \mathcal{G} -action on X and denote this by $\mathcal{G} \curvearrowright X$.

Notice that we may study properties of the symmetries of some mathematical object X without reference to the structure of X in particular.

3.7. Permutations. Take a finite set X with $|X| = n$, then the transformations of X are called **permutations** of the elements of X . In particular, the group of permutations of $X = \{1, 2, \dots, n\}$ is a **symmetric group**, denoted S_n , with **order** $|S_n| = n!$. Thus, by taking any subgroup of S_n we have a **permutation group**. Also note that, for finite sets, *permutation* and *bijective maps* refer to the same operation, namely rearrangement of elements of X . Another way is to consider, a group \mathcal{G} and set X . Then a group action is defined as a group homomorphism φ from \mathcal{G} to the symmetric group of X . That is, the action $\varphi : \mathcal{G} \rightarrow S_n(X)$, assigns a permutation of X to each element of the group \mathcal{G} in the following way:

- From the identity element $e \in \mathcal{G}$ to the identity transformation id_X of X , that is, $\varphi : e \rightarrow id_X$;

- A product of group homomorphisms $\varphi \circ \psi \in \mathcal{G}$ is then the composite of permutations given by φ and ψ in X .

Given that each element of \mathcal{G} is represented as a permutation. Then a group action can also be consider as a permutation representation.

A permutation $\sigma \in S_n$ can be written,

$$\sigma = \begin{pmatrix} 1 & 2 & \cdots & n \\ a_1 & a_2 & \cdots & a_n \end{pmatrix} \text{ where } a_1 = \sigma(1), a_2 = \sigma(2), \cdots .$$

The identity permutation $id_n \in S_n$ is simply,

$$id_n = \begin{pmatrix} 1 & 2 & \cdots & n \\ 1 & 2 & \cdots & n \end{pmatrix}$$

Since $|S_n| = n!$ then the total number of ways n elements maybe permuted is $n!$.

Take any two permutations $\sigma, \pi \in S_n$ then composition is well defined as **functional composition** as follows.

Given,

$$\sigma = \begin{pmatrix} 1 & 2 & \cdots & n \\ a_1 & a_2 & \cdots & a_n \end{pmatrix} \text{ and } \pi = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ b_1 & b_2 & \cdots & b_n \end{pmatrix}$$

then,

$$\begin{aligned} \pi \circ \sigma &= \begin{pmatrix} 1 & 2 & \cdots & n \\ \pi(a_1) & \pi(a_2) & \cdots & \pi(a_n) \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & \cdots & n \\ b_1 & b_2 & \cdots & b_n \end{pmatrix} \end{aligned}$$

A inverse of any permutation $\sigma \in S_n$ is given by,

$$\sigma^{-1} = \begin{pmatrix} 1 & 2 & \cdots & n \\ a_1 & a_2 & \cdots & a_n \end{pmatrix}^{-1} = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ 1 & 2 & \cdots & n \end{pmatrix}$$

3.8. Permutation parity. Consider the algebraic structure:

$$\triangle_n(x_1, \dots, x_n) = \prod_{i < j} (x_i - x_j)$$

TODO..

3.9. Fields. We now may build higher order algebraic structures using the notion of a group.

Definition 3.28 (Field). A **field** \mathbb{F} is a set together with two binary operations, addition and multiplication, such that:

- addition forms a abelian group,
- multiplication forms a abelian quasi-group, i.e. a commutative multiplicative group on the set $\mathbb{F} - \{0\}$,

coupled together with a law of distribution between the two binary operations.

4. EXACT SEQUENCE

An **exact sequence** may either be a finite or infinite sequence of objects and morphisms between them. Such a sequence is constructed so that the image of one morphism equals the kernel of the next.

In particular;

Definition 4.1 (Exact Sequence). Consider the sequence of n group homomorphism between $n + 1$ groups in the following way:

$$\mathcal{G}_0 \xrightarrow{\varphi_1} \mathcal{G}_1 \xrightarrow{\varphi_2} \mathcal{G}_2 \xrightarrow{\varphi_3} \dots \xrightarrow{\varphi_n} \mathcal{G}_n$$

Then the sequence is said to be *exact* if,

$$\ker(\varphi_{k+1}) = \text{im}(\varphi_k)$$

for every $k \in \{1 \dots n\}$. For $n = 3$ the sequence is said to be a **short exact sequence**.

Example 4.2. Suppose we have $\mathcal{K} \trianglelefteq \mathcal{G}$ and that $q : \mathcal{G} \rightarrow \mathcal{G}/\mathcal{K}$ is the quotient mapping. Then,

$$1 \longrightarrow \mathcal{K} \xrightarrow{\subseteq} \mathcal{G} \xrightarrow{q} \mathcal{G}/\mathcal{K} \longrightarrow 1$$

is a short exact sequence.


5. FIRST ISOMORPHISM THEOREM

Definition 5.1. Let \mathcal{G} and \mathcal{H} , and let $\varphi : \mathcal{G} \rightarrow \mathcal{H}$ be a group homomorphism. Then:

- The kernel of φ is a normal subgroup of \mathcal{G} ; $\ker(\varphi) \trianglelefteq \mathcal{G}$,
- The image of φ is a subgroup of \mathcal{H} ; $\text{im}(\varphi) \leq \mathcal{H}$, and
- The image of φ is also isomorphic to the factor group $\mathcal{G}/\ker(\varphi)$; $\text{im}(\varphi) \cong \mathcal{G}/\ker(\varphi)$.

In particular, if φ is epimorphic then $\mathcal{H} \cong \mathcal{G}/\ker(\varphi)$.

We may represent these fundamental relations in the following commutative diagram.



Notice the *exact sequence* that runs from the lower left to the upper right of the commutative diagram.

6. DETERMINANTS

Definition 6.1 (Determinant).

$$\det(A) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{i=1}^n a_{i,\sigma(i)}.$$

Example 6.2. Find the determinant of some matrix $A \in \mathcal{M}_{2,2}(\mathbb{F})$.

Suppose

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

and take the permutation group

$$S_2 = \{\sigma_1, \sigma_2\}$$

with $\sigma_1 = id$ and $\sigma_2 = [1\ 2]$. Then, by definition, we have:

$$\begin{aligned}
 \det(A) &= \sum_{\sigma \in S_2} \operatorname{sgn}(\sigma) \prod_{i=1}^2 a_{i,\sigma(i)} \\
 &= \operatorname{sgn}(\sigma_1)(a_{1,\sigma_1(1)}a_{2,\sigma_1(2)}) + \operatorname{sgn}(\sigma_2)(a_{1,\sigma_2(1)}a_{2,\sigma_2(2)}) \\
 &= +(a_{11}a_{22}) - (a_{12}a_{21}) \\
 &= a_{11}a_{22} - a_{12}a_{21}.
 \end{aligned}$$

Example 6.3. Find the determinant of some matrix $A \in \mathcal{M}_{3,3}(\mathbb{F})$.

Suppose

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

and take the permutation group

$$S_6 = \{\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5, \sigma_6\}$$

with

$$\sigma_1 = [1\ 2\ 3] \sigma_2 = [1\ 3\ 2]$$

$$\sigma_3 = [2\ 1\ 3] \sigma_4 = [2\ 3\ 1]$$

$$\sigma_5 = [3\ 2\ 1] \sigma_6 = [3\ 1\ 2]$$

Then, by definition, we have:

$$\begin{aligned} \det(A) &= \sum_{\sigma \in S_6} \operatorname{sgn}(\sigma) \prod_{i=1}^6 a_{i, \sigma(i)} \\ &= \operatorname{sgn}(\sigma_1)(a_{1, \sigma_1(1)} a_{2, \sigma_1(2)} a_{3, \sigma_1(3)} a_{4, \sigma_1(4)} a_{5, \sigma_1(5)} a_{6, \sigma_1(6)}) \\ &\quad + \operatorname{sgn}(\sigma_2)(a_{1, \sigma_2(1)} a_{2, \sigma_2(2)} a_{3, \sigma_2(3)} a_{4, \sigma_2(4)} a_{5, \sigma_2(5)} a_{6, \sigma_2(6)}) \\ &\quad + \operatorname{sgn}(\sigma_3)(a_{1, \sigma_3(1)} a_{2, \sigma_3(2)} a_{3, \sigma_3(3)} a_{4, \sigma_3(4)} a_{5, \sigma_3(5)} a_{6, \sigma_3(6)}) \\ &\quad + \operatorname{sgn}(\sigma_4)(a_{1, \sigma_4(1)} a_{2, \sigma_4(2)} a_{3, \sigma_4(3)} a_{4, \sigma_4(4)} a_{5, \sigma_4(5)} a_{6, \sigma_4(6)}) \\ &\quad + \operatorname{sgn}(\sigma_5)(a_{1, \sigma_5(1)} a_{2, \sigma_5(2)} a_{3, \sigma_5(3)} a_{4, \sigma_5(4)} a_{5, \sigma_5(5)} a_{6, \sigma_5(6)}) \\ &\quad + \operatorname{sgn}(\sigma_6)(a_{1, \sigma_6(1)} a_{2, \sigma_6(2)} a_{3, \sigma_6(3)} a_{4, \sigma_6(4)} a_{5, \sigma_6(5)} a_{6, \sigma_6(6)}) \end{aligned}$$

Lemma 6.4.

$$\det(A^T) = \det(A).$$

Proof. Consider some σ such that $\sigma_1, \dots, \sigma_n$ is in order so that $\tau = \sigma^{-1}$ and take the matrix A_j^i so that $(A_j^i)^T = A_i^j$.

Then we see,

$$A_j^{\sigma(i)} = A_{\sigma^{-1}(j)}^j = A_{\tau(j)}^j$$

with $\text{sgn}(\tau) = \text{sgn}(\sigma)$. So,

$$\begin{aligned}
 \det(A) &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i=1}^n a_{i, \sigma(i)} \\
 &= \sum_{\tau \in S_n} \text{sgn}(\tau) \prod_{i=1}^n a_{\tau(i), \tau \cdot \sigma(i)} \\
 &= \sum_{\tau \in S_n} \text{sgn}(\tau) \prod_{i=1}^n a_{\tau(i), i} \\
 &= \det(A^T). \quad \square
 \end{aligned}$$

Lemma 6.5.

$$\det(AB) = \det(A) \det(B).$$

Proof. Let $A = [a_j^i]$ and $B = [b_j^i]$ with $A, B \in \mathcal{M}_{nn}(\mathbb{F})$ so that $AB = [\sum_{k=1}^n a_k^i b_j^k]$.

$$\begin{aligned}
 \det(AB) &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i=1}^n \left[\sum_{k=1}^n a_k^i b_{\sigma(i)}^k \right] \\
 &= .. \quad \square
 \end{aligned}$$

7. VECTOR SPACES

Definition 7.1 (Vector space). A **vector space** \mathcal{V} over a field \mathbb{F} is an abelian additive group $(\mathcal{V}, +)$, coupled with a mapping (scalar multiplication) $(\lambda, x) \rightarrow \lambda x$ of $\mathbb{F} \times \mathcal{V}$ into \mathcal{V} which satisfies

- $1 \cdot x = x$,
- $(\lambda + \mu)(x) = \lambda x + \mu x$,
- $\lambda(\mu x) = (\lambda \mu)x$,
- $\lambda(x + y) = \lambda x + \lambda y$,

for $\lambda, \mu \in \mathbb{F}$ and $x, y \in \mathcal{V}$. We note here that the elements of \mathcal{V} are called *vectors* and that the elements of \mathbb{F} are called *scalars*.

Remark. Note that any field \mathbb{F} forms a degenerate vector space over itself.

Example 7.2. For a set X , the set of functions from $X \rightarrow \mathbb{F}$, denoted $\text{Fun}(X, \mathbb{F})$, is a vector space with pointwise addition and scalar multiplication. That is, for some $\varphi, \psi \in \text{Fun}(X, \mathbb{F})$, $x \in X$ and $\lambda \in \mathbb{F}$ we have,

$$(\varphi + \psi)(x) = \varphi(x) + \psi(x)$$

and

$$(\lambda\varphi)(x) = \lambda(\varphi(x)).$$

7.1. Subspaces.

Definition 7.3. A non-empty subset \mathcal{U} of a vector space \mathcal{V} over \mathbb{F} is a *linear subspace* if it is a subgroup of \mathcal{V} and if $\lambda x \in \mathcal{U}$ whenever $\lambda \in \mathbb{F}$ and $x \in \mathcal{U}$ written $\mathcal{U} \leq \mathcal{V}$.

Corollary. A non-empty subset \mathcal{U} of a vector space \mathcal{V} over a field \mathbb{F} is a linear subspace if and only if, for any $u, u' \in \mathcal{U}$ and $\lambda \in \mathbb{F}$ we have, $u + \lambda u' \in \mathcal{U}$.

Example 7.4. Let the set of all real-valued functions on \mathbb{R}^n be denoted by $\mathcal{F}(\mathbb{R}^n)$. Notice that, for any $f, g \in \mathcal{F}(\mathbb{R}^n)$, $\mathbf{x} \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$ we have, $(f + g)(\mathbf{x}) = f(\mathbf{x}) + g(\mathbf{x})$ and $(\lambda f)(\mathbf{x}) = \lambda f(\mathbf{x})$. That is we claim that, $\mathcal{F}(\mathbb{R}^n)$ is a vector space.

Now, observe that the set of all continuous real-valued functions, denoted $\mathcal{C}(\mathbb{R}^n)$ is a subspace of $\mathcal{F}(\mathbb{R}^n)$.

Example 7.5. Given two subspaces $\mathcal{U}, \mathcal{V} \leq \mathcal{W}$ of vector space \mathcal{W} , then $\mathcal{U} \cap \mathcal{V} \leq \mathcal{W}$.

Proof.

Suppose $x, \lambda x' \in \mathcal{U} \cap \mathcal{V}$ with any $\lambda \in \mathbb{F}$. Then,

$$x, \lambda x' \in \mathcal{U} \text{ and } x, \lambda x' \in \mathcal{V}.$$

Now, since both \mathcal{U} and \mathcal{V} are subspaces then,

$$\begin{aligned} x + \lambda x' &\in \mathcal{U} \text{ and } x + \lambda x' \in \mathcal{V} \\ \implies x + \lambda x' &\in \mathcal{U} \cap \mathcal{V}. \end{aligned}$$

□

Lemma 7.6. In general, for a vector space \mathcal{V} and any family of subspaces $\mathcal{U}_i \leq \mathcal{V}$ for every $i \in I$, we have $\bigcap_{i \in I} \mathcal{U}_i \leq \mathcal{V}$.

Example 7.7. Consider the subspaces $\mathcal{V}', \mathcal{V}'', \mathcal{W} \leq \mathcal{V}$. Prove that

$$(\mathcal{V}' \cap \mathcal{W}) + (\mathcal{V}'' \cap \mathcal{W}) \subseteq (\mathcal{V}' + \mathcal{V}'') \cap \mathcal{W}.$$

Proof.

Pick some

$$x \in (\mathcal{V}' \cap \mathcal{W}) + (\mathcal{V}'' \cap \mathcal{W}) : x = u + v$$

where $u \in \mathcal{V}' \cap \mathcal{W}$ and $v \in \mathcal{V}'' \cap \mathcal{W}$.

Then,

$$u \in \mathcal{V}', u \in \mathcal{W}$$

and

$$v \in \mathcal{V}'', v \in \mathcal{W}.$$

Now, since $u, v \in \mathcal{W}$ and that \mathcal{W} is a subspace,

$$x = u + v \in \mathcal{W}.$$

Also, since

$$u \in \mathcal{V}' \text{ and } v \in \mathcal{V}''$$

then

$$u + v \in \mathcal{V}' + \mathcal{V}''.$$

Hence,

$$x \in \mathcal{W} \cap (\mathcal{V}' + \mathcal{V}'').$$

□

Does the reverse inclusion always hold?

Proof.

Pick some $y \in (\mathcal{V}' + \mathcal{V}'') \cap \mathcal{W}$ so that

$$y \in (\mathcal{V}' + \mathcal{V}'') \text{ or } y \in \mathcal{W}.$$

Now, write $y = y' + y''$ with $y' \in \mathcal{V}'$ and $y'' \in \mathcal{V}''$. Observe that,

$$\begin{aligned} & \text{for } y' \in \mathcal{W} \text{ so that } y' \in \mathcal{V}' \cap \mathcal{W} \\ & \nRightarrow y'' \in \mathcal{W} \text{ so that } y'' \in \mathcal{V}'' \cap \mathcal{W} \end{aligned}$$

and so,

$$y = y' + y'' \notin (\mathcal{V}' \cap \mathcal{W}) + (\mathcal{V}'' \cap \mathcal{W}).$$

Hence no. □

Example 7.8. Consider the subspaces $\mathcal{V}', \mathcal{V}'' \leq \mathcal{V}$ and suppose that $\mathcal{V}' \cup \mathcal{V}''$ is also a subspace of \mathcal{V} . Then show that either $\mathcal{V}' \subseteq \mathcal{V}''$ or $\mathcal{V}'' \subseteq \mathcal{V}'$.

Proof.

Let $x \in \mathcal{V}'$ and $y \in \mathcal{V}''$ so that $x \in \mathcal{V}' \cup \mathcal{V}''$ and $y \in \mathcal{V}' \cup \mathcal{V}''$.

Now, since $\mathcal{V}' \cup \mathcal{V}''$ is a subspace then,

$$x + y \in \mathcal{V}' \cup \mathcal{V}''$$

so,

$$x + y \in \mathcal{V}' \text{ or } x + y \in \mathcal{V}''.$$

Case i.) If $x + y \in \mathcal{V}'$ while $x \in \mathcal{V}'$ and that, since \mathcal{V}' is a subspace then, $y \in \mathcal{V}'$. So,

$$\begin{aligned} y \in \mathcal{V}'' &\implies y \in \mathcal{V}' \\ \mathcal{V}'' &\subseteq \mathcal{V}'. \end{aligned}$$

Case ii.) If $x + y \in \mathcal{V}''$ while $y \in \mathcal{V}''$ and that, since \mathcal{V}'' is a subspace then, $x \in \mathcal{V}''$. So,

$$\begin{aligned} x \in \mathcal{V}' &\implies x \in \mathcal{V}'' \\ \mathcal{V}' &\subseteq \mathcal{V}'. \end{aligned}$$

□

Remark. Observe that, in general, unions of subspaces are not necessarily subspaces.

Example 7.9. Consider the subspaces $\mathcal{W}, \mathcal{W}' \leq \mathcal{V}$ of some vector space \mathcal{V} . Show that, if \mathcal{W}'' is any subspace of \mathcal{V} containing \mathcal{W} and \mathcal{W}' then $\mathcal{W} + \mathcal{W}' \subseteq \mathcal{W}''$.

Proof.

If,

$$\mathcal{W} \cup \mathcal{W}' \subseteq \mathcal{W}'' \text{ with } \mathcal{W}'' \leq \mathcal{V}$$

then,

$$\mathcal{W} \subseteq \mathcal{W}'' \text{ and } \mathcal{W}' \subseteq \mathcal{W}''.$$

Now, suppose $x = x' + x''$ with $x' \in \mathcal{W}$ and $x'' \in \mathcal{W}'$. Then,

$$x \in \mathcal{W} \cup \mathcal{W}' \implies x \in \mathcal{W}''.$$

Hence,

$$\mathcal{W} + \mathcal{W}' \subseteq \mathcal{W}''.$$

□

8. DIRECT SUM

Consider two subspaces $\mathcal{U}, \mathcal{V} \leq \mathcal{W}$ of \mathcal{W} . Then \mathcal{U} and \mathcal{V} are said to be **complementary** if every vector $w \in \mathcal{W}$ has a *unique* decomposition $w = u + v : u \in \mathcal{U}$ and $v \in \mathcal{V}$. The vector space \mathcal{W} is then said to be the **internal direct sum** of subspaces \mathcal{U} and \mathcal{V} , written $\mathcal{W} = \mathcal{U} \oplus \mathcal{V}$.

Definition 8.1 (Direct sum). Consider two vector spaces \mathcal{U} and \mathcal{V} . Then the sum $\mathcal{U} + \mathcal{V}$ is called *direct* if and only if $\mathcal{U} \cap \mathcal{V} = \{\mathbf{0}\}$, written $\mathcal{U} \oplus \mathcal{V}$.

Definition 8.2 (Complementary subspaces). Both \mathcal{U} and \mathcal{V} are complementary subspaces of \mathcal{W} if and only if,

- $\mathcal{W} = \mathcal{U} + \mathcal{V}$ and,
- $\mathcal{U} \cap \mathcal{V} = \{\mathbf{0}\}$.

That is, if $\mathcal{U}, \mathcal{V} \leq \mathcal{W}$ we have that $\mathcal{W} = \mathcal{U} \oplus \mathcal{V}$.

Definition 8.3 (External direct sum). Consider two arbitrary vector spaces \mathcal{U} and \mathcal{V} over some field \mathbb{F} . We may define their direct sum constructively by setting $\mathcal{U} \oplus \mathcal{V} = \mathcal{U} \times \mathcal{V} = \{(u, v) : u \in \mathcal{U}, v \in \mathcal{V}\}$ with vector addition and scalar multiplication defined by:

- $(u, v) + (u', v') = (u + u', v + v')$ and
- $\lambda(u, v) = (\lambda u, \lambda v) : \lambda \in \mathbb{F}$.

Consider two vector spaces \mathcal{U} and \mathcal{V} with the linear map $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ and the short exact sequence

$$\mathcal{U} \xrightarrow{\alpha} \mathcal{U} / \ker(\varphi) \xrightarrow{\gamma} \text{im}(\varphi) \xrightarrow{\beta} \mathcal{V}.$$

In particular, observe that the map $\alpha : u \mapsto (u, 0)$ and $\beta : v \mapsto (0, v)$ are injective. While the maps $\alpha' : (u, 0) \mapsto u$ and $\beta' : (0, v) \mapsto v$ are surjective. That is γ is isomorphic. Hence we have the following commutative diagram:

$$\begin{array}{ccccc} \mathcal{U} & & & & \mathcal{V} \\ & \searrow \alpha & & \swarrow \beta & \\ & & \mathcal{U} \oplus \mathcal{V} & & \\ & \swarrow \alpha' & & \searrow \beta' & \\ \mathcal{U} & & & & \mathcal{V} \end{array}$$

Observe that the internal and external direct sums are isomorphic in this case.

We can now write higher dimensional vector spaces are direct sums of one dimensional subspaces by the natural isomorphism outlined above.

Example 8.4. Let $\mathbb{R}_1 = \{(\mathbf{x}_1, \mathbf{0}) : \mathbf{x}_1 \in \mathbb{R}\}$ and $\mathbb{R}_2 = \{(\mathbf{0}, \mathbf{x}_2) : \mathbf{x}_2 \in \mathbb{R}\}$. Then we see that $\mathbb{R}^2 = \mathbb{R}_1 \oplus \mathbb{R}_2 \cong \mathbb{R} \oplus \mathbb{R}$.

More generally, consider the subspace \mathbb{R}_i with vectors of the form

$$\mathbb{R}_i = \{(\mathbf{0}_1, \dots, \mathbf{0}_{i-1}, \mathbf{x}_i, \mathbf{0}_{i+1}, \dots, \mathbf{0}_n) : \mathbf{x}_i \in \mathbb{R}\}.$$

Then we have the following:

$$\begin{aligned} \mathbb{R}^n &= \bigoplus_{i=1}^n \mathbb{R}_i \\ &= \mathbb{R}_1 \oplus \mathbb{R}_2 \oplus \dots \oplus \mathbb{R}_n \\ &\cong \underbrace{\mathbb{R} \oplus \dots \oplus \mathbb{R}}_{n \text{ times}} \\ &= \bigoplus_{i=1}^n \mathbb{R}. \end{aligned}$$

9. INVARIANT SUBSPACES

Definition 9.1 (Characteristic polynomial). Consider some matrix $A \in \mathcal{M}_{n,n}(\mathbb{F})$ with the eigenvector \mathbf{x} and eigenvalue $\lambda \in \mathbb{F}$ relation,

$$A\mathbf{x} = \lambda\mathbf{x} : \mathbf{x} \neq \mathbf{0}$$

then we have,

$$\begin{aligned} A\mathbf{x} - \lambda\mathbf{x} &= \mathbf{0} \\ \Rightarrow (A - \lambda I_n)\mathbf{x} &= \mathbf{0}. \end{aligned}$$

Hence, the characteristic polynomial is given by

$$\chi_A(\lambda) = |A - \lambda I_n|.$$

Theorem 9.2 (Cayley-Hamilton). For any matrix $A \in \mathcal{M}_{n,n}(\mathbb{F})$ with characteristic polynomial $\chi_A(\lambda) : \lambda \in \mathbb{F}$. Then A satisfies $\chi_A(\lambda)$, that is, $\chi_A(A) = 0$.

Example 9.3. Suppose some matrix $A \in \mathcal{M}_{n,n}(\mathbb{F})$ is *idempotent*, that is $A^2 = A$. Then show that each eigenvalue of A is either zero or one.

Proof.

Suppose that,

$$A\mathbf{x} = \lambda\mathbf{x} : \mathbf{x} \neq \mathbf{0}$$

by induction we see that,

$$\begin{aligned} A^2\mathbf{x} &= \lambda^2\mathbf{x} \\ \Rightarrow A\mathbf{x} &= \lambda^2\mathbf{x} \\ &= \lambda\mathbf{x} \\ \Rightarrow \lambda &= \lambda^2 \text{ iff } \lambda \in \{0, 1\}. \end{aligned} \quad \square$$

Example 9.4. Suppose some matrix $A \in \mathcal{M}_{n,n}(\mathbb{F})$ is *nilpotent*, that is $A^k = \mathbf{0}$ for some positive integer k . Then show that all the eigenvalues of A are zero.

Proof.

Suppose that,

$$A\mathbf{x} = \lambda\mathbf{x} : \mathbf{x} \neq \mathbf{0}$$

by induction we see that,

$$\begin{aligned} A^k\mathbf{x} &= \lambda^k\mathbf{x} \\ \Rightarrow \mathbf{0} &= \lambda^k\mathbf{x} \\ \Rightarrow \lambda^k &= 0 \\ \text{and so, } \lambda &= 0 \text{ for every } \lambda \in \mathbb{F}. \end{aligned} \quad \square$$

Definition 9.5 (Eigen Subspace). For some matrix $A \in \mathcal{M}_{n,n}$. Define a eigen-subspace \mathcal{E}_λ for every eigenvalue λ as the space that contains all the eigenvectors for that particular eigenvalue. That is,

$$\mathcal{E}_\lambda(k) = \ker(A - \lambda I_n)^k.$$

Definition 9.6 (Spectrum). The *spectrum* of a linear operator $A \in \mathcal{M}_{n,n}(\mathbb{C})$ is the set of all its eigenvalues, denoted by $\sigma(A)$. That is,

$$\sigma(A) = \{\lambda \in \mathbb{C} : A\mathbf{x} = \lambda\mathbf{x} : \mathbf{x} \neq \mathbf{0}\}.$$

Definition 9.7 (Spectral radius). The *spectral radius* of a linear operator $A \in \mathcal{M}_{n,n}(\mathbb{C})$ is the radius of the smallest disc centered at the origin in the complex plane that includes all the eigenvalues

of A , denoted by $\rho(A)$. That is,

$$\rho(A) = \max\{|\lambda| : \lambda \in \sigma(A)\}.$$

Definition 9.8 (T-invariant subspace). For a vector space \mathcal{V} consider the linear endomorphism $T : \mathcal{V} \rightarrow \mathcal{V}$. Then, for some subspace $\mathcal{W} \leq \mathcal{V}$, \mathcal{W} is said to be T -invariant if $T(\mathcal{W}) \subseteq \mathcal{W}$.

Example 9.9. Show that,

$$W = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 2 \\ 0 \end{pmatrix} \right\} \text{ and } W' = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ -1 \\ 0 \end{pmatrix} \right\}$$

are T -invariant subspaces of the linear map:

$$T = \begin{pmatrix} 4 & -2 & -1 & -1 \\ 3 & -1 & -1 & -1 \\ -2 & 2 & 2 & 0 \\ 1 & -1 & 0 & 1 \end{pmatrix}.$$

Observe that,

$$T(W) = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \\ 2 \\ 1 \end{pmatrix} \right\} \subseteq W$$

and that

$$T(W') = \text{span} \left\{ \begin{pmatrix} 2 \\ 2 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 3 \\ -2 \\ 0 \end{pmatrix} \right\} \subseteq W'$$

and so, both W and W' are T -invariant subspaces.

10. ALGEBRAS

Definition 10.1 (Algebras). A finite-dimensional (associative) *algebra* \mathcal{A} over a field \mathbb{F} is a finite-dimensional vector space over \mathbb{F} equipped with a law of composition, that is, a mapping (multiplication) $(a, b) \rightarrow ab$ from $\mathcal{A} \times \mathcal{A}$ into \mathcal{A} which satisfies

- $(ab)c = a(bc)$ (associativity),

- $a(b + c) = ab + ac$,
- $(a + b)c = ac + bc$,
- $\lambda(ab) = (\lambda a)b = a(\lambda b)$,

for $\lambda \in \mathbb{F}$ and $a, b, c \in \mathcal{A}$.

Remark. A algebra \mathcal{A} is called *unital* if there exists $1 \in \mathcal{A}$, the *identity element*, such that $1a = a1 = a \forall a \in \mathcal{A}$.

Remark. A algebra \mathcal{A} is *commutative* if $ab = ba \forall a, b \in \mathcal{A}$.

11. VECTOR SPACE HOMOMORPHISMS

Definition 11.1 (Vector space Homomorphism). A vector space homomorphism is the structure preserving mapping between two vector spaces. That is, for some vector spaces \mathcal{U} and \mathcal{V} over common field \mathbb{F} with morphism $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ we have,

$$\varphi(u_1 + \lambda u_2) = \varphi(u_1) + \lambda \varphi(u_2) : u_1, u_2 \in \mathcal{U} \text{ and } \lambda \in \mathbb{F}.$$

Remark. A vector space homomorphism preserves linearity and so we typically call the homomorphism a linear map.

Definition 11.2 (Kernel). Consider the linear map $\varphi : \mathcal{U} \rightarrow \mathcal{V}$, the *kernel* of φ , denoted $\ker(\varphi)$ is given by:

$$\ker(\varphi) = \{u \in \mathcal{U} : \varphi(u) = \mathbf{0} \in \mathcal{V}\}.$$

The set or space of elements inside the kernel is thus called the *null space*.

Definition 11.3 (Nullity). The *nullity* is the dimension of the null space or kernel.

Remark. Note that the nullity gives a measure of the injectivity of the linear map φ .

Definition 11.4 (Image). For some morphism $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ the *image* is defined as,

$$\text{img}(\varphi) = \varphi(\mathcal{U}) = \{v \in \mathcal{V} : v = \varphi(u) \text{ for some } u \in \mathcal{U}\}.$$

Definition 11.5 (Preimage). For some morphism $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ the *preimage* is defined as,

$$\varphi^{-1}(\mathcal{V}) = \{u \in \mathcal{U} : \varphi(u) \in \mathcal{V}\}$$

Remark. The preimage can be defined even when no inverse morphism exists.

Theorem 11.6. Let \mathcal{U} and \mathcal{V} be vector spaces over some common field \mathbb{F} and some linear map $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ then

$$\dim(\text{img}(\varphi)) + \dim(\ker(\varphi)) = \dim(\mathcal{U})$$

Definition 11.7 (Injective). If, for the linear map $\varphi : \mathcal{U} \rightarrow \mathcal{V}$, we have

$$\varphi(u_1) = \varphi(u_2) \implies u_1 = u_2 \quad \forall u_1, u_2 \in \mathcal{U}$$

then φ is said to be *injective*.

Definition 11.8 (Surjective). If, for the linear map $\varphi : \mathcal{U} \rightarrow \mathcal{V}$, we have

$$\forall v \in \mathcal{V} \exists u \in \mathcal{U} : \varphi(u) = v$$

then φ is said to be *surjective*.

Definition 11.9 (Bijective). If the linear map $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ is both injective and surjective then φ is said to be *bijective*, or, isomorphic in the context vector space homomorphisms.

11.1. Properties of linear maps. One may note that a linear map can always be represented in the form of a matrix and as such the determinant can be taken given that linear maps are from vector spaces to vector spaces over a common field. Hence, the conceptual question arises, geometrically, what exactly is the determinant? It turns out the determinant is a *measure* of dilation much like how scalar multiplies of the unit scalar dilate it. That is, $3 * 1 = 3$ is the dilation of the unit scalar 1 by a measure of 3. Now, recall that a field is a vector space over itself and so we may use this to generalise our intuition from above. Thus, we may view a linear map as a linear dilation of some vector in one vector space to another vector space and the determinant as the measure of dilation of this mapping.

TODO.. motivate to prove the determinant definition here.. and motivate to the characteristic polynomial and eigen spaces..

12. DUAL SPACES

Recall that; If \mathcal{U} and \mathcal{V} are two vector spaces over a field \mathbb{F} , then the set of all linear maps from $\mathcal{U} \rightarrow \mathcal{V}$, denoted $L(\mathcal{U}, \mathcal{V})$, is a vector space over \mathbb{F} .

Definition 12.1. Let \mathcal{V} be a vector space over \mathbb{F} . The dual space of \mathcal{V} , denoted \mathcal{V}^* , is the set of all linear maps from \mathcal{V} to \mathbb{F} . That is, $\mathcal{V}^* = L(\mathcal{V}, \mathbb{F})$. The elements of \mathcal{V}^* are called *linear functionals* on \mathcal{V} .

Remark. Since $L(\mathcal{V}, \mathbb{F})$ is a vector space over \mathbb{F} then so too is the dual \mathcal{V}^* , with point-wise addition and scalar multiplication. Moreover, if $n = \dim \mathcal{V} < \infty$, then $\dim \mathcal{V}^* = \dim L(\mathcal{V}, \mathbb{F}) = n$.

Example 12.2. Fix $x \in [0, 1]$ and define the *evaluation map* $ev_x : \mathcal{C}([0, 1]) \rightarrow \mathbb{R}$ by $ev_x(f) = f(x)$ for any $f \in \mathcal{C}([0, 1])$. Then $ev_x \in (\mathcal{C}([0, 1]))^*$.

13. NORMED VECTOR SPACE

A norm on a complex vector space \mathcal{V} is a real-valued function $\|\cdot\| : \mathcal{V} \rightarrow \mathbb{R}$ that, to any vector $x \in \mathcal{V}$ associates a real number $\|x\|$, satisfies the following:

- $\|x\| \geq 0$, $\forall x \in \mathcal{V}$ and in particular $\|x\| = 0 \Leftrightarrow x = 0$ (positive definite),
- $\|\alpha x\| = |\alpha| \|x\| \forall x \in \mathcal{V}$, and $\alpha \in \mathbb{C}$ (uniform scaling),
- $\|x + y\| \leq \|x\| + \|y\| \forall x, y \in \mathcal{V}$ (triangle inequality).

13.1. Linear Operators. In functional analysis one considers linear maps between normed vector spaces. In this context such linear mappings are termed *linear operators* or just operators. In this light one is motivated to talk about notions of *continuity* of such operators.

For any normed vector space $(\mathcal{U}, \|\cdot\|_{\mathcal{U}})$ consider the closed unit ball in \mathcal{U} given by $B(\mathcal{U}) = \{x \in \mathcal{U} : \|x\|_{\mathcal{U}} \leq 1\}$. Hence, we may now define boundedness of an operator formally as follows:

Definition 13.1 (Bounded linear operator). A linear operator $T : \mathcal{U} \rightarrow \mathcal{V}$ between normed vector spaces $(\mathcal{U}, \|\cdot\|_{\mathcal{U}})$ and $(\mathcal{V}, \|\cdot\|_{\mathcal{V}})$ is called a *bounded* linear operator if it maps the closed unit ball $B(\mathcal{U})$ in \mathcal{U} into a bounded set (a ball) in \mathcal{V} .

Corollary. A linear operator $T : \mathcal{U} \rightarrow \mathcal{V}$ between normed vector spaces $(\mathcal{U}, \|\cdot\|_{\mathcal{U}})$ and $(\mathcal{V}, \|\cdot\|_{\mathcal{V}})$ is continuous if and only if it is bounded.

14. INNER PRODUCT SPACES

Let V be a complex vector space.

Definition 14.1 (Inner product). An *inner product* in V is a mapping $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$ satisfying

- Hermitian symmetry: $\langle x, y \rangle = \overline{\langle y, x \rangle} \forall x, y \in V$;
- Linearity: $\langle \alpha x_1 + \beta x_2, y \rangle = \alpha \langle x_1, y \rangle + \beta \langle x_2, y \rangle \forall x_1, x_2, y \in V$ and $\alpha, \beta \in \mathbb{C}$;
- Positive definite: $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0$ iff $x = 0$, $\forall x \in V$;

Remark. Note that condition (i) implies that $\langle x, x \rangle \in \mathbb{R}$ and so condition (ii) is consistent.

Remark. Note that condition (ii) defines that the inner product is linear in the first argument. However, the inner product is *conjugate linear* in the second argument as shown by:

Proof.

$$\begin{aligned} \langle x, \alpha y_1 + \beta y_2 \rangle &= \overline{\langle \alpha y_1 + \beta y_2, x \rangle} \\ &= \overline{\langle \alpha y_1, x \rangle + \langle \beta y_2, x \rangle} \\ &= \bar{\alpha} \overline{\langle y_1, x \rangle} + \bar{\beta} \overline{\langle y_2, x \rangle} \\ &= \bar{\alpha} \langle x, y_1 \rangle + \bar{\beta} \langle x, y_2 \rangle. \end{aligned}$$

□

A vector space V together with an inner product $\langle \cdot, \cdot \rangle$ is termed an *inner product space*.

Example 14.2. Consider the vector space $V = M_{mn}(\mathbb{F})$. We wish to show that, for some $X, Y \in V$, that $\text{tr}(X^*Y)$ defines an inner product in V . Hence we have, $\langle X, Y \rangle \doteq \text{tr}(X^*Y) \forall X, Y \in V$.

Proof. First suppose that $\langle X, Y \rangle = \text{tr}(X^*Y)$ and so;

We check that $\text{tr}(X^*Y)$ is Hermitian symmetric:

$$\begin{aligned} \langle X, Y \rangle &= \text{tr}(X^*Y) \\ &= \overline{\text{tr}(\overline{X^*Y})} \\ &= \overline{\text{tr}(YX^*)} \\ &= \overline{\text{tr}((X^*Y)^T)} \\ &= \overline{\text{tr}(Y^*X)} \\ &= \overline{\langle Y, X \rangle}. \end{aligned}$$

Check that $\text{tr}(X^*Y)$ is linear in the first parameters by considering some $\alpha, \beta \in \mathbb{F}$ so that

$$\begin{aligned} \langle \alpha X_1 + \beta X_2, Y \rangle &= \text{tr}((\alpha X_1 + \beta X_2)^*Y) \\ &= \text{tr}((\alpha X_1)^*Y + (\beta X_2)^*Y) \\ &= \text{tr}((\alpha X_1)^*Y) + \text{tr}((\beta X_2)^*Y) \\ &= \bar{\alpha} \text{tr}(X_1^*Y) + \bar{\beta} \text{tr}(X_2^*Y) \\ &= \bar{\alpha} \langle X_1, Y \rangle + \bar{\beta} \langle X_2, Y \rangle. \end{aligned}$$

and check that $\text{tr}(X^*Y)$ is positive definite for $X = Y$,

$$\langle X, X \rangle = \text{tr}(X^*X) \geq 0$$

and when

$$\begin{aligned} X = 0 &\implies X^* = 0 \\ &\implies \text{tr}(X^*X) = 0. \end{aligned}$$

□

TODO..

Example 14.3. Let \mathcal{V} be the vector space of continuous real-valued functions $[0, 1]$ endowed with the standard inner product. If $f(x) = x$ and $g(x) = e^x$ then find $\langle f, g \rangle$ and $\|f\|$.

For $\langle f, g \rangle$ we have,

$$\begin{aligned}
 \langle f, g \rangle &= \int_0^1 f(x) \overline{g(x)} dx \\
 &= \int_0^1 x e^x dx && \text{(by parts we have,)} \\
 &= x e^x \Big|_0^1 - \int_0^1 e^x dx \\
 &= e - (e - 1) = 1.
 \end{aligned}$$

Now, for $\|f\|$ we have,

$$\begin{aligned}
 \|f\| &= \left(\int_0^1 x^2 dx \right)^{\frac{1}{2}} \\
 &= \sqrt{\frac{1}{x^3} \Big|_0^1} \\
 &= \frac{1}{\sqrt{3}}.
 \end{aligned}$$

Theorem 14.4. *Let \mathcal{V} be a complex vector space with inner product $\langle \cdot, \cdot \rangle$. Then the inner product induces a norm $\|\cdot\|$ in \mathcal{V} by the definition*

$$\|x\| = \sqrt{\langle x, x \rangle} \text{ for } x \in \mathcal{V}.$$

Proof.

Notice that $\|\cdot\|$ is trivially positive definite as follows:

$$\|x\| = \sqrt{\langle x, x \rangle} \geq 0 \Leftrightarrow x = 0, \forall x \in \mathcal{V}.$$

By linearity we see that,

$$\begin{aligned}
 \|\alpha x\| &= \sqrt{\langle \alpha x, \alpha x \rangle} \\
 &= \sqrt{\alpha \bar{\alpha} \langle x, x \rangle} \\
 &= \sqrt{|\alpha|^2 \langle x, x \rangle} = |\alpha| \|x\| \forall \alpha \in \mathbb{C} \text{ and } \forall x \in \mathcal{V}.
 \end{aligned}$$

Finally the triangle inequality follows as shown,

$$\begin{aligned}
 \|x + y\|^2 &= \langle x + y, x + y \rangle \\
 &= \|x\|^2 + \|y\|^2 + \langle x, y \rangle + \langle y, x \rangle \\
 &= \|x\|^2 + \|y\|^2 + \langle x, y \rangle + \overline{\langle x, y \rangle} \\
 &= \|x\|^2 + \|y\|^2 + 2\Re[\langle x, y \rangle] \\
 &\leq \|x\|^2 + \|y\|^2 + 2|\langle x, y \rangle| \\
 &\leq \|x\|^2 + \|y\|^2 + 2\|x\|\|y\| \\
 &= (\|x\| + \|y\|)^2 \quad \forall x, y \in \mathcal{V}.
 \end{aligned}$$

□

TODO.. MOVE ME..

Example 14.5. Find a basis for the orthogonal complement of $W = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} \right\} \in \mathbb{R}^4$.

Therefore we do the following:

$$\begin{aligned}
 W^\perp &= \ker \begin{pmatrix} 1 & 1 & 1 & 0 \\ 2 & 2 & 1 & 1 \end{pmatrix} \\
 &= \ker \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix} && \text{(after a row reduction)} \\
 &= \text{span} \left\{ \begin{pmatrix} 0 \\ -1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 1 \\ 1 \end{pmatrix} \right\}.
 \end{aligned}$$