




Condensed Matter Theory



MINGYU XIA | PhD @Westlake University



This page was intentionally left blank.

Contents

Lecture #1 Free electron model	i
Problem Set	i
Lecture #2 Phonon	vii
Lecture #3 Crystal lattice	xv
Lecture #4 X-ray diffraction	xix
Lecture #5 Nearly free electron model	xxii
Lecture #6 Tight binding model	xxvii
Lecture #7 Semiconductor	xxx
Lecture #8 Electrons in a magnetic field	xxxii
Lecture #9 Band structure of graphene	xxxv

Lecture #1 Free electron model

Problem 1.1. Two-charge-carrier Drude model:

Consider a system of two types of charge carriers in the Drude model. The two carriers have the same density (n) and opposite charge (e and $-e$), and their masses and relaxation rates are m_1, m_2 and τ_1, τ_2 , respectively. So their corresponding mobilities are $\mu_1 = \frac{e\tau_1}{m_1}$ and $\mu_2 = \frac{e\tau_2}{m_2}$, respectively.

- Calculate the magnetoresistance $\Delta\rho = \rho(B) - \rho(B=0)$, where B is the magnetic field.
- Calculate the Hall coefficient.
- In an undoped semiconductor, $n = n_0 e^{-\Delta/k_B T}$ describes the temperature dependence of the carrier concentration. What will the temperature dependence of the magnetoresistance and the Hall coefficient be?

Solution (By TA). The EOM

$$m \frac{dv}{dt} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}) - m \frac{v}{\tau} = 0$$

Apply $\mathbf{j} = \hat{\sigma} \mathbf{E} = nq\mathbf{v}$, then

$$\mathbf{E} = \frac{m}{mq^2\tau} \mathbf{j} - \frac{1}{qn} \mathbf{j} \times \mathbf{B} = \begin{pmatrix} \frac{m}{nq^2\tau} & -\frac{B}{qn} & 0 \\ \frac{B}{qn} & \frac{m}{nq^2\tau} & 0 \\ 0 & 0 & \dots \end{pmatrix}$$

The conductivity

$$\hat{\sigma} = \mathbf{E}^{-1} = \begin{pmatrix} \frac{\sigma_0}{1+\omega_c^2\tau^2} & \sigma_0 \frac{\omega_c\tau}{1+\omega_c^2\tau^2} & 0 \\ -\sigma_0 \frac{\omega_c\tau}{1+\omega_c^2\tau^2} & \frac{\sigma_0}{1+\omega_c^2\tau^2} & 0 \\ 0 & 0 & \dots \end{pmatrix}$$

where $\omega_c = \frac{Bq}{m}$. When $B=0$,

$$\rho_{xx}(0) = \frac{1}{\sigma_{10} + \sigma_{20}}, \quad \rho_{xy}(0) = 0$$

When $B \neq 0$

$$\rho_{xx} = \frac{\sigma_{xx}}{\sigma_{xx}^2 + \sigma_{xy}^2}, \quad \rho_{xy} = -\frac{\sigma_{xy}}{\sigma_{xx}^2 + \sigma_{xy}^2}$$

Solution.

- For the two carriers, we have

$$\sigma_{0i} = \frac{ne^2\tau_i}{m_i} = ne\mu_i, \quad \omega_{ci} = \frac{eB}{m_i}, \quad \omega_{ci}\tau_i = \mu_i B$$

The total conductivity tensor is

$$\hat{\sigma} = \hat{\sigma}_1 + \hat{\sigma}_2$$

and the components are

$$\begin{aligned}\sigma_{xx} &= \frac{ne\mu_1}{1 + (\mu_1 B)^2} + \frac{ne\mu_2}{1 + (\mu_2 B)^2} \\ \sigma_{xy} &= \frac{ne\mu_1^2 B}{1 + (\mu_1 B)^2} + \frac{ne\mu_2^2 B}{1 + (\mu_2 B)^2} \\ \sigma_{zz} &= ne(\mu_1 + \mu_2)\end{aligned}$$

The resistance tensor is the inverse

$$\hat{\rho} = \hat{\sigma}^{-1} = \frac{1}{\sigma_{xx}^2 + \sigma_{xy}^2} \begin{pmatrix} \sigma_{xx} & -\sigma_{xy} & 0 \\ \sigma_{xy} & \sigma_{xx} & 0 \\ 0 & 0 & \frac{\sigma_{xx}^2 + \sigma_{xy}^2}{\sigma_{zz}} \end{pmatrix}.$$

Then, the element ρ_{xx} becomes

$$\rho_{xx}(0) = \sigma_{xx}^{-1}(0) = \frac{1}{ne(\mu_1 + \mu_2)}, \quad \text{and} \quad \rho_{xx} = \frac{\sigma_{xx}(B)}{\sigma_{xx}^2(B) + \sigma_{xy}^2(B)}$$

So, the magnetoresistance is

$$\Delta\rho = \rho_{xx}(B) - \rho_{xx}(0) = \frac{1}{ne} \left[\frac{\frac{\mu_1}{1+(\mu_1 B)^2} + \frac{\mu_2}{1+(\mu_2 B)^2}}{\left(\frac{\mu_1}{1+(\mu_1 B)^2} + \frac{\mu_2}{1+(\mu_2 B)^2} \right)^2 + B^2 \left(\frac{\mu_1}{1+(\mu_1 B)^2} - \frac{\mu_2}{1+(\mu_2 B)^2} \right)^2} - \frac{1}{\mu_1 + \mu_2} \right]$$

(b) Since

$$\rho_{xy}(B) = -\frac{\sigma_{xy}(B)}{\sigma_{xx}(B)^2 + \sigma_{xy}(B)^2},$$

substitute it into the definition of Hall coefficient $R_H = \frac{\rho_{xy}}{B}$, that is

$$R_H = -\frac{1}{ne} \cdot \frac{\frac{\mu_1}{1+(\mu_1 B)^2} + \frac{\mu_2}{1+(\mu_2 B)^2}}{\left(\frac{\mu_1}{1+(\mu_1 B)^2} + \frac{\mu_2}{1+(\mu_2 B)^2}\right)^2 + B^2 \left(\frac{\mu_1}{1+(\mu_1 B)^2} - \frac{\mu_2}{1+(\mu_2 B)^2}\right)^2}$$


(c) For simplicity, consider the case B is small. Now, $\Delta\rho$ and R_H becomes

$$\Delta\rho = ne\mu_1\mu_2(\mu_1 + \mu_2) \left(\frac{\mu_1 - \mu_2}{\mu_1 + \mu_2}\right)^2 B^2, \quad \text{and} \quad R_H = -\frac{1}{ne} \frac{\mu_1 - \mu_2}{\mu_1 + \mu_2}$$

So, in this case, the magnetoresistance $\Delta\rho \propto e^{-\Delta/k_B T}$, while the Hall coefficient $R_H \propto e^{\Delta/k_B T}$.

Problem 1.2. Density of States in 2D system:

Consider free electron in a 2D box with a width of a . The allowed wave-numbers k_x and k_y for x and y directions respectively are expressed as $\frac{\pi n_x}{a}$ and $\frac{\pi n_y}{a}$, where $n_x, n_y = 1, 2, 3, \dots$ are integer numbers. The following figure shows the schematic view of the 2D array of allowed quantum states in k -space.



- Determine the area in k -space occupied by one allowed k point.
- Express number of allowed states in the area enclosed by the first quarter circles with the radius of k and $k + dk$, where $k \gg dk$ (Consider the spin degeneracy).
- Express density of allowed states in real space per unit energy.

Solution.

- The spaces of unit k point in k_x -direction and k_y -direction are

$$\Delta k_x = \frac{\pi}{a}, \quad \text{and} \quad \Delta k_y = \frac{\pi}{a}$$

Therefore, the area in k -space occupied by one allowed k point is

$$\Delta k_x \cdot \Delta k_y = \left(\frac{\pi}{a}\right)^2$$

- The area enclosed by the two quarter circles is

$$dA = \frac{1}{4} 2\pi k dk = \frac{\pi k}{2} dk$$

Since every k point occupy an area of $(\pi/a)^2$, so the number of allowed k -points in this area is

$$dN = \frac{2 dA}{(\pi/a)^2} = \frac{ka^2}{\pi} dk$$

Since two electrons can occupy each grid point ($|\uparrow\rangle$ and $|\downarrow\rangle$).

- Derivate it to the energy for a free election, we get

$$dE = d\left(\frac{\hbar^2 k^2}{2m}\right) = \frac{\hbar^2 k}{m} dk, \quad dk = \frac{m dE}{\hbar^2 k^2}.$$

Substitute it into the result in (b): the number of states in dk

$$dN = \frac{ka^2}{\pi} dk = \frac{ma^2}{\pi \hbar^2} dE,$$

means in the whole system of area a^2 , a number dN of states are allowed in energy range dE . So, the density of allowed states in real space per unit energy is

$$g(E) = \frac{1}{a^2} \frac{dN}{dE} = \frac{m}{\pi \hbar^2}.$$

Problem 1.3. Density of states from a general formula:

- (a) In a 3D solid and starting from the general formula, $g(E) = \frac{2V}{(2\pi)^3} \iint \frac{dS_E}{|\nabla_k E|}$, find the expression for the electronic density of states: (i) for free electrons; (ii) for electrons that obey the isotropic dispersion relation: $E = -E_0 - 2\gamma \cos\left(\frac{ka}{2}\right)$, where $E_0, \gamma > 0$.
- (b) After having established the corresponding expression for 2D, find $g(E)$ for (i) and (ii) above.
- (c) Same question for 1D.

Solution (By TA).

$$dV_E = dS_E \cdot dk = dS_E \cdot \frac{dE}{|\nabla_k E|}$$

Then, the volume of the sphere surface

$$V_{\text{site}} = dE \iint \frac{dS}{|\nabla_k E|}$$

Also for

$$dN = 2 \frac{1}{\left(\frac{2\pi}{a}\right)^3}, \quad dN = g(E) dE$$

For 2D case

$$g(E) = 2 \frac{A}{(2\pi)^2} \int \frac{dL}{|\nabla_k E|}$$

1D

$$g(E) = 2 \cdot \frac{L}{2\pi} \frac{1}{|\nabla_k E|}$$

Solution. Firstly, we shall calculate something before: For the isotropic dispersion

$$E = -E_0 - 2\gamma \cos(ka/2),$$

we can obtain

$$\sin\left(\frac{ka}{2}\right) = \sqrt{1 - \left(\frac{E + E_0}{2\gamma}\right)^2}, \quad \text{and} \quad k = \frac{2}{a} \arccos\left(-\frac{E + E_0}{2\gamma}\right).$$

Since $E = \hbar^2 k^2 / 2m$, so $k = \sqrt{2mE} / \hbar$, $\nabla_k E = \hbar^2 k / m$, $\nabla_k E = \gamma a \sin(ka/2)$ for isotropic dispersion.

- (a) In a 3D situation, the general formula becomes

$$g(E) = \frac{2V}{(2\pi)^3} \cdot \frac{4\pi k^2}{\hbar^2 k / m} = \frac{Vmk}{\pi^2 \hbar^2} = \frac{Vm}{\pi^2 \hbar^3} \sqrt{2mE}.$$

For isotropic dispersion, the general formula becomes

$$g(E) = \frac{V k^2}{\pi^2 \gamma a \sqrt{1 - \left(\frac{E + E_0}{2\gamma}\right)^2}}.$$

(b) In a 2D situation, the general formula becomes

$$g(E) = \frac{2A}{(2\pi)^2} \cdot \frac{2\pi k}{\hbar^2 k/m} = \frac{mA}{\pi \hbar^2} = \text{Const.}$$

For isotropic dispersion, the general formula becomes

$$g(E) = \frac{2A}{(2\pi)^2} \cdot \frac{2\pi k}{\gamma a \sin(ka/2)} = \frac{2A}{\pi \gamma a^2} \frac{\arccos\left(-\frac{E+E_0}{2\gamma}\right)}{\sqrt{1 - \left(\frac{E+E_0}{2\gamma}\right)^2}}$$

(c) In 1D situation, $g(E) = \frac{2L}{2\pi} \cdot \frac{2}{dE/dk}$. The general formula becomes

$$g(E) = \frac{2L}{2\pi} \cdot \frac{2}{dE/dk} = \frac{2L}{\pi dE/dk} = \frac{L}{\pi \hbar} \sqrt{\frac{2m}{E}}$$

For isotropic dispersion, the general formula becomes

$$g(E) = \frac{2L}{\pi \gamma a} \frac{1}{\sqrt{1 - \left(\frac{E+E_0}{2\gamma}\right)^2}}.$$

Lecture #2 Phonon

Problem 2.1 (Normal modes of a square lattice). Consider a square lattice of atoms of mass M , in which the nearest neighbors are coupled to each other by springs with constants K . Derive the equation of motion for the mode in which atomic displacements are normal to the crystal plane and find the dispersion of this mode. Analyze the behavior of this mode in the continuum limit.

Solution (By TA). Only consider the vibration perpendicular to the surface

$$M \frac{d^2 U_{n,m}}{dt^2} = -k \sum [u_{n,m} - u_{n\pm 1, m\pm 1}]$$

Substitute the plane wave

$$u_{n,m}(t) = U e^{i(q_x n a + q_y m a - \omega t)}$$

we have

$$-M\omega^2 = 2k[\cos(q_x a) + \cos(q_y a) - 2]$$

For $|q_x a|, |q_y a| \ll \pi$, $\sin qa = qa$.

Solution. Assume the lattice constant a , then the lattice sites become

$$\mathbf{R}_{n,m} = (na, ma), \quad n, m \in \mathbb{Z}.$$

We consider every atom has a single degree of freedom: the out-of-plane scalar displacement $u_{n,m}(t)$. So, the spring connecting site (n, m) and a neighbour (n', m') has extension $\Delta = u_{n',m'} - u_{n,m}$, which brings the force to atom (n, m)

$$F_{n,m} = -K(u_{n,m} - u_{n',m'})$$

due to Hooke's law. Since every atom has four nearest neighbours: $(n \pm 1, m)$, $(n, m \pm 1)$, which bring four forces to the “central atom”. Summing these forces

$$F_{n,m}^{\text{tot}} = - \sum_{\langle n,m;n',m' \rangle} K(u_{n,m} - u_{n',m'}) = -K(4u_{n,m} - u_{n+1,m} - u_{n-1,m} - u_{n,m+1} - u_{n,m-1})$$

According to Newton's second law, we obtain the EOM

$$M\ddot{u}_{n,m} = -K(4u_{n,m} - u_{n+1,m} - u_{n-1,m} - u_{n,m+1} - u_{n,m-1}).$$

Assume the EOM has a plane-wave solution

$$u_{n,m}(t) = A e^{i(k_x n a + k_y m a - \omega t)},$$

then substitute it into the EOM, the LHS is

$$\text{LHS} = M(-\omega^2)u_{n,m}$$

and the RHS is

$$\text{RHS} = -K(4 - e^{ik_x a} - e^{-ik_x a} - e^{ik_y a} - e^{-ik_y a})u_{n,m} = -2K[2 - \cos(k_x a) - \cos(k_y a)]u_{n,m}$$

Finally we have the dispersion of this model

$$\omega^2(\mathbf{k}) = \frac{2K}{M} [2 - \cos(k_x a) - \cos(k_y a)], \quad \omega(\mathbf{k}) = 2\sqrt{\frac{K}{M}} \sqrt{\sin^2\left(\frac{k_x a}{2}\right) + \sin^2\left(\frac{k_y a}{2}\right)}.$$

In the continuum limit, i.e., $|\mathbf{k}|a \ll 1$, then the dispersion of this model becomes linear to $|\mathbf{k}|$

$$\omega(\mathbf{k}) \approx 2\sqrt{\frac{K}{M}} \sqrt{\left(\frac{k_x a}{2}\right)^2 + \left(\frac{k_y a}{2}\right)^2} = c|\mathbf{k}|$$

where $c = a\sqrt{K/M}$.

Problem 2.2 (Localized phonons on an impurity). Consider a linear chain of identical atoms of mass m each equidistant by a .

- Limiting the problem to interactions between nearest neighbors characterized by the force constant β , give the dispersion relation of longitudinal phonons and derive the expression of their group velocity. Express the results as a function of “ a ” and of the velocity of sound, v_s . What is maximum frequency of atomic vibration, ω_M , when $v_s = 5000 \text{ m/s}$?
- An atom in the middle of the chain is replaced by the atom of an impurity with a mass m' (where $m' < m$). Write the displacement equations, u_0 , for this impurity atom and the displacement “ u_n ” for an atom located at a distance “ na ”. Assume that the force constant β between nearest neighbors is the same regardless of the atom considered.
- The impurity atom being lighter than the others has an oscillation frequency ω'_m larger than ω_M and it drags the other neighbor atoms by its motion. We try to describe its effect with the expression of a damped wave of form: $u_n = u_0 e^{-\alpha|x|} e^{i(\omega't - kx)}$. What is the value of the frequency ω'_M , which is compatible with the equations of motion when the wave vector is near the boundary of the first Brillouin Zone? Verify that when $m = m'$, the solution matches that obtained in (a).
- After having numerically evaluated ω'_M and a when $m' = m/2$, sketch the instantaneous position of atoms as a function of x .

Solution (By TA). $v_y = \frac{dm}{dk} \cdot v_s$ is long-wave limit v_s : $\lim_{k \rightarrow 0} v_g = a\sqrt{\frac{\beta}{m}}$, and $\omega_m = \frac{2v_s}{a}$.

$$u_n = u_0 e^{-\alpha na} e^{i(\omega't - kna)}$$

Substitute it into the EOM

$$\begin{cases} -\omega'^2 m' = \beta[e^{ika}(e^{ika} + e^{-ika}) - 2] \\ -\omega'^2 m = \beta[e^{(\alpha+ik)a} + e^{-(\alpha+ik)a} - 2] \end{cases}$$

where $ka = \pi$, $\omega' = \omega'_m$,

$$\begin{aligned} u_n &= u_0 (-1)^n e^{i\omega't} e^{-\alpha a|n|} \\ \begin{cases} m\omega'^2_m = 2\beta(1 + e^{-\alpha a}), \\ m\omega'^2_m = \Delta\beta \text{ch}^2\left(\frac{\alpha a}{2}\right) \end{cases} \end{aligned}$$

Solution.

- Assume the nearest-neighbor spring constant β . The EOM for the n -th atom is

$$m\ddot{u}_n = -\beta(u_n - u_{n+1}) - \beta(u_n - u_{n-1}) = \beta(u_{n+1} - u_{n-1} - 2u_n).$$

We assume the EOM has a plane-wave solution $u_n(t) = A e^{i(kna - \omega t)}$, then substitute it into the EOM

$$-m\omega^2 u_n = -\beta(2 - e^{ika} - e^{-ika})u_n = -\beta(2 - 2\cos(ka))u_n = -4\beta \sin^2\left(\frac{ka}{2}\right),$$

we have the dispersion

$$\omega = 2\sqrt{\frac{\beta}{m}} \left| \sin\left(\frac{ka}{2}\right) \right|.$$

The group velocity and the velocity of sound are

$$v_g(k) = \frac{d\omega}{dk} = a\sqrt{\frac{\beta}{m}} \cos\left(\frac{ka}{2}\right), \quad \text{and} \quad v_s(a) = \lim_{k \rightarrow 0} v_g(k) = a\sqrt{\frac{\beta}{m}}$$

then the group velocity, as well as the frequency can be expressed as

$$v_g(k) = v_s \cos\left(\frac{ka}{2}\right), \quad \text{and} \quad \omega = \frac{2v_s}{a} \left| \sin\left(\frac{ka}{2}\right) \right|.$$

when $k = \pi/a$, the frequency arrives at a maximal value

$$\omega_M = \frac{2v_s}{a} \xrightarrow{v_s=5000 \text{ m/s}} \frac{10000}{a} \text{ s}^{-1}.$$

- (b) The impurity is located at site $n = 0$, with mass m' ; while its neighbors $n = \pm 1$ with mass m . The EOM for the impurity and other atoms are

$$\begin{aligned} m' \ddot{u}_0 &= -\beta(u_1 - u_0) + \beta(u_{-1} - u_0) = \beta(u_1 + u_{-1} - 2u_0) \\ m \ddot{u}_n &= \beta(u_{n+1} + u_{n-1} - 2u_n) \end{aligned}$$

- (c) In (a), near the boundary of the first BZ, i.e. $k = \pi/a$, the frequency is $\omega_1 = 2\sqrt{\beta/m}$. Write the given wave form in terms of site number n

$$u_n = A e^{-\alpha|n|a} e^{i(\omega't - kna)}$$

Substitute it into the two EOMs in (b), and let $n = 1$

$$\begin{aligned} -\omega^2 m' u_0 &= \beta(e^{-\alpha a - ika} + e^{-\alpha a + ika} - 2)u_0 \xrightarrow{k=\pi/a} -2\beta(e^{-\alpha a} + 1)u_0 \\ -\omega^2 m u_1 &= \beta(u_2 + u_0 - 2u_1) = \beta(e^{-\alpha a - ika} + e^{\alpha a + ika} - 2)u_1 \xrightarrow{k=\pi/a} -2\beta(\text{ch}(\alpha a) + 1)u_1 \end{aligned}$$

Derive the two expressions, we have

$$\frac{m'}{m} = \frac{e^{-\alpha a} + 1}{\text{ch}(\alpha a) + 1} = \frac{2}{e^{\alpha a} + 1}, \quad \text{and} \quad e^{-\alpha a} + 1 = \frac{2m}{2m - m'}$$

Substitute it into one of the EOM, we obtain the dispersion

$$\omega_M'^2 = \omega'^2 = \frac{4\beta m}{m'(2m - m')}$$

where $\omega_M'^2 = \omega'^2$ is due to the boundary of the first BZ ($k = \pi/a$). When $m' = m$, the frequency becomes


$$\omega_M'^2 = \frac{4\beta}{m}, \quad \text{and} \quad \omega_M' = 2\sqrt{\frac{\beta}{m}} = \omega_1$$

which matches that obtained in (a).

- (d) Substitute $m' = m/2$ in the result of (c), we have $\omega_M'^2 = \frac{16}{3} \frac{\beta}{m}$. To simplify, we choose the time $t = 0$. Near the first BZ, the displacement of atoms is

$$u_n = (-1)^n A e^{-\alpha|n|a}$$

The plotted figure of atoms is attached as follows (Just show the relative position for reference).



Problem 2.3 (Density of states and specific heat of a monoatomic 1D lattice). Consider a row of N identical atoms of mass m and equidistant by a .

- Limiting the problem to interactions between nearest neighbors with a force constant β , find the expression for the dispersion relation of acoustic longitudinal vibrations propagating along this row. What is the maximum frequency ω_M of atomic vibrations, written in terms of v_s and a ? Compare the value obtained with that of ω_D deduced from the Debye model.
- Find the corresponding expression for the density of states $g(\omega)$.
- Write the expressions for internal energy U and for the heat capacity C_v . Find the limits as these functions approach high temperatures ($\hbar\omega_M \ll k_B T$).
- What is the evolution of C_v at low temperature? Compare the result obtained here with that deduced from the Debye model.

Solution (By TA). (a) $\omega = v_s k$, $\int_0^{\omega_0} g_D(\omega) d\omega = N$, $\omega(k) = 2\sqrt{\beta/m} |\sin(ka/2)|$.

$$g_D(\omega) = \frac{L}{\pi} \frac{dk}{d\omega} = \frac{L}{\pi} \cdot \frac{1}{v_s} = \frac{Na}{\pi v_s}$$

$$\omega_D = \pi v_s / a, \omega_M = 2v_s / a.$$

$$(b) \langle n \rangle = \frac{1}{e^{\hbar\omega/k_B T} - 1}, \langle E \rangle = \langle n \rangle \hbar\omega, u_0 = \frac{1}{2} \hbar\omega.$$

$$U = \int_0^{\omega_M} g(\omega) \left[\frac{1}{2} \hbar\omega + \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} \right] d\omega$$

$$D_v = \frac{dU}{dT}, \text{ for } \hbar\omega \ll k_B T, \text{ the exponential is } \frac{\hbar\omega}{k_B T},$$

$$C_v = k_B \int_0^{\omega_a} g(\omega) d\omega = Nk_B$$

- Only $\omega \rightarrow 0$ contributes to the integration.

$$e^{\hbar\omega/k_B T} \begin{cases} \hbar\omega \gg k_B T \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} = \hbar\omega e^{-\hbar\omega/k_B T}, \\ \hbar\omega \ll k_B T \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} \approx k_B T \end{cases}$$

$$\omega \rightarrow 0, g(\omega) = g(0) = \frac{2N}{\pi\omega_M}.$$

Solution.

- The same to Problem 2.2(a): the dispersion

$$\omega(k) = 2\sqrt{\frac{\beta}{m}} \left| \sin\left(\frac{ka}{2}\right) \right|, v_s = \lim_{k \rightarrow 0} \frac{d\omega}{dk} = a\sqrt{\frac{\beta}{m}}, \quad \text{and} \quad \omega_M \stackrel{k=\pi/a}{=} 2\sqrt{\frac{\beta}{m}} = \frac{2v_s}{a},$$

and the Debye frequency for 1D chain is $\omega_D = v_s k = \frac{\pi v_s}{a}$. Since $\omega_M / \omega_D = 2/\pi < 1$, the maximum frequency in the real dispersion is smaller than in the Debye model.

(b) In 1D chain, the number of states within dk is $2 \cdot \frac{L}{2\pi} dk$. So, the density of states is

$$g(\omega) = \frac{L}{\pi} \frac{dk}{d\omega} = \frac{2L}{\pi a \omega_M} \frac{1}{\sqrt{1 - (\omega/\omega_M)^2}} \stackrel{L=Na}{=} \frac{2N}{\pi \omega_M} \frac{1}{\sqrt{1 - (\omega/\omega_M)^2}}.$$

(c) The phonon thermal occupation

$$n(\omega) = \left[\exp\left(\frac{\hbar\omega}{k_B T}\right) - 1 \right]^{-1}$$

Then the internal energy is

$$U(T) = \int_0^{\omega_M} \hbar\omega n(\omega) g(\omega) d\omega = \int_0^{\omega_M} \frac{\hbar\omega}{e^{\frac{\hbar\omega}{k_B T}} - 1} g(\omega) d\omega.$$

and the heat capacity

$$C_V(T) = \frac{dU}{dT} = \int_0^{\omega_M} \frac{\partial n(\omega)}{\partial T} g(\omega) d\omega$$

At high-temperature limit, i.e., $\hbar\omega_M \ll k_B T$. The expression of internal energy and heat capacity become

$$U(T) = \int_0^{\omega_M} k_B T g(\omega) d\omega = N k_B T, \quad C_V(T) = N k_B.$$

where applied the total number of vibration modes $\int_0^{\omega_M} g(\omega) d\omega = N$.

- (d) At low temperature limit, only small k matters, i.e., the main contributions come from small ω . Then, $g(\omega) = \frac{L}{\pi v_s}$. The internal energy becomes

$$U = \frac{L}{\pi v_s} \int_0^\infty \frac{\hbar\omega}{e^{\frac{\hbar\omega}{k_B T}} - 1} d\omega = \frac{\pi L (k_B T)^2}{6\hbar v_s}$$

and the capacity

$$C_v = \frac{dU}{dT} = \frac{\pi N a k_B^2 T}{3\hbar v_s} \propto T$$

the same conclusion as the Debye model.

Lecture #3 Crystal lattice

Problem 3.1 (Crystallographic restriction theorem). Proof the crystallographic restriction theorem: Consider a crystal (i.e. with translational symmetry) in three dimensions that is invariant under rotations by an angle $2\pi/n$ around an axis. Then the only allowed possibilities are $n = 1, 2, 3, 4, 6$ with the case $n = 1$ being the trivial case with no rotational symmetry.

Solution (By TA). Rotation matrix

$$\mathbf{R} = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The vector

$$A = (\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)$$

after rotation

$$(L_1, L_2, L_3)^T = A^{-1} R A (L_1, L_2, L_3)^T$$

Since $\text{Tr}(A^{-1} R A) = \text{Tr}(R)$, then, $2 \cos \alpha + 1 = \mathbb{Z}$.

Solution. *Proof.* Denote the 3D lattice can be generated by the basis vectors $\mathbf{e}_1$, $\mathbf{e}_2$, and $\mathbf{e}_3$. So, any lattice point vector set should be

$$\mathbf{L} = \{a_1 \mathbf{e}_1 + a_2 \mathbf{e}_2 + a_3 \mathbf{e}_3\}.$$

Obviously, $a_i \in \mathbb{Z}$, i.e., any lattice point can be labeled by integer coordinates (a_1, a_2, a_3) . Consider the 3D rotation matrix (operator)

$$\mathbf{R} = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$


When acting it on the vector $\mathbf{L}$, it should leave the lattice invariant, i.e., the trace must be integers

$$\text{Tr}(\mathbf{R}) = 2 \cos \theta + 1 \in \mathbb{Z}.$$

So we have $2 \cos \theta = 2 \cos(2\pi/n) \in \mathbb{Z}$, i.e., $\cos(2\pi/n) = -1, -1/2, 0, 1/2, 1$, then $n \in \{1, 2, 3, 4, 6\}$. $\square$

Problem 3.2 (2D triangular lattice). Let us consider a two-dimensional triangular lattice as shown in the following figure.

- Identify the primitive unit cell of the two-dimensional triangular lattice. Find the basis vectors.
- Construct the basis vectors of the reciprocal unit cell.



Solution (By TA). $\mathbf{a}_i \mathbf{b}_j = 2\pi \delta_{ij}$, $\mathbf{b}_i = (b_{1i}, b_{2i})$.

$$(\mathbf{a}_1, \mathbf{a}_2)(\mathbf{b}_1, \mathbf{b}_2)^T = 2\pi \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Solution.

(a) Denote the lattice constant a . Then the simplest choice of primitive lattice vectors is

$$\mathbf{a}_1 = a(1, 0), \quad \mathbf{a}_2 = a\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right).$$

(b) Denote the two reciprocal-lattice primitive vectors $\mathbf{b}_1 = (b_{1x}, b_{1y})$, $\mathbf{b}_2 = (b_{2x}, b_{2y})$. Using the relation

$$\mathbf{a}_i \cdot \mathbf{b}_j = 2\pi \delta_{ij}.$$

To expand it

$$\mathbf{a}_1 \cdot \mathbf{b}_1 = 2\pi, \quad \mathbf{a}_2 \cdot \mathbf{b}_1 = 0, \quad \mathbf{a}_1 \cdot \mathbf{b}_2 = 0, \quad \mathbf{a}_2 \cdot \mathbf{b}_2 = 2\pi,$$

we obtain the two reciprocal-lattice primitive vectors

$$\mathbf{b}_1 = \frac{2\pi}{a} \left(1, -\frac{1}{\sqrt{3}}\right), \quad \mathbf{b}_2 = \frac{2\pi}{a} \left(0, \frac{2}{\sqrt{3}}\right).$$

Problem 3.3 (Properties of the reciprocal lattice).

- (a) Show that all reciprocal lattice vectors of the form $\mathbf{G} = h\mathbf{A} + k\mathbf{B} + l\mathbf{C}$ are perpendicular to the planes of the same indices (h, k, l) in real space.
- (b) Show that the distance $d_{h,k,l}$ between two consecutive planes (h, k, l) is inversely proportional to $|\mathbf{G}_{h,k,l}|$.
- (c) Find $d_{h,k,l}$ for a simple cubic lattice and an orthorhombic lattice ($a \neq b \neq c$, $\alpha = \beta = \gamma = \pi/2$).

Solution. “

(a) *Proof.* Substitute the expressions of the reciprocal lattice vectors to $\mathbf{G}$

$$\mathbf{G} = \frac{2\pi}{V} [h(\mathbf{a}_2 \times \mathbf{a}_3) + k(\mathbf{a}_3 \times \mathbf{a}_1) + l(\mathbf{a}_1 \times \mathbf{a}_2)].$$

Since the Miller indices (h, k, l) correspond to a plane that intercepts the crystal axes at $\mathbf{a}_1/h$, $\mathbf{a}_2/k$, $\mathbf{a}_3/l$, then, any vector lying in the (h, k, l) plane can be written as a linear combination of two independent vectors in that plane, i.e.,

$$\mathbf{r}_1 = \frac{\mathbf{a}_2}{k} - \frac{\mathbf{a}_1}{h}, \quad \mathbf{r}_2 = \frac{\mathbf{a}_3}{l} - \frac{\mathbf{a}_2}{k}.$$

Then, the normal to this plane can be written as a cross product

$$\hat{\mathbf{n}}_{hkl} = \mathbf{r}_1 \times \mathbf{r}_2 = \frac{1}{kl}(\mathbf{a}_2 \times \mathbf{a}_3) + \frac{1}{hl}(\mathbf{a}_3 \times \mathbf{a}_1) + \frac{1}{hk}(\mathbf{a}_1 \times \mathbf{a}_2).$$

which is parallel to $\mathbf{G}$, implies that $\mathbf{G}$ is perpendicular to the (h, k, l) plane. □

(b) *Proof.* For the vector pointing to the $n + 1$ and n -th (h, k, l) plane, it satisfies

$$\mathbf{G} \cdot \mathbf{r}_{n+1} = 2\pi(n + 1), \quad \text{and} \quad \mathbf{G} \cdot \mathbf{r}_n = 2\pi n,$$

where $\mathbf{r}_{n+1} - \mathbf{r}_n = d\hat{\mathbf{n}}$. Combine the two relations

$$\mathbf{G} \cdot (\mathbf{r}_{n+1} - \mathbf{r}_n) = \mathbf{G} \cdot \hat{\mathbf{n}}d = 2\pi d.$$

So, we have

$$d = \frac{2\pi}{\mathbf{G} \cdot \hat{\mathbf{n}}} = \frac{2\pi}{|\mathbf{G}_{h,k,l}|},$$

i.e., the distance between two consecutive planes is inversely proportional to $|\mathbf{G}_{h,k,l}|$. $\square$

(c) For a simple cubic lattice

$$\mathbf{a} = (a, 0, 0), \quad \mathbf{b} = (0, b, 0), \quad \mathbf{c} = (0, 0, c).$$

Then, the reciprocal vectors can be expressed as

$$\mathbf{G} = \frac{2\pi h}{a}\hat{\mathbf{x}} + \frac{2\pi k}{b}\hat{\mathbf{y}} + \frac{2\pi l}{c}\hat{\mathbf{z}},$$

and its magnitude

$$|\mathbf{G}| = 2\pi \sqrt{\frac{h^2}{a^2} + \frac{k^2}{b^2} + \frac{l^2}{c^2}}.$$

So, we obtain the distance $d = \frac{2\pi}{|\mathbf{G}|} = \left(\frac{h^2}{a^2} + \frac{k^2}{b^2} + \frac{l^2}{c^2}\right)^{-1/2}$.

Problem 3.4 (Atomic planes and Miller indices: application to lithium). The Bravais lattice of lithium is simple cubic with lattice parameter $a = 3.48 \text{ \AA}$.

- Suppose that the atoms (assumed to be spheres) are placed along the $[111]$, represent the atomic distribution along the following faces (100), (110), (111), and (201).
- For each such 2D structure, state the direction and basis vector, $\mathbf{a}$ and $\mathbf{b}$ of the elementary lattice as well as the value of the angle, γ .
- Find the atomic concentration and the mass density of lithium ($A \approx 7$).

Solution.

- (b) The distribution, basis vector, and the angle of the faces are listed.

2D Structure	Distribution	$\mathbf{a}$	$\mathbf{b}$	γ
(100)	square	$(0, a, 0)$	$(0, 0, a)$	90°
(110)	rectangular	$(a, -a, 0)$	$(0, 0, a)$	90°
(111)	hexagonal	$(a, 0, -a)$	$(0, a, -a)$	60°
(201)	rectangular	$(0, a, 0)$	$(a, 0, -2a)$	90°

- (c) Since the atoms are placed along the $[111]$, then every cubic lattice's corners has an atom, i.e., it is equivalent that there is an atom in every lattice. So, the atomic concentration is

$$n = \frac{1}{V} = \frac{1}{a^3} = 2.3728 \times 10^{28} \text{ atoms/m}^3.$$

The mass density is

$$\rho = nm_{\text{atom}} = \frac{nA}{N_A} = 275.816 \text{ kg/m}^3.$$

Lecture #4 X-ray diffraction

Problem 4.1 (Structure factor for a tri-atomic basis). Consider a linear crystal with lattice constant a in which the basis consists of atom of species A at 0 and two atoms of species B at $1/3$ and $2/3$.

- Find the structure factor of such a crystal.
- X-rays of wavelength $\lambda (= a/4)$ irradiate this crystal at an oblique incidence (45°). Using the Ewald construction in the plane of incidence, show graphically, the diffraction directions and indicate the relative weights of the different diffracted intensities.
- State the physical significance of the vector $\mathbf{k}$ from which the extremity is on a plane passing through the origin of the reciprocal lattice.

Solution (By TA). (a)

$$S_a = \sum_j f_j e^{-i\mathbf{a} \cdot \mathbf{r}_j} = f_A e^{-i\mathbf{G} \cdot 0} + f_B (e^{-i\mathbf{G} \cdot \frac{a}{3}\hat{x}} + e^{-i\mathbf{a} \cdot \frac{2}{3}a\hat{x}}) = f_A + 2f_B \cos\left(\frac{2\pi}{3}h\right) = \begin{cases} f_A + 2f_B, \\ f_A - f_B \end{cases}$$

$$(b) \quad k = \frac{2\pi}{\lambda} = \frac{8\pi}{a},$$

$$k \cos 45^\circ - k \cos \theta = G_x$$

$$\cos \theta = \frac{h+2\sqrt{2}}{4} \in [-1, 1].$$

Solution.

- Substitute $r_i = 0, \frac{a}{3}, \frac{2a}{3}$ into the definition of the structure factor

$$S(h) = f_A e^{i\mathbf{G} \cdot 0} + f_B e^{i\mathbf{G} \cdot \frac{a}{3}} + f_B e^{i\mathbf{G} \cdot \frac{2a}{3}} \stackrel{G=2\pi h/a}{=} f_A + f_B \left(e^{i\frac{2\pi h}{3}} + e^{i\frac{4\pi h}{3}} \right)$$

$$\stackrel{\text{Euler's formula}}{=} \begin{cases} f_A + 2f_B, & h = 3n \\ f_A - f_B, & \text{otherwise} \end{cases}$$

- The wave number

$$k = \frac{2\pi}{\lambda} = \frac{8\pi}{a} = 4b_0,$$

where $b_0 = \frac{2\pi}{a}$ is the reciprocal lattice vector. The incident wavevector is


$$\mathbf{k}_i = \left(k/\sqrt{2}, -k/\sqrt{2} \right)$$

Then, the Ewald circle satisfies

$$\left(x + k/\sqrt{2} \right)^2 + \left(y - k/\sqrt{2} \right)^2 = k^2$$

and we can obtain the two intersection on x -axis: $x_1 = 0$ and $x_1 = -\sqrt{2}k$.

Obviously, $x_1 = 0b_0, n_1 = 0; x_2 = -5.656b_0, n_2 = -5.656$.



Since n is not an integer, no other reciprocal lattice points lie on the Ewald circle except the one on the original point. The intensity of each diffraction order is proportional to $|S(h)|^2$, i.e.,

$$I_h \propto |S(h)|^2 = \begin{cases} (f_A + 2f_B)^2, & h = 3m, \\ (f_A - f_B)^2, & \text{otherwise} \end{cases}$$

- (c) The tip of $\mathbf{k}$ lying on a plane through the origin of the reciprocal lattice means the Bragg condition is satisfied for diffraction.

Problem 4.2. The main experimental methods of X-ray diffraction are

- (a) the Laue method
- (b) the crystal rotation method
- (c) the powder method (or the Debye-Scherrer method)

Describe the corresponding experimental set-up and with the help of the Ewald construction in reciprocal space, deduce the observed diffraction patterns in each case.

Solution.

- (a) Laue Method

- i. Setup: Stationary single crystal, polychromatic X-rays.
- ii. Ewald & Pattern: A continuous range of Ewald sphere radii (due to varying λ) means many reciprocal lattice points lie on some sphere. This produces a pattern of sharp spots, each from a different (hkl) plane and wavelength. The spot arrangement directly reveals the crystal's symmetry.

- (b) Crystal Rotation Method

- i. Setup: Rotating single crystal, monochromatic X-rays.
- ii. Ewald & Pattern: The fixed Ewald sphere is intersected by rotating reciprocal lattice points. This generates discrete spots or arcs on the detector. Measuring the rotation angles and intensities of these spots allows for the determination of the unit cell and atomic structure.

- (c) Powder Method

- i. Setup: Powdered sample (random crystallites), monochromatic X-rays.

- ii. Ewald & Pattern: Each set of (hkl) planes forms a spherical shell in reciprocal space. The Ewald sphere's intersection with these shells creates cones of diffracted beams (Debye cones), which appear as characteristic rings on the detector. The ring positions (2θ angles) provide d -spacings for phase identification and lattice parameter calculation.

Lecture #5 Nearly free electron model

Problem 5.1 (Compare Kronig-Penney model with the nearly free electron model). The Kronig-Penney model allows one to consider both the nearly-free-electron and tight-binding limits. Consider the (repulsive) Dirac-Kronig-Penney model. It gives the following implicit equation for the energy bands

$$\cos(ka) = \cos(qa) + u \frac{\sin(qa)}{qa}$$

where $q = \sqrt{2mE}/\hbar$ and $u = \frac{mU_0a}{\hbar^2} > 0$. Assuming that $u \ll q$, find the solution of above equation near the Bragg points $k \approx \pm \frac{\pi}{a}, \pm \frac{2\pi}{a}, \dots$. Compare your result with that of the nearly-free electron model. (Reminder: the $\approx$ sign means that k is close yet not equal to the wavenumber of the Bragg point.)

Solution. KP model $k = \frac{n\pi}{a} + \delta$.

$$\cos ka = \cos qa + u \frac{\sin qa}{qa}$$

Since

$$\cos(n\pi + \delta a) = \cos n\pi \cos \delta a - \sin n\pi \sin \delta a = (-1)^n \cos \delta a = (-1)^n \left[1 - \frac{(\delta a)^2}{2} \right]$$

The energy

$$E^2 = \frac{\hbar^2}{2m} \left(\frac{n\pi}{a} \right)^2 + \Delta E, \quad q = \frac{\sqrt{2mE}}{\hbar} = \frac{n\pi}{a} + \epsilon$$

Then the cosine and sine

$$\cos(qa) = (-1)^n \left[1 - \frac{(\epsilon a)^2}{2} \right], \quad \frac{\sin(qa)}{qa} = \frac{(-1)^n \epsilon a}{n\pi}$$

We can obtain an equation about ϵ

$$\epsilon^2 - \frac{2U}{an\pi} \epsilon - \delta^2 = 0$$

NFE: Expand into a Fourier mode

$$V(x) = \sum_a V_a e^{iGx} = U_0 \delta(x - na)$$

$$k = \frac{n\pi}{a} + \Delta, \quad k' = \frac{n\pi}{a} - \Delta$$

The wavefunction

$$\psi(x) = a\psi_k^0 + b\psi_{k'}^0$$

where $\psi_k^0 = \frac{1}{\sqrt{L}} e^{ikx}$. The eigenequation

$$(H_0 + H')\psi(x) = E\psi(x)$$

where $H_0 = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + \bar{V}$, $\Delta V = V(x) - \bar{V}$.

$$H_0\psi_k^0 = E_k^0\psi_k^0, \quad H_0\psi_{k'}^0 = E_{k'}^0\psi_{k'}^0$$

$$a(E_k^0 - E + \Delta V)\psi_k^0 + b(E_{k'}^0 - E + \Delta V)\psi_{k'}^0 = 0$$

$$\begin{cases} (E_k^0 - E)a + V_n^*b = 0, \\ V_n^a(E_{k'}^0 - E)b = 0 \end{cases}$$

$$V_n = \langle k|V(x)|k \rangle = \frac{1}{a} \int_0^a U_0 \delta(x) e^{-i2\pi mx/a} dx$$

The Determine should be zero

$$\begin{vmatrix} E_k^0 - E & V_n^* \\ V_n & E_{k'}^0 - E \end{vmatrix} = 0$$

$$E_k^0 = E_{k'}^0,$$

$$E_{\pm} = \frac{1}{2} \{E_k^0 + E_{k'}^0 \pm \sqrt{(E_k^0 - E_{k'}^0)^2 + 4|V_n|^2}\}, \quad E_{\text{gap}} = 2|V_n| = \frac{2U_0}{a}$$

Solution. Near the Bragg points, assume

$$ka = n\pi + \kappa a, \quad \text{and} \quad qa = q_n a + \delta.$$

with $|\kappa|, |\delta| \ll 1$, $q_n = n\pi/a$. Keep the 2nd order of κ and δ , the implicit equation can be written as

$$(-1)^n \left[1 - \frac{1}{2}(\kappa a)^2 \right] = (-1)^n \left(1 - \frac{1}{2}\delta^2 \right) + u \frac{(-1)^n \delta}{n\pi}.$$

Then, we can have the solution of δ

$$\delta = \frac{u}{n\pi} \pm \sqrt{\left(\frac{u}{n\pi} \right)^2 + (\kappa a)^2}.$$

Since $q = \sqrt{2mE}/\hbar$, then we have

$$E = \frac{\hbar^2(q_n + \delta/a)^2}{2m} \cong E_n^0 + \frac{\hbar^2 q_n \delta/a}{m},$$

where $E_n^0 = \frac{\hbar^2 q_n^2}{2m}$. Substitute $q_n = n\pi/a$, and the solution of δ into the expression of E

$$E = E_n^0 + \frac{n\pi \hbar^2 \delta}{m} = E_n^0 + \frac{n\pi \hbar^2}{ma^2} \left[\frac{u}{n\pi} \pm \sqrt{\left(\frac{u}{n\pi} \right)^2 + (\kappa a)^2} \right].$$

Substitute $u = mU_0 a/\hbar^2$ into the expression above

$$E = E_n^0 + \frac{U_0}{a} \pm \sqrt{\left(\frac{U_0}{a} \right)^2 + \left(\frac{n\pi \hbar^2 \kappa}{ma} \right)^2},$$

with the band gap

$$\Delta E_n = E_+ - E_- = 2\sqrt{\left(\frac{U_0}{a} \right)^2 + \left(\frac{n\pi \hbar^2 \kappa}{ma} \right)^2}.$$

When $\kappa a \rightarrow 0$, the energy band and the band gap become

$$E = E_n^0 + \frac{U_0}{a} \pm \frac{U_0}{a}, \quad \text{and} \quad \Delta E_n = \frac{2U_0}{a}.$$

The results perfectly match that of the nearly-free electron model.

Problem 5.2 (Nearly free electron model). Consider 2D electrons subject to a weak periodic potential

$$U(x, y) = U_0 \left(\cos \frac{2\pi x}{a} + \cos \frac{2\pi y}{a} \right).$$

- (a) Find the energy bands near the edges of the first Brillouin zone. (Hint: the edges of the Brillouin zone are the Bragg planes where the eigenstates are doubly degenerate.) Plot the bands and isoenergetic surfaces.
- (b) Repeat the analysis of part (a) for regions near the corners of the first Brillouin zone, where there are four degenerate eigenstates.

Solution (By TA). (a)

$$U(\mathbf{r}) = \sum_{\mathbf{G}} U_{\mathbf{G}} e^{i\mathbf{G}\cdot\mathbf{r}}, \quad U_{\mathbf{G}} = \frac{1}{V_{\text{Cell}}} \int_{\text{Cell}} U(\mathbf{r}) e^{-i\mathbf{G}\cdot\mathbf{r}} d\mathbf{r}$$

$$U = U_0 \left(\cos \frac{2\pi x}{a} + \cos \frac{2\pi y}{a} \right) = \frac{U_0}{2} (e^{\pm i \frac{2\pi x}{a}} + e^{\pm i \frac{2\pi y}{a}})$$

$$\mathbf{G} = (\pm \frac{2\pi}{a}, 0), (0, \pm \frac{2\pi}{a}), U_{\mathbf{G}} = \frac{U_0}{2}.$$

$$k_x = \frac{\pi}{a} + \delta, k_y.$$

$$\begin{vmatrix} E_k^0 - E & \frac{U_0}{2} \\ \frac{U_0}{2} & E_{k'}^0 - E \end{vmatrix} = 0$$

$$E = \frac{\hbar^2}{2m} \left(\frac{\pi^2}{a^2} + q_x^2 + k_y^2 \right) \pm \sqrt{\left(\frac{\pi \hbar^2}{ma} q_x \right)^2 + \left(\frac{U_0}{2} \right)^2}$$

$$k_x = \pm \frac{\pi}{a} + q_x, \quad k_y = \pm \frac{\pi}{a} + q_x.$$

- (b) $\langle k_i | U(\mathbf{r}) | k_j \rangle \neq 0, \mathbf{k}_i - \mathbf{k}_j = (0, \pm \frac{2\pi}{a}), (\pm \frac{2\pi}{a}, 0).$

Solution.

- (a) In reciprocal space, the reciprocal lattice vectors are

$$\mathbf{G}_1 = \left(\frac{2\pi}{a}, 0 \right), \quad \mathbf{G}_2 = \left(0, \frac{2\pi}{a} \right).$$

The potential can be expressed as

$$U(x, y) = \frac{U_0}{2} \left[e^{i\mathbf{G}_1 \cdot \mathbf{r}} + e^{-i\mathbf{G}_1 \cdot \mathbf{r}} + e^{i\mathbf{G}_2 \cdot \mathbf{r}} + e^{-i\mathbf{G}_2 \cdot \mathbf{r}} \right].$$

At the zone edge (e.g., $k_x = \pi/a$), consider a wavevector

$$\mathbf{k} = \left(\frac{\pi}{a} + q_x, k_y \right),$$




with $|q_x| \ll \pi/a$. The two nearly degenerate plane waves are $|\mathbf{k}\rangle$ and $|\mathbf{k} - \mathbf{G}_1\rangle$. Their free-electron energies are

$$E^{(0)}(\mathbf{k}) = \frac{\hbar^2}{2m} \left[\left(\frac{\pi}{a} + q_x \right)^2 + k_y^2 \right],$$

$$E^{(0)}(\mathbf{k} - \mathbf{G}_1) = \frac{\hbar^2}{2m} \left[\left(-\frac{\pi}{a} + q_x \right)^2 + k_y^2 \right].$$

Expanding to first order in q_x

$$E^{(0)} \approx E_0 \pm \alpha q_x, \quad \text{where} \quad E_0 = \frac{\hbar^2}{2m} \left[\left(\frac{\pi}{a} \right)^2 + k_y^2 \right], \quad \alpha = \frac{\hbar^2 \pi}{ma}.$$



The perturbation couples these states with matrix element $\langle \mathbf{k} | U | \mathbf{k} - \mathbf{G}_1 \rangle = U_0/2$. The Hamiltonian in the subspace $\{|\mathbf{k}\rangle, |\mathbf{k} - \mathbf{G}_1\rangle\}$ is

$$H = \begin{pmatrix} E_0 + \alpha q_x & U_0/2 \\ U_0/2 & E_0 - \alpha q_x \end{pmatrix}.$$

Diagonalizing gives the eigenenergies

$$E_{\pm} = E_0 \pm \sqrt{(\alpha q_x)^2 + \left(\frac{U_0}{2} \right)^2} = E_0 + \frac{\hbar^2 q_x^2}{2m} \pm \sqrt{\left(\frac{\hbar^2 \pi}{ma} q_x \right)^2 + \left(\frac{U_0}{2} \right)^2}$$

- (b) The corners of the first Brillouin zone are points like $\mathbf{k}_0 = (\pi/a, \pi/a)$ (the M point). The four degenerate plane waves at this point are

$$\begin{aligned} \mathbf{k}_1 &= \mathbf{k}_0 = (+\pi/a, +\pi/a), \\ \mathbf{k}_2 &= \mathbf{k}_0 - \mathbf{G}_1 = (-\pi/a, +\pi/a), \\ \mathbf{k}_3 &= \mathbf{k}_0 - \mathbf{G}_2 = (+\pi/a, -\pi/a), \\ \mathbf{k}_4 &= \mathbf{k}_0 - \mathbf{G}_1 - \mathbf{G}_2 = (-\pi/a, -\pi/a). \end{aligned}$$

All have the same free-electron energy $E_0 = \frac{\hbar^2}{2m} [(\pi/a)^2 + (\pi/a)^2] = \frac{\hbar^2 \pi^2}{ma^2}$.

In the basis $\{|\mathbf{k}_1\rangle, |\mathbf{k}_2\rangle, |\mathbf{k}_3\rangle, |\mathbf{k}_4\rangle\}$, states are coupled if they differ by $\pm \mathbf{G}_1$ or $\pm \mathbf{G}_2$. The non-zero off-diagonal matrix elements are

$$\langle \mathbf{k}_1 | U | \mathbf{k}_2 \rangle = \langle \mathbf{k}_1 | U | \mathbf{k}_3 \rangle = \langle \mathbf{k}_2 | U | \mathbf{k}_4 \rangle = \langle \mathbf{k}_3 | U | \mathbf{k}_4 \rangle = U_0/2.$$

The Hamiltonian is

$$H = E_0 \mathbb{1} + \frac{U_0}{2} \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}.$$

Diagonalizing this matrix yields eigenvalues

$$E = E_0 + \frac{U_0}{2} \times \{2, -2, 0, 0\} = \{E_0 + U_0, E_0 - U_0, E_0, E_0\}.$$

Thus, at $\mathbf{q} = 0$, the fourfold degeneracy splits into three distinct levels: one non-degenerate band at $E_0 + U_0$, one non-degenerate band at $E_0 - U_0$, and a two-fold degenerate band at E_0 .

Lecture #6 Tight binding model

Problem 6.1 (Tight bindings in the BCC and FCC lattices). We consider identical atoms located at points of a lattice that are body-centered cubic (**BCC: i**) and face-centered cubic (**FCC: ii**).

(a) For each case, determine the dispersion relation for s -electrons, having the general form

$$E = -\alpha - \gamma \sum_s e^{-i\mathbf{k} \cdot \mathbf{R}_s},$$

where α and γ are the positive energies that can be calculated and $\mathbf{R}_s$ represents the vectors that connect an atom located at the origin to each of its nearest neighbors.

(b) Find the expressions for E at the points located at the intersection of the first Brillouin zone with the directions $[100]$, $[110]$, and $[111]$ and thus the value $E(\Gamma)$ to the center of this zone.

(c) Deduce the total width of the corresponding band, ΔE , and find the expression for the effective mass m^* in the vicinity of Γ as well as the shape of constant energy surfaces around this point.

Solution (By TA). $\frac{a}{2}(\pm 1, \pm 1, \pm 1)$

$$\sum_s e^{-i\mathbf{k} \cdot \mathbf{R}_s} = \sum_{u,v,w} \left[-\frac{a}{2} i(uk_x + vk_y + wk_z) \right] = 8 \cos\left(\frac{ak_x}{2}\right) \cos\left(\frac{ak_y}{2}\right) \cos\left(\frac{ak_z}{2}\right)$$

$$\frac{a}{2}(\pm 1, 0, \pm 1), \frac{a}{2}(\pm 1, \pm 1, 0), \frac{a}{2}(0, \pm 1, \pm 1).$$

For BCC $E(\Gamma) = E(0, 0, 0) = -\alpha - 8\gamma$.

$$|\mathbf{k}|^2 = |\mathbf{k} - \mathbf{G}|^2 = |\mathbf{k}|^2 - 2\mathbf{k} \cdot \mathbf{G} + |\mathbf{G}|^2, \quad 2\mathbf{k} \cdot \mathbf{G} = |\mathbf{G}|^2$$

The nearest $\frac{2\pi}{a}(\pm 1, 0, \pm 1)$

$$\left\{ k_x \pm k_y = \pm \frac{2\pi}{a}, \quad k_y \pm k_z = \pm \frac{2\pi}{a}, \quad k_z \pm k_x = \pm \frac{2\pi}{a} \right\}.$$

For $[1, 0, 0]$

$$\begin{cases} k_x = t, \\ k_y = k_z = 0 \end{cases}$$

$(\pm \frac{2\pi}{a}, 0, 0)$, $E = -\alpha + 8\gamma$. For $[1, 1, 0]$

$$\begin{cases} k_x = k_y = t, \\ k_z = 0 \end{cases}$$

$(\frac{\pi}{a}, \frac{\pi}{a}, 0)$, $E = -\alpha$. For $[1, 1, 1]$: $k_x = k_y = k_z = t$, $(\frac{\pi}{a}, \frac{\pi}{a}, \frac{\pi}{a})$, $E = -\alpha$.

For FCC:

$$E(\Gamma) = -\alpha - 12\gamma$$

Consider the nearest $\frac{2\pi}{a}(\pm 1, \pm 1, \pm 1)$.

$$\mathbf{k} \cdot \mathbf{a} = \left(\frac{2\pi}{a}\right)^2 \cdot \frac{3}{2},$$

$$\pm k_x \pm k_y \pm k_z = \frac{3\pi}{a}$$

Sub-nearest: $\frac{2\pi}{a}(\pm 2, 0, 0)$,

$$k_x = \pm \frac{2\pi}{a}, \quad k_y = \pm \frac{2\pi}{a}, \quad k_z = \pm \frac{2\pi}{a}.$$

For $[1, 0, 0]: (\frac{2\pi}{a}, 0, 0)$

$$\begin{cases} k_x = t, \\ k_y = k_z = 0 \end{cases}, \quad E = -\alpha + 4r$$

For $[1, 1, 0]: (\frac{3\pi}{2a}, \frac{3\pi}{2a}, 0)$

$$\begin{cases} k_x = k_y = t, \\ k_z = 0 \end{cases}, \quad E = -\alpha - 2\gamma + 4\sqrt{2}r$$

For $[1, 1, 1]: (\frac{\pi}{a}, \frac{\pi}{a}, \frac{\pi}{a})$

$$k_x = k_y = k_z = t, \quad E = -\alpha$$

$$E_k = E_0 + \frac{\hbar^2}{2m}(\mathbf{k} - \mathbf{k}_0)^2 \quad \text{for } k \rightarrow 0$$

Use some approximation

$$k \rightarrow 0, \quad \cos \frac{ak}{2} = 1 - \frac{a^2 k^2}{8}$$

BCC:

$$E_k = -\alpha - 8\gamma \left(1 - \frac{a^2 k_x^2}{8}\right) (\dots k_y) (\dots k_z) = -\alpha - 8\gamma + \gamma a^2 k^2$$

where $\gamma a^2 = \frac{\hbar^2}{2m^*}$, $m^* = \frac{\hbar^2}{2\gamma a}$.

Solution (i = BCC, ii = FCC). Assume the lattice constant a .

(a_i) Each atom has 8 NNs, i.e.,

$$\{\mathbf{R}_s\} = \left\{ \frac{a}{2}(\pm 1, \pm 1, \pm 1) \right\}$$

Then, the sum in the general form becomes

$$\sum_s e^{-i\mathbf{k} \cdot \mathbf{R}_s} = \sum_{\sigma_x = \pm} \sum_{\sigma_y = \pm} \sum_{\sigma_z = \pm} e^{-i\frac{a}{2}\sigma_x k_x} \cdot e^{-i\frac{a}{2}\sigma_y k_y} \cdot e^{-i\frac{a}{2}\sigma_z k_z} = 8 \cos\left(\frac{ak_x}{2}\right) \cos\left(\frac{ak_y}{2}\right) \cos\left(\frac{ak_z}{2}\right)$$

Hence, the dispersion relation is

$$E(\mathbf{k}) = -\alpha - 8\gamma \cos\left(\frac{ak_x}{2}\right) \cos\left(\frac{ak_y}{2}\right) \cos\left(\frac{ak_z}{2}\right)$$

- (b_i) i. Γ point: $\mathbf{k} = (0, 0, 0)$, $E(\Gamma) = -\alpha - 8\gamma$. iii. $[110]: \mathbf{k} = \frac{2\pi}{a}(\frac{1}{2}, \frac{1}{2}, 0)$, $E(N) = -\alpha$.
 ii. $[100]: \mathbf{k} = \frac{2\pi}{a}(1, 0, 0)$, $E(X) = -\alpha + 8\gamma$. iv. $[111]: \mathbf{k} = \frac{\pi}{a}(1, 1, 1)$, $E(P) = -\alpha$

(c_i) The cos terms in the general form differs between $[-8\gamma, +8\gamma]$. So, the energy gap is

$$\Delta E_{\text{BCC}} = (-\alpha + 8\gamma) - (-\alpha - 8\gamma) = 16\gamma$$

The dispersion relation can be expanded around the Γ point

$$E(\mathbf{k}) = -\alpha - 8\gamma \left[1 - \frac{1}{2} \left(\frac{ak_x}{2} \right)^2 - \frac{1}{2} \left(\frac{ak_y}{2} \right)^2 - \frac{1}{2} \left(\frac{ak_z}{2} \right)^2 \right] \frac{k^2 = k_x^2 + k_y^2 + k_z^2}{2\gamma a^2} - \alpha - 8\gamma + \gamma a^2 k^2$$

Then, the effective mass $m^* = \frac{\hbar^2}{\partial^2 E / \partial k^2} = \frac{\hbar^2}{2\gamma a^2}$.

(a_{ii}) Each atom has 12 NNs, i.e.,

$$\{\mathbf{R}_s\} = \left\{ \frac{a}{2}(\pm 1, \pm 1, 0), \frac{a}{2}(\pm 1, 0, \pm 1), \frac{a}{2}(0, \pm 1, \pm 1) \right\}$$

Then, the sum in the general form becomes

$$\begin{aligned} \sum_s e^{-i\mathbf{k} \cdot \mathbf{R}_s} &= \sum_{\text{cyclic}(x,y,z)} \sum_{\pm} e^{-i\frac{a}{2}(\pm k_i \pm k_j)} \\ &= 4 \left[\cos\left(\frac{ak_x}{2}\right) \cos\left(\frac{ak_y}{2}\right) + \cos\left(\frac{ak_y}{2}\right) \cos\left(\frac{ak_z}{2}\right) + \cos\left(\frac{ak_z}{2}\right) \cos\left(\frac{ak_x}{2}\right) \right] \end{aligned}$$

Hence, the dispersion relation is

$$E(\mathbf{k}) = -\alpha - 4\gamma \left[\cos\left(\frac{ak_x}{2}\right) \cos\left(\frac{ak_y}{2}\right) + \cos\left(\frac{ak_y}{2}\right) \cos\left(\frac{ak_z}{2}\right) + \cos\left(\frac{ak_z}{2}\right) \cos\left(\frac{ak_x}{2}\right) \right]$$

- (b_{ii}) i. Γ point: $\mathbf{k} = (0, 0, 0)$, $E(\Gamma) = -\alpha - 8\gamma$. ii. $[100]$: $\mathbf{k} = \frac{2\pi}{a}(1, 0, 0)$, $E(X) = -\alpha - 4\gamma$.
 iii. $[110]$ point: $\mathbf{k} = \frac{2\pi}{a}(\frac{3}{4}, \frac{3}{4}, 0)$, $E(N) = -\alpha + \gamma(4\sqrt{2} - 2)$.
 iv. $[111]$ point: $\mathbf{k} = \frac{\pi}{a}(1, 1, 1)$, $E(P) = -\alpha - 12\gamma$

(c_{ii}) The cos terms in the general form differs between $[-12\gamma, +4\gamma]$. So, the energy gap is

$$\Delta E_{\text{FCC}} = (-\alpha + 4\gamma) - (-\alpha - 12\gamma) = 16\gamma$$

The dispersion relation can be expanded around the Γ point


$$\begin{aligned} E(\mathbf{k}) &= -\alpha - 4\gamma \sum_{\text{cyclic}(x,y,z)} \left[1 - \frac{1}{2} \left(\frac{ak_i}{2} \right)^2 - \frac{1}{2} \left(\frac{ak_j}{2} \right)^2 \right] \\ &= -\alpha - 12\gamma + 4\gamma \left[\left(\frac{k_x a}{2} \right)^2 + \left(\frac{k_y a}{2} \right)^2 + \left(\frac{k_z a}{2} \right)^2 \right] = -\alpha - 12\gamma + \gamma a^2 k^2 \end{aligned}$$

Then, the effective mass $m^* = \frac{\hbar^2}{\partial^2 E / \partial k^2} = \frac{\hbar^2}{2\gamma a^2}$.

Lecture #7 Semiconductor

Problem 7.1 (Elementary study of an intrinsic semiconductor).

Consider a semiconductor characterized by a constant density of states ($g(E) = C$) in both the valence and conduction band. These bands, separated by E_g , have a respective width of E_V and E_C (see the Figure).



- (a) Knowing that the Fermi level E_F is in the bandgap ($5k_B T < E_F < E_g - 5k_B T$), leading to the conventional simplification for $f(E)$ (The Boltzmann approximation can be applied), find the expression for the number of electrons n in the conduction band and the number of holes h in the valence band respectively as a function of the data and absolute temperature.
- (b) Deduce the expression for the volumetric density of intrinsic carriers n_i at ambient temperature ($k_B T = 25 \text{ meV}$) as well as the position of the Fermi level E_{F_i} .
Numerical applications: $E_g = 0.7 \text{ eV}$, $E_C = E_V = 5 \text{ eV}$, and $C = 2 \times 10^{21} \text{ electrons/cm}^3/\text{eV}$.
- (c) At the same temperature, what is the intrinsic electronic conductivity of this material? Take $\mu_h = \mu_e = 1000 \text{ cm}^2/\text{Vs}$.
- (d) The material is doped with N_0 boron impurities. Find the expressions and the new values of n_d , h_d , σ_d , and E_{F_d} with $N_0 = 10^{12} \text{ impurities per cm}^3$ and next with $N_0 = 10^{14} \text{ impurities per cm}^3$.

Solution.

- (a) The number of electrons in the conduction band

$$n = \int_{E_g}^{E_g+E_C} g(E) f(E) dE = C k_B T e^{-\frac{E_g-E_F}{k_B T}} (1 - e^{-\frac{E_C}{k_B T}}) \xrightarrow{\text{Boltzmann approximation}} \frac{E_C \gg k_B T}{C k_B T e^{-\frac{E_g-E_F}{k_B T}}}.$$

The number of holes in the valence band

$$h = \int_{-E_V}^0 g(E) [1 - f(E)] dE \approx C k_B T e^{-\frac{E_F}{k_B T}} (1 - e^{-\frac{E_V}{k_B T}}) \xrightarrow{\text{Boltzmann approximation}} \frac{E_V \gg k_B T}{C k_B T e^{-\frac{E_F}{k_B T}}}.$$

where $f(E) \approx e^{-\frac{E-E_F}{k_B T}}$, $[1 - f(E)] \approx e^{\frac{E-E_F}{k_B T}}$.

- (b) For the intrinsic carriers, the number of electrons and holes are the same

$$n = C k_B T e^{-\frac{E_g-E_F}{k_B T}} = h = C k_B T e^{-\frac{E_F}{k_B T}}, \quad E_g - E_F = E_F.$$

Then, the Fermi level is at mid-gap, i.e., $E_F = \frac{E_g}{2}$. Substitute it back into the expression of n (or h), we get the intrinsic carrier concentration

$$n_i = C k_B T e^{-\frac{E_g}{2k_B T}}.$$

Apply the parameters, $n_i = 4.158 \times 10^{19} \text{ m}^{-3}$, $E_F = 0.35 \text{ eV}$.

(c) The conductivity

$$\sigma_i = e(n_i\mu_e + n_i\mu_h) = 1.332 \text{ S/m}.$$

(d) Boron is an acceptor, so doping concentration $N_A = N_0$. Now, the number of electrons and holes

$$n = n_i, \quad h = n_i \longrightarrow n_d = n, \quad h_d = n + N_0,$$

and they satisfy the identity: mass-action law

$$n_i^2 = n_d h_d = n(n + N_0).$$

We can solve that

$$n = \frac{\sqrt{N_0^2 + 4n_i^2} - N_0}{2}, \quad n_d = n = \frac{\sqrt{N_0^2 + 4n_i^2} - N_0}{2}, \quad h_d = n + N_0 = \frac{\sqrt{N_0^2 + 4n_i^2} + N_0}{2}.$$

The conductivity becomes

$$\sigma_d = e(n_d\mu_e + h_d\mu_h).$$

From $h = Ck_B T e^{-\frac{E_F}{k_B T}}$, we can obtain the Fermi level

$$E_{F_d} = k_B T \ln \frac{Ck_B T}{h_d}.$$

Apply the parameters (with the previous $n_i = 4.158 \times 10^{19} \text{ m}^{-3}$)

i. $N_0 = 10^{18} \text{ m}^{-3}$.

The numbers of the acceptors and donators

$$\begin{aligned} n_d &= 4.108 \times 10^{19} \text{ m}^{-3}, \\ h_d &= 4.208 \times 10^{19} \text{ m}^{-3}. \end{aligned}$$

The conductivity and the Fermi level

$$\sigma_d = 1.332 \text{ S/m}, \quad E_{F_d} = 0.350 \text{ eV}.$$

ii. $N_0 = 10^{20} \text{ m}^{-3}$.

The numbers of the acceptors and donators

$$\begin{aligned} n_d &= 1.503 \times 10^{19} \text{ m}^{-3}, \\ h_d &= 1.150 \times 10^{20} \text{ m}^{-3}. \end{aligned}$$

The conductivity and the Fermi level

$$\sigma_d = 2.083 \text{ S/m}, \quad E_{F_d} = 0.325 \text{ eV}.$$

Lecture #8 Electrons in a magnetic field

Problem 8.1 (Onsager formula). Suppose that the electron spectrum for a tetragonal lattice can be described by the following equation

$$E_k = \frac{\hbar^2 k_x^2}{2m_{xx}} + \frac{\hbar^2 k_y^2}{2m_{xx}} + \frac{\hbar^2 k_z^2}{2m_{zz}},$$

with $m_{xx} \neq m_{zz}$. Using the Onsager formula for the cyclotron mass, $m_c = \frac{\hbar^2}{2\pi} \frac{\partial A}{\partial E_k}$ with A being the area of an isoenergetic contour perpendicular to the magnetic field. Find m_c for the case when the magnetic field is

(a) Along the z -axis.

(b) In the $x - y$ plane.

Solution.

(a) For $\mathbf{B} \parallel \hat{z}$, the motion of electrons will be perpendicular to $\hat{z}$, i.e., parallel to $x - y$ plane. The equation of motion becomes

$$E_k = \frac{\hbar^2}{2m_{xx}}(k_x^2 + k_y^2),$$

with the equation of the motion boundary in k -space

$$k_{\parallel}^2 \equiv k_x^2 + k_y^2 = \frac{2m_{xx}E_k}{\hbar^2}.$$

Then, we get the radius. The circle area of the isoenergetic contour is

$$A = \pi k_{\parallel}^2 = \frac{2\pi m_{xx}E_k}{\hbar^2}.$$

Substituting to the definition of cyclotron mass, we can obtain

$$m_c = \frac{\hbar^2}{2\pi} \frac{\partial A}{\partial E_k} = \frac{\hbar^2}{2\pi} \frac{2\pi m_{xx}}{\hbar^2} = m_{xx}.$$

(b) Simply taking $\mathbf{B} \parallel \hat{x}$, then orbits lie in the $k_y - k_z$ plane with an ellipse area. The equation of motion becomes

$$E_k = \frac{\hbar^2 k_y^2}{2m_{xx}} + \frac{\hbar^2 k_z^2}{2m_{zz}},$$

with the equation of the motion boundary in k -space

$$\frac{k_y^2}{2m_{xx}E/\hbar^2} + \frac{k_z^2}{2m_{zz}E/\hbar^2} = 1.$$

Then we get the semi-axes. The ellipse area of the isoenergetic contour is

$$A = \pi \sqrt{\frac{2m_{xx}E}{\hbar^2}} \sqrt{\frac{2m_{zz}E}{\hbar^2}} = \frac{2\pi E}{\hbar^2} \sqrt{m_{xx}m_{zz}}$$

Substituting to the definition of cyclotron mass, we can obtain

$$m_c = \frac{\hbar^2}{2\pi} \frac{\partial A}{\partial E_k} = \sqrt{m_{xx}m_{zz}}.$$

Problem 8.2 (Integer Quantum Hall Effect). The Hall resistance R_{xy} of a two-dimensional electron gas is measured as a function of magnetic field B at a very low temperature. A series of plateaus are observed at values $R_{xy} = \frac{h}{\nu e^2}$, where ν is an integer. Simultaneously, the longitudinal resistance R_{xx} shows peaks and drops to zero.

- (a) Explain physically why R_{xx} drops to zero whenever R_{xy} is on a plateau.
- (b) A sample has an electron density of $n_s = 4.0 \times 10^{11} \text{ cm}^{-2}$.
 - i. Calculate the filling factor ν at $B = 2.0 \text{ T}$.
 - ii. What is the expected value of R_{xy} on the plateau nearest to this field?
 - iii. How many Landau levels are completely filled at this plateau?
- (c) Why is the IQHE used as a resistance standard? Which two fundamental constants are involved?

Solution.

- (a) When the Hall resistance R_{xy} is on a plateau, the Fermi energy lies in a mobility gap between Landau levels. All extended states in the bulk are absent, making the bulk insulating. Current flows only along dissipationless chiral edge states.

At the edges, backscattering is prevented by spatial separation: even if impurities or imperfections exist along an edge, they may cause small forward deviations in an electron's path but cannot reverse its direction, so no resistance is produced. With no extended states in the bulk to allow dissipation and no backscattering at the edges, the longitudinal voltage drop vanishes, giving $R_{xx} = 0$.

- (b) i. From the degeneracy of Landau level $n_s = \nu \frac{eB}{h}$, we can obtain the filling factor

$$\nu = \frac{n_s h}{eB} = \frac{4.0 \times 10^{15} \text{ m}^{-2} \times 6.626 \times 10^{-34} \text{ J} \cdot \text{s}}{1.602 \times 10^{-19} \text{ C} \times 2.0 \text{ T}} = 8.271.$$

- ii. Substituting the values into the expression of the Hall resistance

$$R_{xy} = \frac{R_K}{\tilde{\nu}} = \frac{h}{\tilde{\nu} e^2} = 3226.60 \Omega.$$

where $\tilde{\nu}$ is the nearest integer of ν .

- iii. The filling factor $\tilde{\nu} = 8$ directly corresponds to the number of completely filled Landau levels.
- (c) IQHE provides a quantized Hall resistance

$$R_H = \frac{h}{e^2},$$

which precisely quantized and only based on two fundamental constants: The Planck constant h and the elementary charge e , so it is used as a resistance standard.

Problem 8.3 (Lifshitz-Kosevich formula). The Lifshitz-Kosevich formula describes the amplitude of the Shubnikov-de Haas (SdH) oscillations. The amplitude ΔR is proportional to the factor: $R_T = \frac{\lambda(T)}{\text{sh } \lambda(T)}$, where $\lambda(T) = 2\pi^2 \frac{k_B T m^*}{\hbar e B}$.

- (a) A researcher measures the SdH oscillation amplitude ΔR at a fixed magnetic field B as a function of temperature T . How can they use this data to determine the effective mass m^* of the charge carriers?
- (b) Another researcher fixes the temperature T and measures the SdH amplitude ΔR as a function of $1/B$. They find that the amplitude decreases exponentially with $1/B$. What physical parameter does this decay reveal, and what does a rapid decay imply about the quality of the sample?

Solution.

- (a) Take $\alpha = \frac{2\pi^2 k_B T m^*}{\hbar e B}$, then $\lambda(T) = \alpha T$. Since at $T \gg 1$, the amplitude ΔR satisfies


$$\Delta R \propto R_T = \frac{\alpha T}{\text{sh}(\alpha T)} \rightarrow \frac{1}{2} \alpha T e^{-\alpha T}, \quad \Delta R = \tilde{C} T e^{-\alpha T}$$

Take the logarithm to the two sides

$$\ln \Delta R = \ln \tilde{C} + \ln T - \alpha T, \quad \ln \left(\frac{\Delta R}{T} \right) = -\alpha T + C,$$

where $C = \ln \tilde{C}$.

So, one can take the data at high temperature, and use some software tool to convert the original data from T - ΔR to T - $\ln \left(\frac{\Delta R}{T} \right)$, then plot the new data set like the right figure.



By fitting the slope $k = -\alpha$ that fits the figure, one can obtain the effective mass

$$m^* = \frac{\alpha \hbar e B}{2\pi^2 k_B} = -\frac{k \hbar e B}{2\pi^2 k_B}$$

- (b) Similarly to part (a), we take $T \gg 1$. Since ΔR is proportional to $R_T R_D R_S$, and if we only consider the magnetic variable B , then $\Delta R \propto R_T R_D$. To be more specific,

$$\Delta R \propto \frac{2\pi^2 k_B T m^*}{\hbar e} \cdot \frac{1}{B} \cdot e^{-\frac{2\pi^2 k_B T m^*}{\hbar e} \cdot \frac{1}{B}} \cdot e^{-\frac{\pi m^*}{e \tau_q} \cdot \frac{1}{B}} = \tilde{C} \cdot \frac{1}{B} \cdot e^{-(\alpha + \beta/\tau_q) \cdot \frac{1}{B}}$$

where the coefficients α and β can be easily known since we have already obtained the effective mass. Taking the logarithm to both sides, we can obtain

$$\ln(\Delta R) = C - \ln B - \left(\alpha + \frac{\beta}{\tau_q} \right) \frac{1}{B},$$

Since the effective mass m^* has been determined from (a), so the coefficients α and β are known, and the slope directly reveals the quantum scattering rate $1/\tau_q$.

A rapid exponential decay corresponds to a short τ_q , meaning a short mean free path and therefore significant disorder: an indicator of poor sample quality.

Lecture #9 Band structure of graphene

Problem 9.1. In the calculation of the monolayer graphene band structure using tight-binding model, if we keep the contribution of the next-nearest hopping t' .

- Write down the dispersion relation.
- Expand the energy spectrum around the Dirac point up to second order. Search the literature and check what's trigonal warping effect is in a monolayer.

Solution.

- The Hamiltonian in second quantization form for the graphene with the next-nearest (NN) hopping is

$$\mathcal{H} = -t \sum_{\langle i,j \rangle} (\hat{a}_i \hat{b}_j^\dagger + \text{H.c.}) - t' \sum_{\langle\langle i,j \rangle\rangle} (\hat{a}_i \hat{a}_j^\dagger + \hat{b}_i \hat{b}_j^\dagger + \text{H.c.}) = \mathcal{H}_t + \mathcal{H}_{t'},$$

where i and j label the nearest neighbor, i.e., the bond A_i, B_i .

The fermion operators a_i and $a_i^\dagger$ annihilate/create an electron at site A , also for b_i and $b_i^\dagger$ at site B . The vectors pointing from A_i to the 3 nearest B -type sites and 6 next-nearest A -type sites are

$$\begin{aligned} \delta_1 &= a(1, 0), & \delta_2 &= a\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), & \delta_3 &= a\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right); \\ a_1 &= a\left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right), & a_2 &= a(0, \sqrt{3}), & a_3 &= a\left(-\frac{3}{2}, \frac{\sqrt{3}}{2}\right), \\ a_4 &= a\left(-\frac{3}{2}, -\frac{\sqrt{3}}{2}\right), & a_5 &= a(0, -\sqrt{3}), & a_6 &= a\left(\frac{3}{2}, -\frac{\sqrt{3}}{2}\right). \end{aligned}$$

Assume there are N atoms in total, then there are $N/2$ A -type or B -type sites in total.


Applying the Fourier transformation to the fermion operators

$$\begin{aligned} \hat{a}_i &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}_i} \hat{a}_{\mathbf{k}}, & \hat{b}_{i+\delta} &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k},j} e^{i\mathbf{k}\cdot(\mathbf{r}_i+\delta_j)} \hat{b}_{\mathbf{k}}, \\ \hat{a}_i^\dagger &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}_i} \hat{a}_{\mathbf{k}}^\dagger, & \hat{b}_{i+\delta}^\dagger &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k},j} e^{-i\mathbf{k}\cdot(\mathbf{r}_i+\delta_j)} \hat{b}_{\mathbf{k}}^\dagger. \end{aligned}$$

Consider the first term of the Hamiltonian, it can be expressed as

$$\begin{aligned} \mathcal{H}_t &= -t \sum_{i \in A} \sum_{j=1}^3 \left(\frac{1}{N} \sum_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}_i} \hat{a}_{\mathbf{k}}^\dagger \sum_{\mathbf{k}'} e^{i\mathbf{k}'\cdot(\mathbf{r}_i+\delta_j)} \hat{b}_{\mathbf{k}'} + \text{h.c.} \right) \\ &= -t \frac{1}{N} \sum_{\mathbf{k}, \mathbf{k}'} \sum_{i \in A} e^{i(\mathbf{k}'-\mathbf{k})\cdot\mathbf{r}_i} \sum_{j=1}^3 e^{i\mathbf{k}'\cdot\delta_j} \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}'} + \text{h.c.} = -t \sum_{\mathbf{k}} \sum_{j=1}^3 e^{i\mathbf{k}\cdot\delta_j} \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} + \text{h.c.} \end{aligned}$$

where $\sum_{i \in A} e^{i(\mathbf{k}'-\mathbf{k})\cdot\mathbf{r}_i} = N\delta_{\mathbf{k}',\mathbf{k}}$ is applied, and multiple sum will not occur here since we only sum over every A -type site.



Denote $\Delta(\mathbf{k}) = -t \sum_{j=1}^3 e^{i\mathbf{k} \cdot \delta_j}$, this part becomes

$$\mathcal{H}_t = \sum_{\mathbf{k}} \Delta(\mathbf{k}) \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} + \sum_{\mathbf{k}} \Delta^*(\mathbf{k}) \hat{b}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}}$$

Similarly for the second part: Denote $\epsilon(\mathbf{k}) = -t' \sum_{m=1}^6 e^{i\mathbf{k} \cdot \mathbf{a}_m}$, one can obtain (The Hermit conjugate is applied, but the factor $\frac{1}{2}$ is also applied to prevent a twice sum between the same NN bond.)

$$\mathcal{H}_{t'} = \sum_{\mathbf{k}} \epsilon(\mathbf{k}) \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \sum_{\mathbf{k}} \epsilon(\mathbf{k}) \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}.$$

Here the $\epsilon(\mathbf{k})$ should not taken Hermit conjugate, since the hopping A_i-A_j and B_i-B_j are the same physical process. Hence, the total Hamiltonian is

$$\begin{aligned} \mathcal{H} &= \mathcal{H}_t + \mathcal{H}_{t'} = \sum_{\mathbf{k}} \left(\Delta(\mathbf{k}) \hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} + \Delta^*(\mathbf{k}) \hat{b}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \epsilon(\mathbf{k}) \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}} + \epsilon^*(\mathbf{k}) \hat{b}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}} \right) \\ &= \sum_{\mathbf{k}} \left(\hat{a}_{\mathbf{k}}^\dagger \hat{b}_{\mathbf{k}}^\dagger \right) \begin{pmatrix} \epsilon(\mathbf{k}) & \Delta(\mathbf{k}) \\ \Delta^*(\mathbf{k}) & \epsilon(\mathbf{k}) \end{pmatrix} \begin{pmatrix} \hat{a}_{\mathbf{k}} \\ \hat{b}_{\mathbf{k}} \end{pmatrix} = \sum_{\mathbf{k}} \Psi^\dagger \mathbf{H}_{2 \times 2} \Psi \end{aligned}$$

By taking $\det(\mathbf{H}_{2 \times 2} - E\mathbf{1}) = 0$, one can obtain the dispersion relation

$$(\epsilon(\mathbf{k}) - E)^2 = \Delta(\mathbf{k}) \Delta^*(\mathbf{k}), \quad E_{\pm} = \pm \sqrt{\Delta(\mathbf{k}) \Delta^*(\mathbf{k})} + \epsilon(\mathbf{k})$$

To obtain the exact expression, expanding $\Delta(\mathbf{k})$ and $\epsilon(\mathbf{k})$

$$\begin{aligned} \Delta(\mathbf{k}) &= -t \left[e^{i\mathbf{k} \cdot \mathbf{a}(1,0)} + e^{i\mathbf{k} \cdot \mathbf{a}\left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right)} + e^{i\mathbf{k} \cdot \mathbf{a}\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right)} \right] \\ &= -t \left(e^{ik_x a} + e^{-\frac{1}{2}ik_x a} \cdot e^{\frac{\sqrt{3}}{2}ik_y a} + e^{-\frac{1}{2}ik_x a} \cdot e^{-\frac{\sqrt{3}}{2}ik_y a} \right) \\ &= -t e^{-ik_x a} \left[1 + 2 e^{\frac{3}{2}ik_x a} \cos\left(\frac{\sqrt{3}}{2}k_y a\right) \right], \\ \Delta(\mathbf{k}) \Delta^*(\mathbf{k}) &= t^2 \left[3 + 2 \cos(\sqrt{3}k_y a) + 4 \cos\left(\frac{3}{2}k_x a\right) \cos\left(\frac{\sqrt{3}}{2}k_y a\right) \right], \\ &\quad \underbrace{\hspace{10em}}_{f(\mathbf{k})} \\ \epsilon(\mathbf{k}) &= -t' \left[e^{ika\left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right)} + e^{ika(0, \sqrt{3})} + e^{ika\left(-\frac{3}{2}, \frac{\sqrt{3}}{2}\right)} \right. \\ &\quad \left. + e^{ika\left(-\frac{3}{2}, -\frac{\sqrt{3}}{2}\right)} + e^{ika(0, -\sqrt{3})} + e^{ika\left(\frac{3}{2}, -\frac{\sqrt{3}}{2}\right)} \right] \\ &= -t' \left[e^{\sqrt{3}ik_y a} + 2 e^{\frac{\sqrt{3}}{2}ik_y a} \cos\left(\frac{3}{2}k_x a\right) + 2 e^{-\frac{\sqrt{3}}{2}ik_y a} \cos\left(\frac{3}{2}k_x a\right) + e^{-\sqrt{3}ik_y a} \right] \\ &= -t' \left[2 \cos(\sqrt{3}k_x a) + 4 \cos\left(\frac{\sqrt{3}}{2}k_y a\right) \cos\left(\frac{3}{2}k_x a\right) \right] = -t' f(\mathbf{k}) \end{aligned}$$

Substituting them back to the dispersion relation, we can obtain

$$E_{\pm} = \pm t \sqrt{3 + f(\mathbf{k})} - t' f(\mathbf{k})$$

with $f(\mathbf{k})$ defined above.

(b) Take the Dirac point $\mathbf{K} = \frac{2\pi}{3\sqrt{3}a}(\sqrt{3}, 1)$, then calculate $f(\mathbf{K} + \mathbf{q})$ ($|\mathbf{q}| \ll 1$).

$$\begin{aligned} f(\mathbf{K} + \mathbf{q}) &= 2 \cos(\sqrt{3}K_y a + \sqrt{3}q_y a) \\ &\quad + 4 \cos\left(\frac{3}{2}K_x a + \frac{3}{2}q_x a\right) \cos\left(\frac{\sqrt{3}}{2}K_y a + \frac{\sqrt{3}}{2}q_y a\right) = -3 + \frac{9}{4}|\mathbf{q}|^2 a^2, \end{aligned}$$

where the identities for the trigonometric functions and the approximations $\cos(x) \sim 1 - \frac{1}{2}x^2$, $\sin x \sim x$ are used. Substituting $f(\mathbf{K} + \mathbf{q})$ to $E_{\pm}$

$$E_{\pm} = \pm t \sqrt{3 - 3 + \frac{9}{4}|\mathbf{q}|^2 a^2} - t' \left(-3 + \frac{9}{4}|\mathbf{q}|^2 a^2 \right) + \mathcal{O}(|\mathbf{q}|^3) = \pm \frac{3t}{2}|\mathbf{q}|a + 3t' - \frac{9}{4}t'|\mathbf{q}|^2 a^2 + \mathcal{O}(|\mathbf{q}|^3).$$

Problem 9.2. Graphene has the record room-temperature electron mobility, mainly due to two reasons. The first is the high Debye temperature (~ 2100 K) that suppresses phonon scattering arising from the in-plane covalent sp^2 bonding between carbon atoms. The second one is related to its unique linear Dirac band, resulting in massless Dirac fermions ($m^* = 0$) near the Dirac point. However, if we apply the definition of the effective mass $m^* = \hbar^2 \left(\frac{d^2 E(k)}{dk^2} \right)^{-1}$ to the band dispersion of graphene ($E_{\pm}(k) = \pm \hbar v_F k$), we will get diverging m^* and non-differentiable Dirac point. Please explain this contradiction.

Solution. The contradiction arises from applying different definitions of “mass” to graphene’s unique band structure. The description of “massless Dirac fermions” comes from the analogy to relativistic particles.

From a massless special relativity particle like a photon to the electrons in the graphene, their dispersion relations have the comparasion

$$E = pc \implies E = \pm \hbar v_F k = p v_F$$

with v_F playing the role of c . Hence, electrons near the Dirac point behave as if they have zero rest mass.

However, the conventional effective mass $m^* = \hbar^2 (d^2 E / dk^2)^{-1}$ is defined under the parabolic-band approximation $E \propto k^2$. For graphene’s linear dispersion, $d^2 E / dk^2 = 0$ away from $k = 0$, leading to $m^* \rightarrow \infty$ mathematically. At $k = 0$ the dispersion is non-differentiable, so the formula breaks down.

Physically, this infinite effective mass reflects that the electron’s speed is constant at v_F : An external force parallel to its momentum cannot change its speed, much like a photon’s speed is fixed at c . Thus, the “massless” label refers to the relativistic analogy, while the divergent m^* indicates the inapplicability of the parabolic effective mass concept to a linear band.