

HEALTHCARE PROFESSIONAL TRAINING

# PARACETAMOL FRESENIUS

10 mg/ml IV infusion

---

Short-term treatment of mild to moderate pain or fever when oral administration is unsuitable.

# Dose by weight — and count every paracetamol source

Adults >50 kg

**1 g**

One 100 ml vial per dose

Up to 4 times daily

MAX 4 G / 24 H

>33 kg to ≤50 kg

**15** mg/kg

Equivalent to 1.5 ml/kg

Up to 4 times daily

MAX 60 MG/KG; NOT >3 G / 24 H

>10 kg to ≤33 kg

**15** mg/kg

Equivalent to 1.5 ml/kg

Up to 4 times daily

MAX 60 MG/KG; NOT >2 G / 24 H

**15-minute IV infusion for all patients**

Minimum interval is 4 hours; use 6 hours if creatinine clearance ≤30 ml/min. Use non-oedematous body weight and include all concomitant paracetamol-containing medicines.

# Prevent overdose; respond early if it occurs

## BEFORE & DURING USE

### Reduce avoidable risk

- Contraindicated in hypersensitivity and severe hepatocellular insufficiency or active liver disease.
- Use oral paracetamol as soon as the oral route is available.
- Stop at first rash or hypersensitivity sign.

## OVERDOSE RESPONSE

### Treat as a medical emergency

- Early symptoms may be mild; liver injury can be delayed.
- Hospitalise, assay plasma paracetamol, and start N-acetylcysteine promptly.
- Monitor hepatic tests and significant overdose for 96 hours.