

INTERNAL HCP TRAINING

# Paracetamol Fresenius IV

10 mg/ml solution for intravenous  
infusion

---

Short-term treatment of mild to moderate pain and fever when oral administration is unsuitable.

Scheduling status: S3 | Based on approved PI: 26 Nov 2021

## DOSE BY WEIGHT

# Start with non-oedematous body weight

All doses are given by  
15-minute IV infusion.

**Adults >50 kg**

**1 g** / 100 ml

Up to 4 times daily

Maximum  
**4 g in 24 hours**

**>33 kg to ≤50 kg**

**15 mg/kg**

Equivalent to 1.5 ml/kg, up to 4 times daily

Maximum  
**60 mg/kg; not >3 g/day**

**>10 kg to ≤33 kg**

**15 mg/kg**

Equivalent to 1.5 ml/kg, up to 4 times daily

Maximum  
**60 mg/kg; not >2 g/day**

### MINIMUM INTERVAL

**4 hours between doses**

### SEVERE RENAL IMPAIRMENT

**Use at least 6 hours**

### 100 ML VIAL

**Only if patient >33 kg**

# Prevent overdose and infusion errors

## Before prescribing or administering

- Check weight, liver risk, renal function and all paracetamol medicines.
- Do not exceed recommended dose; include every paracetamol source.
- Use oral paracetamol as soon as the oral route is available.

## Prepare, monitor, escalate

- Inspect vial; use immediately after opening and discard unused solution.
- Do not mix with other medicines; monitor glass vial infusions for air embolism.
- Stop at first rash or hypersensitivity sign; overdose needs urgent hospital care and NAC.

**Contraindicated in severe hepatocellular insufficiency or active liver disease, including alcoholic hepatitis.**