

UNIVERSITY OF CALIFORNIA
SANTA CRUZ

**VOICE QUALITY AND LARYNGEAL COMPLEXITY IN
SANTIAGO LAXOPA ZAPOTEC**

A dissertation submitted in partial satisfaction of the
requirements for the degree of

DOCTOR OF PHILOSOPHY

in

LINGUISTICS

by

Mykel Loren Brinkerhoff

May 2025

The Dissertation of Mykel Loren Brinkerhoff
is approved:

Professor Grant McGuire, Chair

Professor Jaye Padgett

Professor Ryan Bennett

Professor Marc Garellek

Peter Biehl
Vice Provost and Dean of Graduate Studies

Copyright © by

Mykel Loren Brinkerhoff

2025

Contents

List of Figures	vi
List of Tables	ix
Abstract	xi
Dedication	xii
Acknowledgments	xiii
1 Introduction	1
1.1 What is Voice Quality	1
1.2 Voice Quality and Tone	2
1.3 Voice Quality and Phonation	2
1.4 Interactions between Voice Quality and Tone	2
1.4.1 Why Esposito & Khan 2020 matters to my research	2
1.4.2 Connecting the two	4
1.5 Esposito & Khan 2020	5
1.5.1 Type I languages	8
1.5.2 Type II languages	9
1.5.3 Type III languages	10
1.5.4 Type IV languages	12
2 Vowels and suprasegmentals in Santiago Laxopa Zapotec	16
2.1 Introduction	16
2.2 Vowels in Santiago Laxopa Zapotec	17
2.3 Phonation in Santiago Laxopa Zapotec	21
2.3.1 Breathy vowels in SLZ	22
2.3.2 Checked vowels in SLZ	23

2.3.3	Rearticulated vowels in SLZ	24
2.3.4	Laryngealization's status in Zapotec languages	27
2.4	Tonal contrasts in Santiago Laxopa Zapotec	29
2.5	Interactions between tone and voice quality	29
3	On using Residual H1* for voice quality research	32
3.1	Introduction	32
3.2	Santiago Laxopa Zapotec	36
3.3	Methods	38
3.3.1	Elicitation	38
3.3.2	Data Processing	39
3.3.3	Statistical modeling	41
3.4	Results	43
3.4.1	H1*–H2*	44
3.4.2	Residual H1*	44
3.4.3	Model Comparison	45
3.4.4	GAMM analysis and model comparisons	47
3.5	Conclusion	52
4	The acoustic space of voice quality in Santiago Laxopa Zapotec	54
4.1	Introduction	54
4.2	Methods	55
4.2.1	Participants	55
4.2.2	Recordings	55
4.2.3	Acoustic measuring	56
4.2.4	Data processing	57
4.2.5	Statistical analysis	59
4.3	Results	62
4.3.1	Acoustic space of voice quality	62
4.3.2	Acoustic correlates of voice quality	69
4.4	Discussion	75
4.5	Conclusion	79
5	Trees reveal the importance of measures in SLZ	81
5.1	Introduction	81
5.2	What are Decision Trees	82
5.3	Decision trees in linguistics	84
5.4	Growing a forest of decision trees	86
5.4.1	Bagging trees	87
5.4.2	Random Forests	88
5.4.3	How to interpret the results	89

5.5	Random Forests in SLZ	93
5.5.1	Methods	94
5.5.2	Parameter selection	98
5.6	Results	99
5.7	Discussion of the results	102
5.7.1	Comparing the MDS and bagging trees	103
5.7.2	Importance of duration	103
5.7.3	Importance of A1*	108
5.7.4	Importance of H1*–A1*	111
5.7.5	Importance of Residual H1*	114
5.7.6	Importance of the HNR <1500 Hz	116
5.7.7	Importance of Strength of Excitation	117
5.8	Conclusion	120
6	Testing laryngeal complexity in SLZ	122
6.1	Introduction	122
6.2	What is Laryngeal Complexity?	124
6.2.1	Phasing and recoverability	125
6.2.2	Implicational hierarchy of laryngealization	133
6.3	Previous analyses of laryngeal complexity	136
6.4	Analysis of laryngeal complexity	139
6.4.1	Methods	139
6.5	Results	145
6.5.1	Fundamental frequency	146
6.5.2	Harmonic-to-noise ratio	149
6.5.3	Strength of excitation	152
6.6	Discussion	155
6.7	Conclusion	159
7	Conclusion	160
7.1	The Laryngeal Articulator Model	160
7.2	Modeling laryngeal complexity	162
7.3	Alternative accounts	162
7.3.1	Articulatory Phonology account	162
7.3.2	Q-theory account	162
7.3.3	Radical CV Phonology account	162
Bibliography		163
A Some Ancillary Stuff		194

List of Figures

2.1	Photo of Santiago Laxopa taken by Beto Diaz, a resident.	17
2.2	Vowel space of Santiago Laxopa Zapotec. The ellipses around each vowel mean represents 1 standard deviation. The scale of the axes are in barks with their corresponding Hz values.	19
2.3	Vowel space for SLZ speakers by gender. The ellipses around each vowel mean represents 1 standard deviation. The scale of the axes are in barks with their corresponding Hz values.	19
2.4	Breathy vowel in the word <i>yah</i> ‘metal; rifle’	23
2.5	Checked vowel in the word <i>yu'</i> ‘earth’	24
2.6	FSR’s rearticulated vowels in <i>za’ɑ</i> ‘corncob’ and <i>xa’ag</i> ‘topil’	26
2.7	RD’s rearticulated vowels in <i>za’ɑ</i> ‘corncob’ and <i>xa’ag</i> ‘topil’	27
3.1	H1*–H2* across the duration of the vowel. Points represent the mean of each measure across the ten intervals. The error bars around each point represent a 95% confidence interval. A line was plotted over each to show how the acoustic measure functions across the ten intervals.	45
3.2	Residual H1* across the duration of the vowel. Points represent the mean of each measure across the ten intervals. The error bars around each point represent a 95% confidence interval. A line was plotted over each to show how the acoustic measure functions across the ten intervals.	46
3.3	GAMM smooths for H1*–H2* across the duration of the vowel.	48
3.4	GAMM difference plots for H1*–H2* across the duration of the vowel.	49
3.5	GAMM smooths and difference plots for residual H1* across the duration of the vowel.	51
3.6	GAMM smooths and difference plots for residual H1* across the duration of the vowel.	52
4.1	Scree plot showing the stress for each dimension for the MDS analysis.	61

4.2	Two-dimensional MDS solution showing the first and second dimensions.	63
4.3	Two-dimensional MDS solution showing the first and third dimensions.	64
4.4	Two-dimensional MDS solution showing the first and fourth dimensions.	65
4.5	Two-dimensional MDS solution showing the second and third dimensions.	66
4.6	Two-dimensional MDS solution showing the second and fourth dimensions.	67
4.7	Two-dimensional MDS solution showing the third and fourth dimensions.	68
4.8	A diagram showing the relationship between breathy, modal, and creaky phonation types from Gordon & Ladefoged (2001).	68
4.9	A diagram showing the relationship between breathy, modal, and creaky phonation types. Based on Gordon & Ladefoged (2001).	77
4.10	A two-dimensional representation of the acoustic space of voice quality in SLZ. The horizontal axis represents the spectral slope of the signal, while the vertical axis represents the amount of noise or energy in the signal.	79
5.1	Classification tree of phonation categories from Keating et al. (2023). Abbreviations used in this figure are: HNR05_means002: harmonics-to-noise ratio over the frequency range from 0 Hz to 500 Hz for the middle third of each vowel; SHR_means002: subharmonic-to-harmonic ratio for the middle third of each vowel; H1H2c_means002: H1* – H2* for the middle third of each vowel; B: breathy, M: modal, and C: creaky phonation categories.	85
5.2	Variable importance plot from Tagliamonte & Baayen (2012). The plot on the left shows the impurity importance (i.e., Gini index) and the plot on the right shows the permutation importance. The y-axis shows the different predictors for both plots.	92
5.3	Plot showing the percent of inaccurately classified phonation types as a function of the number of trees ran. The different colored lines indicate the different m_{try} values.	100
5.4	Variable importance plots showing the impurity importance and permutation importance of each acoustic measure.	101
5.5	Plot showing the distribution of duration across the different voice qualities in SLZ.	106
5.6	Plot showing the distribution of A1* across the different voice qualities in SLZ.	109

5.7	Plot showing the distribution of H1*–A1* across the different voice qualities in SLZ. Each point represents the mean of the ten equally spaced intervals across the duration of the vowel and the error bars represent a 95% confidence interval.	113
5.8	Plot showing the distribution of residual H1* across the different voice qualities in SLZ.	115
5.9	Plot showing the distribution of HNR < 1500 Hz across the different voice qualities in SLZ.	117
5.10	Plot showing the distribution of Strength of Excitation across the different voice qualities in SLZ.	119
6.1	A diagram showing the relationship between breathy, modal, and creaky phonation types. Based on Gordon & Ladefoged (2001).	129
6.2	Schematic representation of the characteristics of [ha˥] sequences. Adaptation of figure from Silverman (1997a).	132
6.3	Schematic representation of the characteristics of [a˥h] sequences. Adaptation of figure from Silverman (1997a).	132
6.4	Schematic representation of the characteristics of [aha˥] sequences. Adaptation of figure from Silverman (1997a).	133
6.5	Model fit for f_0 . Each line represents the predicted values from the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval.	147
6.6	Plot of the difference between modal and each of the non-modal phonation types.	148
6.7	Model fit for HNR. Each line represents the predicted values from the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval.	150
6.8	Plot of the difference between modal and each of the non-modal phonation types.	151
6.9	Model fit for SoE. Each line represents the predicted values from the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval.	153
6.10	Plot of the difference between modal and each of the non-modal phonation types.	154
7.1	The Laryngeal Articulator Model from Esling et al. (2019). This model shows the interactions between the laryngeal articulators (labeled circles). Synergetic interactions are shown with solid lines, while anti-synergetic interactions are shown with dotted lines.	161

List of Tables

1.1	Language types based on the lexically contrastive nature of <i>f0</i> /tone/pitch accents (rows) and voice quality on vowels (columns) and their interactions (Esposito & Khan 2020)	2
1.2	Language types based on the lexically contrastive nature of <i>f0</i> /tone/pitch accents (rows) and voice quality on vowels (columns) and their interactions (Esposito & Khan 2020)	8
1.3	Updated language types based on the lexically contrastive nature of <i>f0</i> /tone/pitch accents (rows) and voice quality on vowels (columns) and their interactions	15
2.1	Vowel qualities in Santiago Laxopa Zapotec.	18
2.2	Pillai scores and Bhattacharyya's Affinity for /o/ and /u/ in SLZ.	20
2.3	Layngealized Vowels in Yalálag Zapotec	25
2.4	SLQZ tone and phonation interactions (Chávez-Péón 2010).	30
2.5	SLZ tone and voice quality combinations.	31
3.1	Distribution of tone and voice quality in the wordlist	39
3.2	Model comparison between H1*–H2* and Residual H1* in distinguishing Santiago Laxopa Zapotec voice quality.	47
3.3	AIC for the H1*–H2* and residual H1* models.	47
4.1	Correlations for each acoustic measure to the four dimensions (NMDS1, NMDS2, NMDS3, NMDS4). The four largest correlations in each dimension are bolded.	70
5.1	Correlations for each acoustic measure to the four dimensions (NMDS1, NMDS2, NMDS3, NMDS4). The four largest correlations in each dimension are bolded.	104
5.2	Possible phonetic correlates of register. From Brunelle & Kirby (2016).	110

6.1	Implicational hierarchy of laryngeal complexity. The symbols h and ? represent laryngealization. The symbol V represents where the modal vowel is located in relation to the laryngealization. Modified from Silverman (1997a).	134
6.2	Distribution of the number of syllables containing the combination of tone and voice quality in the wordlist.	143
6.3	Implicational hierarchy of laryngeal complexity. The symbols h and ? represent laryngealization. The symbol V represents where the modal vowel is located in relation to the laryngealization. Modified from Silverman (1997a).	157

Abstract

Voice Quality and Laryngeal Complexity in

Santiago Laxopa Zapotec

by

Mykel Loren Brinkerhoff

This dissertation provides a detailed description and analysis of the SLZ voice quality system, a minority language spoken by about 1000 people in the municipality of Santiago Laxopa, and its interactions with the tonal system of the language. Standard assumptions about the interaction between tone and voice quality in Otomanguean languages proposed by Silverman (1997a,b), where nonmodal phonation is realized in only a portion of the vowel and tone is realized on a modal portion, do not fully hold in Santiago Laxopa Zapotec. Instead, speakers routinely produce nonmodal phonation throughout the entire vowel for breathy vowels. The only time phasing is observed is with the two types of creaky voice that occur in the language: rearticulated and checked. Rearticulated vowels have a period of creakiness in the middle of the vowel, whereas checked vowels have creakiness at the end. Although this creakiness is pronounced in distinct locations, non-modal phonation remains throughout the entire vowel. These results were confirmed through statistical modeling.

...

Dedicated to my family,

Betsy and Maelyn,

I wouldn't be here without you.

Acknowledgments

Acknowledgements in dissertations and theses are always so awkward and sometimes difficult to write. This is because there are no words that can adequately express the gratitude that you feel towards those who have helped you along the way. But I will try my best.

I would like to thank my advisor, Professor Grant McGuire, for his guidance and support throughout my time at the graduate program at the University of California, Santa Cruz. Thanks to him, I have learned and grown as a researcher and as a person. Thanks to his guidance, encouragement, and support, I have been able to complete this dissertation.

This dissertation would also not have been possible without the help of my committee members, Professors Jaye Padgett, Ryan Bennett, and Marc Garellek. Their feedback and advice have been invaluable in shaping my research and my writing. Thank you for your time and effort in reading my dissertation and for your thoughtful comments. I am grateful for your support and encouragement.

Besides my committee members, I would like to also thank the first linguistic Banana slugs I had the pleasure of knowing. Abby and Aaron Kaplan, both alumni from the linguistic program here at UC Santa Cruz, introduced me to the world of phonetics and phonology while I was an undergraduate student at the University of Utah and by their own love of the field, they inspired me to pursue a career in linguistics. I am

grateful for their support while an undergraduate and for helping me take the first steps in my career as a linguist.

Finally, I would like to thank my wife, Betsy, and my daughter, Maelyn, for their patience and understanding. I could not have done this without them. Without their love and support, I would not have been able to complete this dissertation.

Chapter 1

Introduction

1.1 What is Voice Quality

Voice quality describes the state of the larynx during phonation, when the vocal folds are set in motion. Languages make use of voice quality for paralinguistic purposes, such as conveying indexation of “biological, psychological, and social characteristics of the speaker” (Laver 1968) and racial identity (Podesva 2016).

Voice quality is also used linguistically. In English, it is often the case that we use creaky voice to indicate that we are at the end of an utterance (e.g., Garellek 2013). In many other languages, voice quality is used as part of the phonological system. Most famously, Gujarati has a phonemic contrast between breathy and modal voice in vowels (e.g., Fischer-Jørgensen 1968, Esposito & Khan 2012, Khan 2012, Esposito et al.

2019).

Esposito & Khan (2020)

1.2 Voice Quality and Tone

1.3 Voice Quality and Phonation

1.4 Interactions between Voice Quality and Tone

1.4.1 Why Esposito & Khan 2020 matters to my research

- Esposito & Khan (2020) is primarily important for my research because of their typology of the interactions between tone and phonation, which are summarized in Table 1.1.

Table 1.1: Language types based on the lexically contrastive nature of f_0 /tone/pitch accents (rows) and voice quality on vowels (columns) and their interactions (Esposito & Khan 2020)

	No VQ contrast		VQ contrast	
No tone contrast	No contrasts (Type I)		f_0 not a cue (Type IIa)	f_0 is a cue (Type IIb)
Tone contrast	VQ not a cue (Type IIIa)	VQ is a cue (Type IIIb)	Orthogonal (Type IVa)	Fused (Type IVb)

- Another aspect of my research into the tone and phonation interactions is why we see gaps/restrictions for which tones and phonations are allowed to combine.
 - This is important because of the gaps I observe in SLZ where some we never see breathy with high tone and we do not see checked vowels with rising tone.
 - * This is true for nominals.
 - * I have not looked at verbs and whether or not we see this same gap.
 - I expect that the gap is also present in the verbal paradigms similar to what Uchihara (2016) observed where breathy phonation fails to appear in some parts of the paradigm.
 - Especially with the Potential Aspect, which is always realized as a high tone on the verbal root.
- These four types of languages offer an excellent way for me to characterize what we see cross-linguistically in the interactions between tone and phonation.
- This primarily is useful for me as a way to zero in on which languages I need to look at.
 - This means that I need to focus my search into types IIb, IIIb, and IV languages.

1.4.2 Connecting the two

- The connection here between Silverman (1997a) and the typology is that these both have to deal with the interactions between tone and phonation.
- Silverman offers an account for what we expect to see when a language has both tone and phonation.
 - It additionally offers three motivating factors for interactions between tone and phonation.
 - These three factors also offer ways to account for the patterns that we observe and don't observe in the typology
- The typology allows us to see more generally what combinations we have and don't have.
- If there are any cross-linguistic gaps that we observe, then do Silverman's (1997a) three factors of (i) sufficient acoustic distance, (ii) sufficient articulatory compatibility, and (iii) optimal auditory salience provide the answers to how and why.

1.5 Esposito & Khan 2020

- Esposito & Khan (2020) is a paper that explains the various cross-linguistic patterns that exist in the world's languages for phonation.
- They describe that for the majority of the world's languages there is only modal voice.
- However, a subset of the languages have non-modal phonation.
- They state that languages can either associate phonation, which they define as production of sound by the vocal folds, with consonants or vowels.
 - For consonants this takes the form of breathy or creaky voice (4ff).
- I am going to ignore their discussion of phonation associated with consonants for the most part and focus on what they have to say about vowels.
 - According to them only five languages contrast phonation on both vowels and consonants.
 - These are !Xóõ, Ju'hoansi, Wa, White Hmong, and Gujarati
- They describe that there are two main avenues for researching phonation's production.
 - This is done through acoustic measurements and electroglottography

- The most common way that this is done is with spectral silt measures which reflect the open quotient.
 - The open quotient is the proportion of the glottal cycle during which the glottis is open (Holmberg et al. 1995).
- Other measures that reflect the periodicity of the signal are also consulted such as HNR and CPP. Typically nonmodal phonation is also aperiodic and will have a lower score compared to the modal.
- They also state that “H1-H2 may be a (near-)universal acoustic measure of phonation” (8).
- They further say that this measure works the best the most often but there are several exceptions where other H1-X measures are more robust at capturing the phonation contrasts in the language.
 - As discussed in Chai & Garellek (2022), what they are actually trying to do is get at the amplitude differences in H1 as first observed by Fischer-Jørgensen (1968) and Laver (1968).
 - I think this is where, I and other researchers fell into a fallacy where we think that just the spectral-tilt is what matters instead of realizing what those measures were trying to accomplish which was to normalize H1 for comparison.

- Esposito & Khan (2020) further discuss the role that EGG play.
- Two things that I think are relevant for my dissertation is the discussion around localization of non-modal phonation and how phonation relates to tone.
- For the discussion around localization, they keep in line with Silverman (1997a) who says that phonation is located in certain portions of the vowel.
- Localization of non-modal phonation is also important in languages with complex “clusters” (i.e., phonetic sequences) of phonation type, which involve at least one non-modal phonation localized to a portion of the vowel.
 - !Xóõ distinguishes clusters of breathy-creaky, pharyngealized-creaky, and breathy-pharyngealized phonation in addition to modal, breathy, creaky, and pharyngealized voice
- In terms of tone and phonation interactions, Esposito & Khan (2020) claim that there are four different types of interactions based on whether or not tone and phonation is contrastive in the language.
- Table 1.2 shows each of the different interactions between tone and phonation. In this table each row represents whether or not tone/ f_0 /pitch accent is contrastive and each column represents whether or not phonation is contrastive.
- Each cell represent each of the four different types of languages (I-IV).

- Types II-IV are further subdivided based on whether or not tone and phonation are cues for each other.

Table 1.2: Language types based on the lexically contrastive nature of f_0 /tone/pitch accents (rows) and voice quality on vowels (columns) and their interactions (Esposito & Khan 2020)

	No VQ contrast	VQ contrast	
No tone contrast	No contrasts (Type I)	f_0 not a cue (Type IIa)	f_0 is a cue (Type IIb)
Tone contrast	VQ not a cue (Type IIIa)	VQ is a cue (Type IIIb)	Orthogonal (Type IVa)
			Fused (Type IVb)

1.5.1 Type I languages

- Type I languages are those languages that lack tonal contrasts and phonation contrasts.
- According to Esposito & Khan (2020) these are the languages that belong to this type:
 - most Australian Aboriginal languages,
 - most Austronesian languages,
 - most Indo-European languages
 - Afro-Asiatic languages,

- Standard Khmer,
- Turkic languages
- They do not go into much detail about these languages

1.5.2 Type II languages

- Type II languages are languages that lack tonal contrasts but do have phonation contrasts on the vowel.
- This is further subdivided into two subtypes (IIa and IIb) based on whether or not changes in f_0 are a cue to the phonation.
- Type IIa languages are those where f_0 is not a cue.
- According to Esposito & Khan (2020), these languages are quite rare and only two languages are known to exhibit this behavior:
 - Danish (Grønnum, Vazquez-Larruscaín & Basbøll 2013)
 - Gujarati (Khan 2012)
- Type IIb languages are those where f_0 is a cue for the different phonation types.
- These languages are sometimes called “register languages”
- These are languages such as:

- Chanthaburi Khmer
 - Chong
 - Javanese
 - Kedang
 - Mon
 - Suai
 - Wa
- In type IIb languages the way in which f_0 is a cue is language specific.
- They describe that breathy vowels in Wa and Kedang both begin with a lower f_0 than their clear/modal counterparts
- However, breathy vowels in Chanthaburi Khmer have a higher f_0 than their clear counterparts.

1.5.3 Type III languages

- Type III languages are characterized by languages that have a tonal contrast but lack a phonation contrast.
- These languages are further subdivided into two subtypes based on whether or not phonation is a cue for the tonal contrasts (Type IIIa vs. Type IIIb)

- In Type IIIa languages, tone is contrastive and phonation is not a cue for those tonal contrasts.
- These are languages like:
 - Japanese,
 - Navajo
 - Punjabi
 - Manange
 - Most W. African languages
 - Swedish
 - Central Thai
- Type IIIb languages are those that have lexical tones and no lexical vowel phonation. However, subsets of the tone categories have been reported to have optional phonation which helps cue the tonal contrasts.
- For example, tone 3 of Mandarin Chinese which is usually accompanied with creak (Kuang 2017).
- These are languages such as:
 - Cantonese Yue,
 - Khmu' Rawk

- Mandarin
- Pakphanang Thai
- Ph.Penh Khmer
- Yueyang Xiang

1.5.4 Type IV languages

- Type IV languages are those languages that have both a contrast in tone and phonation and is also subdivided into two subtypes (IVa and IVb).
- Type IVa are those languages where tone and phonation are completely orthogonal to each other. This means that all tone contrasts can appear with any of the phonation contrasts.
- Type IVa languages include:
 - Dinka,
 - Mazatec language,
 - Mpi
 - Yalálag Zapotec
 - Yi languages such as Bo

- Type IVb languages are often called “register languages” because tone and phonation are so closely tied to one another that it is impossible to state whether or not they have tonal contrasts or phonation contrasts.
- In these languages suprasegmental categories (i.e., tone) are consistently produced with both a specific f_0 as well as a specific phonation.
- Examples for this include the behavior where specific tones only arise with specific phonation types. This is most commonly found among languages in Mainland Southeast Asia.
- Esposito & Khan (2020) additionally describe that most Zapotec languages also fall into this type of language.
 - I do not agree with the strong version of this statement. The evidence that they cite for this comes from Esposito (2010) where she describes that SAV Zapotec syllables fall into one of four categories: breathy phonation with falling tone, creaky phonation with low falling tone, modal phonation with a high tone, and modal phonation with rising tone
 - It is true that some phonation types are closely tied to one tonal pattern in Zapotec languages.
 - For example, in Isthmus Zapotec (Pickett, Villalobos & Marlett 2010) is described as having five tonal melodies (H, L, LH, HL, and LHL) and three

phonation types (modal, checked, and laryngealized).

- In monosyllabic nouns L and LH can surface with any of the three phonation types but the combinations of H and HL with any of the phonation types do not appear to be present in the lexicon
- When we consider disyllabic nouns we see the following combinations:
 - * L appears with any of the phonation types.
 - * H appears with modal and laryngealized
 - * LH appears with modal and checked
 - * HL appears with modal and checked
 - * LHL only appears with modal.
- This is also true in SLZ where the only combination of tone and phonation that we fail to see is H with breathy phonation and MH with checked.
- This leads me to believe that maybe it would be better to think of Zapotec languages as a third subtype of languages where some of the tone and phonation contrasts are orthogonal and other tone and phonation contrasts are fused.
- In other words a hybrid between Type IVa and Type IVb which I would call mixed languages or Type IVc.
- This is illustrated in a new updated table in Table 1.3.

Table 1.3: Updated language types based on the lexically contrastive nature of f_0 /tone/pitch accents (rows) and voice quality on vowels (columns) and their interactions

	No VQ contrast	VQ contrast	
No tone contrast	No contrasts (Type I)	f_0 not a cue (Type IIa)	f_0 is a cue (Type IIb)
Tone contrast	VQ not a cue (Type IIIa)	VQ is a cue (Type IIIb)	Orthogonal (Type IVa)
		Fused (Type IVb)	Mixed (Type IVc)

Chapter 2

Vowels and suprasegmentals in Santiago Laxopa Zapotec

2.1 Introduction

Santiago Laxopa Zapotec (SLZ; *Dille'xhunh Laxup* [diɬe'ɬunh l:aɬsup^h]) is a Northern Zapotec language spoken by approximately 1000 people in the municipality of Santiago Laxopa, Ixtlán, Oaxaca, Mexico and in diaspora communities throughout Mexico and the United States (Adler & Morimoto 2016, Adler et al. 2018, Foley, Kalivoda & Toosarvandani 2018, Foley & Toosarvandani 2022). According to Smith-Stark (2007), SLZ is part of the macro variety of Cajonos Zapotec, which also includes Zoogocho Zapotec, Yatzachi Zapotec, Yalálag Zapotec, Tabaá Zapotec, Lachirioag Za-

potec, and several other varieties spoken in the Sierra Norte of Oaxaca, Mexico.



Figure 2.1: Photo of Santiago Laxopa taken by Beto Diaz, a resident.

2.2 Vowels in Santiago Laxopa Zapotec

SLZ exhibits a four-vowel inventory; see Table 2.1. This type of vowel inventory is very common among Sierra Norte Zapotecs. Most varieties have the vowels /i/, /e/, /a/, and /o/ (Nellis & Hollenbach 1980, Jaeger & Van Valin 1982, Butler H. 1997, Avelino 2004, Long & Cruz 2005, Sonnenschein 2005).

The vowel /o/ is marginal in SLZ's lexicon, only appearing in a few lexical items

Table 2.1: Vowel qualities in Santiago Laxopa Zapotec.

	front	central	back
high	i		u~o
mid	e		
low		a	

such as the diminutive classifier *do'*. Instead, this vowel is replaced by /u/ in most cases. However, this difference is not universal among all speakers in the community. For the most part older speakers exhibit the vowel /o/ in their speech, while younger speakers tend to replace it with /u/. Most speakers, when asked, classify the two back rounded vowels as the same phoneme and view them as a dialectal feature between the different pueblos. For example, in neighboring San Bartolomé Zoogocho the /u/ vowel is very marginal and has led Sonnenschein (2005) to describe the language as having only four vowels. It is interesting to note that everywhere that SLZ has the vowels /u/ or /o/, Zoogocho only has /o/. Further evidence for this comes from plotting the vowels along the first two formants. As shown in Figure 2.2, the vowels /o/ and /u/ occupy nearly identical vowel spaces.

Additional evidence for the overlap of /o/ and /u/ can be measured with a combination of Pillai scores (Pillai 1955, Hay, Warren & Drager 2006, Nycz & Hall-Lew 2014) and Bhattacharyya's Affinity (Bhattacharyya 1943, Johnson 2015, Warren 2018, Strelluf 2018). Both of these measures show what degree of overlap exists between




Figure 2.2: Vowel space of Santiago Laxopa Zapotec. The ellipses around each vowel mean represents 1 standard deviation. The scale of the axes are in barks with their corresponding Hz values.




Figure 2.3: Vowel space for SLZ speakers by gender. The ellipses around each vowel mean represents 1 standard deviation. The scale of the axes are in barks with their corresponding Hz values.

two different items in some space. Their use in linguistics has been used mainly to show the process of complete and partial mergers between vowels, such as the NEAR-SQUARE vowel merger in New Zealand English (Hay, Warren & Drager 2006). The Pillai scores and Bhattacharyya's Affinity show that the vowels /o/ and /u/ are nearly identical in their vowel space; see Table 2.2.

Table 2.2: Pillai scores and Bhattacharyya's Affinity for /o/ and /u/ in SLZ.

	Pillai score	Bhattacharyya's Affinity
All speakers	0.157	0.892
Females	0.138	0.890
Males	0.224	0.858

In interpreting these results, Pillai scores range from 0 to 1, with 0 indicating overlap and 1 indicating complete no overlap. Bhattacharyya's Affinity ranges from 0 to 1, with 0 indicating no overlap and 1 indicating complete overlap. The results show that the overlap between /o/ and /u/ is not complete, but it is also not completely separating. This is consistent with the observations made by myself and other researchers for this variety (Toosarvandani, p.c.).

In summary, we can conclude that SLZ is similar to other Northern Zapotec varieties in having a four-vowel inventory. The vowel /o/ is marginal in the lexicon and is often replaced by /u/ in younger speakers. The vowels /o/ and /u/ occupy nearly identical vowel spaces, and the overlap between the two vowels is not complete but

is also not completely separating.

2.3 Phonation in Santiago Laxopa Zapotec

One of the defining characteristics of the Oto-Manguean languages is their use of contrastive phonation (Campbell 2017a,b). This fact is so common and widespread that it has been reconstructed for the proto-language (see Rensch 1976). Additionally, the use of contrastive phonation in Oto-Manguean languages led Silverman (1997a,b) to create the term *laryngeal complexity* to describe how phonation is realized in these languages and the development of theories of how phonation interacts with tone.

This use of contrastive phonation is prevalent in the Zapotec languages (see Ariza-García 2018 for an overview and typology of the phonation contrasts in the Zapotec language family), with SLZ being no exception. SLZ has a four-way voice quality contrast: modal, breathy, and two types of laryngealization which are checked and rearticulated. These contrasts are exemplified in the near-minimal quadruple in (1).

(1) Four-way near minimal phonation contrast

- a. *yag* [çax-ɬ] ‘tree; wood; almúd (unit of measurement ~4kg)’
- b. *yah* [çə-ɬ] ‘metal; rifle; bell’
- c. *cha'* [tʃa'ɬ] ‘cooking pot’
- d. *ya'a* [ça'a-ɬ] ‘market’

In representing the checked and rearticulated vowels, I follow Avelino (2010) and Uchihara (2016) in representing laryngealization as a superscript glottal stop in the IPA transcriptions (i.e., [a[’]] or [a[’]a]). This allows for a standardized way to discuss these phonations without referencing their variable pronunciation, see Sections 2.3.2 and 2.3.3 for a more detailed discussion of checked and rearticulated phonations respectively.

2.3.1 Breathy vowels in SLZ

SLZ is also unique in regards to its voice quality contrasts because it is a Northern Core Zapotec that has developed breathy voice, which has not been described in any of the neighboring Sierra Norte varieties (Nellis & Hollenbach 1980, Jaeger & Van Valin 1982, Butler H. 1997, Avelino 2004, Sonnenschein 2005, Long & Cruz 2005). However, breathy voice does appear in other Zapotec languages most frequently in Central Valley Zapotecs (Munro & Lopez 1999, Esposito 2004, 2010, Uchihara 2016, Ariza-García 2018). Breathy voice is characterized by aperiodicity in the harmonic structure of throughout the vowel or a portion thereof depending on the speaker.

An example of breathy voice is shown in Figure 2.4, which is the spectrogram and waveform for the word *yah* [çaɬ] ‘metal; rifle; bell’ produced by a female speaker in their 50s. In this figure, we see that the breathy voice on the vowel shows a large amount of noise as evidenced by the aperiodicity in the waveform and the haziness

seen in the spectrogram.




Figure 2.4: Breathy vowel in the word *yah* ‘metal; rifle’

As will be discussed in Chapters 5 and 6, breathy voice is also characterized as having a longer duration than modal voice and having a lower fundamental frequency (f_0) than the other phonation types with the same tone. This behavior is consistent with the behavior of breathy voice in other languages (e.g., Blankenship 1997, Brunelle 2012, Brunelle & Kirby 2016, Hillenbrand & Houde 1996).

2.3.2 Checked vowels in SLZ

SLZ shares with other Zapotec variates two types of laryngealization: checked and rearticulated. These two types of laryngealization are characterized by the presence

of a glottal occlusion or some form of creaky voice that is produced in a portion of the vowel. Checked vowels are typically produced with a glottal stop at the end of the vowel (i.e., /aʔ/) or a period of creakiness at the end of the vowel (i.e., /aɑʔ/).

As seen in Figure 2.5, which was produced by a male speaker in his mid-twenties, the checked vowel in the word *yu'* [çu'ɿ] ‘earth’ is characterized by a rapid decrease in amplitude and aperiodicity at the end of the vowel.




Figure 2.5: Checked vowel in the word *yu'* ‘earth’

2.3.3 Rearticulated vowels in SLZ

Rearticulated vowels are the term used by Oto-Mangueanists to describe the class of laryngealized vowels which have a glottal occlusion or some sort of glottalization in

the middle of the vowel (Ariza-García 2018) and are distinct from other uses of the term such as rearticulated vowels in Mayan languages (Baird 2011). Linguists have also used other terms such as interrupted, laryngealized, broken, creaky, et cetera (Adler & Morimoto 2016, Ariza-García 2018, Avelino 2010, Long & Cruz 2005, Sonnenschein 2005). I will be using the term rearticulated to refer to these vowels throughout the dissertation.

These vowels are also highly variable in their production, hence the large number of terms used to refer to these vowels. For some speakers, they are produced with a full glottal stop in the middle of the vowel (i.e., /aʔa/), while others produce a vowel with apparent modal voice but with a drop in amplitude (similar to what Gerfen & Baker 2005 found for some Mixtec varieties), while others produce a creaky voice throughout the entire vowel. Some speakers produce a combination of these unique productions. This is also consistent with the findings of Avelino (2010) who found that the rearticulated vowels in Yalálag Zapotec had at least four different pronunciations among his consultants, as shown in Table 2.3.

Table 2.3: Layngealized Vowels in Yalálag Zapotec

/V'V/	[V?V]
	[VVV]
	[VV:V̄]
	[VVV̄]

Figures 2.6 and 2.7 show how two different speakers produce these vowels. Fig-

ure 2.6 shows how a female speaker in her 50s produces this vowel. Figure 2.6a shows the rearticulated vowel in the word *za'a* [za'a-ɿ] ‘corncob’. In this spectrogram and waveform FSR produces a modal vowel that is clearly divided by a glottal stop. This is the typical pronunciation used by this speaker. However, when the rearticulated vowel cooccurs with a low tone, like in the word *xa'ag* [ʂa'ax-ɿ] ‘topil’, FSR produces the vowel with creaky voice, as seen in the spectrogram in Figure 2.6b.




Figure 2.6: FSR’s rearticulated vowels in *za'a* ‘corncob’ and *xa'ag* ‘topil’

However, RD, a male in their mid-20s, produces the rearticulated vowels consistently with creaky voice. This is shown in Figures 2.7a and 2.7b which are the same words as found in Figure 2.6.




Figure 2.7: RD's rearticulated vowels in *za'a* 'corncob' and *xa'ag* 'topil'

2.3.4 Laryngealization's status in Zapotec languages

The status of laryngealization in Zapotec languages has been a topic of much debate. Some, such as Avelino (2010) and Esposito (2004) have argued that are not actually phonation types but involve a glottal stop consonant, either as a syllabic onset or coda. It is indeed logical that this could be the case. However, much work has shown that this cannot be true in Zapotec for the laryngealized vowels, which have a glottal closure in their center. Chávez-Péón (2010) summarizes this work and offers six reasons why these “glottal stops” are a vocalic feature, not a consonant in laryngealized vowels. The first reason has to do with the distribution of the glottal stop. If it is indeed a coda or onset, then we would expect it could also occur word-initially or in consonant clusters. This does not happen in SLZ or other Zapotec languages (Jaeger & Van Valin 1982). Another point that he raises is that when linguists ask na-

tive language consultants about how many syllables or beats the word has, they treat laryngealized vowels as if they only had a single beat.¹ These and other points raised by Chávez-Péón are presented in (2).

(2) Summary: glottal stop as a vocalic feature in laryngealized vowels (adapted from Chávez-Péón 2010).

- a. /ʔ/ defective distribution (not in onset, not in clusters)
- b. Laryngealized vowels have the same tonal sequences as single vowels
- c. Monosyllabic tendency of the majority of languages (roots = 1σ)
- d. *VʔC_{fortis}, predicted by bimoraicity of laryngealized vowels
- e. Same vowel quality, i.e., one vowel gesture (diphthongs a minority)
- f. Perceived as single syllables by native speakers (? ≠ sufficient consonantal barrier, i.e., syllable boundary)

One point that he does not mention is that if we assume the glottal stop is a coda consonant, then we would also expect to see the other phonation types being able to cooccur with this coda consonant. Much work still needs to be done from a phonological perspective. Treating /ʔ/ and /h/ as a vocalic feature or as a consonant is worth further study, but in the present work I assume that these sounds are vocal characteristics and contribute to the phonation contrasts following the traditional interpretation

¹This is something Maya Wax Cavallaro and I tested with one of our consultants. The only times they said one of these laryngealized vowels was not a single beat was when they had different vowel qualities on either side of the glottal closure (e.g., *bi'a* [bi'a] 'fly (insect)').

of these sounds.

2.4 Tonal contrasts in Santiago Laxopa Zapotec

One of the most well known features of all Oto-Manguean languages is the fact that they are tonal languages and exhibit a large range of tonal systems (Pike 1948, Rensch 1976, Josserand 1983, Silverman 1997a, Beam de Azcona 2007, DiCanio 2010, 2012a, Elliott, Edmondson & Cruz 2016, Campbell 2017a,b, Lillehaugen 2019, Eischens 2022). SLZ has a five-way tonal contrast which consists of three level tones (high, mid, low) and two contour tones (rising and falling).

Brinkerhoff, Duff & Wax Cavallaro (2022)

2.5 Interactions between tone and voice quality

Most previous work on the interaction of tone and phonation has been focused on the languages of East and Southeast Asia (e.g., Masica 1976, Thurgood 2002, Yip 2002, Enfield 2005, Michaud 2012, Brunelle & Kirby 2016). What has been found in these descriptions is that certain tones and phonations are codependent. For example, Smalley (1976) and Ratliff (1992) both describe White Hmong's -g tone as being a mid-low tone with breathy phonation, and Mandarin's tone 3 is often associated with creaky phonation (Hockett 1947). Brunelle (2009) found that creaky phonation plays

an important role in producing certain tones. Additionally, work on S'gaw Karen has found that two tones are only differentiated by some form of non-modal phonation (Boehm p.c.).

However, there have been some observations—especially in Mesoamerica—that tone and phonation are independent of each other (e.g., Silverman 1997a, Garellek & Keating 2011). This means that tone can independently occur with any phonation type. This has also been extensively described in multiple Zapotecan languages (e.g., Avelino 2004, 2010, Chávez-Péón 2010, Campbell 2011, Villard 2015, López Nicolás 2016)

Chávez-Péón (2010) describes the tone and phonation interactions in San Lucas Quiaviní Zapotec (SLQZ), a central valley variety of Zapotec. The distribution of tone and phonation is found in Table 2.4. We see that in SLQZ, both low- and falling-tones have the full range of possible combinations. However, we see gaps in the high-tone for breathy and rising tones that can only occur with modal phonation.

Table 2.4: SLQZ tone and phonation interactions (Chávez-Péón 2010).

	Modal	Breathy	Creaky	Interrupted
High	✓	–	✓	✓
Low	✓	✓	✓	✓
Falling	✓	✓	✓	✓
Rising	✓	–	–	–

Based on elicitation data collected from 2020-2022, SLZ has a more expansive distribution of tone and phonation when compared to SLQZ but seems to be very similar

to other Northern Zapotec varieties (e.g., Avelino 2004). The distribution of SLZ tonal and phonation combinations are given in Table 2.5.

Table 2.5: SLZ tone and voice quality combinations.

	Modal	Breathy	Checked	Rearticulated
High	✓	–	✓	✓
Mid	✓	✓	✓	✓
Low	✓	✓	✓	✓
High-Low	✓	✓	✓	✓
Mid-High	✓	✓	–	✓

One of the striking things in this is the lack of high tone with breathy phonation. This gap is interesting because of the long-time association of high pitch with breathiness (a good overview—of this association and other phonation types—is found in Esling et al. 2019). This gap is common across the Zapotecan languages that have breathy voice (Campbell p.c.). Regarding breathy phonation in SLQZ, one of the Valley Zapotec varieties, Uchihara (2016) offers some convincing evidence that the phonation originated in syllables with low tone and then spread to other tones via analogy. Investigating the origin of breathy voice in SLZ would be important in understanding how breathy vowels originated in the Zapotecan family where breathy voice is rare typologically (Ariza-García 2018). Such a study is beyond the scope of this paper.

Chapter 3

On using Residual H1* for voice quality research

3.1 Introduction

Languages use voice quality distinctions to convey phonemic distinctions (Garellek 2019) and to convey paralinguistic information by “indexing the biological, psychological, and social characteristics of the speaker” (Laver 1968, Podesva 2016). Voice quality contrasts have been studied extensively by examining their correlates in the acoustic signal (e.g., Esposito & Khan 2020), resulting in a large and complex literature on acoustic correlates of phonation differences (see Garellek 2019).

⁰The majority of the chapter was previously published as Brinkerhoff & McGuire (2025). This chapter adds to Brinkerhoff & McGuire by including discussion of a generalized additive mixed model (Hastie & Tibshirani 1986, Wood 2017) analysis of H1*–H2* and residual H1*.

One measure that Fischer-Jørgensen established, the fundamental's relative strengthive amplitudes of the first harmonic and second harmonic. As established by Fischer-Jørgensen, the relative strength of the fundamental is a correlated measure of breathy voice in contrast with modal voice Fischer-Jørgensen (1968). In order to normalize the amplitude of the fundamental and counteract some of the effects of high-pass filtering and differences in sound pressure in the signal, she proposed that you could subtract the amplitude of a higher harmonic, in this case, the second harmonic (H2), from the amplitude of the fundamental (H1). Since its introduction, $H1^* - H2^*$ has been used in many studies to measure not only breathy voice but other voice quality contrasts as well (Garellek 2019, Chai & Garellek 2022).

Despite the large amount of evidence in support of $H1^* - H2^*$, it is not without its problems. At the fundamental level, it is not clear that $H1^* - H2^*$ adequately measures the strength of the fundamental. Sundberg (2022) found that H1 and H2 are affected differently by subglottal pressure, compromising some of the original reasoning behind the use of $H1^* - H2^*$ from Fischer-Jørgensen (1968). Similarly, in a comprehensive overview of the main concerns with $H1^* - H2^*$ as a phonation type measure, Chai & Garellek (2022) found that in addition to the issues mentioned above, errors in measuring $H1^* - H2^*$ are uncomfortably high. This is mainly due to the need to precisely measure two different harmonic amplitudes; when there are errors in calculating H1, this, in turn, leads to errors in calculating H2 (Arras 1998). An example of this type

of error propagation is that errors in measuring the fundamental frequency, which is especially common with non-modal phonation, are introduced into measuring harmonics because they are based on the fundamental. Despite algorithms correcting for vowel height, a common error that occurs is when a high fundamental frequency co-occurs with a low first formant (Chai & Garellek 2022). This situation causes errors in tracking the fundamental frequency and the first formant. A final issue that can occur when measuring the harmonics is in contexts where the vowel is nasalized. Simpson (2012) shows that in these nasalized contexts, the first nasal pole (P_0) can increase the amplitude of H_2 and, when the fundamental frequency is high, H_1 increases instead.

This collection of errors leads Chai & Garellek (2022) to propose a new measure, residual H_1^* . This measure is calculated by first regressing H_1 on energy and then subtracting the product of energy and the energy factor from H_1 . Chai and Garellek argue that this measure better reflects the initial purpose of using $H_1^*-H_2^*$. Furthermore, they find that residual H_1^* : (i) provides better differentiation between phonation types in !Xóõ; (ii) was more robust for measuring creak in Mandarin with respect to different utterance positions; and (iii) has a stronger relationship to the open quotient than $H_1^*H_2^*$ based on a comparison using electroglottogram data.

The contributions Chai & Garellek (2022) make are very intriguing and have the potential to alter how spectral analyses of voice quality are performed. However, they are not convincing on their own. If residual H_1^* is to be widely adopted in linguistics,

speech pathology, and other speech sciences, then it requires considerable evidence of its effectiveness in acoustic studies. This paper offers additional evidence for residual H1*'s effectiveness.

Given the promising nature of this measure, we tested residual H1* with data from Santiago Laxopa Zapotec. Although Chai & Garellek (2022) evaluated residual H1* for !Xóõ and Mandarin, both of which contain tone and voice quality, they do not factor tone into their analysis.¹ This is where testing the effectiveness of residual H1* in Zapotec languages is beneficial. Zapotec languages are often described as laryngeally complex (Silverman 1997a, Ariza-García 2018). Laryngeal complexity is defined as a language that allows contrastive tone and contrastive voice quality that are unrestricted in their interactions. This laryngeal complexity between tone and phonation types presents a unique challenge for acoustic analysis and testing of the residual H1* measure.

Other Zapotec languages have also been studied for voice quality research. For example, Esposito (2010) found that in Santa Ana del Valle Zapotec there was a biological sex difference in which acoustic measures best capture the voice quality contrasts similar to observations from Klatt & Klatt (1990). Arellanes Arellanes (2010) found that the realization of laryngealization in a variety of San Pablo Güilá Zapotec is highly variable and depends on the position of laryngealization in the phrase. This

¹!Xóõ is most like Zapotec in that it has three phonologized phonation categories. It also has tone, which to our understanding is not restricted by phonation and which is not analyzed in Chai & Garellek (2022). The Mandarin case is less relevant as its phonation is prosodic and tonally linked.

was also found to be the case in Betaza Zapotec (Crowhurst, Kelly & Teodocio 2016). Barzilai & Riestenberg (2021) found that tone and phrasal position also played a role in how voice quality is realized in San Pablo Macuiltianguis Zapotec. These studies show that Zapotec languages have much to contribute to our understanding of voice quality.

We find that residual H1* can adequately capture differences in voice quality and is a more robust measure of voice quality than H1*–H2*, adding credence to the use of this measure instead of H1*–H2* in voice quality research. The remainder of this paper is organized as follows. Section 3.2 provides a brief overview of the Santiago Laxopa Zapotec language. Section 3.3 describes the methods used in the data collection, data processing, and statistical modeling used in this study. Section 3.4 presents the results of the study. Section 3.5 concludes the paper.

3.2 Santiago Laxopa Zapotec

Santiago Laxopa Zapotec is a Northern Zapotec language of the Oto-Manguean language family (Adler & Morimoto 2016, Adler et al. 2018, Foley, Kalivoda & Toosarvandani 2018, Foley & Toosarvandani 2022, Sichel & Toosarvandani 2020a,b, Brinkhoff, Duff & Wax Cavallaro 2021, 2022). It is spoken by 981 people in the municipality of Santiago Laxopa, Ixtlán, Oaxaca, Mexico and a small number of other speakers from the diaspora in Mexico and the United States.

Santiago Laxopa Zapotec exhibits a four-vowel inventory (that is, /i/, /e/, /a/, and /u/), which is further distinguished by a four-way contrast in voice quality. This variety is unique because it is a Northern Zapotec that has developed a breathy voice (/V̥/) in addition to the two types of laryngealization that characterize the rest of the Zapotec languages, namely checked and articulated. Checked vowels are defined as a modal vowel that ends with a period of creaky voice or a glottal closure (/VV̥/ or /V'V̥/). Rearticulated vowels are also defined as a modal vowel that also has a period of creakiness or glottal closure but aligned to the middle of a vowel (/V̥VV/ or /V'V̥V/). This makes an otherwise modal vowel appear as if it is interrupted by this laryngealization.² This difference in laryngeal timing is one of the key differences between these phonations (see Ariza-García 2018 for a detailed typological study of voice quality distinctions in Zapotec languages).

Santiago Laxopa Zapotec is also tonal with three level tones (H, M, and L) and two contours (MH and HL) appearing in nominals (Brinkerhoff, Duff & Wax Cavallaro 2022).³ The language has a complex interaction between tone and phonation types. Every tone can appear with every phonation type, with two exceptions being that breathy voice cannot appear with the high tone and checked voice cannot appear with the rising contour tone. It is unclear whether these are accidental gaps or have

²This is different from how rearticulated vowels are defined in other languages, where they are a modal vowel followed by an “echo vowel” which may or may not have a glottal closure intervening between the modal and “echo” vowel (see Baird 2011). A fuller description of how these vowels are realized in one variety of Zapotec and the terms used are found in Avelino (2010).

³The tonal system of Santiago Laxopa Zapotec for verbs and other lexical categories is still being evaluated.

a phonetic underpinning.

These interactions between voice quality and tone present a rich environment for testing the reliability of voice quality measures in laryngeally complex languages.

3.3 Methods

3.3.1 Elicitation

Ten native speakers of SLZ (five female; five male) participated in a wordlist elicitation. Elicitation was performed in the pueblo of Santiago Laxopa, Ixtlán, Oaxaca, Mexico during the summer of 2022 using the built-in microphone of a Zoom H4n handheld recorder (16-bit, 44.5 kHz).

The wordlist consisted of 72 items repeated three times in isolation and the carrier sentence *Shnia' X chonhe lhas* [ʃn:ia' X tʃone ras] "I say X three times". This sentence was chosen to minimize any effect of the phrasal position (see Crowhurst, Kelly & Teodocio 2016: for a study about phrasal effects on voice quality in Zapotec). Additionally, there are no co-articulatory effects on voice quality found with this frame and the items in the wordlist. Between these 72 words, there were 11 words with breathy voice, 9 with rearticulated voice, 10 with checked voice, and 42 with modal. Thirteen of the 72 words were disyllabic and eight of the thirteen contained the same voice quality in each syllable. Of those 13 disyllabic words, only five contained different

voice qualities in each of the syllables. This resulted in 85 different syllables.⁴

The token selection for the wordlist was done in consultation with a native speaker. Similarly to Barzilai & Riestenberg (2021), we did not balance the tones with the different voice qualities. The frequency in which certain tones co-occur with certain voice qualities is uneven in the language, making it difficult to control for. The distribution of tones and voice quality across the 85 syllables used in this study are presented in Table 3.1. This imbalance was taken into account by including Tone as a fixed effect in our statistical models in Section 3.3.3.

Table 3.1: Distribution of tone and voice quality in the wordlist

	High	Mid	Low	Rising	Falling
Modal	14	9	15	2	10
Breathy	—	—	11	—	2
Checked	1	—	9	—	
Rearticulated	1	—	4	—	4

3.3.2 Data Processing

Each vowel of the target words in the carrier sentence condition was labeled following Garellek (2020) for where the vowel began and ended. Each vowel in the word list was annotated for speaker, word, vowel, tone, voice quality, and utterance number.

⁴There is ongoing debate about the status of strong syllables in Zapotec. The majority of evidence for the presence of stress comes from non-Northern Zapotec varieties (e.g., Chávez-Péón 2008, Mock 1988). At this time, evidence for stress in Northern Zapotec remains to be seen.

This labeling was conducted for each vowel in the target word from the elicitation list of the carrier sentences.

These vowels were then extracted and fed into VoiceSauce for acoustic measurement (Shue et al. 2011). The formants were measured using Snack (Sjölander 2004), while the fundamental frequency (f_0) was measured using the STRAIGHT algorithm (Kawahara, Chevigne & Patterson 1998). Spectral slope measures were corrected for formants and bandwidths (Hanson 1997, Iseli, Shue & Alwan 2007). Each vowel was measured with ten equal time intervals, resulting in 22890 data points in total.

The data were cleaned of outliers following the same steps as taken by Chai & Garellek (2022) in their study. The $H1^*$, $H1^*-H2^*$, and f_0 values were z-scored by speaker to reduce the variation between the speakers and provide a way to directly compare the different measures on the same scale. Data points with an absolute z-score value greater than 3 were considered outliers and excluded from the analyses. Within each vowel category, we calculated the Mahalanobis distance in the F1-F2 panel. Each data point with a Mahalanobis distance greater than 6 was considered an outlier and excluded from the analysis. Using the Mahalanobis distance allows us to compare the data points to the mean of the F1-F2 panel for each vowel category. The larger the Mahalanobis distance is the more deviant the data point is from the mean which in turn means that the data point was improperly tracked. This is comparable to what was done in Seyfarth & Garellek (2018), Chai & Ye (2022), and Garellek &

Esposito (2023).

Time points whose f_0 , F1, or F2 values were outliers were also excluded from H1* and H1*–H2* analyzes because H1* and H1*–H2* are calculated based on f_0 , F1, and F2. Energy was excluded if it had a zero value and then logarithmically transformed to normalize its right-skewed distribution. Afterward, the resulting logarithmically transformed data was z-scored, and any data point with a z-score greater than 3 was excluded. This outlier removal resulted in 1918 datapoints being removed.

After removing the outliers, we calculated residual H1* for the remaining data points following Chai & Garellek (2022). First, a linear mixed effects model was generated with the z-scored H1* as the response variable and the z-scored energy as the fixed effect. The uncorrelated interaction of the z-scored energy by speaker was treated as random. The energy factor resulting from this linear mixed-effects model was extracted. Finally, the z-scored H1* had the product of the z-scored energy and the energy factor subtracted from it, giving us the residual H1* measure.

The measures were then orthogonally coded according to their position in the vowel (first, middle, and third) and for tone (H, M, L, R, F) for statistical modeling.

3.3.3 Statistical modeling

Two linear mixed-effects regression models were fitted, one for the normalized H1*–H2* and residual H1*. Each model had the tone and the interaction between

voice quality and position in the vowel as fixed effects. Vowel and interaction between the speaker, the word, and the repetition were treated as random intercepts.⁵

The tone and the interaction between voice quality and position in the vowel were selected as fixed effects for several reasons. The first is that five unique tones appeared in the data and it is well established that tone interacts with voice quality in different ways (see Esposito & Khan 2020 and Garellek 2019 for discussion). By treating tone as a fixed effect in our model, we can account for these interactions. The interaction between voice quality and position in the vowel as a fixed effect was included to account for the temporal differences between the two different laryngealizations, checked and rearticulated vowels. Checked vowels in Zapotec languages have a glottal occlusion or a short period of creaky voice located at the right edge of the vowel. This is in contrast to rearticulated vowels, where there is a glottal occlusion or creaky voice in the middle of the vowel. Because this difference between checked and rearticulated vowels is temporal in nature, we can account for this difference through the interaction of voice quality and position in the vowel.⁶

The interaction between speaker, word, and repetition was treated as a random intercept because this allows us to consider that each speaker said each word on the elicitation list three times. This intercept accounts for not only the intra-speaker variability, but also the inter-speaker variability during each time the word was uttered.

⁵Measure ~ Phonation * Position + Tone + (1|Speaker : Word : Repetition) + (1|Vowel)

⁶Tone and voice quality are closely linked. By including only the positional interaction with voice quality, we can avoid collinear interactions that appear when we try to include tone in the interaction.

Treating a vowel as a random intercept allows us to capture the fact that each voice quality occurred with different vowels during elicitation.

Additional statistical modeling was performed with two generalized additive mixed models (GAMM) using the `mgcv` package in R (Wood 2017) and plotted using the `tidygam` package (Coretta 2024). Phonation was treated as a fixed effect to model the main effect of phonation on the acoustic measures. A smooth term for time (i.e., measurement number) was included to model the nonlinear relationship between time. A second smooth term was included to allow time to vary by phonation type. Random effects for speaker and the interaction between speaker and phonation were included to account for variability in the data due to individual differences between speakers and individual differences in how they produce the phonation types. A tensor product interaction between time and repetition was also included as a random effect to account for the variability in the data due to the different repetitions the speakers were asked to complete. The inclusion of tone and word did not improve the model fit and, therefore, were excluded from the final model.

3.4 Results

In interpreting the results, there are certain expectations about how these measures should capture the voice quality contrasts. As discussed in Garellek (2019), it is expected that breathy vowels should have a higher spectral slope than modal vow-

els across the two measures in keeping with observations about breathy vowels in other languages. Because both checked and rearticulated vowels make use of creaky voice, they should have a lower spectral slope than modal vowels. Additionally, because there is a temporal distinction between checked and rearticulated with where the creakiness appears, this should also be captured in the two measures.

3.4.1 $H1^* - H2^*$

Figure 3.1 shows the mean $H1^* - H2^*$ values for each voice quality at each of the ten vowel intervals. We see that the breathy, checked, and rearticulated values are lower than the modal values at each of the first nine intervals. In the final intervals, breathy and rearticulated are essentially equal to the modal value. In contrast, checked's value remains lower than the modal's value throughout the entire vowel. This measure does not match the expectations discussed above.

3.4.2 Residual $H1^*$

Figure 3.2 shows the mean residual $H1^*$ values for each voice quality at each of the ten vowel intervals. In contrast to Figure 3.1, we see that breathy has a higher residual $H1^*$ measure than modal throughout the duration of the vowel, which is consistent with other observations for breathy voice (Fischer-Jørgensen 1968). Checked and rearticulated both have lower values than the modal at each of the 10 intervals.




Figure 3.1: $H1^* - H2^*$ across the duration of the vowel. Points represent the mean of each measure across the ten intervals. The error bars around each point represent a 95% confidence interval. A line was plotted over each to show how the acoustic measure functions across the ten intervals.

In addition, it shows that the checked voice has a lower residual $H1^*$ value than the rearticulated voice at intervals 8 through 10. The rearticulated voice has a lower residual $H1^*$ value than the checked voice at intervals 1 through 7, showing the temporal distinction between these two voice qualities. This measure complies with the expectations discussed above.

3.4.3 Model Comparison

To assess the robustness of the models, we compared the residual $H1^*$ linear mixed-effects model to the $H1^* - H2^*$ linear mixed-effects model. This was done using two methods: direct comparison of the outputs of the two models in the same way as




Figure 3.2: Residual $H1^*$ across the duration of the vowel. Points represent the mean of each measure across the ten intervals. The error bars around each point represent a 95% confidence interval. A line was plotted over each to show how the acoustic measure functions across the ten intervals.

Chai & Garellek (2022) and the Akaike Information Criterion (AIC).

Table 3.2 compares the linear mixed effects models for $H1^* - H2^*$ and residual $H1^*$. In comparing these models, we find that the residual $H1^*$ model performed better than the $H1^* - H2^*$ model in distinguishing voice quality contrasts in Santiago Laxopa Zapotec. This is supported by the larger absolute value of the coefficient estimate, the lower standard error, and the higher t-value of the residual $H1^*$ to distinguish breathy, checked, and rearticulated vowels from modal vowels.

Table 3.3 shows the results of the AIC comparison between the $H1^* - H2^*$ and residual $H1^*$ models. The residual $H1^*$ model had a lower AIC than the $H1^* - H2^*$ model, indicating that the residual $H1^*$ model is a better fit for the data than the $H1^* - H2^*$

Table 3.2: Model comparison between H1*–H2* and Residual H1* in distinguishing Santiago Laxopa Zapotec voice quality.

Voice Quality Contrast	Model	β	Std. Error	t-value	p-value
Breathy vs Modal	H1*–H2*	0.04631	0.03806	1.21680	0.22372
	Res. H1*	0.23625	0.02866	8.24177	<0.001 ***
Checked vs Modal	H1*–H2*	-0.11880	0.03476	-3.41793	<0.001 ***
	Res. H1*	-0.40099	0.02621	-15.30098	<0.001 ***
Rearticulated vs Modal	H1*-H2	-0.09175	0.04588	-1.99968	0.04560 *
	Res. H1*	-0.44162	0.03450	-12.80027	<0.001 ***

model. Although AIC comparison is usually performed on nested models, it is still a useful tool for comparing non-nested models (Burnham & Anderson 2004b, Burnham, Anderson & Huyvaert 2011, Burnham & Anderson 2004a).

Table 3.3: AIC for the H1*–H2* and residual H1* models.

Model	AIC	Δ AIC
H1*–H2* model	43443.33	11182.76
Residual H1* model	32260.57	0

3.4.4 GAMM analysis and model comparisons

The GAMM analysis shows that there are nonlinear effects present in the data. This was expected due to the dynamic nature of voice quality in SLZ. Due to the nature of GAMM analyses, it is important to visually inspect the results to see how well the model fits the data. The model that best fits the data is the one that shows the clearest

distinction between the different voice qualities.

Figure 3.3 shows the predicted values for $H1^* - H2^*$ from the GAMM. We see that the different voice qualities are very difficult to observe clearly. We see that modal and breathy voice occupy the same space. However, checked and rearticulated voice are more separated from modal, with both appearing lower in the graph, which is expected because of creaky voice. We also observe that the rearticulated voice begins to increase throughout the final third of the vowel until it is higher than the modal voice.




Figure 3.3: GAMM smooths for $H1^* - H2^*$ across the duration of the vowel.

The difference plots in Figure 3.4 show three panels for a comparison between the modal voice and each of the other voice qualities. The leftmost panel shows that the difference between modal and breathy voice never reaches significance. This means

that the breathy voice and the modal voice are essentially the same according to the model. The middle panel shows the difference between modal and checked voice. This shows that checked voice is significantly different from modal voice throughout the duration of the vowel, except for a short period of time right before measurement number 10. The rightmost panel shows the difference between modal and rearticulated voice. This panel shows that rearticulated voice is significantly different from modal voice for the first two-thirds of the vowel. However, in the final third of the vowel, rearticulated voice is not significantly different from modal voice.




Figure 3.4: GAMM difference plots for $H1^* - H2^*$ across the duration of the vowel.

The residual $H1^*$ GAMM showed a clearer distinction between the different voice qualities than $H1^* - H2^*$. This is especially clear in the smooth plot of the predicted values in Figure 3.5. In this plot, we see that beginning at measurement number 3

breathy voice is higher than modal voice. This is expected because the breathy voice is characterized by a higher spectral slope than the modal voice. We also see that the checked and rearticulated voice are lower than modal voice, which is also expected because of the presence of creaky voice, which is characterized by a lower spectral slope than modal voice. Additionally, we see that the temporal difference between checked and rearticulated voice is present in the smooth plot. The checked voice is higher than the rearticulated voice for the first two-thirds of the vowel, but then the checked voice becomes lower than the rearticulated voice in the final third of the vowel. This is expected because checked voice is characterized by a glottal occlusion at the end of the vowel, while rearticulated voice is characterized by a glottal occlusion in the middle of the vowel. Furthermore, this plot suggests that rearticulated voice is predominately more creaky than checked voice in the first two thirds of the vowel. This suggests that the glottal occlusion in the middle of the vowel predominantly affects the first half of the vowel. This difference might be due to the need to distinguish the rearticulated vowel from the checked vowel more effectively.

Figure 3.6 shows the difference plots for residual H1*. The leftmost panel shows that the difference we observed in Figure 3.5 between the modal and the breathy voice is significant for the last two thirds of the vowel. The middle panel shows that the difference observed between the modal and checked voice is significant across the entire duration of the vowel. The rightmost panel shows that the difference we observe




Figure 3.5: GAMM smooths and difference plots for residual $H1^*$ across the duration of the vowel.

between modal and rearticulated voice is significant for almost the entire duration of the vowel except for measurement number 10.

In comparing the two GAMM analyses, we find that the residual $H1^*$ model provides a clearer distinction between the different voice qualities than the $H1^* - H2^*$ model. This is further evidence that residual $H1^*$ is a more robust measure of voice quality than $H1^* - H2^*$ in Santiago Laxopa Zapotec. Additionally, the GAMM analysis shows that there appear to be some amount of phasing in the data, which the linear models were unable to capture. This is especially clear in the residual $H1^*$ model, where the difference plots show that the breathy voice is only significantly different from the modal in the final two thirds of the vowel. This suggests that the breathy




Figure 3.6: GAMM smooths and difference plots for residual $H1^*$ across the duration of the vowel.

voice aligns with the end of the vowel. This is similar to the predictions made by Silverman (1997a,b) about the alignment of nonmodal phonation in laryngeally complex languages. This will be further discussed in Chapters 6.

3.5 Conclusion

In conclusion, we find that residual $H1^*$ is a more robust measure of voice quality than $H1^* - H2^*$ in Santiago Laxopa Zapotec. This is supported by the results of the linear mixed-effects models, which show that residual $H1^*$ is better at distinguishing breathy, checked, and rearticulated vowels from modal vowels. This is further supported by the AIC comparison, which shows that the residual $H1^*$ model is a better

fit for the data than the H1*–H2* model. These results lend credence to the claims of Chai & Garellek (2022) and support using residual H1* instead of H1*–H2* in voice quality research, especially in laryngeally complex languages.

However, the results neither suggest nor support the notion that residual H1* should be the sole measure used to determine phonation quality. Instead, we suggest that residual H1* should be included as one of the measures that researchers should consult during their acoustic studies.

Chapter 4

The acoustic space of voice quality in Santiago Laxopa Zapotec

4.1 Introduction

This chapter studies the acoustic dimension of voice quality in Santiago Laxopa Zapotec (SLZ) using a Multidimensional Scaling (MDS) analysis of acoustic data. MDS is a statistical method that reduces the dimensionality of a dataset and visualizes the relationships between data points. This study uses MDS to visualize the acoustic space of voice quality in SLZ. This analysis provides information on the acoustic correlates of voice quality in SLZ and contributes to our understanding of the phonetic properties of this under-documented language.

This study is based on the work conducted by Keating et al. (2023) on the acoustic space of voice quality in 11 languages. However, this study focuses on a single language, SLZ, and provides a detailed analysis of the acoustic properties of voice quality in this language. The results of this study will contribute to our understanding of the phonetic properties of SLZ and how the acoustic properties of voice quality in this language compare with other languages.

4.2 Methods

4.2.1 Participants

This study uses data collected from 10 native speakers of SLZ during the summer of 2022. Participants were recruited from the community of Santiago Laxopa, Oaxaca, Mexico. All participants were native speakers of SLZ. The participants were between 18 and 60 years old and consisted of five males and five females.

4.2.2 Recordings

Participants were asked to perform a word list elicitation task consisting of 72 words. These words were selected to elicit the entire range of types of voice quality in SLZ, including modal voice, the two kinds of creaky (i.e., checked and rearticulated), and breathy voice. The words were selected based on previous research conducted

as part of the Zapotec Language Project at the University of California, Santa Cruz (*Zapotec Language Project – University of California, Santa Cruz* 2022). Because participants were not literate in SLZ, the word list was prompted for them by asking them “How do you say [word in Spanish]?” by myself and another researcher in Zapotec. Participants were asked to respond with the desired word in the carrier phrase *Shnia’ [WORD] chonhe lhas* “I say [WORD] three times.” which was repeated three times. These utterances were recorded in a quiet environment using a Zoom H4n digital recorder. The recordings were saved as 16-bit WAV files with a sampling rate of 44.1 kHz.

4.2.3 Acoustic measuring

The resulting audio files were then processed in Praat to isolate the vowel portion of each word. The onset of the vowel was set to the second glottal pulse after the onset, and the offset of the vowel was set to the last glottal pulse before the decrease in amplitude at the end of the vowel (Garellek 2020). The vowel was then extracted and saved as a separate file for analysis.

These vowels were fed into VoiceSauce (Shue et al. 2011) to generate the acoustic measures for the studies discussed in this dissertation. Because many acoustic measures are based on the fundamental frequency, this measure was calculated using the STRAIGHT algorithm from (Kawahara, Cheveigne & Patterson 1998) to esti-

mate the fundamental frequency in millisecond (ms) intervals. Once the fundamental frequency is calculated, VoiceSauce then uses an optimization function to locate the harmonics of the spectrum, finding their amplitudes.

VoiceSauce then uses the Snack Sound toolkit (Sjölander 2004) to find the frequencies and bandwidths of the first four formants, also at millisecond intervals. The amplitudes of the harmonics closest to these formant frequencies are located and treated as the amplitudes of the formants. These formant frequencies and bandwidths are used to correct the harmonic amplitudes for the filtering effects of the vocal tract, using Iseli, Shue & Alwan's 2007 extension of the method employed by Hanson (1997). Each vowel was measured across ten equal time intervals, resulting in 22890 data points in total. These measures were then z-scored by speaker to reduce the variation between speakers and provide a way to compare the different measures directly on similar scales.

4.2.4 Data processing

Data points with an absolute z-score value greater than three were considered outliers and excluded from the dissertation analyzes. The Mahalanobis distance was calculated in the F1-F2 panel within each vowel category. Each data point with a Mahalanobis distance greater than six was considered an outlier and excluded from the analysis. Using the Mahalanobis distance allows us to compare the data points

to the mean of the F1-F2 panel for each vowel category. The larger the Mahalanobis distance is the more deviant the data point is from the mean which in turn means that the data point was improperly tracked. This is comparable to what was done in Seyfarth & Garellek (2018), Chai & Ye (2022), and Garellek & Esposito (2023).

Energy was excluded if it had a zero value and then logarithmically transformed to normalize its right-skewed distribution. Afterward, the resulting log-transformed energy was z-scored and any data point with a z-score greater than three was excluded. This outlier removal resulted in 1918 data points being removed.

All data points were then z-scored by speaker to reduce the variation between speakers and provide a way to compare the different measures directly on the same scale.

The residual H1^{*} was then calculated for the remaining data points following Chai & Garellek (2022). First, a linear mixed effects model was generated with the z-scored H1^{*} as the response variable and the z-scored energy as the fixed effect. The uncorrelated interaction of the z-scored energy by speaker was treated as random. The energy factor resulting from this linear mixed-effects model was extracted. Finally, the z-scored H1^{*} had the product of the z-scored energy and the energy factor subtracted from it to produce the residual H1^{*} measure.

Once these steps were completed, the mean of each combination of phonation and speaker was taken for the fourth to seventh interval of the vowel. This is similar to

what Keating et al. (2023) did by taking the middle of the vowel for their analysis. This choice minimizes the effect of the onset and offset of the vowel on the acoustic measures, which are more likely to be affected by the surrounding consonants and should give us the most accurate representation of the vowel quality. Because z-scores were used, this resulted in negative measures, which presents a problem for MDS analyses. To correct for this, I added the absolute value of the minimum z-score to each measure. This results in a dataset that still preserves the relative differences in the scores while providing a dataset that is all positive for the MDS analysis.

4.2.5 Statistical analysis

Multidimensional scaling analysis (MDS) is a type of dimensionality reduction to visualize the relationships between data points (Kruskal & Wish 1978). Using MDS is especially effective when many variables could contribute to the data. In the case of voice quality, this is especially warranted.

As shown in Kreiman et al. (2014, 2021) and Garellek (2020), voice quality is psychoacoustically complex and a single measure is not enough to capture the full range of voice quality. Instead, multiple measures are required that function as cues for the different types of voice quality. For example, a vowel characterized as having a breathy voice has an elevated spectral-slope and a lower harmonics-to-noise ratio than a modal voice. A creaky voice has a lowered spectral-slope and a lowered harmonics-to-noise

ratio.

Because MDS analyses that contain many variables can result in rather unmeaningful results, I chose to focus on the speaker by phonation interaction. This allows us to see how speakers differ in their production of the different voice qualities. This choice to focus on speaker by phonation means that each speaker's production of each of the four phonation contrasts is represented as a single point in the MDS plot (e.g., one point for the first speaker's modal voice, one for their checked voice, one for their rearticulated voice, and one for their breathy voice). This is similar to what Keating et al. (2023) did in their study of the acoustic space of voice quality in 11 languages, except that they compared the language by voice quality interaction. Both of these interactions show us similar information. The analysis of speaker-by-voice quality shows us the acoustic space within a language, while the analysis using language-by-voice quality shows us the acoustic space between languages.

The MDS analysis was conducted using the `metaMDS` function in the `vegan` package (Oksanen et al. 2025) in the R programming language (R Core Team 2024). The Manhattan distance was used to estimate the differences between the speaker-by-phonation pairs. Because the distances are non-Euclidean, the MDS analysis was conducted using the nonmetric option.

This algorithm resulted in a solution that involves several different dimensions. The number of dimensions retained directly affects how well the original data is cap-

tured. Too many dimensions and the data are overfitted; too few, and the data are underfitted. To determine the number of dimensions to retain, I used a scree plot to plot the stress of each dimension. As shown in Figure 4.1, most of the data are captured in the first four dimensions. These four dimensions were retained for the analysis.




Figure 4.1: Scree plot showing the stress for each dimension for the MDS analysis.

4.3 Results

4.3.1 Acoustic space of voice quality

The results of the MDS analysis show that the acoustical space is represented primarily by a three-dimensional space.¹ In all subsequent plots, breathy voice is represented by orange, checked voice with blue, rearticulated voice with green, and modal voice with black. In each of the plots, the modal voice is generally more densely packed than the non-modal voice qualities. This is likely due to the fact that the modal voice represents approximately 60% of the data, while the non-modal voice qualities represent approximately 40% of the data.

Figure 4.2 shows the first dimension plotted against the second dimension. In this plot, we observe that breathy voice is located in the top left of the plot, modal voice is located in the bottom center of the plot, and the two types of creaky voices are located to the right of the plot, with checked voice located at the extreme right of the plot, and rearticulated voice located closer to the center. From this plot, we see that the first dimension separates breathy, modal, and creaky voices. The second dimension separates the modal voice, bottom of the plot, from the non-modal voice qualities, top of the plot.

Figure 4.3 compares the first dimension with the third dimension. In this plot, we

¹ A 3D plot showing the acoustic space can be found at https://www.mlbrinkerhoff.me/files/3d_plot.html.




Figure 4.2: Two-dimensional MDS solution showing the first and second dimensions.

observe that the breathy voice is located in the bottom left of the plot, modal voice is located in the very center of the plot, and the two types of creaky voices are located in the top right of the plot. It should be noted that the distribution of the different voice qualities follows a line from bottom left to top right in the plot. This suggests that the first and third dimensions capture similar information about voice quality in SLZ.

Figure 4.4 shows the first dimension plotted against the fourth dimension. This plot is very similar to Figure 4.2, with the only difference being that the fourth dimension moves the modal voice from the bottom-center of the plot to almost the exact center of the plot.

Figure 4.5 shows the second dimension plotted against the third dimension. This




Figure 4.3: Two-dimensional MDS solution showing the first and third dimensions.

plot is essentially the same as Figure 4.3, except that the coordinates are flipped.

Figure 4.6 shows the second dimension plotted against the fourth dimension. This plot shows that the modal voice and the non-modal voice are separated into two different clusters, with modal voice located to the extreme left of the plot and the nonmodal voice qualities located to the right of the modal grouping. Again, as first seen in Figure 4.4, the fourth dimension centralizes the modal voice, but no discernible pattern is observed for the other phonations.

Figure 4.7 shows the third dimension plotted against the fourth dimension. This plot is very similar to Figure 4.4 with the exception that along the third dimension checked and rearticulated voice have swapped places. Checked voice is more central-




Figure 4.4: Two-dimensional MDS solution showing the first and fourth dimensions.

ized in the plot, whereas rearticulated voice is located more to the right of the plot. Again, we observe that the modal voice is located in the center of the plot.

4.3.1.1 Interim summary for dimension plots

The plots of the MDS analysis show that the acoustic space of voice quality in SLZ is primarily represented by a three-dimensional space. The first dimension and third dimensions are very similar in that they capture a continuum from breathy, to modal, and finally creaky voice. This is similar to what Keating et al. (2023) found in their study for the second dimension.

This continuum from breathy to modal to creaky is very similar to the open-

NMDS Plot with Duration added (Dimension 2 x Dimension 3)




Figure 4.5: Two-dimensional MDS solution showing the second and third dimensions.

quotient model of voice quality proposed by Gordon & Ladefoged (2001), as illustrated in Figure 4.8. Because of this similarity to the open-quotient model, the first and third dimensions are likely capturing the open-quotient of the glottis.

From the plots involving the second dimension, we see that this dimension separates the modals from the non-modal ones. This is similar to what we observe with the various harmonics-to-noise ratios and the cepstral peak prominence (CPP) which are all measures of the amount of noise present in the signal across various bandwidths (de Krom 1993, Hillenbrand & Houde 1996, Blankenship 2002, Ferrer Riesgo & Nöth 2020). In addition to the amount of noise, it also captures the strength of the vocal fold vibration, similar to what Garellek et al. (2021) found in their study of glot-




Figure 4.6: Two-dimensional MDS solution showing the second and fourth dimensions.

tal consonants and nonmodals. In their study, they found that the modal voice had the highest strength of vocal fold vibration (measured by the Strength of Excitation), while the non-modal voice had a lower strength of vocal fold vibration. This suggests that the second dimension is to capture the amount of aperiodic noise in the signal, the strength of the vocal fold vibration, or both.

The fourth dimension is less clear about what it is potentially capturing. In all of the plots involving the fourth dimension, we see that modal voice is always located near the center of the plot, while the nonmodal voice qualities are located around this




Figure 4.7: Two-dimensional MDS solution showing the third and fourth dimensions.



Figure 4.8: A diagram showing the relationship between breathy, modal, and creaky phonation types from Gordon & Ladefoged (2001).

point depending on the patterns from the other dimensions. This suggests that the fourth dimension possibly captures something about modal voice.

The rest of this chapter will focus on how the different acoustic measures contribute to the different dimensions of the MDS analysis. This will be followed by a general discussion of both the MDS analysis dimensions and the acoustic measures that are correlated with these dimensions. This discussion will focus on how the re-

sults of this study relate to previous work on voice quality and the implications of these results for our understanding of the acoustic space of voice quality.

4.3.2 Acoustic correlates of voice quality

Looking at the visual representation of the dimensions is only part of the full story. In order to fully understand what is occurring, we need to determine how the acoustic measures contribute to each of the different dimensions. There are two ways to do this: (i) looking at the amount of weight each acoustic measure contributes to the different dimensions, and (ii) looking at how correlated the different acoustic measures are to the different dimensions (Kruskal & Wish 1978, Hastie, Tibshirani & Friedman 2009). Both of these methods are useful in understanding the shape of the data. In the following discussion, I will take the second approach and look at the correlations between the different acoustic measures and the different dimensions. This is done to make the resulting discussion easier to follow.

Table 4.1 shows the correlation score, computed by the `cor` function in R, for each acoustic measure and dimension. The four largest correlations in each dimension are in bold. The choice to bold the four largest correlations was arbitrary and was done to make it easier to discuss. Although only the four largest correlations are bolded, there are several correlations that share similar correlation values. These cases will be discussed as needed.

Table 4.1: Correlations for each acoustic measure to the four dimensions (NMDS1, NMDS2, NMDS3, NMDS4). The four largest correlations in each dimension are bolded.

Acoustic Measure	NMDS1	NMDS2	NMDS3	NMDS4
H1*–H2*	-0.221	-0.339	0.031	0.314
H2*–H4	-0.437	0.239	-0.689	-0.364
H1*–A1*	-0.828	0.048	-0.459	0.044
H1*–A2*	-0.855	-0.067	-0.343	0.114
H1*–A3*	-0.809	-0.218	-0.297	0.126
H4*–H2k*	-0.452	-0.598	0.294	0.366
H2k*–H5k*	0.152	0.023	0.101	0.057
residual H1*	-0.290	-0.443	-0.722	0.084
H2*	-0.157	-0.555	-0.679	0.114
H4*	0.295	-0.778	0.078	0.479
A1*	0.756	-0.549	0.092	0.124
A2*	0.779	-0.476	-0.103	0.086
A3*	0.735	-0.416	-0.211	0.093
CPP	-0.590	-0.606	0.209	-0.179
HNR < 500 Hz	-0.513	-0.792	0.152	-0.202
HNR < 1500 Hz	-0.275	-0.799	0.323	-0.290
HNR < 2500 Hz	-0.327	-0.714	0.391	-0.348
HNR < 3500 Hz	-0.446	-0.644	0.393	-0.356
Strength of Excitation	-0.013	-0.741	-0.238	0.145
SHR	0.144	-0.176	0.122	-0.597
Energy	-0.080	-0.793	-0.015	0.341
Duration	-0.622	0.539	0.257	0.030

4.3.2.1 Dimension 1

From Table 4.1, we observe that the first dimension is negatively correlated with the acoustic slope measures of H1*–A1*, H1*–A2*, and H1*–A3* ($r^2 \approx -0.825$ for all three of these measures). These measures are all types of spectral slope measures which attempt to normalize the amplitude of the formant corrected H1 against the

amplitudes of the formant correct harmonic closest to the first three formants. It has been noted that these measures are correlated with the open-quotient of the glottis (see Garellek 2019, 2022 for an overview of these measures). Because these measures all capture the same information and they are all highly correlated with the first dimension, we can conclude that the first dimension is primarily concerned with the spectral slope of the signal (i.e., the open-quotient of the glottis).

The first dimension is also positively correlated with the amplitude of the first three formants (that is, $A1^*$, $A2^*$, and $A3^*$). These acoustic measures also share correlations of a similar value ($r^2 \approx 0.75$). It is also worth noting that these amplitude measures are all used in the normalization of the spectral slope measures (i.e., $H1^*-A1^*$, $H1^*-A2^*$, and $H1^*-A3^*$). I believe that this fact suggests that the first dimension is primarily concerned with the spectral slope of the signal, but is also factoring in the amplitude of the formants.

4.3.2.2 Dimension 2

The second dimension is strongly correlated with the harmonics-to-noise ratio (HNR) measures. The HNR measures $HNR < 500$ Hz, $HNR < 1500$ Hz, and $HNR < 2500$ Hz are all negatively correlated with the second dimension ($r^2 \approx -0.79$). These measures are all measures of the amount of noise in the signal and are used to indicate the amount of periodicity in the signal (e.g., whether something is modal or non-modal). This suggests that the second dimension is concerned with the amount of

noise in the signal.

Furthermore, there are strong negative correlations with energy, which is the root mean squared energy of the acoustic signal, and the Strength of Excitation (SoE), which is defined as “the instant of significant excitation of the vocal-tract system during production of speech” and represents “the relative amplitude of impulse-like excitation” (Mittal, Yegnanarayana & Bhaskararaao 2014: p. 1934). In other words, the SoE correlates with how strongly the vocal folds vibrate during the signal with modal voice showing the strongest amount of voicing and nonmodal phonation the least. This suggests that the second dimension is also concerned with strength of the vocal-fold vibration.

The last measure that shows a strong correlation with the second dimension is the amplitude of the fourth harmonic ($H4^*$). This measure is negatively correlated with the second dimension ($r^2 \approx -0.78$). This measure is typically used to help normalize the amplitude of the first formant and is used in the calculation of the spectral slope measures (e.g., $H2^* - H4^*$).

Together, given the correlations of the HNR measures, energy, and SoE, we can conclude that the second dimension is primarily concerned with the periodicity of the signal (i.e., the amount of noise in the signal) and the strength of the vocal fold vibration.

4.3.2.3 Dimension 3

The third dimension is negatively correlated with the two spectral slope measures of residual H1* and H2*–H4* ($r^2 \approx -0.7$). Residual H1* is a measure that has been shown to be robust in capturing the strength of the first harmonic (Chai & Garellek 2022, Brinkerhoff & McGuire 2025), which was the original goal of using higher harmonics to normalize H1 (Fischer-Jørgensen 1968). H2*–H4* has also been shown to be another spectral slope measure that can capture the differences in amplitude between different phonation types (Garellek 2019, Garellek et al. 2016, Kreiman et al. 2014, 2021). This suggests that the third dimension is primarily concerned with the spectral slope of the signal, similar to what we observed in the first dimension.

In addition to these measures, the amplitude of the second harmonic (H2*) was also found to be negatively correlated with the third dimension ($r^2 \approx -0.69$). This measure is typically used to help normalize the amplitude of the first harmonic and is used in the calculation of the spectral slope measures (e.g., H1*–H2*, H2*–H4*, etc.). Due to its use in spectral slope measures and that H2*–H4* is also negatively correlated with the third dimension, we can conclude that H2* also contributes to the spectral slope of the signal. This further supports the idea that the third dimension is primarily concerned with the spectral slope of the signal.

The last acoustic measure also correlated with the third dimension is the spectral slope measure of H1*–A1* ($r^2 \approx -0.46$). Even though the correlation is not as strong

as the other acoustic measures, it still shows that the third dimension corresponds to the spectral slope of the signal (i.e., the open-quotient of the glottis).

4.3.2.4 Dimension 4

In the fourth dimension, we observe that the correlations are less clear than in the previous three dimensions. The strongest correlations in this dimension with the Subharmonic-to-Harmonic ratio (SHR; $r^2 \approx -0.60$) describe the relative strength of any subharmonics (interharmonics) in the signal (Sun 2002). The subharmonics in the signal correspond to alternating periods in the time domain (that is, period doubling), which typically occurs in creaky voice and with broader laryngeal constrictions (see Herbst 2021 for concerns about this acoustic measure).

The positive correlation with H4* ($r^2 \approx 0.48$) suggests that the fourth dimension may also capture some information about the amplitude of the higher harmonics. This is also true with the positive correlation with H4*–H2k* ($r^2 \approx 0.37$) and H2*–H4* ($r^2 \approx 0.36$), which are both measures that capture the spectral slope of the signal. This suggests that the fourth dimension may capture information about the amplitude of the harmonics in the signal.

4.3.2.5 Interim summary for acoustic correlates

Based on the correlations observed in Table 4.1, we can summarize the acoustic correlates of each dimension as follows: Dimension 1 captures the spectral slope of

the signal, primarily through the spectral slope measures $H1^* - A1^*$, $H1^* - A2^*$, and $H1^* - A3^*$. This dimension appears to be related to the open-quotient of the glottis. Dimension 2 captures the amount of noise in the signal and the strength of vocal fold vibration, primarily through the HNR measures ($HNR < 500$ Hz, $HNR < 1500$ Hz, and $HNR < 2500$ Hz), Energy, and Strength of Excitation. This dimension separates modal from nonmodal voice quality, which is what periodicity is concerned with. Dimension 3 captures the spectral slope of the signal as well, just as Dimension 1 does, primarily through residual $H1^*$, $H2^* - H4^*$, and $H2^*$. This dimension also appears to be related to the open-quotient of the glottis. Finally, Dimension 4 captures the periodicity in the signal through SHR and also appears to capture some information about the spectral slope, as evidenced by the amplitude of the higher harmonics through $H4^*$ and $H4^* - H2k^*$. However, it is not entirely clear whether this dimension is primarily concerned with the spectral slope of the signal or the periodicity in the signal.

4.4 Discussion

The results of this study show that the voice quality in SLZ occupies an acoustic space, but also shows that this space is similar to what Keating et al. (2023) found in their study of phonation in 11 languages. However, the results of this study differ from Keating et al. (2023). Instead of a two-dimensional space like in Keating et al. (2023), SLZ's voice quality occupies a three-dimensional space. Despite these differences, the

behavior of the dimensions is similar to that in Keating et al. (2023).

In the analysis presented in this chapter, we see that the first and third dimensions are primarily concerned with a spectral slope continuum from positive spectral slope to negative spectral slope which correlates to breathy voice to modal voice and finally to creaky voice. As extensive research has shown, the spectral slope of the signal is closely related to the open-quotient of the glottis, or in other words, how open or closed the glottis is during phonation (see Garellek (2022) for an overview of this history). This continuum was also found to exist in the second dimension of Keating et al.'s 2023 MDS analysis. Based on Keating et al. (2023) and the analysis presented in this chapter, at least one of the dimensions of any acoustic space related to voice quality must correspond to the spectral slope of the signal.

As mentioned in Section 4.3.1.1, the first and third dimensions of this chapter's analysis and Keating et al.'s second dimension, bear a striking resemblance to the voice quality model proposed by Gordon & Ladefoged (2001). In Gordon & Ladefoged's model, voice quality is described as being a single continuum based on how open or closed the glottis is during speech. The more open the glottis, the more breathy the phonation will be. The more closed the glottis, the more creaky the phonation will be. This model also claims that laryngeal consonants [h] and [?] also exist along this continuum and represent the extreme ends of this continuum. This model can be visually represented as a line with [h] on one end and [?] on the other end. The

various voice qualities exist between these two extremes, as represented in Figure 4.9.




Figure 4.9: A diagram showing the relationship between breathy, modal, and creaky phonation types. Based on Gordon & Ladefoged (2001).

As mentioned above, the measures that correlate the most with the first and third dimensions are several spectral slope measures (i.e., H1^{*}–A1^{*}, H1^{*}–A2^{*}, H1^{*}–A3^{*}, residual H1^{*}, and H2^{*}-H4^{*}). These measure correlations provide further support that the first and third dimensions of the MDS analysis are concerned with the spectral slope of the signal, which in turn is closely correlated with the state of the glottis during phonation (Holmberg et al. 1995, Kreiman, Gerratt & Antoñanzas-Barroso 2007, Garellek et al. 2016, Garellek 2019, Chai & Garellek 2022).

The second dimension of this chapter's analysis is concerned with dividing the acoustic space into modal and nonmodal voice qualities. This split between modal and nonmodal is similar to the split that exists for periodicity and strength of voicing. The more modal the voice quality is, the greater the amount of periodicity or voicing that we observe in the signal. I propose that this dimension is concerned with how harmonic the signal is, or in other words, how much noise is present in the acoustic signal. This is similar to what Keating et al. (2023) found in their study, where the first dimension of their analysis was primarily concerned with making the same split

in the acoustic space. Based on these results, I propose that another dimension in the acoustic space of voice quality must correspond to the amount of noise in the signal. This proposal is further supported by the fact that the second dimension of this chapter's analysis is primarily correlated with the harmonics-to-noise ratios, energy, and SoE measures, which are all measures of the amount of noise or the amount of energy present in the acoustic signal.

The last dimension of the MDS analysis is less clear in what it is capturing. The correlation with the subharmonics-to-harmonic ratio (SHR) suggests that this dimension might be capturing whether or not there is period doubling which is a common feature in creaky voice. However, there is no easily describable pattern that emerges with this measure.

The results of this study show that the voice quality acoustic space in SLZ is primarily represented by a three-dimensional space. However, it can primarily be broken down into two aspects that the acoustic space is attempting to capture: (i) the spectral slope of the signal and (ii) the amount of noise in the signal. This can be represented as a primarily two-dimensional space with higher dimensions adding additional information related to these two primary concerns. This can be visualized as a two-dimensional space with the first dimension representing the spectral slope of the signal and the second dimension representing the amount of noise in the signal, as shown in Figure 4.10.




Figure 4.10: A two-dimensional representation of the acoustic space of voice quality in SLZ. The horizontal axis represents the spectral slope of the signal, while the vertical axis represents the amount of noise or energy in the signal.

4.5 Conclusion

Although the discussion has predominately been about the correlations of the measures that contribute to the different dimensions, it is important to note that the measures are not independent of each other. Instead, all of the measures contribute to the acoustic space of voice quality in SLZ to some extent or another. Just because a measure has a low correlation does not mean that it does not contribute to the acoustic space. Rather than thinking of the measures as independent of each other, it is better to think of them as a group of measures that work together to create the acoustic space

of voice quality in SLZ. This is especially true given the fact that the MDS analysis is a reduction of the data to a few dimensions. This analysis offers a snapshot of the voice quality acoustic space in SLZ, but is not the full picture.

Furthermore, as will be discussed in Chapter 5, another way in which we can determine which measures are the most important is by performing a classification and regression tree analysis (Breiman et al. 1986, Breiman 1996, 2001).

Chapter 5

Trees reveal the importance of measures in SLZ

5.1 Introduction

The MDS analysis presented in Chapter 4 provides us an understanding of the acoustic shape that voice quality takes in SLZ. This shape is defined by different dimensions that correspond to the glottal configuration needed to produce each voice quality and the amount of noise in the signal. The MDS analysis also provides us with a way to visualize how the different voice qualities occupy an acoustic space. Finally, the chapter discussed how different acoustic measures are correlated with different dimensions of the acoustic space. This provides us with a potential avenue to explore

which measures contribute to our understanding of the different voice qualities in SLZ. However, the analysis does not tell us which measures are the most important in making the splits between the different voice qualities. This is where decision trees are helpful as they provide a way to cut through the noise and reveal which measures play the most important role in dividing that acoustic space.

In this chapter, I present an analysis using a flavor of decision trees called Random Forests (Breiman et al. 1986, Breiman 2001). Random Forests are a type of ensemble learning method that combines multiple decision trees that are built on the same data. This allows us to take advantage of the strengths of decision trees while minimizing the weaknesses that are present in classical decision trees (see Hastie, Tibshirani & Friedman 2009, Boehmke & Greenwell 2019, James et al. 2021 for a discussion of the strengths and weaknesses of the different types of decision trees).

In this chapter, I show that only a small number of acoustic measures are needed to classify the different voice qualities and make the splits in the acoustic space.

5.2 What are Decision Trees

Decision trees are a statistical tool that helps to reveal which predictors divide the space under investigation. Essentially, this is done by stratifying or segmenting the predictor space into some number of simpler regions. The rules that divide the space into these regions are based on some aspect of the predictors (see Hastie, Tibshirani &

Friedman 2009, James et al. 2021 for explanations of the statistics and how to perform these analyzes in R).

These trees can be used for both regression and classification. In the case of regressions, it splits the predictor space into regions and calculates how the item under discussion behaves in each region. This process of splitting into regions and calculating how something responds in that region continues until some stopping rule is applied, which is usually defined to some number of terminal nodes. This resulting tree is rather large and is then pruned based on the cost-complexity pruning to a subset of itself. This subsetted tree is the tree that has minimized its cost-complexity criterion for all potential subsets. That is, it balances the trade-off between the complexity of the tree and its fit to the data.

In the case of classification, the algorithms that result in a tree are very similar to those used for regression trees. The main difference in the algorithm comes from what is used to split the nodes and how the tree is pruned. Instead of predicting a continuous outcome like with regression trees, classification trees predict a categorical outcome. The predictor space is divided into regions and within each region, the majority class is assigned as the predicted class for that region. This process continues until a stopping rule is applied, similar to regression trees. The resulting tree can also be pruned to avoid overfitting, using a cost-complexity criterion.

Decision trees are easy to interpret and visualize, making them an ideal choice

for understanding the structure of data and how the different predictors interact with data (Hastie, Tibshirani & Friedman 2009, James et al. 2021).

5.3 Decision trees in linguistics

The use of decision trees in linguistics is not new. One of the first uses was done by Tagliamonte & Baayen (2012), where they illustrated the use of decision trees in investigating which sociolinguistic factors were the most important in the use of *was* versus *were* in York English.

Recently, decision trees were used to show which acoustic measures were important in making the split in the acoustic space for voice quality (Keating et al. 2023). Keating et al. performed a simple decision tree analysis to supplement their MDS analysis of voice quality in 11 languages. The results of this analysis are shown in Figure 5.1. Decision trees like the one in Figure 5.1, show the binary splits that are made in the space and what predictor, and the value of that predictor, makes that split. In the case of Keating et al.'s (2023) tree, the first split is made on the harmonics-to-noise ratio over the frequency range from 0 to 500 Hz for the middle third of each vowel. This split is made at the z-score of 0.48. If the predictor value is greater than or equal to 0.48, the dominant voice quality of that region is modal. If $\text{HNR} < 500 \text{ Hz}$ value is less than 0.48, the region needs to be further split.

The next split in the region is made on the subharmonic-to-harmonic ratio for the




Figure 5.1: Classification tree of phonation categories from Keating et al. (2023).

Abbreviations used in this figure are: HNR05_means002: harmonics-to-noise ratio over the frequency range from 0 Hz to 500 Hz for the middle third of each vowel; SHR_means002: subharmonic-to-harmonic ratio for the middle third of each vowel; H1H2c_means002: $H1^* - H2^*$ for the middle third of each vowel; B: breathy, M: modal, and C: creaky phonation categories.

middle third of each vowel. If the value of this predictor is less than 0.0024, the voice quality is classified as breathy. If the value is greater than or equal to 0.0024, the region needs to be split further.

The final split in the region is made on the $H1^* - H2^*$ for the middle third of each vowel. If the value of this predictor is less than 0.45, the voice quality is classified as creaky. If the value is greater than or equal to 0.45, the voice quality is classified as modal. This tree shows that using only three acoustic measures, one can classify the voice quality of the data. This is a powerful tool for understanding the importance of different acoustic measures in the acoustic space.

5.4 Growing a forest of decision trees

However, simple decision trees suffer from two main disadvantages. The first is that decision trees can suffer from high variance. In other words, the tree can be very sensitive to small changes in the data on which it was trained. The second disadvantage is that decision trees do not have the same predictive accuracy as other regression or classification models (see Hastie, Tibshirani & Friedman 2009 for a discussion). The following subsections explain two methods that have been proposed to overcome these disadvantages. The first method is called Bagging trees and the second method is called Random Forests. As will be explained in the following sections, these two methods are essentially the same except for one key difference, the num-

ber of predictors that are used. Bagging trees use all predictors in each split, whereas Random Forests use a random subset of predictors in each split.

5.4.1 Bagging trees

One way to overcome the disadvantages of simple decision trees is to make use of a technique called bootstrap aggregating, or bagging (Breiman 1996). This means that instead of growing a single tree with all of the predictors (i.e., variables in your data) in the data like in simple decision trees, we grow many trees on random samples of the data until we reach a given number of trees using the full set of predictors. Once these trees are grown, we then average across the trees to get a more stable prediction of how the regions are split and what predictors are most important in making those splits. This averaging across the trees helps to explain the variance in the data and improve the predictive accuracy of the model. However, this comes at the cost of interpretability.

In decision trees, we usually represent the splits in the data as a tree. When we use bagging, because of the large number of trees that are grown, it is impossible to represent the results in this way. Instead of using a tree, we use variable importance measures to understand which predictors are most important in dividing the data. There are two measures that are commonly used to understand the importance of variables in tree bagging: the residual sum of squares (RSS) for regression trees and

the Gini index, which is a measure of *purity*, for classification. In regression trees, the amount of RSS that decreases due to the splits over a given predictor is recorded and averaged across all the trees. In the classification trees, the total amount that the Gini index decreases for each predictor is recorded and averaged across all trees. The higher the value of the RSS, or Gini index, the more important that predictor is in making the splits in the data. These are then graphed to show the importance of each predictor in the data with the most important predictors at the top of the graph.

The exact number of trees to be grown is not known *a priori*. Instead, the number of trees is determined by the user and is usually determined by the number of trees needed to stabilize the prediction. This is done by comparing multiple models that were built with different numbers of trees and determining which number of trees produces the most stable prediction. This is done by comparing the predictions of the different models and calculating the variance of the predictions across the different models. The model that produces the most stable prediction is the one chosen.

5.4.2 Random Forests

However, subsequent research by Breiman (2001), showed that bagging sometimes would not stabilize and that they would overfit the data. To remedy this problem, Breiman proposed that instead of growing multiple trees that consider all predictors every time, we should only consider a random subset of predictors in each split. This

means that bagging and Random Forests are essentially the exact same thing except for the number of predictors that are considered at each split. This is a very important distinction, as it allows us to grow trees that are more stable and less likely to overfit the data.

Research has shown that generally speaking the number of predictors that must be considered in each split (that is, m_{try}) is usually the square root of the total number of predictors in the data for classification trees and one-third of the total number of predictors for regression trees (Breiman 2001, Sandri & and Zuccolotto 2008, Hastie, Tibshirani & Friedman 2009, Janitza & Hornung 2018, Boehmke & Greenwell 2019, James et al. 2021). This means that if we have 81 predictors in our data, we would only consider ≈ 9 predictors at each split for classification trees ($\sqrt{81} = 9$) and ≈ 27 predictors for regression trees ($\frac{81}{3} = 27$). However, this is not a hard and fast rule, and the number of predictors that are considered at each split can be tuned to improve the performance of the model. This is done by comparing the predictions of the different models and calculating the variance of the predictions across the different models. The model that produces the most stable prediction is the one chosen.

5.4.3 How to interpret the results

The benefit of using Bagging and Random Forests is that they are able to produce more stable predictions than simple decision trees. This is because they are able to

average across the predictions of multiple trees instead of growing a single tree. This allows us to take advantage of the strengths of decision trees while minimizing the weaknesses that are present in classical decision trees.

However, the benefits of using these methods come at the cost of interpretability. In classical decision trees, the results of the analysis are presented as a tree plot, similar to the tree in Figure 5.1. This allows us to see how the different predictors are used to split the data. This is not possible in bagging and random forests due to the large number of trees that are grown and then used to average the predictions. Instead, we used variable importance plots to show how important a variable is across all trees in making the data splits. This is done by calculating a measure of importance for each variable and then plotting the results. In the case of random forests, the most common measure of importance is called the Gini Index, which is a measure of how pure the split is generally. The Gini index is calculated for each variable and then averaged across all trees. The higher the Gini index, the more important that variable is in making the splits in the data. This is similar to how we interpret the results of a classical decision tree, where the most important predictors are at the top of the tree and the least important predictors are at the bottom. In bagging and random forests, we are not concerned with the exact value of the Gini index, but rather with the mean decrease in the Gini index. This is because the Gini index is a measure of how pure the split is and not a measure of how important the variable is in making the splits in

the data. The mean decrease in the Gini index is a measure of how much the variable contributes to the overall purity of the split. For bagging, this is the only measure that is considered.

There is some evidence that the Gini index is not always the best measure for random forests. For example, Strobl et al. (2007) showed that the Gini index can be biased in some cases with random forests. To get around this fact, interpretation of random forests also frequently consult the permutation importance in addition to the Gini Index. The permutation importance measures the change in the model's prediction accuracy when the values of a variable are randomly permuted. This means that the variable is shuffled and the model is re-evaluated. The difference in accuracy between the original model and the permuted model is then used to measure the importance of that variable. The higher the difference, the more important that variable is in splitting the data.

Tagliamonte & Baayen (2012) made use of both the Gini index and the importance of permutation to evaluate the importance of the different sociolinguistic predictors in their analysis. The results of their analysis are shown in Figure 5.2. The Gini index is shown on the left, and the permutation importance is shown on the right. The y-axis shows the different predictors for both plots. We observe in this graph that for the impurity importance (i.e., Gini index), the most important predictors are those that are at the top of the graph. The most important predictor is the one that has the largest

Gini index, in this case the Individual variable. The second most important predictor is the Age variable, etc. The permutation importance is shown on the right and is interpreted in the same way. However, they kept the ordering of the predictors the same across the two plots. The most important predictor according to the permutation importance is the one that has the largest value in this case, the Individual variable.

The second most important predictor is the Proximate1 variable, etc.




Figure 5.2: Variable importance plot from Tagliamonte & Baayen (2012). The plot on the left shows the impurity importance (i.e., Gini index) and the plot on the right shows the permutation importance. The y-axis shows the different predictors for both plots.

Looking at both the impurity and permutation importance, we can get a better understanding of which predictors are the most important in making the splits in the space. For Tagliamonte & Baayen (2012), this meant that the Individual variable was the most important predictor in making the splits in the data. The other predictors

were also important, but we now see differences between the two plots in which are important. In general, the researcher should look at both the impurity importance and the permutation importance to gain an understanding of which predictors are the most important in making the splits in the data. If there is a overlap between the two plots, then we can be more confident that the predictor is important in making the splits in the data. If there is a large difference between the importance of a predictor between the two plots, then we should be cautious about that predictor's importance.

5.5 Random Forests in SLZ

In this section, I will present the results of a Random Forest analysis that was performed on the SLZ data. The goal of this analysis is to understand which acoustic measures are the most important in achieving the separations in the acoustic space of the SLZ. The data collection and analysis methods are similar to those used in the MDS analysis presented in Chapter 4. The main difference is that I will be using a Random Forests analysis instead of an MDS analysis. The results of this analysis will be presented in the following sections.

5.5.1 Methods

5.5.1.1 Participants

This study uses data collected from 10 native speakers of SLZ during the summer of 2022. Participants were recruited from the community of Santiago Laxopa, Oaxaca, Mexico. All participants were native speakers of SLZ. The participants were between 18 and 60 years old and consisted of five males and five females.

5.5.1.2 Recordings

Participants were asked to perform a word list elicitation task consisting of 72 words. These words were selected to elicit the entire range of types of voice quality in SLZ, including modal voice, the two kinds of creaky (i.e., checked and rearticulated), and breathy voice. The words were selected based on previous research conducted as part of the Zapotec Language Project at the University of California, Santa Cruz (*Zapotec Language Project – University of California, Santa Cruz 2022*). Because participants were not literate in SLZ, the word list was prompted for them by asking them “How do you say [word in Spanish]?” by myself and another researcher in Zapotec. Participants were asked to respond with the desired word in the carrier phrase *Shnia’ X chonhe lhas* [ʃn̥ia’ X tʃone ras] “I say X three times.” which was repeated three times. These utterances were recorded in a quiet environment using a Zoom H4n handheld digital recorder. The recordings were saved as 16-bit WAV files with a sampling rate

of 44.1 kHz.

5.5.1.3 Acoustic measuring

The resulting audio files were then processed in Praat to isolate the vowel portion of each word. The onset of the vowel was set to the second glottal pulse after the onset, and the offset of the vowel was set to the last glottal pulse before the decrease in amplitude at the end of the vowel (Garellek 2020). The vowel was then extracted and saved as a separate file for analysis.

These vowels were fed into VoiceSauce (Shue et al. 2011) to generate the acoustic measures for the studies discussed in this dissertation. Because many acoustic measures are based on the fundamental frequency, this measure was calculated using the STRAIGHT algorithm from (Kawahara, Chevigne & Patterson 1998) to estimate the fundamental frequency in millisecond (ms) intervals. Once the fundamental frequency is calculated, VoiceSauce then uses an optimization function to locate the harmonics of the spectrum, finding their amplitudes.

VoiceSauce then uses the Snack Sound toolkit (Sjölander 2004) to find the frequencies and bandwidths of the first four formants, also at millisecond intervals. The amplitudes of the harmonics closest to these formant frequencies are located and treated as the amplitudes of the formants. These formant frequencies and bandwidths are used to correct the harmonic amplitudes for the filtering effects of the vocal tract, using Iseli, Shue & Alwan's 2007 extension of the method employed by Hanson (1997). Each

vowel was measured across ten equal time intervals, resulting in 22890 data points in total. These measures were then z-scored by speaker to reduce the variation between speakers and provide a way to compare the different measures directly on similar scales.

5.5.1.4 Data processing

Data points with an absolute z-score value greater than three were considered outliers and excluded from the analysis. The Mahalanobis distance was calculated in the F1-F2 panel within each vowel category. Each data point with a Mahalanobis distance greater than six was considered an outlier and excluded from the analysis. Using the Mahalanobis distance allows us to compare the data points to the mean of the F1-F2 panel for each vowel category. The larger the Mahalanobis distance is the more deviant the data point is from the mean which in turn means that the data point was improperly tracked. This is comparable to what was done in Seyfarth & Garellek (2018), Chai & Ye (2022), and Garellek & Esposito (2023).

Energy was excluded if it had a zero value and then log-transformed to normalize its right-skewed distribution. Afterward, the resulting log-transformed Energy was z-scored, and any data point with a z-score greater than three was excluded. This outlier removal resulted in 1918 data points being removed.

All data points were then z-scored by speaker to reduce the variation between speakers and provide a way to compare the different measures directly on the same

scale.

Residual H1* for the remaining data points following Chai & Garellek (2022). First, a linear mixed effects model was generated with the z-scored H1* as the response variable and the z-scored energy as the fixed effect. The uncorrelated interaction of the z-scored energy by speaker was treated as random. The energy factor resulting from this linear mixed-effects model was extracted. Finally, the z-scored H1* had the product of the z-scored energy and energy factor subtracted from it to produce the residual H1* measure.

Once these steps were completed, the mean of each combination of phonation and speaker was taken for the fourth to seventh interval of the vowel. This is similar to what Keating et al. (2023) did by taking the middle of the vowel for their analysis. This choice minimizes the effect of the onset and offset of the vowel on the acoustic measures, which are more likely to be affected by the surrounding consonants and should give us the most accurate representation of the vowel quality. Because z-scores were used, this resulted in negative measures, which presents a problem for MDS analyses. To correct for this, I added the absolute value of the minimum z-score to each measure. This results in a dataset that still preserves the relative differences in the scores.

5.5.2 Parameter selection

In order to determine the correct number of trees and the number of predictors to use at each split, I followed the methods outlined in Boehmke & Greenwell (2019) and James et al. (2021) for parameter selection. The data was split into a training set and a test set. The training set was used to train the model and the test set was used to evaluate the performance of the model. The training set consisted of 70% of the data and the test set consisted of 30% of the data. The split was done so that the distributions of the different voice qualities was the same between the training and test sets. The training set was then used to tune the parameters of the model. The parameters that were tuned were the number of trees, the number of predictors to use at each split, the amount to sample from the training set, and whether to sample with replacement. The values for the parameters were chosen based on the results of a grid search.

Using a grid search allowed me to systematically search through the different combinations of parameters to find the best combination for the model. Another reason was that this allowed me to determine if Bagging (i.e., using all of the predictors at each split) or Random Forests (i.e., using a random subset of the predictors at each split) was the best model for the data. The model whose parameters resulted in the most accurate model was chosen as the final model.

Figure 5.3, shows the results of the grid search in relation to the number of trees and the number of predictors to use at each split. The x-axis shows the number of

trees that were used in the model and the y-axis shows the percentage of incorrectly classified out-of-bag tokens. The lower the out-of-bag error percentage, the better the model is at predicting unseen data. The colored lines show the different values of m_{try} that were used in the model. When $m_{try} = 22$, the model uses all the predictors in each split and is therefore a bagging model. The other potential values for m_{try} are the values corresponding to 5%, 15%, 25%, and 40% of the total number of predictors. Furthermore, the default values of $m_{try} = \sqrt{22}$ and $m_{try} = \frac{22}{3}$ for classification and regression trees were considered, these values were recommended by Boehmke & Greenwell (2019).

Figure 5.3 shows that the model with $m_{try} = 22$ (i.e., bagging) is not the best model for the data. This is because it has a very high percentage of out-of-bag error compared to the other models. The model that performed the best was the model that trained on 300 trees and whose $m_{try} = 5$ had the lowest percentages of out-of-bag errors. Furthermore, it was found that sampling only 80% of the replacement data produced the best model.¹

5.6 Results

The results of the Random Forest analysis are shown in Figure 5.4. The plot on the left shows the impurity importance which uses the mean decrease in the Gini index

¹The full grid search results can be found online at my website: mlbrinkerhoff.me.




Figure 5.3: Plot showing the percent of inaccurately classified phonation types as a function of the number of trees ran. The different colored lines indicate the different m_{try} values.

to measure the importance of each predictor. The Gini index is a measure of how pure the split is and is calculated for each variable and then averaged across all trees. The higher the Gini index, the more important that variable is in making the splits in the data. The plot on the right shows the permutation importance, which measures the change in the model's prediction accuracy when the values of a variable are randomly permuted (i.e., the variable is shuffled and the model is reevaluated). The difference in accuracy between the original model and the permuted model is then used to measure the importance of that variable. The higher the difference, the more important

that variable is in splitting the data. The y-axis shows all 21 different predictors for both plots. However, the predictors in each plot are ranked in descending order of importance.




Figure 5.4: Variable importance plots showing the impurity importance and permutation importance of each acoustic measure.

From Figure 5.4, we can see that the two most important predictors for classifying the different phonations are: (i) duration and (ii) HNR < 1500 Hz. The reason for this is that both of these measures appear as the two highest predictors in terms of impurity and permutation. Each of the predictors suggested by the model will be discussed in more detail below.

The rest of the variables require some discussion to determine their importance. This is because the next several predictors in both plots are not consistent. In the impurity importance plot, we see that Strength of Excitation and residual H1* are the next most important measures. However, their importance decreases when we consider permutation. Instead of being the third and fourth most important impurity predictors, they are now the ninth and eleventh most important permutation predictors. Because both measures are sifted only slightly and are in the upper half of both plots, we can assume that these acoustic measures still play an important role to some extent.

Furthermore, we see that A1* is relatively consistent between the two plots. In the impurity plot on the left it is the sixth most important predictor and in the permutation plot on the right it is the fifth most important predictor. This suggests that A1* plays an important role in classifying the different phonations. The same reasoning holds for H1*–A1*, which is the seventh most important predictor in both plots.

5.7 Discussion of the results

This section will discuss the results of the Random Forest analysis in two parts. The first part will discuss the results of the Random Forest analysis in relation to the MDS analysis presented in Chapter 4. The second part will discuss certain acoustic measures that were shared by the analyses and those that were uniquely chosen by

the Random Forest analysis.

5.7.1 Comparing the MDS and bagging trees

There was a large amount of overlap between which measures were correlated with the dimensions of the MDS analysis and the important predictors found by the Random Forest analysis. The correlations for the MDS analysis were found in Chapter 4 and are shown in Table 4.1. They are repeated here as Table 5.1 for convenience.

The rest of this section will discuss each of the measures where there was overlap between which measures showed a correlation to a MDS dimension and the Random Forest results. Furthermore, I will discuss duration and A1* due to how important the Random Forest model showed these measures. These measures are: (i) duration, (ii) A1*, (iii) H1* – A1*, (iv) residual H1*, (v) HNR < 1500 Hz, and (vi) Strength of Excitation. Each of these measures will be discussed in the following subsections in the order described above.

5.7.2 Importance of duration

In the Random Forest analysis, duration was found to be one of the most important predictors in classifying the different phonations. This importance is reflected in the fact that it is the most important predictor in the impurity importance plot and the second most important predictor in the permutation importance plot.

Table 5.1: Correlations for each acoustic measure to the four dimensions (NMDS1, NMDS2, NMDS3, NMDS4). The four largest correlations in each dimension are bolded.

Acoustic Measure	NMDS1	NMDS2	NMDS3	NMDS4
H1*–H2*	-0.221	-0.339	0.031	0.314
H2*–H4	-0.437	0.239	-0.689	-0.364
H1*–A1*	-0.828	0.048	-0.459	0.044
H1*–A2*	-0.855	-0.067	-0.343	0.114
H1*–A3*	-0.809	-0.218	-0.297	0.126
H4*–H2k*	-0.452	-0.598	0.294	0.366
H2k*–H5k*	0.152	0.023	0.101	0.057
residual H1*	-0.290	-0.443	-0.722	0.084
H2*	-0.157	-0.555	-0.679	0.114
H4*	0.295	-0.778	0.078	0.479
A1*	0.756	-0.549	0.092	0.124
A2*	0.779	-0.476	-0.103	0.086
A3*	0.735	-0.416	-0.211	0.093
CPP	-0.590	-0.606	0.209	-0.179
HNR < 500 Hz	-0.513	-0.792	0.152	-0.202
HNR < 1500 Hz	-0.275	-0.799	0.323	-0.290
HNR < 2500 Hz	-0.327	-0.714	0.391	-0.348
HNR < 3500 Hz	-0.446	-0.644	0.393	-0.356
Strength of Excitation	-0.013	-0.741	-0.238	0.145
SHR	0.144	-0.176	0.122	-0.597
Energy	-0.080	-0.793	-0.015	0.341
Duration	-0.622	0.539	0.257	0.030

Duration frequently plays an important role in Zapotec phonation contrasts. Many descriptions of Zapotec languages have shown that duration is often an important correlate for checked vowels (Ariza-García 2018). This has also been reported by Chai (2025) for Yateé Zapotec, a variety of Zapotec spoken in the Sierra Norte region of Oaxaca, Mexico, and a close relative of SLZ. Chai (2025) found that duration was the most

important percept for distinguishing between Yateé Zapotec's checked and rearticulated vowels. Chai also found that the longer the duration of the vowel, the more likely the speakers of Yateé Zapotec were to classify the vowel as rearticulated. Similar results for checked vowels have been reported for other Zapotec varieties (Arellanes Arellanes 2009, 2010, Chávez-Péón 2010, López Nicolás 2016, Merrill 2008), other members of the Oto-Manguean family (e.g., Campbell 2014), and crosslinguistically (Gao & Kuang 2022, Chai & Ye 2022).

In their study on contextual enhancement effects on phonation contrasts in San Pablo Macuitianguis Zapotec, Barzilai & Riestenberg (2021) found that duration played a role in distinguishing between the different types of phonation, regardless of the phrasal context. They found that modal vowels were the longest and checked vowels the shortest. In regards to rearticulated vowels, they claim that they do not function as a separate phonation type but rather are a sequence of a checked vowel followed by a modal vowel. Despite this reanalysis, we would then expect the duration of the rearticulated vowel to be longer, given that they are a sequence of two vowels.

In SLZ, the duration distribution between the different phonation types is shown in Figure 5.5. The y-axis shows the duration of each of the different phonation types in z-scores, again, z-scores were used to minimize individual speaker variation. The x-axis shows each of the four different phonations. The plot shows that the breathy voice has the longest duration and the rearticulated voice has the longest duration,

followed by the checked voice. The shortest duration is found in the modal voice.




Figure 5.5: Plot showing the distribution of duration across the different voice qualities in SLZ.

These results are somewhat surprising because of the generalization that checked vowels are vowels that are abruptly stopped at the end of the vowel. This is not the case in SLZ. The checked vowels in SLZ are indeed the shortest of the three non-modal phonations. However, they are not the shortest when modal is included. This is likely due to the fact that checked vowels are often limited to final open word syllables in SLZ, and there is a general trend in Zapotec phonology for these syllables to be longer or minimally bimoraic (e.g., Chávez-Péón 2010, Nellis & Hollenbach 1980, Uchihara & Pérez Báez 2016).

When we consider the behavior of nonmodal phonation cross-linguistically, we see that the behavior of duration is consistent with what is generally found. For example, Gordon & Ladefoged (2001) and Esposito & Khan (2020) both report that non-modal voice qualities are typically longer than modal voice. One of the reasons for this is that by lengthening the vowel, the speaker is able to increase the amount of time that the glottal source is able to produce the desired voice quality, which increases the likelihood that the listener will be able to perceive the desired voice quality.

The behavior of duration warrants further investigation in SLZ. If checked vowels are indeed longer than modal vowels, then this would suggest that the checked vowels in SLZ are not the same as the checked vowels in other Zapotec languages. Furthermore, vowels have been claimed to undergo a process of vowel lengthening in Zapotec languages depending on the type of syllable it is in and whether the coda is a fortis or lenis consonant (Nellis & Hollenbach 1980, Uchihara & Pérez Báez 2016).

If the vowel is in an open syllable or a closed syllable with a lenis coda, then the vowel is lengthened. If the vowel is in a closed syllable with a fortis coda, then the vowel is not lengthened. If this behavior is true for SLZ, this means that the duration of the vowel depends not only on the type of phonation but also on the type of syllable in which it is placed. This requires further investigation to determine if the duration is affected by the phonation and type of syllable.

5.7.3 Importance of A1*

The measure that was found to be the most important for classifying the SLZ phonation contrasts was A1*. This measure captures the amplitude of the harmonic closest to the first formant. This measure is typically not found in voice quality research except as a way to normalize the amplitude of the first harmonic (i.e., H1*). This goes back to Fischer-Jørgensen (1968) who used this as one of the ways to correct for the high-pass filtering, in addition to the widely used H1*–H2* measure. However, A1* as a standalone measure is not typically used in voice quality research. Therefore, what we are to make of this measure's importance in SLZ is not clear as is its behavior in regards to the different phonations.

When we look at how A1* is distributed across the different voice qualities in SLZ, as seen in Figure 5.6, we see that the modal vowels are found at the top of the chart with the other phonation contrasts located lower on the chart. The phonation that is found at the bottom of the chart is a breathy voice.

The pattern of behavior for A1* is very similar to what is found in descriptions of the behavior of the frequency of F1 in breathy voice contexts. For example, differences in the frequency of F1 are typically used to distinguish register differences in Southeast Asian languages (Brunelle 2005, 2012, Brunelle & Kirby 2016).

As described in Brunelle & Kirby (2016), many of the languages of Southeast Asia have what is called register. The linguistic term register is defined as “the redundant




Figure 5.6: Plot showing the distribution of $A1^*$ across the different voice qualities in SLZ.

use of pitch, voice quality, vowel quality, and durational differences to distinguish (typically two) contrastive categories" (Brunelle & Kirby 2016: p. 193), and was first used by Henderson (1952) to describe the categorical contrasts found in Khmer. The characteristics that define the higher and lower registers are found in Table 5.2.

From this table, the lower register is what interests us here. The lower register is associated with breathy voice and a lowered first formant. There is evidence that breathy voice frequently has a lower first formant than modal voice in paralinguistic settings for English (Lotto, Holt & Kluender 1997). However, these studies do not discuss the amplitude of the first formant but rather the frequency of the first formant.

Table 5.2: Possible phonetic correlates of register. From Brunelle & Kirby (2016).

Higher Register	Lower Register
Higher pitch	Lower pitch
Tense/Modal voice	Lax/Breathy voice
Monophthongs/shorter vowels	Diphthongs/longer vowels
Raised F1/lower vowels/[+ATR]	Lowered F1/higher vowels/[-ATR]
Plain stops/shorter VOT	Aspirated stops/longer VOT

Another comparison can be found in the research on nasality, where this measure is discussed extensively either alone or in association with the nasal pole (e.g., Chen 1997, Delvaux 2009, Macmillan et al. 1999, Pruthi & Espy-Wilson 2004, Schwartz 1968, Stevens 2000, Styler 2015, 2017). These studies discuss how in nasalized contexts the amplitude of the first formant is typically found to be lower than in oral contexts, again similar to what we see in Figure 5.6.

There is a large body of research that discusses that nasality is closely associated with breathy voice or glottal consonants, in a phenomenon called *rhinoglottophilia* (Matisoff 1975, Ohala 1975, Ohala & Ohala 1993, Bennett 2016). In Blevins & Garrett (1993) and Matisoff (1975), this association is attributed to the acoustic and perceptual similarities between nasalization and breathy voice. In one study, Garellek, Ritchart & Kuang (2016) showed that nasalization in three different Yi languages was associated with a slender voice. The authors suggested that breathy voice during nasalization can arise from misperception or as a type of phonetic enhancement.

In terms of what is going on in SLZ, it is not clear if the lowering of A1* for breathy

voice can be attributed to the same observations as discussed above. I suggest that there are three different possibilities to explain what is going on in SLZ. First; because SLZ does not have phonemic nasalization, it is possible that speakers of SLZ are using nasalization as a way to phonetically enhance the contrast of breathy voice. Essentially, the reverse of what was reported by Garellek, Ritchart & Kuang (2016). This possibility could be tested acoustically by performing an experiment to detect nasal airflow during breathy vowels. The second possibility is that the same measures that work for detecting nasality can also be used to detect breathy voice. This second possibility could be easily tested by examining whether A1* and the other measures for nasality work in other breathy voice contexts cross-linguistically. The third possibility is that the lowering of A1* is a result of subglottal resonances, which is much harder to test than the other two possibilities.

5.7.4 Importance of H1*–A1*

The Random Forest analysis found H1*–A1* to be one of the most important predictors in classifying the different phonations. This is because of its high ranking in both the impurity and permutation importance plots. In the impurity plot, it is the seventh most important predictor, and in the permutation plot, it is the sixth most important predictor.

The acoustic measure H1*–A1* is a measure of the difference between the am-

plitude of the first harmonic and the amplitude of the harmonic closest to the first formant. This acoustic measure falls into the category of spectral slope measures. As discussed in Chapter 3, these measures capture the differences in the amplitude of the harmonics of the spectrum. The higher the values of the spectral slopes, the more breathy the phonation is (Fischer-Jørgensen 1968). The lower the spectral slope values, the more creaky the phonation is.

The distribution of $H1^* - A1^*$ across the different phonation types is shown in Figure 5.7. The y-axis is the z-scored $H1^* - A1^*$ values and the x-axis shows the ten equally spaced intervals throughout the duration of the vowel.

The plot shows that breathy voice has the highest $H1^* - A1^*$ score, which is consistent with what we would expect from a spectral slope measure. Things are more complicated when we look at the other non-modal phonations. It is expected that checked voice and rearticulated voice have a lower spectral slope than modal voice because of their association with creaky voice. However, this is not the case in Figure 5.7.

The plot shows that the checked voice has a spectral slope that is identical to that of the modal voice. Additionally, we see that the rearticulated voice has a spectral slope that is higher than the modal voice except at measurement intervals three and four, where it is identical to the modal voice. This behavior for a rearticulated voice is not too surprising, especially at the end of the vowel.




Figure 5.7: Plot showing the distribution of $H1^* - A1^*$ across the different voice qualities in SLZ. Each point represents the mean of the ten equally spaced intervals across the duration of the vowel and the error bars represent a 95% confidence interval.

One reason for this is that when listening to the rearticulated vowels, the portion of the vowel after the glottal occlusion sounds somewhat breathy. One of the possible explanations for this is that in order to produce the glottal occlusion, the speaker is tightly constricting the vocal folds, which causes either a glottal stop or creaky voice. The speaker then relaxes the vocal folds in order to produce the modal voice which stereotypically occurs after this glottal occlusion. However, one way for the speaker to quickly relax the vocal folds is to open them as much as possible to reach the ideal position for the modal voice. This general behavior of opening the vocal folds

to produce a modal voice is similar to what is found in a breathy voice. This could be a possible explanation for why the rearticulated voice has a higher $H1^* - A1^*$ score than modal voice.

5.7.5 Importance of Residual $H1^*$

Another spectral slope measure that was found to be important in the Random Forest and MDS analyses was residual $H1^*$. This measure is similar to $H1^* - A1^*$ in that it captures the differences in the amplitude of the harmonic of the spectrum. However, residual $H1^*$ is a measure that is calculated after removing the effect of energy on $H1^*$.

As residual $H1^*$ is a type of spectral slope measure, it is expected to show similar behavior to other spectral slope measures. That is, a breathy voice should be associated with a higher score than a modal voice. Since the two types of laryngealization are closely associated with creaky voice, they should be associated with a lower score than modal voice. Additionally, there is a temporal difference between the two types of laryngealization, it is expected that checked voice will have a lower score toward the end of the vowel and rearticulated voice will have a lower score in the middle of the vowel.

In Figure 5.8, we see that the predictions are correct. Breathy voice is associated with a higher score than the modal voice. The two types of laryngealization are associated with a lower score than the modal voice.




Figure 5.8: Plot showing the distribution of residual $H1^*$ across the different voice qualities in SLZ.

A further analysis of the results of this measure using generalized additive (mixed) models (GAM(M)s; Hastie & Tibshirani 1986, Wood 2017, Sóskuthy 2017, Wieling 2018) shows that the temporal behavior of the two types of laryngealization is also correct. Checked voice is associated with a lower score toward the end of the vowel, and rearticulated voice is associated with a lower score in the middle of the vowel. Additionally, the results of the GAMM show that for the first three time points there is no significant difference between the breathy and modal voice. The suggestion is that the breathy voice is more closely associated with the latter part of the vowel than with the beginning. The full results of the GAMM for residual $H1^*$ are discussed in

5.7.6 Importance of the HNR <1500 Hz

The acoustic measure HNR < 1500 Hz is a harmonics-to-noise ratio measure that is calculated over the frequency range from 0 Hz to 1500 Hz and is a measure of the amount of noise in the signal, or in other words, it is a measure of periodicity. In these measures, modal voice is associated with a higher score than nonmodal phonations. This is because modal voice is associated with a higher degree of periodicity than non-modal phonations because a-periodicity is a defining feature of nonmodal phonations (e.g., Hillenbrand & Houde 1996, Blankenship 1997, Kent & Ball 1999).

As seen in Figure 5.9, this is exactly what we see in SLZ. Modal voice is associated with a higher score than non-modal phonations. Among nonmodal phonations, breathy and rearticulated voice have a higher HNR < 1500 Hz than checked voice.

The fact that breathy and rearticulated voice has a higher HNR < 1500 Hz than checked voice is not surprising. As discussed in Chapter 2, rearticulated and breathy voices are associated with more modal-like qualities than checked voice. In most instances of rearticulated voice, the vowel is produced with modal voice except for a small portion of the middle of the vowel where the vowel is produced with creaky voice or a glottal occlusion. Breathy voice typically appears very periodic but with a high degree of noise in the signal. Checked voice on the other hand, is associated




Figure 5.9: Plot showing the distribution of HNR < 1500 Hz across the different voice qualities in SLZ.

with a high degree of aperiodicity in the signal from the creaky voice that is produced at the end of the vowel. This is why checked voice has the lowest HNR < 1500 Hz of nonmodal phonations.

This measure will be very important in Chapter 6, where I will use it to model laryngeal complexity.

5.7.7 Importance of Strength of Excitation

Strength of Excitation (SoE) is a measure that is defined as “the instant of significant excitation of the vocal-tract system during production of speech [and rep-

resents] the relative amplitude of impulse-like excitation” (Mittal, Yegnanarayana & Bhaskararao 2014: p. 1934). This measure is typically associated with the amplitude of the voicing, or in other words, the degree of voicing during production (Murty & Yegnanarayana 2008, Mittal, Yegnanarayana & Bhaskararao 2014). This measure has been shown to be an effective measure for showing the effects of laryngealization on the amplitude of voicing (Garellek et al. 2021) and has been shown to be effective in distinguishing the different phonations in the Oto-Manguean languages (Chai, Fernández & Mendez 2023, Weller et al. 2023a,b, 2024).

Following Garellek et al. (2021), it is expected that the modal voice will have the highest SoE score. It is also expected that, because the breathy, rearticulated, and checked voice will have a lower SoE score. The reason for this expectation is that according to Garellek et al. (2021), there is a strong tendency for all types of laryngealization to have a dampening effect on voicing. This is evidenced by the lower SoE scores for these phonations. The score for SoE ranges from 0 to 1, where 0 is no voicing and 1 is full voicing.

In Figure 5.10, we observe that the modal voice has the highest SoE score. The three non-modal phonations are all lower. We can ignore the SoE values at measurement numbers one and ten because they represent the beginning and end of the vowel and are more likely to be affected by the coarticulatory effects of the surrounding consonants. However, we can see that breathy voice has a higher SoE score than

checked voice and rearticulated voice. Additionally, we see that between checked and rearticulated voice, we see that checked voice has a higher SoE score in the first half of the vowel and rearticulated voice has a higher SoE score in the second half of the vowel. This is consistent with what we would expect from the behavior of the phasing difference between checked and rearticulated vowels.




Figure 5.10: Plot showing the distribution of Strength of Excitation across the different voice qualities in SLZ.

One thing to note about the SoE measures is that even though nonmodal phonations are associated with a lower SoE score than modal voice, the values for these

measures are still extremely high. This likely indicates that even though laryngealization is present in these phonations, the speakers are only weakly laryngealizing the vowel. This is consistent with the claims made by Silverman (1997a,b) that laryngeally complex languages that do not show a phasing relationship between modal and non-modal phonation must weakly produce non-modal phonation. The claims about laryngeal complexity will be discussed in more detail in Chapter 6.

5.8 Conclusion

In summary, we find that the results of the Random Forest and the MDS analyses showed a lot of overlap between which measures were correlated and/or important for classifying SLZ’s four phonation types. In both analyses, A1*, H1*–A1*, residual H1*, HNR < 1500 Hz, and SoE were found to be important acoustic measures. We saw that each of these measures was able to capture some aspect of the phonation types. It also shows that the spectral slope measures and harmonics-to-noise ratio measures are generally the most important measures for classifying the different phonation types. This is consistent with previous work on the acoustics of phonation (e.g., Garellek 2019). We also saw that SoE plays an important role in classifying the different phonation types. This contributes to the growing body of literature that shows that SoE is an important measure for classifying the different phonation types (Chai, Fernández & Mendez 2023, Garellek et al. 2021, Weller et al. 2023a,b, 2024).

Interestingly, the results of the Random Forest analysis showed that duration was the most important predictor in classifying the different phonations. This is consistent with what has been found in other Zapotec languages (e.g., Barzilai & Riestenberg 2021, Chai 2025, Chávez-Péón 2010). However, when we plotted the duration, it showed that modal vowels had the shortest duration instead of checked vowels having the shortest duration. The effect of duration needs to be further investigated in SLZ, especially with respect to differences in syllable type and the claimed effects of vowel lengthening in the various Zapotec languages (e.g., Chávez-Péón 2010, Merrill 2008, Nellis & Hollenbach 1980, Pickett, Villalobos & Marlett 2010, Uchihara & Pérez Báez 2016).

Chapter 6

Testing laryngeal complexity in SLZ

6.1 Introduction

This chapter investigates the role that laryngeal complexity plays in Santiago Laxopa Zapotec. Laryngeal complexity is used to describe the contrastive use of tone and phonation in many languages, especially in the Oto-Manguean languages (BlankenSHIP 1997, 2002, Silverman 1997a,b). As mentioned in Chapter 2, SLZ is a laryngeally complex language, because of its contrastive use of both tone and phonation. This means that SLZ can be used to test the predictions of laryngeal complexity, with respect to the phasing and recoverability of tone and phonation.

The goal of this chapter is to show that laryngeal complexity is present in SLZ as evidenced by the phasing of nonmodal phonation in several acoustic measures. This

is done by looking at the acoustic properties of SLZ phonation using a combination of Strength of Excitation (SoE), Harmonic-to-Noise Ratio (HNR) < 1500 Hz and f_0 perturbations. It will be shown that there is a clear phasing between modal and nonmodal phonation in vowels that are described as breathy, checked, and rearticulated.

These measures were selected based on the results of the MDS analysis in Chapter 4 and the Random Forest analysis in Chapter 5. Both MDS and Random Forest analyses showed that SoE and HNR < 1500 Hz were the most important measures to distinguish between the different types of phonation. SoE has previously been used to show that there is a clear phasing between modal and rearticulated vowels in San Sebastián del Monte Mixtec (Weller et al. 2023a,b, 2024).

HNR < 1500 Hz has not been used to date to show phasing between modal and nonmodal phonation. As a harmonics-to-noise ratio measure, HNR < 1500 Hz is a measure of the noise in the signal. This means that it can be used to determine if there is aperiodicity in the signal, which is one of the defining characteristics of nonmodal phonation (Ladefoged & Maddieson 1996). This means that HNR < 1500 Hz can be used to determine if there is a clear phasing between modal and nonmodal phonation in these vowels.

Additionally, f_0 perturbation has been used previously to show that there is phasing between modal and nonmodal phonation in several Oto-Manguean languages (Garellek & Keating 2011, DiCanio 2012a, Kelterer & Schuppler 2020).

The remainder of this chapter will be organized as follows. First, in Section 6.2, I will provide a brief overview of laryngeal complexity and phasing and recoverability of tone and phonation in Section 6.2. In Section 6.3 I will discuss previous analyses of laryngeal complexity. In Section 6.4 I will discuss the methods used to analyze laryngeal complexity in SLZ. In Section 6.5 I will present the results of the analysis. Finally, in Section 6.6 I will discuss the implications of these results for our understanding of laryngeal complexity in SLZ.

6.2 What is Laryngeal Complexity?

Laryngeal complexity is defined as the contrastive use of tone and phonation within the same syllabic nucleus (Blankenship 1997, 2002, Silverman 1997a,b). This use of contrastive tone and phonation is one of the defining characteristics of the Oto-Manguean languages (Silverman 1997a). However, it is not limited to just these languages. It has also been used to describe the behavior of tone or pitch in languages outside of the Oto-Manguean languages; such as the Tibeto-Burman languages of Mpi and Tamang (Silverman 1997a,b), the Mayan language Yucatec Mayan (Frazier 2013), and to describe the behavior of coarticulatory pitch and phonation in the Germanic language Danish (Frazier 2013, Peña 2022, 2024).

According to Silverman (1997a,b), even though tone and phonation are apparently allowed to cooccur in the same syllabic nucleus, they are not allowed to be realized

at the same time. Silverman argues that this is because they are in competition with each other over the same articulatory gestures and perceptual resources. This means that tone and phonation must be performed in a temporally ordered fashion. This is what Silverman refers to as phasing. Phasing is the idea that the two components of laryngeal complexity, tone and phonation, are temporally ordered with respect to one another. This means that one portion of the vowel is realized with modal phonation with the acoustic correlates of tone and the other portion of the vowel is realized with non-modal phonation with the acoustic correlates of the said phonation. This phasing is also closely linked to the second main concept of laryngeal complexity, recoverability. Recoverability is the idea that the listener must be able to recover the underlying phonation and tone of the signal. These two concepts will be discussed in more detail in Section 6.2.1.

6.2.1 Phasing and recoverability

According to Silverman (1997a,b), one of the defining aspects of laryngeal complexity is the concept of phasing and recoverability. Under this idea, in laryngeally complex vowels the phonation and tone are phased with respect to one another in a way that lends itself to a listener's ability to recover the underlying phonation and tone. In practical terms this means that laryngeally complex vowels are composed of two components: a modal voice portion of the vowel where tone is realized, and a

nonmodal voice portion of the vowel where phonation is realized. For the researcher, that means that there are two distinct portions of the vowel that can be analyzed separately and that need to be analyzed temporally rather than spectrally (Silverman 1997a: p. 237).

For example, in the Oto-Manguean language Jalapa Mazatec, breathiness or creakiness is realized only on the first portion of the vowel, either as a full laryngeal consonant or as a laryngeal feature on the vowel (Silverman 1997a: p. 238). The second portion of the vowel is realized as a modal voice vowel with one of the three tones belonging to the tonemes of the language. This means that the breathiness or creakiness is phased with respect to the tone.

Silverman argues that there are three principles that help explain why laryngeal complexity should be temporally ordered or phased: (i) sufficient acoustic distance, (ii) sufficient articulatory compatibility, and (iii) optimal auditory salience.

6.2.1.1 Sufficient acoustic distance

Silverman (1997a) argues that sufficient acoustic distance is necessary for the recovery of phonation and tone. As Silverman explained, listeners do not rely on the fundamental frequency alone to perceive pitch. Instead, listeners use the harmonic spacing and the pulse period in the signal to perceive the pitch (Ritsma 1967, Remez & Rubin 1993). For modal phonation, this means that the harmonic spacing and pulse periods are present and encode a salient pitch value. However, during non-modal

phonation, the harmonic spacing and pulse periods are often obscured or not present.

For breathy voice, this means that there is a general weakening of the harmonic structure that makes it difficult for the listener to recover the pitch (Silverman 1997b). Creaky voice on the other hand had obscured the pulse periods due to its aperiodic and unstable glottal vibration (Ladefoged & Maddieson 1996). This is what was observed in Mazatec where the harmonic structure is gone and the pulses are indiscernible (Kirk, Ladefoged & Ladefoged 1993). Furthermore, pitch perception is rendered indiscernible when pulse periods are varied by 10% or more (Rosenberg 1966).

These observations lead Silverman (1997a) to conclude that if a period glottal wave is either obscured (as with breathy voice) or not present (as with creaky voice), the acoustic signal cannot encode a salient pitch value. This means that the phonation and tone must be phased with respect to each other to allow the listener to recover the underlying phonation and tone. I will return to this point in Section 6.6.

6.2.1.2 Sufficient articulatory compatibility

Another important point for Silverman's theory about laryngeal complexity has to deal with the articulatory compatibility of the phonation and tone. One of the guiding ideas behind this principle is that there is a principle of least effort in biological motor systems such as speech production (Lindblom 1983). According to Lindblom (1983), speech gestures can be thought of as distinct motor goals in our speech production system. These goals are achieved by the speaker through coordination of the articu-

lators with the least amount of effort. This means that the gestures are coordinated in such a way that they are compatible with one another. This manifests itself through either sequencing or coarticulation of the gestures.

A good example of this comes from nasalization. In nasal contexts, the velum is lowered to allow air to pass through the nasal cavity, creating a nasal sound. This velum-lowering gesture is compatible with the gestures needed in the oral cavity to produce different vowel qualities. This is what leads to the production of nasal vowels in languages like French or Portuguese. Additionally, this lowering also occurs in languages that do not have contrastive nasal vowels, such as English, where the velum is lowered in anticipation of a nasal consonant. This lowering of the velum is compatible with the gestures needed to produce the vowels in the oral cavity (e.g., Ohala 1975, Chen 1997, Styler 2015).

According to Silverman's theory of laryngeal complexity, this idea of articulatory compatibility is also driving the need to phase phonation and tone. In the case of laryngeal complexity, this is because it is assumed that both tone and phonation are produced by the larynx, more specifically the vocal folds and the glottis. This comes from early work on phonation and tone. For tone, Ohala (1978) showed that pitch was controlled primarily by tensing or laxing of the vocal folds, which changed the rate at which the vocal folds vibrate. For phonation, it was similarly shown that the amount of vocal folds were held open or closed determined the type of phonation that

was produced (Ladefoged & Maddieson 1996). This aspect of phonation was shown in

Figure 4.8 and here is repeated as Figure 6.1.




Figure 6.1: A diagram showing the relationship between breathy, modal, and creaky phonation types. Based on Gordon & Ladefoged (2001).

For Silverman's 1997 theory of laryngeal complexity, the articulatory mechanisms for tone and phonation are exactly the same, which leads to a need to phase the two in order to optimally make use of the same articulatory gestures. However, there is a growing body of literature that shows that tone and phonation are much more complex and rely on the entire larynx, not just the vocal folds (see Esling et al. 2019 for a detailed discussion of this). This matter will be discussed again in Section 6.6.

6.2.1.3 Optimal auditory salience

The last important point for Silverman's theory of laryngeal complexity is the idea of optimal auditory salience. Research into the behavior of nerve responses to acoustic signals has shown that the auditory system is sometimes more and other times less recoverable depending on the signal, which led Bladon (1986) to propose two major principles for auditory phonetics: (i) the on/off response asymmetry and (ii) short-term adaptation.

The on/off response asymmetry principle has origins in the work of Tyler et al. (1982). This principle states that “spectral changes whose response in the auditory nerve is predominantly an onset of firing are much more perceptually salient than those producing an offset. (Silverman 1997a: p. 249)”. This means that changes in the signal that cause an increase in the firing rate in the auditory nerve are easier to perceive when this firing starts compared to when it ends.

The second principle, short-term adaptation, states that “after a rapid onset of auditory nerve discharge at a particular frequency, there is a decay to a moderate level of discharge, even though the same speech sound is continuing to be produced (Silverman 1997a: p. 250)” and is based on the work of Delgutte (1982). This means that the auditory nerve is less sensitive to changes in the signal after a rapid onset of firing, which in turn means that the auditory nerve is less able to recover the underlying phonation and tone when the signal is not changing.

For Silverman, these two principles are important because they help provide an acoustic explanation for why laryngeal complexity should be phased. By phasing tone and phonation, the listener is able to recover the signal more easily. This is because when there is a change in the signal, the auditory nerve has the greatest chance of recovering the signal. In regards to tone and phonation, this means that when there is a change from tone’s modal phonation to nonmodal phonation or from nonmodal phonation to tone’s modal phonation, the listener is able to perceive the difference bet-

ter than if the two were produced simultaneously. Furthermore, these two principles suggest that some phasing relationships are more perceptually salient than others. Based on these two principles, things are easier to perceive if they are produced first or if there is a large change in the signal.

To illustrate this point graphically, Silverman provides a series of figures that show the relationship between the articulatory, acoustic, and perceptual components of laryngeal complexity. These figures are shown in Figures 6.2, 6.3, and 6.4.

Figure 6.2 shows the characteristics of the sequence of [ha˥], where breathiness is first produced followed by a modal vowel with a high tone. The figure shows that when the breathiness is first produced, the auditory nerve response is high followed by a sharp decline throughout the rest of the breathiness. As the energy increases when the modal vowel with high tone is produced, there is a sharp increase in the auditory nerve response which is followed by a gradual decline. This means that it is easier for the listener to perceive the transition and to recover the underlying phonation and tone.

For Figure 6.3, the sequence is reversed. In this case, the high-tone modal vowel is produced first followed by breathiness. The figure shows that when the high tone modal vowel is produced first, the auditory nerve response is at its highest, followed by a sharp decline through the rest of the modal vowel portion. Because there is no energy increase associated with breathiness, there is a continual decrease in the




Figure 6.2: Schematic representation of the characteristics of [ha˥] sequences. Adaptation of figure from Silverman (1997a).

auditory nerve response. This means that it is not as easy as prevocalic for the listener to perceive the transitions and recover the underlying phonation and tone.




Figure 6.3: Schematic representation of the characteristics of [a˥h] sequences. Adaptation of figure from Silverman (1997a).

With regard to the third type of sequence, [aha˥], the figure shows that the auditory nerve response is at its highest when the modal vowel is produced first. This is

followed by a decline similar to what we see in Figure 6.3. However, because there is a high-tone modal vowel, we see a sharp increase in auditory nerve response. Again, this is not as optimal as having the breathiness produced first.




Figure 6.4: Schematic representation of the characteristics of [aha̯] sequences. Adaptation of figure from Silverman (1997a).

In summary, Silverman claims that due to this auditory response asymmetry, we see that there is a difference in the auditory nerve response depending on the sequencing of the phonation and tone. For Silverman, this implies that there is an implicational hierarchy between the different types of sequence found in laryngeal complexity. This implicational hierarchy will be discussed in more detail in Section 6.2.2.

6.2.2 Implicational hierarchy of laryngealization

Another aspect of Silverman's 1997 laryngeal complexity theory is that there is an implicational hierarchy in the phasing and ordering of phonation and tone. This hier-

archy is based on the way laryngealization appears in three Oto-Manguean languages. In this implicational hierarchy, laryngealization can only appear in three ways: prevocalic, postvocalic, or interrupted. In the prevocalic case, the laryngealization appears before the vowel. In the post-vocalic case, the laryngealization appears after the vowel. In the interrupted case, laryngealization interrupts a vowel and appears in the middle.

According to Silverman (1997a), if a language has interrupted laryngealization, it must also have post-vocalic laryngealization. If a language has postvocalic laryngealization, it must also have prevocalic laryngealization. In support of his claims, Silverman (1997a) provides data from three Oto-Manguean languages: Jalapa Mazatec, Comaltepec Chinantec, and Copala Trique. These languages are shown in Table 6.1.

Table 6.1: Implicational hierarchy of laryngeal complexity. The symbols h and ? represent laryngealization. The symbol V represents where the modal vowel is located in relation to the laryngealization. Modified from Silverman (1997a).

Language	Prevocalic	Postvocalic	Interrupted
Jalapa Mazatec	hV[], ?V[]	—	—
Comaltepec Chinantec	hV[], ?V[]	Vh[], V?[]	—
Copala Trique	hV[], ?V[]	Vh[], V?[]	VhV[], V?V[]

For many descriptions of languages with laryngeal complexity, the implicational hierarchy seems to hold. This is certainly the case in the other Trique languages (DiCanio 2008, 2010, 2012a,b, 2014, DiCanio et al. 2020, Elliott, Edmondson & Cruz 2016, Hollenbach 1984). However, as mentioned in Frazier (2013), it is not clear how accu-

rate or robust this implicational hierarchy is actually. The reason for this is because it is not always clear whether something is a laryngeal consonant or a laryngeal feature in the vowel. In fact, Silverman (1997a,b) treats laryngeal consonants and laryngeal features in vowels as the same thing. For example, in many Trique languages, the laryngealization is realized as a laryngeal consonant (DiCanio 2008, 2010, 2012a,b, 2014, DiCanio et al. 2020, Elliott, Edmondson & Cruz 2016, Hollenbach 1984), but in Jalapa Mazatec, the laryngealization is realized as a laryngeal feature on the vowel (Kirk, Ladefoged & Ladefoged 1993, Garellek & Keating 2011).

Another issue with the implicational hierarchy is that it is based on only three languages. This is a rather small sample size and does not capture the full range of variation in the Oto-Manguean languages. For example, in many Mixtec languages, laryngealization is understood to be a feature of the vowel rather than a consonant (e.g., Cortés, Mantenuto & Steffman 2023, Eischens 2022, Gerfen 1999, Gerfen & Baker 2005). Additionally, in many of these languages, the laryngealization can appear either in the middle of the vowel, which Silverman (1997a) calls interrupted, or at the end of the vowel (e.g., Cortés, Mantenuto & Steffman 2023, Eischens 2022). This is directly in contrast to the implicational hierarchy, which states that if a language has interrupted laryngealization, it must also have post-vocalic laryngealization and pre-vocalic laryngealization.

Not only is the violation of the implicational hierarchy the case in Mixtecan lan-

guages, it is also the case in other branches of the Oto-Manguean languages. It is often the case that Zapotec languages have only interrupted and post-vocalic vowels or just interrupted vowels (see Ariza-García 2018 for a typology of phonation in Zapotec languages). For example, Avelino (2004, 2010) argues that Yalálag Zapotec only has interrupted laryngealization as a vowel feature and postvocalic laryngealization with a laryngeal consonant. However, there is no prevocalic laryngealization in the language. This is in direct contrast to the implicational hierarchy of laryngeal complexity. This is also true for laryngealization in Santa Ana del Valle Zapotec (Esposito 2004, 2012).

6.3 Previous analyses of laryngeal complexity

Previous analyses of laryngeal complexity fall into two categories: (i) descriptive studies and (ii) instrumental studies. In most descriptive studies, the focus has been on describing the patterns of tone and voice quality and how they interact with each other. For example, Frazier (2013) describes the phonetic properties of tone and voice quality in Yucatec Mayan. In this study, Frazier describes how Yucatec Mayan is one of the few Mayan languages that has developed tonal contrasts. Additionally, it has a series of vowels that have a high tone with glottalized that variably surface as rearticulated vowels¹ or as a vowel with creaky voice. Frazier notes that for most speakers

¹This is very similar to rearticulated vowels in SLZ, where a vowel has a period of glottalization in the middle of the vowel that is often realized as a glottal stop. This is different from how Baird (2011)

these vowels show clear evidence of phasing between the tone and the voice quality. In these vowels, the first portion of the vowel is always modal and is produced with the high tone. The second portion of the vowel is produced with creaky voice, which greatly obscures the pitch.

Not only is this the case in Yucatec Mayan, but it is also the case in languages that primarily have a phonation contrast with tone/pitch as a secondary cue like Danish (Fischer-Jørgensen 1989, Grønnum, Vazquez-Larruscaín & Basbøll 2013, Peña 2022, 2024). In Danish, there is a phonation contrast between modal voice and a type of creaky voice that is called *stød*. Research has shown that *stød* is also associated with a secondary cue of a heightened f_0 (Fischer-Jørgensen 1989, Grønnum, Vazquez-Larruscaín & Basbøll 2013). Peña (2022, 2024) showed that even though Danish is not traditionally classified as a laryngeally complex language it still shows evidence of phasing between the primary and secondary cues to *stød*, with the primary cue of phonation being produced in the second half of the syllabic rhyme and the secondary cue of pitch being produced in the first half of the syllabic rhyme.

In contrast to descriptive studies, instrumental studies have focused on the acoustic properties of laryngeal complexity, mainly how laryngealization affects f_0 and the harmonic structure of the vowel. For example, Garellek & Keating (2011) found that in Jalapa Mazatec, the laryngealization causes f_0 perturbation in the first part of the vowel. Garellek & Keating conclude that this is evidence for the phasing between

describes “broken” or “rearticulated” vowels in K’ichee, another Mayan language.

laryngealization and tone, as predicted by Silverman's 1997 theory of laryngeal complexity. Additionally, this same phenomenon of f_0 perturbation has been shown to be the case with laryngealization in some varieties of Trique, again showing that there is phasing between modal and nonmodal phonation (DiCanio 2012a). For these previous studies, the main evidence for laryngeal complexity has been the perturbation of the f_0 signal in the portion of the vowel affected by nonmodal phonation.

In more recent studies, researchers have also looked at other measures besides f_0 to determine if there is phasing. For example Weller et al. (2023a,b, 2024) have investigated laryngeal complexity in San Sebastián del Monte Mixtec using a combination of f_0 and Strength of Excitation (SoE), a measure that correlates with the strength of voicing, measures. They found that there is a clear phasing between modal and non-modal phonation in the rearticulated vowels of the language.

However, these accounts offer only a limited window into the question of laryngeal complexity. Because there has been a focus on determining whether or not there are f_0 perturbations in the signal, these previous studies have missed the opportunity to look at the full range of acoustic properties. Weller et al. (2023a,b, 2024) have done an excellent job in showing that looking at SoE, in addition to f_0 , we can better understand the phasing between tone and voice quality. However, it is still necessary to look at other acoustic properties of the signal to get a full understanding of laryngeal complexity.

The class of harmonic-to-noise ratio measures is one of those classes that promises to provide this better understanding. Harmonic-to-noise ratio measures are a class of measures that look at the ratio of the harmonic energy to the noise energy in the signal. This class of measures has been particularly helpful in determining whether there is aperiodicity in the signal (de Krom 1993, Ferrer Riesgo & Nöth 2020, Garellek 2019). It is well understood that aperiodicity is one of the defining characteristics of nonmodal phonation (Ladefoged & Maddieson 1996). This means that harmonic-to-noise ratio measures can be used to determine whether there is aperiodicity in the signal and if there is a clear phasing between modal and nonmodal phonation.

6.4 Analysis of laryngeal complexity

6.4.1 Methods

6.4.1.1 Participants

This study uses data collected from 10 native speakers of SLZ during the summer of 2022. Participants were recruited from the community of Santiago Laxopa, Oaxaca, Mexico. All participants were native speakers of SLZ. The participants were between 18 and 60 years old and consisted of five males and five females.

6.4.1.2 Recordings

Participants were asked to perform a word list elicitation task consisting of 72 words. These words were selected to elicit the entire range of types of voice quality in SLZ, including modal voice, the two kinds of creaky (i.e., checked and rearticulated), and breathy voice. The words were selected based on previous research conducted as part of the Zapotec Language Project at the University of California, Santa Cruz (*Zapotec Language Project – University of California, Santa Cruz 2022*). Because participants were not literate in SLZ, the word list was prompted for them by asking them “How do you say [word in Spanish]?” by myself and another researcher in Zapotec. Participants were asked to respond with the desired word in the carrier phrase *Shnia’ X chonhe lhas [ʃn:ia’ X tʃone ras]* “I say X three times.” which was repeated three times. These utterances were recorded in a quiet environment using a Zoom H4n handheld digital recorder. The recordings were saved as 16-bit WAV files with a sampling rate of 44.1 kHz.

6.4.1.3 Acoustic measuring

The resulting audio files were then processed in Praat to isolate the vowel portion of each word. The onset of the vowel was set to the second glottal pulse after the onset, and the offset of the vowel was set to the last glottal pulse before the decrease in amplitude at the end of the vowel (Garellek 2020). The vowel was then extracted

and saved as a separate file for analysis.

These vowels were fed into VoiceSauce (Shue et al. 2011) to generate the acoustic measures for the studies discussed in this dissertation. Because many acoustic measures are based on the fundamental frequency, this measure was calculated using the STRAIGHT algorithm from (Kawahara, Chevigné & Patterson 1998) to estimate the fundamental frequency in millisecond (ms) intervals. Once the fundamental frequency is calculated, VoiceSauce then uses an optimization function to locate the harmonics of the spectrum, finding their amplitudes.

VoiceSauce then uses the Snack Sound toolkit (Sjölander 2004) to find the frequencies and bandwidths of the first four formants, also at millisecond intervals. The amplitudes of the harmonics closest to these formant frequencies are located and treated as the amplitudes of the formants. These formant frequencies and bandwidths are used to correct the harmonic amplitudes for the filtering effects of the vocal tract, using Iseli, Shue & Alwan's 2007 extension of the method employed by Hanson (1997). Each vowel was measured across ten equal time intervals, resulting in 22890 data points in total. These measures were then z-scored by speaker to reduce the variation between speakers and provide a way to compare the different measures directly on similar scales.

6.4.1.4 Data processing

Data points with an absolute z-score value greater than three were considered outliers and excluded from the analysis. The Mahalanobis distance was calculated in the F1-F2 panel within each vowel category. Each data point with a Mahalanobis distance greater than six was considered an outlier and excluded from the analysis. Using the Mahalanobis distance allows us to compare the data points to the mean of the F1-F2 panel for each vowel category. The larger the Mahalanobis distance is, the more deviant the data point is from the mean, which in turn means that the data point was improperly tracked. This is comparable to what was done in Seyfarth & Garellek (2018), Chai & Ye (2022), and Garellek & Esposito (2023).

Energy was excluded if it had a zero value and then logarithmically transformed to normalize its right-skewed distribution. Afterward, the resulting log-transformed energy was z-scored and any data point with a z-score greater than three was excluded. This outlier removal resulted in 1918 data points being removed.

All data points were then z-scored by speaker to reduce the variation between speakers and provide a way to compare the different measures directly on the same scale.

Residual H1* for the remaining data points following Chai & Garellek (2022). First, a linear mixed effects model was generated with the z-scored H1* as the response variable and the z-scored energy as the fixed effect. The uncorrelated interaction of

the z-scored energy by speaker was treated as random. The energy factor resulting from this linear mixed-effects model was extracted. Finally, the z-scored H1* had the product of the z-scored energy and energy factor subtracted from it to produce the residual H1* measure.

As mentioned in Chapter 3, the distribution of tone and phonation was not equal in all combinations of the two. This means that certain combinations of tone and phonation were overrepresented in the data set. As seen in Table 3.1, which is repeated here as Table 6.2, the distribution of tone and phonation is not equal across all combinations.

Table 6.2: Distribution of the number of syllables containing the combination of tone and voice quality in the wordlist.

	High	Mid	Low	Rising	Falling
Modal	14	9	15	2	10
Breathy	—	—	11	—	2
Checked	1	—	9	—	—
Rearticulated	1	—	4	—	4

Modal voice was the only phonation that occurred across all possible tones, and low tone was the only tone that occurred across all possible phonations. This means that the data set is not balanced and that there are more data points for certain combinations of tone and phonation than others. This is a problem because it can lead to biased results in the analysis of laryngeal complexity. To correct for this, only the

phonations that occurred in low tone were used for the analysis of laryngeal complexity. By limiting the analysis to only low tone, we ensure that the dataset accurately captures the effects of the interaction of phonation on a specific tone. Additionally, this will allow us to ignore the potential confounding factor of how creaky voice is not uniformly produced when it appears with different tones (Keating, Garellek & Kreiman 2015).

6.4.1.5 Statistical analysis

Generalized additive mixed models (GAMM; Hastie & Tibshirani 1986, Wood 2017) are a type of statistical model similar to generalized linear mixed models, but allow for smoothing functions of continuous predictors. This allows for the modeling of non-linear relationships between the predictors and the response variable, especially over time. Using GAMMs allows us to capture the non-linear relationships between the predictors and the response variable, which is important for understanding how phonation affects the acoustic measures over time.

The GAMMs were fitted to the data using the `mgcv` package in R (Wood 2017) and plotted using the `tidygam` package (Coretta 2024). Three GAMMs were fitted for each of the acoustic measures: f_0 , HNR < 1500 Hz, and SoE as the response variable. These three measures were chosen based on previous research, with respect to f_0 , and the results of the MDS and Random Forest analyses. Phonation was treated as a fixed effect to model the main effect of phonation on the acoustic measures. A smooth term

for time (i.e., measurement number) was included to model the nonlinear relationship between time. A second smooth term was included to allow time to vary by phonation type. Random effects for speaker and the interaction between speaker and phonation were included to account for variability in the data due to individual differences between speakers and individual differences in how they produce the phonation types. A tensor product interaction between time and repetition was also included as a random effect to account for the variability in the data due to the different repetitions the speakers were asked to complete.

By modeling the GAMMs in this way, each model is able to capture four things: (i) the main effect of phonation on the acoustic measures, (ii) the nonlinear relationships in time both overall and specific for each phonation type, (iii) speaker variability and its interaction with phonation, and (iv) the variability in the data due to the different repetitions the speakers were asked to complete. This allows for a more accurate modeling of the data and a better understanding of how phonation affects the acoustic measures under consideration.

6.5 Results

This section presents the results of the three GAMM analyses. For each measure, I present the GAMM model fit and the model difference plot. The model fit plot shows the predicted values from the GAMM for each of the phonations over time. The model

difference plot shows the difference between the modal phonation and each of the other phonations. This allows us to see how each of the non-modal voice qualities differs from modal phonation over time.

6.5.1 Fundamental frequency

Figure 6.5 shows the model fit for f_0 . Each line represents the predicted values of the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval. The fit of the model shows that the modal voice (the black solid line) has the highest predicted values for f_0 . The nonmodal phonations have lower predicted values, with breathy voice (the orange dashed line) having the lowest predicted values. Checked and rearticulated voice (the blue and green dashed lines, respectively) have predicted values similar to each other and lie between modal and breathy voices. Beginning at measurement number 8 we see that checked voice and rearticulated voice begin to separate from each other. For checked voice, the predicted values begin to increase while for rearticulated voice the predicted values begin to decrease until measurement number 10.

The observations from the model fit are only part of the story when evaluating GAMMs. The other piece of evidence that we need is whether the differences that we observe in the fit of the model are significant. This is done through difference plots that compare the values for a given comparison (in this case modal versus one of the




Figure 6.5: Model fit for f_0 . Each line represents the predicted values from the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval.

nonmodal phonations) and return whether or not the difference is significant or not. In these plots, the red line indicates that the difference that we observe between modal and the other phonation types is significant. The blue line indicates that the difference between modal and the non-modal phonation is not significant. This method of evaluation is the most common method for evaluating GAMMs (see Sóskuthy 2021 for a discussion about evaluating GAMMs).

Figure 6.6 shows the corresponding difference plot for f_0 . The leftmost panel

shows the difference plot between modal and breathy voice. The red line indicates that the differences that we observed between modal and breathy voice in Figure 6.5 are significant throughout the length of the vowel. The center panel shows that the differences between modal and checked voice is not significant at any point during the vowel and in terms of f_0 there is no difference between these two phonations. The rightmost panel shows that the difference between modal and rearticulated voice is also not significant for the first part of the vowel. However, beginning at about measurement number 7, we see that the differences are significant.




Figure 6.6: Plot of the difference between modal and each of the non-modal phonation types.

This suggests that in terms of f_0 , breathy voice is significantly different from modal voice throughout the duration of the vowel. This is consistent with the descriptions for breathy voice cross-linguistically where one of its acoustic cues is a lower f_0 than modal vowels (e.g., Hillenbrand & Houde 1996). However, rearticulated vowels are only significantly different from modal vowels for the final third of the vowel. The results suggest that in terms of f_0 perturbation only rearticulated vowels show phasing.

6.5.2 Harmonic-to-noise ratio

The model fit for $\text{HNR} < 1500 \text{ Hz}$ is shown in Figure 6.7. In the fit of the model, we see that modal voice has the highest predicted values for $\text{HNR} < 1500 \text{ Hz}$, which is consistent with previous research that has shown that modal voice has higher HNR than non-modal phonation types (e.g., Blankenship 1997, 2002, de Krom 1993, Garellek 2012, 2019, Gerratt & Kreiman 2001). Additionally, we see that the three non-modal phonations are all very similar to each other. However, checked vowels show a dipping at the end of the vowel, indicating more noise in the signal.

Figure 6.8 shows the difference plots for each of the nonmodal phonations. Similarly to Figure 6.6, we see that the leftmost panel shows the difference between modal and breathy voice. This difference is significant from measurement number 4 to halfway between measurements 8 and 9.

The center and leftmost panels show that the difference between modal and the




Figure 6.7: Model fit for HNR. Each line represents the predicted values from the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval.

nonmodal phonations is significant across the entire length of the vowel. This indicates that checked and rearticulated vowels are associated with greater aperiodicity throughout the length of the vowel.

From HNR < 1500 Hz, we see that all three nonmodal phonations have more noise in their signal than modal vowels. As I discussed in Section 5.7.6, HNR measures are typically associated with aperiodicity in the signal. If there is phasing between modal phonation and nonmodal phonation as predicted by Silverman (1997a), we would ex-




Figure 6.8: Plot of the difference between modal and each of the non-modal phonation types.

pect to see that these phonations should have periods where they exhibit more modal-like properties or are not significantly different from modal phonation, which is what we observe with only a breathy voice. Breathy vowels show a period of greater aperiodicity than modal vowels in the second half of the vowel, especially if we ignore the interval between 9 and 10 due to coarticulation with the surrounding consonants. Furthermore, this indicates that there is a phasing relationship in breathy vowels where the first half of the vowel is modal and the second half is breathy.

6.5.3 Strength of excitation

As discussed in Section 5.7.7, SoE is an acoustic measure that correlates with the strength of voice. Figure 6.9 shows the fit of the model for SoE. In the model fit, we see that modal voice has the highest predicted values for SoE, which is consistent with Garellek et al. (2021) where modal vowels are associated with a higher SoE than non-modal phonation because of the greater strength in voicing.

Breathy voice is identical to the modal voice until just after measurement number 2 and then begins to decrease but becomes identical to the modal vowels shortly after measurement number 9. Checked vowels vowels are similar to breathy vowels in that they are identical to modal vowels until just after measurement number 2. However, checked vowels have a sharp decrease in SoE throughout the vowel until measurement number 10. However, rearticulated vowels begin lower than the other three phonations but increase in SoE until they are identical to the modal voice between measurements 8 and 9.

In evaluating whether or not the observed differences from the model fit are significant or not, we can again turn to difference plots to help us determine which differences were significant or not. Figure 6.10 shows the difference plots for each of the nonmodal phonations.

The leftmost panel shows the difference between modal and breathy voice. The differences we observed between modal and breathy vowels were only significant be-

Model fit for SoE




Figure 6.9: Model fit for SoE. Each line represents the predicted values from the GAMM for each of the phonations over time. The shaded area represents the 95% confidence interval.

tween measurement numbers 4 and 8. This is very similar to what we observed in Figure 6.8 for HNR < 1500 Hz. For breathy vowels, we can ignore the interval between measurements 9 and 10 because of the coarticulation with the surrounding consonants.

The center panel shows the difference between modal and checked voice. We observe that checked vowels are significantly different from modal vowels from measurement number 3 to the end of the vowel. However, we can consider the interval

between measurements 9 and 10 because checked vowels have a phontactic restriction in the nominal domain that prevents them from occurring in closed syllables. The only time they can accure in a closed syllables is when a morpheme boundary intervenes, such as in the word *beku'*=*nh* [beku'=ŋ] ‘the dog’ where the checked vowel occurs at a morpheme boundary.

The rightmost panel shows the difference between modal and rearticulated voice. The differences we observed between modal and rearticulated vowels were only significant between measurement numbers 1 and just after 7.




Figure 6.10: Plot of the difference between modal and each of the non-modal phonation types.

From the SoE measure, we see that the strength of voicing is not drastically different between modal and non-modal phonation. This suggests that non-modal phonations are not as strongly articulated in SLZ when compared to non-modal phonations in other languages (e.g., Garellek et al. 2021, Garellek, López-Francisco & Amith 2025, Weller et al. 2023a,b, 2024). This weak articulation was predicted by Silverman and is consistent with the idea that laryngealization is not as strongly articulated in Zapotec languages as it is in other Oto-Manguean languages (Herrera Zendejas 2000).

6.6 Discussion

As an Oto-Manguean language, it is not surprising that SLZ exhibits laryngeal complexity. As a Zapotec language, SLZ is expected to offer counterarguments to some of the claims made by Silverman (1997a,b) in relation to the implicational hierarchy. The reason for this is that Zapotec languages are known to typically only have interrupted and post-vocalic laryngealization (Arellanes Arellanes 2009, 2010, Ariza-García 2018, Avelino 2004, 2010, Esposito 2004, 2010). This means that Zapotec languages offer a strong counter-argument against the implicational hierarchy because they lack any type of prevocalic laryngealization. Furthermore, according to Herrera Zendejas (2000), Zapotec languages do not have a phasing relationship between modal and non-modal phonation. This is because these languages have achieved “equilibrium” between tone and laryngealization.

The results of this study show that these expectations for Zapotec languages do not hold in SLZ. Instead, the results, when taken in aggregate, show that SLZ exhibits a phasing relationship between modal and nonmodal phonation in support of Silverman's 1997, 1997 claims about laryngeal complexity. This is supported by the way breathy and checked voice shows a phasing pattern with modal phonation precedes a period of laryngealization, or using Silverman's 1997 vocabulary a post-vocalic laryngealization. Rearticulated vowels, while on the surface appear to belong to the interrupted, do not actually show this. Instead, these vowels begin with non-modal phonation and then end with modal phonation. This classifies them as belonging to the Silverman's 1997 prevocalic laryngealization.

The patterning of nonmodal phonation in prevocalic and postvocalic laryngealization agrees with the implicational hierarchy for laryngeal complexity. As discussed above in Section 6.2.2, the implicational hierarchy states that if a language has laryngealization, it should follow that if it has interrupted laryngealization, it should also have post-vocalic laryngealization, and if it has post-vocalic laryngealization, it should also have prevocalic laryngealization. This can be seen in the table presented above in Table 6.1, which is repeated here as Table 6.3.

As stated previously, the results of this study support this implicational hierarchy for SLZ. SLZ has both prevocalic laryngealization with its rearticulated vowels and postvocalic laryngealization with its checked vowels and breathy vowels. However,

Table 6.3: Implicational hierarchy of laryngeal complexity. The symbols h and ? represent laryngealization. The symbol V represents where the modal vowel is located in relation to the laryngealization. Modified from Silverman (1997a).

Language	Prevocalic	Postvocalic	Interrupted
Jalapa Mazatec	hV[], ?V[]	—	—
Comaltepec Chinantec	hV[], ?V[]	Vh[], V?[]	—
Copala Trique	hV[], ?V[]	Vh[], V?[]	VhV[], V?V[]

this does not mean that this implicational hierarchy does not need further investigation. As I discussed in Section 6.2.2, the implicational hierarchy is based on a small number of languages and does not take into account the full range of laryngeal complexity. This means that while the implicational hierarchy is a good starting point, it needs to be tested against more languages to see if it holds. Additionally, the implicational hierarchy needs to be refined to make it clear whether it is concerned with just laryngeal consonants, laryngealization on vowels, or both. It is also not clear whether the implicational hierarchy is a universal property of laryngeal complexity or if it is just a property of the languages that have been studied so far. Additionally, the implicational hierarchy needs to be further tested to see if it holds even against languages that are not laryngeally complex like Oto-Manguean languages. This is important because it will help us to better understand the nature of tone- and voice-quality interactions cross-linguistically.

The results of the study also suggest that, in keeping with Silverman's 1997 claims,

laryngealization in SLZ appears to be more weakly implemented as evidenced by the rather narrow difference in SoE between modal and nonmodal phonation. This is only possible if the amplitude of voicing is not entirely different between modal and non-modal phonation.

This study also shows that to get a complete picture of laryngeal complexity in a language we need to look at more than just f_0 perturbation. Looking at harmonics-to-noise ratio measures and measures about voicing in addition to f_0 , we can get a better understanding of the phasing relationship between modal and nonmodal phonation. This is important because it allows us to see how these different measures interact with each other and how they can be used to better understand laryngeal complexity in a language. This is also necessary because the nature of voice quality being a multidimensional phenomenon (e.g., Kreiman et al. 2014, 2021). If we only focus our attention on one acoustic phenomenon, such as f_0 perturbation, we will miss out on the full range of acoustic properties present in the signal. This is especially true for laryngeal complexity, where there are multiple dimensions to consider. By looking at a wider range of acoustic properties, we can gain a better understanding of how these different measures interact with each other and how they can be used to better understand laryngeal complexity in a language.

6.7 Conclusion

In conclusion, the results of this study show that Silverman's 1997 claims about laryngeal complexity continue to hold after almost 30 years. We see that SLZ exhibits a phasing relationship between modal and nonmodal phonation in its laryngeally complex vowels. Evidence for this phasing was only possible by looking for f_0 perturbations and through the use of acoustic measures for harmonics-to-noise and Strength of Excitation. This, further, shows that laryngeal complexity is also a multidimensional phenomenon and needs to be carefully considered in light of multiple acoustic measures. If only one acoustic measure is used, such as f_0 perturbation, we will miss out on the full range of acoustic properties present in the signal.

Furthermore, this study shows that the implicational hierarchy for laryngeal complexity holds for SLZ. With its three phonation types, exhibit both prevocalic and postvocalic laryngealization. This also shows that interrupted laryngealization is not actually present in this language. This is important because it shows that the implicational hierarchy is a valid way to think about laryngeal complexity and how it interacts with the quality of tone and voice. However, further work is needed to establish whether this hierarchy holds for all tone and phonation interactions or if they only hold for laryngeally complex languages.

Chapter 7

Conclusion

7.1 The Laryngeal Articulator Model

Esling (2005), Esling et al. (2019), Moisik, Czaykowska-Higgins & Esling (2021),
Moisik et al. (2015), and Moisik & Esling (2014)




Figure 7.1: The Laryngeal Articulator Model from Esling et al. (2019). This model shows the interactions between the laryngeal articulators (labeled circles). Synergetic interactions are shown with solid lines, while anti-synergetic interactions are shown with dotted lines.

7.2 Modeling laryngeal complexity

7.3 Alternative accounts

7.3.1 Articulatory Phonology account

7.3.2 Q-theory account

7.3.3 Radical CV Phonology account

Bibliography

- Adler, Jeff, Steven Foley, Jed Pizarro-Guevara, Kelsey Sasaki & Maziar Toosarvandani. 2018. The derivation of verb initiality in Santiago Laxopa Zapotec. In Jason Merchant, Line Mikkelsen, Deniz Rudin & Kelsey Sasaki (eds.), *A reasonable way to proceed: Essays in honor of Jim McCloskey*, 31–49. Santa Cruz, Berkeley, Chicago: University of California.
- Adler, Jeff & Maho Morimoto. 2016. Acoustics of phonation types and tones in Santiago Laxopa Zapotec. *The Journal of the Acoustical Society of America* 140(4). 3109–3109.
<https://doi.org/10.1121/1.4969713>.
- Arellanes Arellanes, Francisco. 2009. *El sistema fonológico y las propiedades fonéticas del zapoteco de San Pablo Güilá: descripción y análisis formal*. Mexico City, Mexico: El Colegio de México Doctoral thesis.
- Arellanes Arellanes, Francisco. 2010. Dos ‘Grados’ De Laringización Con Pertinencia Fonológica En El Zapoteco De San Pablo Güilá. In Esther Herrera Zendejas (ed.), *Entre cuerdas y velo*, 1st edn., vol. 9 (Estudios Fonológicos de Lenguas Otomangues),

- 85–122. Mexico City, Mexico: El Colegio de México. <https://doi.org/10.2307/j.ctv6jmx5b.7>.
- Ariza-García, Andrea. 2018. Phonation types and tones in Zapotec languages: A synchronic comparison. *Acta Linguistica Petropolitana* XIV(2). 485–516. <https://doi.org/10.30842/alp2306573714220>.
- Arras, Kai Oliver. 1998. *An Introduction to Error Propagation: Derivation, Meaning and Examples of Equation CY= FX CX FXT*. Technical Report EPFL-ASL-TR-98-01 R3. Lausanne, Switzerland: Swiss Federal Institute of Technology Lausanne (EPFL).
- Avelino, Heriberto. 2004. *Topics in Yalálag Zapotec, with particular reference to its phonetic structures*. Los Angeles, CA: University of California, Los Angeles dissertation. 315 pp.
- Avelino, Heriberto. 2010. Acoustic and Electroglossographic Analyses of Nonpathological, Nonmodal Phonation. *Journal of Voice* 24(3). 270–280. <https://doi.org/10.1016/j.jvoice.2008.10.002>.
- Baird, Brandon. 2011. Phonetic and phonological realizations of 'broken glottal' vowels in K'ichee'. In Kirill Shklovsky, Pedro Mateo Pedro & Jessica Coon (eds.), *Proceedings of FAMLi 1: Formal Approaches to Mayan Linguistics* (MIT Working Papers in Linguistics 63). Cambridge, MA: MIT Press.
- Barzilai, Maya L. & Katherine J. Riestenberg. 2021. Context-dependent phonetic enhancement of a phonation contrast in San Pablo Macuiltianguis Zapotec. *Glossa: a*

journal of general linguistics 6(1). 1–36. <https://doi.org/10.5334/gjgl.959>.

Beam de Azcona, Rosemary G. 2007. Problems in Zapotec Tone Reconstruction. *Annual Meeting of the Berkeley Linguistics Society* 33(2). 3. <https://doi.org/10.3765/bls.v33i2.3497>.

Bennett, Ryan. 2016. Mayan phonology. *Language and Linguistics Compass* 10(10). 469–514. <https://doi.org/10.1111/lnc3.12148>.

Bhattacharyya, A. 1943. On a measure of divergence between two statistical populations defined by their probability distribution. *Bulletin of the Calcutta Mathematical Society* 35. 99–110.

Bladon, Anthony. 1986. Phonetics for hearers. In Graham McGregor (ed.), *Language for Hearers*, 1. ed (Language and Communication Library 8), 1–24. Oxford: Pergamon Press.

Blankenship, Barbara. 1997. *The time course of breathiness and laryngealization in vowels*. Los Angeles, CA: University of California, Los Angeles dissertation.

Blankenship, Barbara. 2002. The timing of nonmodal phonation in vowels. *Journal of Phonetics* 30(2). 163–191. <https://doi.org/10.1006/jpho.2001.0155>.

Blevins, Juliette & Andrew Garrett. 1993. The Evolution of Ponapeic Nasal Substitution. *Oceanic Linguistics* 32(2). 199–236. <https://doi.org/10.2307/3623193>.

Boehmke, Brad & Brandon M. Greenwell. 2019. *Hands-On Machine Learning with R*. New York: Chapman and Hall/CRC. 484 pp. <https://doi.org/10.1201/9780367816377>.

Breiman, Leo. 1996. Bagging predictors. *Machine Learning* 24(2). 123–140. <https://doi.org/10.1007/BF00058655>.

Breiman, Leo. 2001. Random Forests. *Machine Learning* 45(1). 5–32. <https://doi.org/10.1023/A:1010933404324>.

Breiman, Leo, Jerome H. Friedman, Richard A. Olshen & Charles J. Stone. 1986. *Classification and regression trees*. Boca Raton, Fla.: Taylor and Francis. 358 pp.

Brinkerhoff, Mykel Loren, John Duff & Maya Wax Cavallaro. 2021. Downstep in Santiago Laxopa Zapotec and the prosodic typology of VSO languages. In *Manchester Phonology Meeting*. Manchester, England.

Brinkerhoff, Mykel Loren, John Duff & Maya Wax Cavallaro. 2022. Tonal patterns and their restrictions in Santiago Laxopa Zapotec.

Brinkerhoff, Mykel Loren & Grant McGuire. 2025. Using residual H1* for voice quality research. *JASA Express Letters* 5(2). 025501. <https://doi.org/10.1121/10.0035881>.

Brunelle, Marc. 2005. *Register in Eastern Cham: Phonological, phonetic and sociolinguistic approaches*. Ithaca, NY: Cornell University dissertation. 355 pp.

Brunelle, Marc. 2009. Tone perception in Northern and Southern Vietnamese. *Journal of Phonetics* 37(1). 79–96. <https://doi.org/10.1016/j.wocn.2008.09.003>.

Brunelle, Marc. 2012. Dialect experience and perceptual integrality in phonological registers: Fundamental frequency, voice quality and the first formant in Cham. *The Journal of the Acoustical Society of America* 131(4). 3088–3102. <https://doi.org/10.1121/1.3693651>.

Brunelle, Marc & James Kirby. 2016. Tone and Phonation in Southeast Asian Languages. *Language and Linguistics Compass* 10(4). 191–207. <https://doi.org/10.1111/lnc3.12182>.

Burnham, Kenneth P. & David R. Anderson (eds.). 2004a. *Model Selection and Multimodel Inference*. New York, NY: Springer. <https://doi.org/10.1007/b97636>.

Burnham, Kenneth P. & David R. Anderson. 2004b. Multimodel Inference: Understanding AIC and BIC in Model Selection. *Sociological Methods & Research* 33(2). 261–304. <https://doi.org/10.1177/0049124104268644>.

Burnham, Kenneth P., David R. Anderson & Kathryn P. Huyvaert. 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology* 65(1). 23–35. <https://doi.org/10.1007/s00265-010-1029-6>.

Butler H., Inez M. 1997. *Diccionario zapoteco de Yatzachi: Yatzachi el Bajo, Yatzachi el Alto, Oaxaca*. 1. ed (Serie de vocabularios y diccionarios indígenas "Mariano Silva y Aceves" no. 37). Tucson, AZ: Instituto Lingüístico de Verano. 528 pp.

Campbell, Eric W. 2011. Zenzontepec Chatino aspect morphology and Zapotecan verb classes. *International Journal of American Linguistics* 77(2). 219–246.

Campbell, Eric W. 2014. *Aspects of the Phonology and Morphology of Zenzontepec Chatino, a Zapotecan Language of Oaxaca, Mexico*. Austin, TX: University of Texas at Austin dissertation.

Campbell, Eric W. 2017a. Otomanguean historical linguistics: Exploring the subgroups. *Language and Linguistics Compass* 11(7). e12244. <https://doi.org/10.1111/lnc3.12244>.

Campbell, Eric W. 2017b. Otomanguean historical linguistics: Past, present, and prospects for the future. *Language and Linguistics Compass* 11(4). e12240. <https://doi.org/10.1111/lnc3.12240>.

Chai, Yuan. 2025. Perception of checked and rearticulated phonations: Effect of duration and glottalization phasing.

Chai, Yuan, Adrián Fernández & Briseida Mendez. 2023. Phonetics of glottalized phonations in Yateé Zapotec. In Radek Skarnitzl & Jan Volín (eds.), *Proceedings of the 20th International Congress of Phonetic Sciences – ICPhS 2023*, 1751–1755. International Phonetic Association.

Chai, Yuan & Marc Garellek. 2022. On H1–H2 as an acoustic measure of linguistic phonation type. *The Journal of the Acoustical Society of America* 152(3). 1856–1870.
<https://doi.org/10.1121/10.0014175>.

Chai, Yuan & Shihong Ye. 2022. Checked Syllables, Checked Tones, and Tone Sandhi in Xiapu Min. *Languages* 7(1). 1–27. <https://doi.org/10.3390/languages7010047>.

Chávez-Peón, Mario E. 2008. Phonetic cues to stress in a tonal language: Prosodic prominence in San Lucas Quiaviní Zapotec. In Susie Jones (ed.), *Proceedings of the 2008 annual conference of the Canadian Linguistic Association*. 13. Vancouver, Canada: University of British Columbia.

Chávez-Peón, Mario E. 2010. *The interaction of metrical structure, tone, and phonation types in Quiaviní Zapotec*. University of British Columbia dissertation. <https://doi.org/10.14288/1.0071253>.

Chen, Marilyn Y. 1997. Acoustic correlates of English and French nasalized vowels. *The Journal of the Acoustical Society of America* 102(4). 2360–2370. <https://doi.org/10.1121/1.419620>.

Coretta, Stefano. 2024. *Tidygam: Tidy Prediction and Plotting of Generalised Additive Models*. manual.

Cortés, Félix, Iara Mantenuto & Jeremy Steffman. 2023. San Sebastián del Monte Mixtec. *Journal of the International Phonetic Association*. 1–22. <https://doi.org/10.1017/S0025100322000226>.

Crowhurst, Megan J., Niamh E. Kelly & Amador Teodocio. 2016. The influence of vowel laryngealisation and duration on the rhythmic grouping preferences of Zapotec speakers. *Journal of Phonetics* 58. 48–70. <https://doi.org/10.1016/j.wocn.2016.06.001>.

Delgutte, Bertrand. 1982. Some correlates of phonetic distinctions at the level of the auditory nerve. In Rolf Carlson & Björn Granström (eds.), *The representation of speech in the peripheral auditory system: proceedings of the Symposium on the Representation of Speech in the Peripheral Auditory System held in Stockholm, Sweden on May 17-19, 1982*, 131–150. Amsterdam: Elsevier Biomedical Press.

Delvaux, Véronique. 2009. Perception du contraste de nasalité vocalique en français. *Journal of French Language Studies* 19(1). 25–59. <https://doi.org/10.1017/S0959269508003566>.

DiCanio, Christian. 2008. *The Phonetics and Phonology of San Martín Itunyoso Trique*. Berkeley, CA: University of California, Berkeley dissertation.

DiCanio, Christian. 2010. Itunyoso Trique. *Journal of the International Phonetic Association* 40(2). 227–238. <https://doi.org/10.1017/S0025100310000034>.

- DiCanio, Christian. 2012a. Coarticulation between tone and glottal consonants in Itunyoso Trique. *Journal of Phonetics* 40(1). 162–176. <https://doi.org/10.1016/j.wocn.2011.10.006>.
- DiCanio, Christian. 2012b. The phonetics of fortis and lenis consonants in Itunyoso Trique. *International Journal of American Linguistics* 78(2). 239–272. <https://doi.org/10.1086/664481>.
- DiCanio, Christian. 2014. Cue weight in the perception of Trique glottal consonants. *The Journal of the Acoustical Society of America* 135(2). 884–895. <https://doi.org/10.1121/1.4861921>.
- DiCanio, Christian, Basileo Martínez Cruz, Benigno Cruz Martínez & Wilberto Martínez Cruz. 2020. Glottal toggling in Itunyoso Triqui. *Phonological Data and Analysis* 2(4). 1–28. <https://doi.org/10.3765/pda.v2art4.3>.
- Eischens, Benjamin. 2022. *Tone, Phonation, and the Phonology-Phonetics Interface in San Martín Peras Mixtec*. UC Santa Cruz.
- Elliott, A. Raymond, Jerold A. Edmondson & Fausto Sandoval Cruz. 2016. Chicahuaxtla Triqui. *Journal of the International Phonetic Association* 46(3). 351–365. <https://doi.org/10.1017/S0025100315000389>.
- Enfield, Nick J. 2005. Areal linguistics and mainland Southeast Asia. *Annual Review of Anthropology* 34. 181–206.

- Esling, John H. 2005. There Are No Back Vowels: The Laryngeal Articulator Model. *Canadian Journal of Linguistics/Revue canadienne de linguistique* 50(1–4). 13–44. <https://doi.org/10.1017/S0008413100003650>.
- Esling, John H., Scott R. Moisik, Allison Benner & Lise Crevier-Buchman. 2019. *Voice Quality: The Laryngeal Articulator Model*. 1st edn. (Cambridge Studies in Linguistics 162). Cambridge University Press. <https://doi.org/10.1017/9781108696555>.
- Esposito, Christina M. 2004. Santa Ana del Valle Zapotec phonation. *UCLA Working Papers in Phonetics* (103). 71–105.
- Esposito, Christina M. 2010. Variation in contrastive phonation in Santa Ana Del Valle Zapotec. *Journal of the International Phonetic Association* 40(2). 181–198. <https://doi.org/10.1017/S0025100310000046>.
- Esposito, Christina M. 2012. An acoustic and electroglottographic study of White Hmong tone and phonation. *Journal of Phonetics* 40(3). 466–476. <https://doi.org/10.1016/j.wocn.2012.02.007>.
- Esposito, Christina M. & Sameer ud Dowla Khan. 2012. Contrastive breathiness across consonants and vowels: A comparative study of Gujarati and White Hmong. *Journal of the International Phonetic Association* 42(2). 123–143. <https://doi.org/10.1017/S0025100312000047>.

Esposito, Christina M. & Sameer ud Dowla Khan. 2020. The cross-linguistic patterns of phonation types. *Language and Linguistics Compass* 14(12). <https://doi.org/10.1111/lnc3.12392>.

Esposito, Christina M., Sameer ud Dowla Khan, Kelly H. Berkson & Max Nelson. 2019. Distinguishing breathy consonants and vowels in Gujarati. *Journal of South Asian Languages and Linguistics* 6(2). 215–243. <https://doi.org/10.1515/jsall-2019-2011>.

Ferrer Riesgo, Carlos A. & Elmar Nöth. 2020. What Makes the Cepstral Peak Prominence Different to Other Acoustic Correlates of Vocal Quality? *Journal of Voice* 34(5). 806.e1–806.e6. <https://doi.org/10.1016/j.jvoice.2019.01.004>.

Fischer-Jørgensen, Eli. 1968. Phonetic Analysis of Breathy (Murmured) Vowels in Gujarati. *Annual Report of the Institute of Phonetics University of Copenhagen* 2. 35–85. <https://doi.org/10.7146/aripuc.v2i.130674>.

Fischer-Jørgensen, Eli. 1989. Phonetic Analysis of the Stød in Standard Danish. *Phonetica* 46(1–3). 1–59. <https://doi.org/10.1159/000261828>.

Foley, Steven, Nick Kalivoda & Maziar Toosarvandani. 2018. Forbidden clitic clusters in Zapotec. In Daniel Edmiston, Marina Ermolaeva, Emre Hakgüder, Jackie Lai, Kathryn Montemurro, Brandon Rhodes, Amara Sankhagowit & Michael Tabatowski (eds.), *Proceedings of the Fifty-third Annual Meeting of the Chicago Linguistic Society*, 87–102.

- Foley, Steven & Maziar Toosarvandani. 2022. Extending the Person-Case Constraint to Gender: Agreement, Locality, and the Syntax of Pronouns. *Linguistic Inquiry* 53(1). 1–40. https://doi.org/10.1162/ling_a_00395.
- Frazier, Melissa. 2013. The phonetics of Yucatec Maya and the typology of laryngeal complexity. *STUF - Language Typology and Universals* 66(1). 7–21. <https://doi.org/10.1524/stuf.2013.0002>.
- Gao, Xin & Jianjing Kuang. 2022. Phonation Variation as a Function of Checked Syllables and Prosodic Boundaries. *Languages* 7(3). 171. <https://doi.org/10.3390/languages7030171>.
- Garellek, Marc. 2012. The timing and sequencing of coarticulated non-modal phonation in English and White Hmong. *Journal of Phonetics* 40(1). 152–161. <https://doi.org/10.1016/j.wocn.2011.10.003>.
- Garellek, Marc. 2013. *Production and perception of glottal stops*. Los Angeles, CA: University of California, Los Angeles dissertation.
- Garellek, Marc. 2019. The phonetics of voice. In William F. Katz & Peter F. Assmann (eds.), *The Routledge handbook of phonetics* (Routledge Handbooks in Linguistics), 75–106. Abingdon, Oxon ; New York, NY: Routledge.
- Garellek, Marc. 2020. Acoustic Discriminability of the Complex Phonation System in !Xóõ. *Phonetica* 77(2). 131–160. <https://doi.org/10.1159/000494301>.

Garellek, Marc. 2022. Theoretical achievements of phonetics in the 21st century: Phonetics of voice quality. *Journal of Phonetics* 94. 101155. <https://doi.org/10.1016/j.wocn.2022.101155>.

Garellek, Marc, Yuan Chai, Yaqian Huang & Maxine Van Doren. 2021. Voicing of glottal consonants and non-modal vowels. *Journal of the International Phonetic Association*. 1–28. <https://doi.org/10.1017/S0025100321000116>.

Garellek, Marc & Christina M. Esposito. 2023. Phonetics of White Hmong vowel and tonal contrasts. *Journal of the International Phonetic Association* 53(1). 213–232. <https://doi.org/10.1017/S0025100321000104>.

Garellek, Marc & Patricia Keating. 2011. The acoustic consequences of phonation and tone interactions in Jalapa Mazatec. *Journal of the International Phonetic Association* 41(2). 185–205. <https://doi.org/10.1017/S0025100311000193>.

Garellek, Marc, Osbel López-Francisco & Jonathan D. Amith. 2025. At the margins: phonology and phonetics of Zongozotla Totonac glottalization. *Laboratory Phonology* 16(1). <https://doi.org/10.16995/labphon.10925>.

Garellek, Marc, Amanda Ritchart & Jianjing Kuang. 2016. Breathy voice during nasal-
ity: A cross-linguistic study. *Journal of Phonetics* 59. 110–121. <https://doi.org/10.1016/j.wocn.2016.09.001>.

- Garellek, Marc, Robin Samlan, Bruce R. Gerratt & Jody Kreiman. 2016. Modeling the voice source in terms of spectral slopes. *The Journal of the Acoustical Society of America* 139(3). 1404–1410. <https://doi.org/10.1121/1.4944474>.
- Gerfen, Chip. 1999. *Phonology and phonetics in Coatzospan Mixtec* (Studies in Natural Language and Linguistic Theory v. 48). Dordrecht ; Boston: Kluwer Academic. 304 pp.
- Gerfen, Chip & Kirk Baker. 2005. The production and perception of laryngealized vowels in Coatzospan Mixtec. *Journal of Phonetics* 33(3). 311–334. <https://doi.org/10.1016/j.wocn.2004.11.002>.
- Gerratt, Bruce R. & Jody Kreiman. 2001. Toward a taxonomy of nonmodal phonation. *Journal of Phonetics* 29(4). 365–381. <https://doi.org/10.1006/jpho.2001.0149>.
- Gordon, Matthew & Peter Ladefoged. 2001. Phonation types: a cross-linguistic overview. *Journal of Phonetics* 29(4). 383–406. <https://doi.org/10.1006/jpho.2001.0147>.
- Grønnum, Nina, Miguel Vazquez-Larruscaín & Hans Basbøll. 2013. Danish Stød: Laryngealization or Tone. *Phonetica* 70(1–2). 66–92. <https://doi.org/10.1159/000354640>.

Hanson, Helen M. 1997. Glottal characteristics of female speakers: Acoustic correlates.

The Journal of the Acoustical Society of America 101(1). 466–481. <https://doi.org/10.1121/1.417991>.

Hastie, Trevor & Robert Tibshirani. 1986. Generalized Additive Models. *Statistical Science* 1(3). 297–310. <https://doi.org/10.1214/ss/1177013604>.

Hastie, Trevor, Robert Tibshirani & Jerome Friedman. 2009. *The Elements of Statistical Learning* (Springer Series in Statistics). New York, NY: Springer. <https://doi.org/10.1007/978-0-387-84858-7>.

Hay, Jennifer, Paul Warren & Katie Drager. 2006. Factors influencing speech perception in the context of a merger-in-progress. *Journal of Phonetics*. Modelling Sociophonetic Variation 34(4). 458–484. <https://doi.org/10.1016/j.wocn.2005.10.001>.

Henderson, Eugénie J. A. 1952. The Main Features of Cambodian Pronunciation. *Bulletin of SOAS* 14(1). 149–174. <https://doi.org/10.1017/S0041977X00084251>.

Herbst, Christian T. 2021. Performance Evaluation of Subharmonic-to-Harmonic Ratio (SHR) Computation. *Journal of Voice* 35(3). 365–375. <https://doi.org/10.1016/j.jvoice.2019.11.005>.

Herrera Zendejas, Esther. 2000. Amuzgo and Zapotec: Two More Cases of Laryngeally Complex Languages. *Anthropological Linguistics* 42(4). 545–563. <http://www.jstor.org/stable/30028566>.

Hillenbrand, James & Robert A. Houde. 1996. Acoustic Correlates of Breathy Vocal Quality: Dysphonic Voices and Continuous Speech. *Journal of Speech, Language, and Hearing Research* 39(2). 311–321. <https://doi.org/10.1044/jshr.3902.311>.

Hockett, Charles F. 1947. Peiping Phonology. *Journal of the American Oriental Society* 67(4). 253–267. <https://doi.org/10.2307/596062>.

Hollenbach, Barbara. 1984. *The phonology and morphology of tone and laryngeals in Copala Trique*. University of Arizona.

Holmberg, Eva B., Robert E. Hillman, Joseph S. Perkell, Peter C. Guiod & Susan L. Goldmann. 1995. Comparisons Among Aerodynamic, Electroglossographic, and Acoustic Spectral Measures of Female Voice. *Journal of Speech, Language, and Hearing Research* 38(6). 1212–1223. <https://doi.org/10.1044/jshr.3806.1212>.

Iseli, Markus, Yen-Liang Shue & Abeer Alwan. 2007. Age, sex, and vowel dependencies of acoustic measures related to the voice source. *The Journal of the Acoustical Society of America* 121(4). 2283–2295. <https://doi.org/10.1121/1.2697522>.

Jaeger, Jeri J. & Robert D. Van Valin. 1982. Initial Consonant Clusters in Yateé Zapotec.

International Journal of American Linguistics 48(2). 125–138. <https://doi.org/10.1086/465724>.

James, Gareth, Daniela Witten, Trevor Hastie & Robert Tibshirani. 2021. *An introduction to statistical learning: with applications in R*. Second edition (Springer Texts in Statistics). New York: Springer. 1 p. <https://doi.org/10.1007/978-1-0716-1418-1>.

Janitza, Silke & Roman Hornung. 2018. On the overestimation of random forest's out-of-bag error. *PLOS ONE* 13(8). e0201904. <https://doi.org/10.1371/journal.pone.0201904>.

Johnson, Daniel Ezra. 2015. Quantifying Overlap with Bhattacharyya's affinity and other measures!

Josserand, Judy Kathryn. 1983. *Mixtec Dialect History*. New Orleans, LA: Tulane University dissertation. 727 pp.

Kawahara, Hideki, Alain De Cheveigne & Roy D. Patterson. 1998. An instantaneous-frequency-based pitch extraction method for high-quality speech transformation: revised TEMPO in the STRAIGHT-suite. In *5th International Conference on Spoken Language Processing (ICSLP 1998)*, paper 0659-. ISCA. <https://doi.org/10.21437/ICSLP.1998-555>.

Keating, Patricia, Marc Garellek & Jody Kreiman. 2015. Acoustic properties of different kinds of creaky voice. In *Proceedings of the 18th International Congress of Phonetic Sciences*.

Keating, Patricia, Jianjing Kuang, Marc Garellek, Christina M. Esposito & Sameer ud Dowla Khan. 2023. A cross-language acoustic space for vocalic phonation distinctions. *Language* 99(2). 351–389. <https://doi.org/10.1353/lan.2023.a900090>.

Kelterer, Anneliese & Barbara Schuppler. 2020. Phonation type contrasts and tone in Chichimec. *The Journal of the Acoustical Society of America* 147(4). 3043–3059. <https://doi.org/10.1121/10.0001015>.

Kent, Raymond D. & Martin J. Ball (eds.). 1999. *Voice quality measurement*. San Diego: Singular Pub. Group. 492 pp.

Khan, Sameer ud Dowla. 2012. The phonetics of contrastive phonation in Gujarati. *Journal of Phonetics* 40(6). 780–795. <https://doi.org/10.1016/j.wocn.2012.07.001>.

Kirk, Paul L, Jenny Ladefoged & Peter Ladefoged. 1993. Quantifying acoustic properties of modal, breathy and creaky vowels in Jalapa Mazatec. In Anthony Mattina & Timothy Montler (eds.), *American Indian linguistics and ethnography in honor of Laurence C. Thompson*, 435–450. Ann Arbor, MI: University of Michigan Press.

- Klatt, Dennis H. & Laura C. Klatt. 1990. Analysis, synthesis, and perception of voice quality variations among female and male talkers. *The Journal of the Acoustical Society of America* 87(2). 820–857. <https://doi.org/10.1121/1.398894>.
- Kreiman, Jody, Bruce R. Gerratt & Norma Antoñanzas-Barroso. 2007. Measures of the Glottal Source Spectrum. *Journal of Speech, Language, and Hearing Research* 50(3). 595–610. [https://doi.org/10.1044/1092-4388\(2007/042\)](https://doi.org/10.1044/1092-4388(2007/042)).
- Kreiman, Jody, Bruce R. Gerratt, Marc Garellek, Robin Samlan & Zhaoyan Zhang. 2014. Toward a unified theory of voice production and perception. *Loquens* 1(1). e009. <https://doi.org/10.3989/loquens.2014.009>.
- Kreiman, Jody, Yoonjeong Lee, Marc Garellek, Robin Samlan & Bruce R. Gerratt. 2021. Validating a psychoacoustic model of voice quality. *The Journal of the Acoustical Society of America* 149(1). 457–465. <https://doi.org/10.1121/10.0003331>.
- de Krom, Guus. 1993. A Cepstrum-Based Technique for Determining a Harmonics-to-Noise Ratio in Speech Signals. *Journal of Speech, Language, and Hearing Research* 36(2). 254–266. <https://doi.org/10.1044/jshr.3602.254>.
- Kruskal, Joseph & Myron Wish. 1978. *Multidimensional Scaling*. SAGE Publications, Inc. <https://doi.org/10.4135/9781412985130>.
- Kuang, Jianjing. 2017. Covariation between voice quality and pitch: Revisiting the case of Mandarin creaky voice. *The Journal of the Acoustical Society of America* 142(3). 1693–1706. <https://doi.org/10.1121/1.5003649>.

Ladefoged, Peter & Ian Maddieson. 1996. *The sounds of the world's languages* (Phonological Theory). Oxford, OX, UK ; Cambridge, Mass., USA: Blackwell Publishers. 425 pp.

Laver, John D. M. 1968. Voice Quality and Indexical Information. *British Journal of Disorders of Communication* 3(1). 43–54. <https://doi.org/10.3109/13682826809011440>.

Lillehaugen, Brook Danielle. 2019. Otomanguean Languages. In *The Routledge Handbook of North American Languages*. Routledge.

Lindblom, Björn. 1983. Economy of Speech Gestures. In Peter F. MacNeilage (ed.), *The Production of Speech*, 217–245. New York, NY: Springer New York. https://doi.org/10.1007/978-1-4613-8202-7_10.

Long, Rebecca & Sofronio Cruz. 2005. *Diccionario zapoteco de San Bartolomé Zoogocho, Oaxaca*. 2nd edition (Serie de vocabularios y diccionarios indígenas "Mariano Silva y Aceves" no. 38). Coyoacán, D.F. [Mexico]: Instituto Lingüístico de Verano. 531 pp.

López Nicolás, Oscar. 2016. *Estudios de la fonología y gramática del zapoteco de Zochina*. Mexico City, Mexico: Ciudad de México: Centro de Investigaciones y Estudios Superiores en Antropología Social Doctoral thesis.

Lotto, A. J., L. L. Holt & K. R. Kluender. 1997. Effect of Voice Quality on Perceived Height of English Vowels. *Phonetica* 54(2). 76–93. <https://doi.org/10.1159/000262212>.

- Macmillan, Neil A., John Kingston, Rachel Thorburn, Laura Walsh Dickey & Christine Bartels. 1999. Integrality of nasalization and F1. II. Basic sensitivity and phonetic labeling measure distinct sensory and decision–rule interactions. *The Journal of the Acoustical Society of America* 106(5). 2913–2932. <https://doi.org/10.1121/1.428113>.
- Masica, Colin P. 1976. *Defining a linguistic area: South Asia*. Chicago: The University of Chicago Press.
- Matisoff, James A. 1975. Rhinoglottophilia: The mysterious connection between nasal-ity and glottality. In Charles A. Ferguson, Larry M. Hyman & John J. Ohala (eds.), *Nasálfest: Papers from a symposium on nasals and nasalization*, 265–287. Stanford: Stanford University Language Universals Project.
- Merrill, Elizabeth D. 2008. Tilquiapan Zapotec. *Journal of the International Phonetic Association* 38(01). <https://doi.org/10.1017/S0025100308003344>.
- Michaud, Alexis. 2012. The Complex Tones of East/Southeast Asian Languages: Current Challenges for Typology and Modelling. 8.
- Mittal, Vinay Kumar, B. Yegnanarayana & Peri Bhaskararao. 2014. Study of the effects of vocal tract constriction on glottal vibration. *The Journal of the Acoustical Society of America* 136(4). 1932–1941. <https://doi.org/10.1121/1.4894789>.
- Mock, Carol. 1988. Pitch accent and stress in Isthmus Zapotec. In Harry van der Hulst (ed.), *Autosegmental studies on pitch accent*, 197–223. Dordrecht: Foris.

- Moisik, Scott R., John H Esling, Lise Crevier-Buchman, Angélique Amelot & Philippe Halimi. 2015. *Multimodal imaging of glottal stop and creaky voice: Evaluating the role of epilaryngeal constriction*. Scottish Consortium (ed.). Vol. paper 247.
- Moisik, Scott R., Ewa Czaykowska-Higgins & John H. Esling. 2021. Phonological potentials and the lower vocal tract. *Journal of the International Phonetic Association* 51(1). 1–35. <https://doi.org/10.1017/S0025100318000403>.
- Moisik, Scott R. & John H. Esling. 2014. Modeling the Biomechanical Influence of Epilaryngeal Stricture on the Vocal Folds: A Low-Dimensional Model of Vocal–Ventricular Fold Coupling. *Journal of Speech, Language, and Hearing Research* 57(2). S687–S704. https://doi.org/10.1044/2014_JSLHR-S-12-0279.
- Munro, Pamela & Felipe H. Lopez. 1999. *Di’csyonaary x:tèe’n dii’zh sah Sann Lu’uc (San Lucas Quiaviní Zapotec dictionary/Diccionario Zapoteco de San Lucas Quiavini)*. 2 vols. Los Angeles: (UCLA) Chicano Studies Research Center Publications.
- Murty, K. Sri Rama & B. Yegnanarayana. 2008. Epoch Extraction from Speech Signals. *IEEE Transactions on Audio, Speech, and Language Processing* 16(8). 1602–1613. <https://doi.org/10.1109/TASL.2008.2004526>.
- Nellis, Donald G. & Barbara E. Hollenbach. 1980. Fortis versus Lenis in Cajonos Zapotec Phonology. *International Journal of American Linguistics* 46(2). 92–105. <https://doi.org/10.1086/465639>.

Nycz, Jennifer & Lauren Hall-Lew. 2014. Best practices in measuring vowel merger.

Proceedings of Meetings on Acoustics 20(1). 060008. <https://doi.org/10.1121/1.4894063>.

Ohala, John J. 1975. Phonetic explanations for nasal sound patterns. In Charles A. Ferguson, Larry M. Hyman & John J. Ohala (eds.), *Nasalfest: Papers from a symposium on nasals and nasalization*, 289–316. Stanford: Stanford University Language Universals Project.

Ohala, John J. 1978. Production of Tone. In Victoria A. Fromkin (ed.), *Tone: A linguistics survey*, 5–39. New York: Academic Press. <https://doi.org/10.1016/B978-0-12-267350-4.50006-6>.

Ohala, John J. & Manjari Ohala. 1993. The phonetics of nasal phonology: Theorems and data. In Marie K. Huffman & Rena A. Krakow (eds.), *Nasals, Nasalization, and the Velum*, vol. 5 (Phonetics and Phonology), 225–249. San Diego: Academic Press. <https://doi.org/10.1016/B978-0-12-360380-7.50013-2>.

Oksanen, Jari, Gavin L. Simpson, F. Guillaume Blanchet, Roeland Kindt, Pierre Legendre, Peter R. Minchin, R.B. O'Hara, Peter Solymos, M. Henry H. Stevens, Eduard Szoecs, Helene Wagner, Matt Barbour, Michael Bedward, Ben Bolker, Daniel Borcard, Gustavo Carvalho, Michael Chirico, Miquel De Caceres, Sebastien Durand, Heloisa Beatriz Antoniazi Evangelista, Rich FitzJohn, Michael Friendly, Brennan Furneaux, Geoffrey Hannigan, Mark O. Hill, Leo Lahti, Dan McGlinn, Marie-

- Helene Ouellette, Eduardo Ribeiro Cunha, Tyler Smith, Adrian Stier, Cajo J.F. Ter Braak, James Weedon & Tuomas Borman. 2025. *Vegan: Community Ecology Package*. manual.
- Peña, Jailyn M. 2022. Stød Timing and Domain in Danish. *Languages* 7(1). 50. <https://doi.org/10.3390/languages7010050>.
- Peña, Jailyn M. 2024. *The production and perception of stød in Danish*. New York City, NY: New York University dissertation.
- Pickett, Velma B., María Villalobos Villalobos & Stephen A. Marlett. 2010. Isthmus (Juchitán) Zapotec. *Journal of the International Phonetic Association* 40(3). 365–372. <https://doi.org/10.1017/S0025100310000174>.
- Pike, Eunice Victoria. 1948. Problems in Zapotec Tone Analysis. *International Journal of American Linguistics* 14(3). 161–170. <https://doi.org/10.1086/463998>.
- Pillai, K. C. S. 1955. Some New Test Criteria in Multivariate Analysis. *The Annals of Mathematical Statistics* 26(1). 117–121. <https://doi.org/10.1214/aoms/117728599>.
- Podesva, Robert J. 2016. Stance as a Window into the Language-Race Connection: Evidence from African American and White Speakers in Washington, DC. In H. Samy Alim, John R. Rickford & Arnetha F. Ball (eds.), *Raciolinguistics: How Language Shapes Our Ideas About Race*, 203–219. Oxford: Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780190625696.003.0012>.

Pruthi, Tarun & Carol Y. Espy-Wilson. 2004. Acoustic parameters for automatic detection of nasal manner. *Speech Communication* 43(3). 225–239. <https://doi.org/10.1016/j.specom.2004.06.001>.

R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. manual. Vienna, Austria: R Foundation for Statistical Computing.

Ratliff, Martha Susan. 1992. *Meaningful tone: a study of tonal morphology in compounds, form classes and expressive phrases in White Hmong* (Special Report no. 27, 1992). DeKalb, Ill: Northern Illinois University, Center for Southeast Asian Studies. 269 pp.

Remez, Robert E. & Philip E. Rubin. 1993. On the intonation of sinusoidal sentences: Contour and pitch height. *The Journal of the Acoustical Society of America* 94(4). 1983–1988. <https://doi.org/10.1121/1.407501>.

Rensch, Calvin R. 1976. *Comparative Otomanguean phonology* (Language Science Monographs 14). Bloomington: Indiana University Press.

Ritsma, Roelof J. 1967. Frequencies Dominant in the Perception of the Pitch of Complex Sounds. *The Journal of the Acoustical Society of America* 42(1). 191–198. <https://doi.org/10.1121/1.1910550>.

Rosenberg, Aaron E. 1966. Pitch Discrimination of Jittered Pulse Trains. *The Journal of the Acoustical Society of America* 39. 920–928. <https://doi.org/10.1121/1.1909973>.

Sandri, Marco & Paola and Zuccolotto. 2008. A Bias Correction Algorithm for the Gini Variable Importance Measure in Classification Trees. *Journal of Computational and Graphical Statistics* 17(3). 611–628. <https://doi.org/10.1198/106186008X344522>.

Schwartz, Martin F. 1968. The Acoustics of Normal and Nasal Vowel Production. *The Cleft Palate Journal* 5(2). 125–140.

Seyfarth, Scott & Marc Garellek. 2018. Plosive voicing acoustics and voice quality in Yerevan Armenian. *Journal of Phonetics* 71. 425–450. <https://doi.org/10.1016/j.wocn.2018.09.001>.

Shue, Yen-Liang, Patricia Keating, Chad Vicenik & Kristine Yu. 2011. VoiceSauce: A program for voice analysis. In *Proceedings of the 17th International Congress of Phonetic Sciences (ICPhS XVII)*, 1846–1849. Hong Kong.

Sichel, Ivy & Maziar Toosarvandani. 2020a. Pronouns and Attraction in Sierra Zapotec. In Andrew Hedding & Morwenna Hoeks (eds.), *Syntax and semantics at Santa Cruz, Volume IV* (Syntax & Semantics at Santa Cruz (SASC) 4). Santa Cruz, CA: Linguistics Research Center.

Sichel, Ivy & Maziar Toosarvandani. 2020b. The featural life of nominals. lingbuzz/005523. Silverman, Daniel. 1997a. Laryngeal complexity in Otomanguean vowels. *Phonology* 14(2). 235–261. <https://doi.org/10.1017/S0952675797003412>.

Silverman, Daniel. 1997b. *Phasing and recoverability* (Outstanding Dissertations in Linguistics). New York: Garland Pub. 242 pp.

Simpson, Adrian P. 2012. The first and second harmonics should not be used to measure breathiness in male and female voices. *Journal of Phonetics* 40(3). 477–490.
<https://doi.org/10.1016/j.wocn.2012.02.001>.

Sjölander, Kåre. 2004. *Snack Sound Toolkit*. Stockholm, Sweden: KTH.

Smalley, William A. 1976. The problems of consonants and tone: Hmong (Meo, Miao). In W. A. Smalley (ed.), *Phonemes and orthography: Language planning in ten minority languages of thailand*, vol. 43 (Pacific Linguistics, Series C), 85–123. Canberra: Australian National University.

Smith-Stark, Thomas. 2007. Algunas isoglosas zapotecas. In *Clasificación de las lenguas indígenas de México: Memorias del III Coloquio Internacional de Lingüística Mauricio Swadesh*, 69–133. Mexico City: UNAM: Instituto de Investigaciones Antropológicas.

Sonnenschein, Aaron H. 2005. *A descriptive grammar of San Bartolomé Zoogocho Zapotec*. Munich: Lincom Europa.

Sóskuthy, Márton. 2017. *Generalised Additive Mixed Models for Dynamic Analysis in Linguistics: A Practical Introduction*. arXiv: [1703.05339 \[stat\]](https://arxiv.org/abs/1703.05339). Pre-published.

- Sóskuthy, Márton. 2021. Evaluating generalised additive mixed modelling strategies for dynamic speech analysis. *Journal of Phonetics* 84. 101017. <https://doi.org/10.1016/j.wocn.2020.101017>.
- Stevens, Kenneth N. 2000. *Acoustic phonetics*. 1. paperback ed (Current Studies in Linguistics Series 30). Cambridge, Mass.: MIT Press. 607 pp.
- Strelluf, Christopher. 2018. Chapter 3: The Low Back Vowel(s). *The Publication of the American Dialect Society* 103(1). 43–64. <https://doi.org/10.1215/00031283-7318737>.
- Strobl, Carolin, Anne-Laure Boulesteix, Achim Zeileis & Torsten Hothorn. 2007. Bias in random forest variable importance measures: Illustrations, sources and a solution. *BMC Bioinformatics* 8(1). 25. <https://doi.org/10.1186/1471-2105-8-25>.
- Styler, Will. 2015. *On the acoustical and perceptual features of vowel nasality*. University of Colorado at Boulder dissertation.
- Styler, Will. 2017. On the acoustical features of vowel nasality in English and French. *The Journal of the Acoustical Society of America* 142(4). 2469–2482. <https://doi.org/10.1121/1.5008854>.
- Sun, Xuejing. 2002. Pitch determination and voice quality analysis using Subharmonic-to-Harmonic Ratio. *2002 IEEE International Conference on Acoustics, Speech, and*

- Signal Processing* 1. I-333-I-336. <https://doi.org/10.1109/ICASSP.2002.5743722>.
- Sundberg, Johan. 2022. Objective Characterization of Phonation Type Using Amplitude of Flow Glottogram Pulse and of Voice Source Fundamental. *Journal of Voice* 36(1). 4–14. <https://doi.org/10.1016/j.jvoice.2020.03.018>.
- Tagliamonte, Sali A. & R. Harald Baayen. 2012. Models, forests, and trees of York English: *Was/were* variation as a case study for statistical practice. *Language Variation and Change* 24(2). 135–178. <https://doi.org/10.1017/S0954394512000129>.
- Thurgood, Graham. 2002. Vietnamese and tonogenesis: Revising the model and the analysis. *Diachronica* 19(2). 333–363. <https://doi.org/10.1075/dia.19.2.04thu>.
- Tyler, Richard S., Quentin Summerfield, Elizabeth J. Wood & Mariano A. Fernandes. 1982. Psychoacoustic and phonetic temporal processing in normal and hearing-impaired listeners. *The Journal of the Acoustical Society of America* 72(3). 740–752. <https://doi.org/10.1121/1.388254>.
- Uchihara, Hiroto. 2016. Tone and registrogenesis in Quiaviní Zapotec. *Diachronica* 33(2). 220–254. <https://doi.org/10.1075/dia.33.2.03uch>.

- Uchihara, Hiroto & Gabriela Pérez Báez. 2016. Fortis/lenis, glides and vowels in Quiav-
iní Zapotec. *Glossa: A Journal of General Linguistics* 1(1). 27. <https://doi.org/10.5334/gjgl.13>.
- Villard, Stéphanie. 2015. *The Phonology and Morphology of Zacatepec Eastern Chatino*. Austin, TX: University of Texas at Austin dissertation.
- Warren, Paul. 2018. Quality and quantity in New Zealand English vowel contrasts. *Journal of the International Phonetic Association* 48(3). 305–330. <https://doi.org/10.1017/S0025100317000329>.
- Weller, Jae, Jeremy Steffman, Félix Cortés & Iara Mantenuto. 2023a. Interactions of tone, glottalization, and word shape in San Sebastián Del Monte Mixtec.
- Weller, Jae, Jeremy Steffman, Félix Cortés & Iara Mantenuto. 2023b. Lexical tone and vowel duration in San Sebastián del Monte Mixtec. *The Journal of the Acoustical Society of America* 153. A293. <https://doi.org/10.1121/10.0018900>.
- Weller, Jae, Jeremy Steffman, Félix Cortés & Iara Mantenuto. 2024. Voice Quality and Tone in San Sebastián del Monte Rearticulated and Modal Vowels.
- Wieling, Martijn. 2018. Analyzing dynamic phonetic data using generalized additive mixed modeling: A tutorial focusing on articulatory differences between L1 and L2 speakers of English. *Journal of Phonetics* 70. 86–116. <https://doi.org/10.1016/j.wocn.2018.03.002>.

- Wood, Simon N. 2017. *Generalized Additive Models: An Introduction with R*. 2nd edn. (Texts in Statistical Science Series). Boca Raton, FL: Chapman and Hall/CRC. <https://doi.org/10.1201/9781315370279>.
- Yip, Moira Jean Winsland. 2002. *Tone* (Cambridge Textbooks in Linguistics). Cambridge ; New York: Cambridge University Press. 341 pp.
- Zapotec Language Project — University of California, Santa Cruz*. 2022. <https://zapotec.ucsc.edu/> (6 December, 2022).

Appendix A

Some Ancillary Stuff

Ancillary material should be put in appendices, which appear after the bibliography.