

A Refined Mechanistic Model for Trajectory-Dependent Biological Response in Spaceflight

Corresponding Author: Mymana Takual

Email: mymanatakual4@gmail.com

Date: June 2025 (revised)

Abstract

This document presents a mathematical hypothesis for systemic biological changes during spaceflight, focusing on reversible dynamics such as telomere length. The model posits a trajectory-dependent biomechano-chemical response, where changes in gravitational acceleration (Δg) act as a primary driver, triggering adaptive cascades via mechanotransduction. Gravity is framed as an inertial compressional load, equivalent in free fall scenarios. The equation integrates non-linear adaptation to a new equilibrium, baseline aging, gravitational stress, and radiation damage, providing a unified, testable framework for space medicine. Aligned with NASA Twins Study data, it offers predictions for multi-transition missions like Artemis or Mars voyages.

The Foundational Concept: Gravity as an Inertial Compressional Load

Gravity imparts a constant, background compressional load on biological structures through inertial forces, manifesting as weight ($\text{mass} \times g$). On Earth ($1g$), cells experience a static preload, creating λ mechanical pre-stress in the cytoskeleton and nuclear lamina. Entry into microgravity ($0g$) removes this load, leading to a "relaxation" of cellular structures. This perturbation is detected as a primary signal, initiating systemic adaptive responses to re-establish equilibrium. The equivalence of free fall scenarios underscores that the response is driven by the effective absence of weight, not gravity per se, aligning with general relativity's principles.

The Mathematical Model: A Unified Framework

The dynamics of a biological state variable $S(t)$ (e.g., telomere length, normalized to its Earth baseline S_0) are described by:

$$\ln \frac{S(t)}{S_0} = [\alpha - \beta \Delta g] \cdot (1 - e^{-t/\tau_{aging}}) - \lambda R t$$

Where :

Symbole	Meaning	Unit /Value
$(S(t))/(S_0)$	Relative biological state	dimensionless
α	Aging magnitude (total decline as $t \rightarrow \infty$)	dimensionless (≈ 1.345)
β	Mechanotransduction coupling constant (static gain)	dimensionless (≈ 0.145 , range 0.12–0.18)

Δg	Change in gravitational acceleration ($g_{\text{new}} - g_{\text{old}}$)	m/s^2
τ_{aging}	Long-term adaptation time constant	≈ 39.2 years (from Ye et al.)
λ	Radiation sensitivity constant	$(\text{Gy} \cdot \text{day})^{-1}$
R	Radiation dose rate	Gy/day
t	Time in the new environment	days / years
ϵ	Gaussian noise term (biological variability)	mean 0, SD σ
σ	Standard deviation of noise	≈ 0.15 (from Ye et al. scatter)

Important notes:

- Radiation damage is cumulative and largely irreversible; therefore it is modeled as a separate linear term $-\lambda R t$, not multiplied by the adaptation function.
- The signed Δg captures directionality: a negative Δg (e.g., Earth $\rightarrow$ space) increases $S(t)$ (lengthening), a positive Δg (space $\rightarrow$ Earth) decreases $S(t)$ (shortening).
- β is defined as a static gain. A differential form $\beta = (dS/dt)/(d\Delta g/dt)$ is conceptually possible for gradual transitions but is not used in the present model.
- Two distinct time constants exist: τ_{aging} (≈ 39.2 years) for long-term adaptation; acute mechanotransduction (e.g., YAP/TAZ nuclear shuttling) would require a much smaller constant ($\tau_{\text{mechanics}} \approx 30$ min – 8 hours), but the present work focuses on aging and spaceflight adaptation, so only τ_{aging} appears in the main equation.

Dimensional analysis:

The left side $\ln(S/S_0)$ is dimensionless. On the right, α and $\beta\Delta g$ are constructed to be dimensionless, $\lambda R t$ is dimensionless because λ has units $(\text{Gy} \cdot \text{day})^{-1}$ and R has Gy/day , and ϵ is dimensionless by definition. Thus the equation is dimensionally consistent.

3. Interpretation and Alignment with Observations

3.1 Earth Aging (Plateau Prediction)

On Earth, $\Delta g = 0$ and radiation effects are small, so the model reduces to:

$$\ln[S(t)/S_0] = -\alpha \cdot (1 - e^{-t/\tau_{\text{aging}}}) + \epsilon$$

Fitting to the large meta-analysis of Ye et al. (2023) gives $\alpha = 1.345$ and $\tau_{\text{aging}} = 39.2$ years. The saturating function $(1 - e^{-t/\tau_{\text{aging}}})$ explains why telomere shortening is fast early in life, gradually decelerates, and reaches a plateau after approximately 60 years. This is a testable prediction that already aligns with existing data.

3.2 Spaceflight (Reversibility and Directionality)

For the Earth-to-space transition ($\Delta g = -9.8 \text{ m/s}^2$, short duration so $(1 - e^{-t/\tau_{\text{aging}}}) \approx t/\tau_{\text{aging}}$), the equation becomes approximately linear in time. Using the NASA Twins Study report of $\sim 14.5\%$ telomere lengthening for Scott Kelly, we obtain $\beta \approx 0.145$. This value is not a universal constant; it is a point estimate for healthy humans, with an expected range of 0.12–0.18.

For the return to Earth ($\Delta g = +9.8 \text{ m/s}^2$), the same β gives a negative driver, predicting rapid shortening – which was indeed observed post-flight. Thus the model captures reversibility and directionality in a single, simple equation.

3.3 Cancer as a Perceptual Shift (Hypothesis)

If a cancer cell exhibits a hyperactive mechanotransduction coupling constant ($\beta_{\text{cancer}} \gg \beta_{\text{normal}}$), its effective perceived gravity becomes $g_{\text{eff}} = g / \beta$. Even on Earth ($g = 9.8 \text{ m/s}^2$), such a cell behaves as if it were in microgravity. This could explain the uncontrolled, aggressive phenotype of cancer cells and why microgravity makes them even more aggressive. This remains a testable hypothesis: we predict that β_{cancer} should be significantly larger than β_{normal} (e.g., > 0.5) and that pharmacological inhibition of the mechanotransduction pathway (e.g., YAP/TAZ inhibitors) would reduce cancer cell aggressiveness.

4. Prediction for Future Missions

Mission Phase	Trajectory	$\Delta g \text{ (m/s}^2\text{)}$	Prediction $S(t)$ response
Short suborbital	Earth $\rightarrow$ 0g $\rightarrow$ Earth	-9.8 then $+9.8$	Rapid lengthening, then rapid shortening; minimal plateau
ISS (6 months)	Earth $\rightarrow$ 0g (prolonged)	-9.8	Quick rise to a plateau (lengthening), then slow erosion by radiation ($-\lambda R t$)
Mars transit	Earth $\rightarrow$ 0g $\rightarrow$ 0.38g	-9.8 then $+6.1$	Lengthening in transit, mild shortening after landing on Mars
Round trip to Mars	Full cycle	multiple signed Δg	Step-wise adaptation; incomplete reversibility due to radiation damage

For long-duration missions ($t \gg \tau_{\text{aging}}$), the reversible adaptation term saturates (i.e., $(1 - e^{-t/\tau_{\text{aging}}}) \approx 1$). The cumulative radiation damage then dominates the time-dependent change, leading to a slow linear decline described by $-\lambda R t$. Thus, after the initial adaptation, telomere length gradually decreases due to radiation, while the gravity-driven part remains constant.

5. Generality of the Framework (Brief Note)

The same mathematical structure can be applied to other age-related biomarkers by redefining $S(t)$ appropriately. For example:

- Muscle mass: Different muscle groups show different atrophy rates, explained by different α values while sharing the same τ_{aging} .
- Bone mineral density: The two-phase behavior (stable then declining) is naturally captured by the same equation.

These examples demonstrate the generality of the framework, but the primary validation remains focused on telomere dynamics, for which quantitative data (Ye et al., 2023; NASA Twins Study) are available.

6. Experimental Design and Falsifiability

6.1 Testable Predictions

- Non-linear adaptation: If telomere shortening continues linearly without plateauing, the $(1 - e^{-t/\tau_{\text{aging}}})$ term is invalid.
- Directionality: If the sign of Δg does not dictate the direction of change (e.g., lengthening upon return to Earth), the model fails.
- Magnitude dependence: In partial gravity (e.g., Moon, Mars), responses must scale with $|\Delta g|$.
- Reversibility: After a Δg flip, changes must reverse (accounting for radiation damage).

6.2 Stochastic Extension for Biological Variability

To account for the inherent scatter in real biological data, a Gaussian noise term ϵ is added:

$$\ln[S(t)/S_0] = [-\alpha - \beta \Delta g] \cdot (1 - e^{-t/\tau_{\text{aging}}}) - \lambda R t + \epsilon$$

Here, ϵ has zero mean and standard deviation σ . The value of σ is not a free parameter; it is estimated from experimental data (e.g., from the scatter in Ye et al., 2023, $\sigma \approx 0.15$). This addition makes the model stochastic and realistic, acknowledging that real data will always scatter around the theoretical curve.

Limitations and Future Work

The model has several limitations that should be addressed by future experiments:

Adaptation time constant for spaceflight: The value $\tau_{\text{aging}} = 39.2$ years was derived from Earth aging data (Ye et al., 2023). Whether the same time constant applies to spaceflight adaptation remains an assumption. If spaceflight adaptation is faster or slower, the value of β would need to be recalculated.

Independent parameter derivation: The parameters α and β were obtained from two independent datasets (Ye et al. and NASA Twins Study). Their mutual consistency within a single equation has not been directly tested; this is a prediction of the model.

Single-subject estimate for β : The value $\beta \approx 0.145$ [The NASA Twins Study reported a 14.5% increase in telomere length during spaceflight. Using the proposed model, this observed change corresponds to a mechanotransduction coupling constant of $\beta \approx 0.145$ (estimated from the relation $\ln(1.145) = \beta \times 9.8 \times t/\tau$). This is not a direct measurement of β , but a model-based estimate derived from the reported telomere elongation data.] comes from a single astronaut (Scott Kelly). The proposed range 0.12–0.18 is a plausible biological interval, not an empirically measured distribution. Population studies are needed to determine the true range of β in healthy humans.

Mechanistic chain not fully established: The molecular chain $\Delta g \rightarrow$ mechanotransduction $\rightarrow$ YAP/TAZ $\rightarrow$ telomerase $\rightarrow$ telomere length is partially supported by the literature but not fully proven. The model aggregates the net effect into the phenomenological parameters β and τ_{aging} ; it does not require every molecular detail to be known.

Cancer hypothesis remains speculative: The prediction that cancer cells have a hyperactive β has not yet been tested experimentally. We propose specific experiments (e.g., measuring β in cancer cell lines under clinorotation) to validate or falsify this hypothesis.

Despite these gaps, the model provides a falsifiable, testable framework that can be refined as new experimental data become available.

Application to Lunar Missions

For the Moon ($g \approx 1.62 \text{ m/s}^2$), the transition from Earth gives $\Delta g \approx -8.2 \text{ m/s}^2$. The model predicts a positive driver and telomere lengthening, but less than in full microgravity. Short Apollo missions showed minimal pathological issues, consistent with the idea that such changes are adaptive and reversible rather than permanent damage. Longer Artemis missions will provide an opportunity to test the predicted partial-gravity scaling.

Conclusion

This model elevates qualitative observations into a quantitative framework, viewing spaceflight biology as adaptive responses to physical stressors like inertial load changes. It unifies Earth aging, spaceflight effects, and cancer under a single testable equation. The model is falsifiable, includes a stochastic term for biological variability, and makes specific predictions for future lunar and Mars missions. The limitations are explicitly acknowledged, and paths for future experimental validation are proposed.

References

- Garrett-Bakelman, F. E., et al. (2019). The NASA Twins Study: A multidimensional analysis of a year-long human spaceflight. *Science*, 364(6436), eaau8650.
- Ye, Q., et al. (2023). Telomere length and chronological age across the human lifespan: A systematic review and meta-analysis of 414 study samples including 743,019 individuals. *Ageing Research Reviews*, 90, 102031.
- Makarova, J. A., et al. (2025). Methods for telomere length measurement: An update on current technologies and emerging approaches. *Frontiers in Molecular Biosciences*, 12, 1725112.
- (2025). Microgravity Therapy as Treatment for Decelerated Aging and Successful Longevity. *International Journal of Molecular Sciences*, 26(13), 6544.
- Wang, X., et al. (2023). Distinct roles of telomerase activity in age-related chronic diseases: An update literature review. *Biomedicine & Pharmacotherapy*, 167, 115553.