

Perceptron (파설트론) _ 신경망의 기원 - 다쉬 산로 입력받아 하내의 산로 걸려 bias) b; 편형 (원. 또가 얼마나 쉽게 04) 7(j : 11년선호 활성화 되내). XΙ (in Put data, node) y; ; 弱链 7(1 (out Put data) Wi : 생에 닦 영향격 数配 を指 培田 住園 (= W; プ 具名 がは) 다는 신호를 함께는 해결 ex) and. Nand. OR HOLE AND 到驻 3叶) $1\left(\chi_{1}.w_{1}+\chi_{2}.w_{2}.7b\right)=\left(\frac{1}{2}\chi_{1}w_{1}\right)$ $0\left(\chi_{1}.w_{1}+\chi_{2}.w_{2}.6b\right)$ ANd gate Nand gate $\chi_1 | \chi_2 |$ $4 = (1 (x_1 \cdot w_1 + x_2 \cdot w_2 + 6 > 0)$ (Q (x1.W1 +X2.W2 +b <0) 4= (1 (x1.w1 + x2.W2 > b) 10 (x, w, +x w2 5b) linear y(x1,x2,b) = $(x_1 \cdot w_1 + x_2 \cdot w_2 - p 70)$ (71.W. +x2.W2-b 40) wieght infut data

ex) And gate				
(J1. 72)	å (w.·W ₂ .b)			
(9.9)	$\bigcirc \cdot \omega_1 + 1 \cdot \omega_2 + b = -\circ \cdot 2$	⟨0 ⇒ 9		
(1.0)	1. W1 + O. W2 tb = -0.17	(0 >9		
(0.1)	$Q \cdot w_1 + Q \cdot w_2 + b = -1.2$	<0 ⇒0		
(1.1)	1. W1 + 1. W2+b = 0.3	>0 ⇒1		

Node

W1=05 W2=1 b=-12

1321 5271 J

문제점: 이라는 가중이. 프랑을 직접 고려 해 주나하지?

but There is a Problem

How about XOB gate?

〈给 工出 野

linear.

马姆 强强刑

$$\frac{d}{dt} = -\frac{w_1}{w_2} \chi - \frac{b}{w_2}$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$


$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$


$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$


$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$

$$= -\alpha \cdot \chi + b \qquad (a_3b = (cnstant))$$





2(1)



 $\frac{1}{\sqrt{(w_1 \cdot w_2 \cdot b)}} = \begin{cases}
4 & (x_1 \cdot w_1 + x_2 \cdot w_2 + b > 0) \\
0 & (x_1 \cdot w_1 + x_2 \cdot w_2 + b \leq 0)
\end{cases}$


그런 XOR의 Perception 의 샾 어떻게 구지?

	च्या विश्वास्त्र अस्ति । अस्त			
4	< X0P	l gat		>
(0.1)	α_{l}	χ ₂	1 4	
	0	()	9	
	1	Q	1	
	, <u> </u>	1	1	
(0.0)	1	1	9	

직선으로 구분이 안된다! > linear 32 25 94. ら un linear 로 가능。

 $\geq (w_1^*.w_2^*.b_1^*) = (1 (y_1.w_1^* + y_2.w_2^* + b_1^*) \circ (y_1.w_1^* + y_2.w_2^* + b_1^*) \circ)$

but) Perceptions 中 中國 如果



(2, w, +16, w) · W"+ (x, w, +12, w) +6 >0 $(\chi_1, w_1 + \chi_2, w_2 + b), w,'' > -(\chi_1, w_1 + \chi_2 w_2 + b)$

