

---

# High\_albedo\_beta0\_50m\_6parameters

```
clear all
close all
lat = ncread('data/high_albedo/beta0_50m.atmos.nc','lat');
tsurf = ncread('data/high_albedo/beta0_50m.atmos.nc','t_surf');
cwvp = ncread('data/high_albedo/beta0_50m.atmos.nc','WVP');
swdn_toa = ncread('data/high_albedo/beta0_50m.atmos.nc','swdn_toa');
swup_toa = ncread('data/high_albedo/beta0_50m.atmos.nc','swup_toa');
cldcvr = ncread('data/high_albedo/beta0_50m.atmos.nc','tot_cld_amt');
netrad_toa = ncread('data/high_albedo/
beta0_50m.atmos.nc','netrad_toa');
icecvr = ncread('data/high_albedo/beta0_50m.ice.nc','ice_cover');
ylat = ncread('data/high_albedo/beta0_50m.ice.nc','yt');

tsurf_ann=squeeze(mean(reshape(tsurf,[90 12 50]),2));
cwvp_ann = squeeze(mean(reshape(cwvp,[90 12 50]),2));
swdn_toa_mn = squeeze(mean(reshape(swdn_toa,[90 12 50]),2));
swup_toa_mn = squeeze(mean(reshape(swup_toa,[90 12 50]),2));
cldcvr_mn = squeeze(mean(reshape(cldcvr,[90 12 50]),2));
netrad_toa_mn = squeeze(mean(reshape(netrad_toa,[90 12 50]),2));
icecvr_mn = squeeze(mean(reshape(icecvr,[90 12 50]),2));

tsurf_ann_cel = tsurf_ann-273.15;
plan_alb = swup_toa_mn./swdn_toa_mn;
initial_cwvp = 32.02;
cwvp_ann_calc = cwvp_ann./initial_cwvp;

year = [1:1:50];
tsurf_ann_gm = zeros(1,50);
cwvp_ann_in = zeros(1,50);
plan_alb_tri = zeros(1,50);
cldcvr_mn_tri = zeros(1,50);
netrad_toa_mn_tri = zeros(1,50);
icecvr_mn_tri = zeros(1,50);

for t=1:size(tsurf_ann_cel,2)
    tsurf_ann_gm(t)=sum(cos(lat*pi/180).*tsurf_ann_cel(:,t))/
sum(cos(lat*pi/180));
    cwvp_ann_in(t) =
    sum(cos(lat*pi/180).*cwvp_ann_calc(:,t)).*100/sum(cos(lat*pi/180));
    plan_alb_tri(t)= sum(cos(lat*pi/180).*plan_alb(:,t)).*100/
sum(cos(lat*pi/180));
    cldcvr_mn_tri(t)= sum(cos(lat*pi/180).*cldcvr_mn(:,t))./
sum(cos(lat*pi/180));
    netrad_toa_mn_tri(t) =
    sum(cos(lat*pi/180).*netrad_toa_mn(:,t))./sum(cos(lat*pi/180));
    icecvr_mn_tri(t) = sum(cos(ylat*pi/180).*icecvr_mn(:,t)).*100/
sum(cos(ylat*pi/180));
end
```

```
yyaxis left;
plot (year,tsurf_ann_gm,'k','LineWidth',2);
%ax = axes;
axis ([0 50 -70 20 ]);
set (gca, 'YLim',[-70 20],'ycolor','k');
set (gca,'YTick',(-70:10:20));
set(gca,'XMinorTick','on');
set(gca,'YMinorTick','on');

ylabel('Surface Temperature (^{\circ}C)','color','k','FontSize',15);
xlabel('Year','color','k','FontSize',15);
text(10.5,-36,'\leftarrow Surface Temperature (^{\circ}C)');

hold on
yyaxis right
ylabel('Per Cent Change','color','k','FontSize',15);
set (gca, 'YLim',[0 100],'ycolor','k');
set (gca,'YTick',(0:20:100));
set(gca,'YMinorTick','on');
plot (year, cwvp_ann_in,'color',[139 203 218]./255,'LineWidth',2);
text(30,3,'\downarrow Column Water Vapor (%)' );
hold on
yyaxis right
plot(year, plan_alb_tri,'color',[225 132 68]./255,'LineWidth',2);
hline = findobj(gcf, 'type', 'line');
set(hline(1),'LineStyle','-');
text(30,77,'\downarrow Planetary Albedo (%)');
hold on
yyaxis right
plot(year, cldcvr_mn_tri,'color',[167 200 103]./255,'LineWidth',2);
hline = findobj(gcf, 'type', 'line');
set(hline(1),'LineStyle','-');
text(14,3,' \downarrow Cloud Cover (%)');
hold on
yyaxis left
plot(year, netrad_toa_mn_tri,'k','LineWidth',2);
text(25,-12,'\uparrow TOA Net Flux (W/m^2)');
hold on
yyaxis right;
plot (year,icecvr_mn_tri,'r','LineWidth',2);
hline = findobj(gcf, 'type', 'line');
set(hline(1),'LineStyle','-');
text(13,87,'\leftarrow Sea Ice Cover (%)');
title('Global Annual Mean Change');
```

*Published with MATLAB® R2017b*