
A
CGM groups agree over time, not with clinical profiles
CGM groups, days 1 to 3 vs days 4 to 6 (held-out)
CGM groups vs C1/C2/C3, days 1 to 3
CGM groups vs C1/C2/C3, days 4 to 6
0.42
kappa 0.63 [0.52, 0.74]
0.030
0.036
Permutation null, 97.5th percentile
Threshold
Refit stability
0.0
0.2
0.4
0.6
0.8
1.0
Adjusted Rand index (ARI)
B
Glucose level is more reproducible than timing
3-day windows
4-day windows
Level
Mean glucose
TIR 70 to 180
TAR > 180
TAR > 250
Variability
SD
MAGE
CV
TBR < 70
Timing and daily pattern
Autocorrelation 1 h
Daily profile PC1
Daily profile PC2
Circadian share
Daily profile PC3
Autocorrelation 3 h
Daily profile PC4
Autocorrelation 24 h
0.0
0.2
0.4
0.6
0.8
1.0
ICC between windows
C
The lower variability of C2 is stable across windows
Change between windows
−0.02 [−0.97, 0.89]
1
0
−1
−2
−3
−4
−5
CV, C2 minus other profiles (percentage points)
−2.33
−2.31
−2.38
Days 1 to 3
Days 4 to 6
Full record