
Reverse direction: grouping by CGM first, non-insulin T2D (n = 458)
A
CGM groups vs clinical profiles
ARI 0.039 [0.020, 0.068]
CGM group (extended, full record)
G1 (n=188)
G2 (n=182)
G3 (n=88)
85
66
37
82
28
72
25
17
46
C1
C2
C3
Clinical profile
B
Forward (CGM across C1 to C3) vs reverse (clinical across G1 to G3)
Mean glucose, 48 h
Time in range, 48 h
CV, 48 h
Mean glucose, full
Time in range, full
CV, full
Age
BMI
C-peptide
TG/HDL
Waist-to-hip
0.06
Forward
Reverse
0.00 0.02 0.04 0.06 0.08 0.10 0.12 0.14
Eta squared [95% CI]
HbA1c (reverse, expected)
0.30 0.35 0.40 0.45 0.50
C
Predicting C1 to C3 from CGM
(grey: permutation null; dotted: chance)
HbA1c plus CGM
CGM features
HbA1c alone
Cross-validated score
0.7
0.6
0.5
0.4
0.3
0.65
Balanced accuracy
(permutation p = 0.005)
Macro AUROC
(permutation p = 0.005)
D
Pooled 1,591: CGM clusters vs diagnosis and vs profile
Within-stratum null mean
Diagnosis (4 strata)
Stratum and profile (10)
C1 to C3, non-insulin T2D only
ARI [95% CI]
0.20
0.15
0.10
0.05
0.00
k = 2
k = 3
k = 4