
A Overlap between clinical and CGM groups
ARI 0.039 (95% CI 0.020 to 0.068)
G1 (n = 188)
G2 (n = 182)
G3 (n = 88)
CGM group
85 45%
66 35%
37 20%
82 45%
28 15%
72 40%
25 28%
17 19%
46 52%
C1
C2
C3
Clinical profile
60
40
20
0
Within CGM group (%)
B Differences across the two groupings
Full record
48 h
CGM outcomes across clinical profiles
Mean glucose
Time in range
Coefficient of variation
Clinical variables across CGM groups
Age
BMI
C-peptide
TG/HDL ratio
Waist-to-hip ratio
0.00 0.05 0.10 0.15
Rank-based effect size (η²)
HbA1c
0.30 0.35 0.40 0.45 0.50
η², HbA1c (separate scale)
C Prediction of clinical profiles
Chance 1/3
Chance 0.50
HbA1c
CGM
HbA1c + CGM
0.3 0.4 0.5
Balanced accuracy
0.5 0.6 0.7
Macro AUROC
Macro AUROC gain of HbA1c + CGM over HbA1c:
+0.055 (95% CI 0.017 to 0.094)
D Agreement across glycemic strata
Adjusted Rand index
0.20
0.15
0.10
0.05
0.00
2 3 4
Number of CGM groups
Enrolment stratum
Stratum + clinical profile
Within-stratum permutation mean
Clinical profile within non-insulin T2D