

¹ Dr. Cipto Mangunkusumo National General Hospital, Jakarta, Indonesia · ² Occupational Medicine Residency Program, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

International consensus indicators achieved

Framework Scott's three-tiered model;² 25-item Indonesian SVES;³ 11 consensus indicators¹

OSH Unit: Environmental Health and Hospital Occupational Safety and Health Installation (IKL-K3RS) · PFA: Psychological First Aid · SVEST: Second Victim Experience and Support Tool



Construct	Mean	SD	Range
Psychological distress	1.21	0.42	0-4
Physical distress	1.19	0.41	0-4
Colleague support	3.12	0.41	1-4
Supervisor support	3.12	0.41	1-4
Institutional support	3.19	0.42	1-4
Non-work-related support	3.68	0.42	1-4
Professional self-efficacy	3.12	0.41	1-4

- Lost workdays, SV vs total **NEW**

4. Scott SD, Hirschinger LE, Cox KR, McCoig M, Brandt J, Hall LW. The natural history of recovery for the healthcare provider "second victim" after adverse patient events. *Qual Saf Health Care*. 2009;18(5):325-30.
5. Mira J, Carillo I, Tella S, Vanhaecht K, Panella M, Seys D, et al. The European Researchers' Network Working on Second Victim (ERNST) policy statement on the second victim phenomenon for increasing patient safety. *Public Health Rev*. 2024;45:160717.
6. Burlinson JD, Quillivan RR, Scott SD, Johnson S, Hoffman JM. The effects of the second victim phenomenon on work-related outcomes: connecting self-reported caregiver distress to turnover intentions and absenteeism. *J Patient Saf*. 2021;17(3):195-9.