

Ghost Cycles of the Syracuse Map: 2-Adic Periodic Orbits and the Exceptional Set

Adam McKenna (adam@mysticflounder.ai)

March 2026

Abstract

We study the Syracuse map $S(n) = (3n + 1)/2^{v_2(3n+1)}$ on odd integers through its transfer operator $\mathcal{L}$ on $C(\mathbb{Z}_2^{\text{odd}})$ and its finite transfer matrices P_k . We prove $\|\mathcal{L}\| = 2/3$ and $\rho(\mathcal{L}) \leq 1/2$, and establish two independent obstructions to spectral gap methods: $\mathcal{L}$ does not preserve any Hölder or Lipschitz space on $(\mathbb{Z}_2^{\text{odd}}, |\cdot|_2)$, and $\mathcal{L}$ is unbounded on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{Q}_2)$ with $\|P_k\|_{2\text{-adic}} = 2^{k+O(1)} \rightarrow \infty$, closing the Mahler/Amice program for 2-adic function spaces. A third obstruction, proved by non-equicontinuity, shows $\mathcal{L}$ is also not compact on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{R})$. All three obstructions trace to a common root cause: the weight 2^{-v} is archimedeanly small but 2-adically large. Our central result is that “ghost cycles” — extra modular cycles beyond the fixed point $\{1\}$ — are not transient artifacts: exhaustive cycle enumeration through $k = 36$ and algebraic analysis through $k = 200$ show they are projections of true 2-adic periodic orbits with negative rational elements in all computed cases (for case-(a) ghost types with $D < 0$). Case-(a) ghosts persist at arithmetic progressions of levels, making the exceptional set E infinite with density $\geq 4\%$. A census identifies 88+ materializing ghost types through cycle length $L = 12$, organized into families by excess $e = V - L$, with record spectral radius $\rho \geq 2^{-16/15} \approx 0.4774$. We propose four replacement conjectures. Conjecture~4 (negative rationality) asserts that all orbit elements of $D < 0$ ghost types are negative rationals — verified through $L = 12$ for 5,996 cases and proved unconditionally for all concentrated patterns via an explicit closed-form formula — providing strong evidence that the entire high-spectral-radius regime ($\rho > 1/3$, equivalently $D < 0$) consists of purely negative 2-adic orbits with no positive-integer elements, as Conjecture~4 asserts. All computations are reproducible from the accompanying open-source repository.

Keywords: Collatz conjecture, Syracuse map, transfer operator, ghost cycles, 2-adic periodic orbits, spectral radius, exceptional set, Mahler basis, p -adic analysis

MSC 2020: 11B83, 37P20, 47B38, 11S80

1 Introduction

The Collatz conjecture, concerning the iteration $n \mapsto (3n + 1)/2^{v_2(3n+1)}$ on odd integers, remains one of the most prominent open problems in number theory. We study the map through its transfer matrices P_k on odd residues modulo 2^k and the associated transfer operator $\mathcal{L}$ on the odd 2-adic integers $\mathbb{Z}_2^{\text{odd}}$.

Our central discovery is that the “ghost cycles” — extra modular cycles beyond the fixed point $\{1\}$ — are not transient artifacts of modular reduction. They are the modular projections of true periodic orbits of S on the 2-adic integers, and case-(a) ghosts (those whose rational orbit matches the prescribed valuation pattern exactly; defined in Section 7) persist at arithmetic progressions of

levels, making the exceptional set E infinite with positive density. This falsifies an initial conjecture, made during the development of this work, that E has density zero and $\rho_k \rightarrow 1/4$.

Our contributions. We present: (1) the transfer operator framework on $C(\mathbb{Z}_2^{\text{odd}})$, including $\|\mathcal{L}\| = 2/3$, $\rho(\mathcal{L}) \leq 1/2$, and a six-part spectral theorem; (2) a proof that $\mathcal{L}$ does not preserve $\text{Lip}_1(\mathbb{Z}_2^{\text{odd}})$ or any Hölder space, obstructing the Lasota–Yorke approach (Theorem~2); (3) a proof that $\mathcal{L}$ is unbounded on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{Q}_2)$ with $\|P_k\|_{2\text{-adic}} = 2^{k+O(1)}$, closing the Mahler/Amice program for nuclearity or compactness in the 2-adic setting (Theorem~3); a third archimedean obstruction — non-compactness of $\mathcal{L}$ on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{R})$, proved by non-equicontinuity — is established in Section~12 (Proposition~6); (4) exhaustive cycle enumeration through $k = 36$ ($2^{35} \approx 3.4 \times 10^{10}$ residues), extending prior searches; (5) identification of ghost cycles as 2-adic periodic orbits with negative rational elements in all computed cases, together with a persistence theorem for case-(a) ghosts; (6) a census of 88+ materializing ghost types (distinct v -patterns across 18 denominator values) through cycle length $L = 12$, organized into families by excess $e = V - L$; (7) density $\delta(E) \geq 4\%$ unconditionally and spectral radius $\rho(\mathcal{L}) \geq 2^{-16/15} \approx 0.4774$ from known ghosts; (8) four replacement conjectures for the density of E , the spectral radius of $\mathcal{L}$, negative rationality of ghost orbits, and the universal case-(a) property; and (9) an unconditional proof (Theorem~6) that all concentrated ghost types $(1, \dots, 1, e+1)$ with $D < 0$ have purely negative rational orbit elements, establishing Conjecture~4 for this family and the universal case-(a) property for concentrated patterns as a byproduct.

The framework extends to the parametric family $S(n) = (xn + y)/2^{v_2(xn+y)}$; see the companion paper for the 2-adic local constancy of transfer matrices in the multiplier x .

All code and data are available at <https://github.com/mysticflounder/collatz>.

Related work. Comprehensive surveys of the Collatz problem and its history are given by Lagarias (1985, 2021). The transfer matrix approach to Collatz-type maps originates with Matthews and Watts (1985), who studied growth rates via Markov chain models, and Wirsching (1998), who developed the dynamical systems perspective systematically. Lagarias and Weiss (1992) introduced stochastic models that predict the heuristic contraction rate. Tao (2022) proved that almost all orbits attain almost bounded values using a probabilistic approach different from transfer matrices. The cycle equation (our Theorem~4) appears in Steiner (1977). Dhiman and Pandey (2026) independently use the term “ghost cycles” for 2-adic periodic orbits of the $3x+1$ map; our work differs in computing the density of exceptional levels and classifying ghost persistence. The Baker–Wüstholz (1993) bounds on linear forms in logarithms, as refined by Laurent (2008), provide unconditional results on ghost cycle lengths (our Propositions~4–5). Kontorovich and Lagarias (2010) study the density of the exceptional set for generalized Collatz maps; our density results (Section 9) address the same question for the standard map using ghost cycle periodicity rather than stochastic models. Assani (2024) treats the Collatz map as a non-singular transformation. Mori (2024) and Neklyudov (2024) study operator-theoretic approaches to Collatz, working on different function spaces; our analysis identifies two independent obstructions (Theorems~2 and~3) that apply to all such approaches using 2-adic regularity or 2-adic function spaces.

Outline. Section 2 defines the objects of study. Section 3 develops the transfer operator on $\mathbb{Z}_2^{\text{odd}}$, establishing the operator norm, spectral radius bound, and the spectral theorem. Section 4 proves the Lasota–Yorke obstruction. Section 5 proves the 2-adic unboundedness obstruction, closing the Mahler/Amice program in the 2-adic setting. Section 6 presents results of exhaustive enumeration of the exceptional set through $k = 36$. Section 7 establishes ghost cycles as 2-adic periodic orbits, including the cycle equation, the case-(a)/(b) classification, and the persistence

theorem together with Baker–Wüstholz bounds. Section 8 presents the census of materializing ghost types organized by family. Section 9 gives the density and spectral radius results together with replacement conjectures, and an unconditional proof of negative rationality for concentrated patterns (Theorem~6). Section 10 presents eigenvalue spectra. Section 11 describes computational methodology. Section 12 discusses the results and directions for future work.

2 Definitions and Setup

Definition 1 (Syracuse map). *The **Syracuse map** on odd positive integers is*

$$S(n) = \frac{3n+1}{2^{v_2(3n+1)}},$$

where $v_2(m)$ denotes the 2-adic valuation of m (the largest power of 2 dividing m). Since $3n+1$ is even for odd n , $v_2(3n+1) \geq 1$ and $S(n)$ is always an odd integer.

Definition 2 (Modular Syracuse map). *At resolution $k \geq 2$, let $R_k = \{1, 3, 5, \dots, 2^k - 1\}$ be the set of $N = 2^{k-1}$ odd integers less than 2^k . The **modular Syracuse map** $S_k : R_k \rightarrow R_k$ is defined by*

$$S_k(j) = S(j) \bmod 2^k, \quad j \in R_k.$$

Definition 3 (Transfer matrix). *The **transfer matrix** P_k is the $N \times N$ matrix encoding S_k with contraction weights. For each odd residue $j \in R_k$, let $v_j = v_2(3j+1)$ and $t_j = S_k(j)$. Then P_k has a single nonzero entry 2^{-v_j} in column j at the row corresponding to t_j . All other entries are zero.*

Since each column has exactly one nonzero entry, the functional graph of P_k decomposes into cycles and trees. Each cycle of length L with valuations $v_1, \dots, v_L$ and total valuation $V = \sum v_i$ contributes eigenvalues that are the L th roots of $\prod 2^{-v_i} = 2^{-V}$.

Definition 4 (Spectral radius). *The **spectral radius** of P_k is*

$$\rho_k = \max_{\text{cycles}} 2^{-V/L},$$

where the maximum is over all cycles in the functional graph. This equals the linear-algebraic spectral radius $\max |\lambda_i|$ because P_k has the special structure of a weighted permutation matrix restricted to its recurrent classes.

Definition 5 (Exceptional set). *The **exceptional set** is*

$$E = \{k \geq 3 : P_k \text{ has a cycle other than the fixed point } \{1\}\}.$$

At non-exceptional levels, $\rho_k = 1/4$ (from the fixed point $\{1\}$ with $v = 2$). At exceptional levels, $\rho_k > 1/4$.

Definition 6 (Ghost cycle). *A **ghost cycle** at level k is any cycle of S_k in R_k other than the fixed point $\{1\}$. Equivalently, $k \in E$ if and only if S_k has at least one ghost cycle. As shown in Section 7, every ghost cycle is the modular projection of a genuine periodic orbit of S on $\mathbb{Z}_2^{\text{odd}}$ whose elements are negative rationals.¹*

¹The term "ghost cycle" was used independently by Zöbelein (2024) in a different sense: patterns in the k_a parametrization of Collatz sequences that satisfy cycle conditions within a larger non-cycling sequence, not 2-adic periodic orbits. Our usage follows Dhiman and Pandey (2026).

Proposition 1 (Valuation distribution). *For odd n chosen uniformly at random modulo 2^k with k large,*

$$P(v_2(3n+1) = j) = 2^{-j}, \quad j = 1, 2, 3, \dots$$

The mean valuation is $\mathbb{E}[v] = 2$.

Proof. For odd n , the value $3n+1$ is even. The residue $3n+1 \bmod 2^{j+1}$ is uniformly distributed over even residues as n ranges over odd residues modulo 2^{j+1} (since multiplication by 3 is a bijection on $\mathbb{Z}/2^{j+1}\mathbb{Z}$ restricted to odd residues). The event $v_2(3n+1) = j$ holds if and only if $3n+1 \equiv 2^j \pmod{2^{j+1}}$, which occurs with probability $1/2^j$ among odd n modulo 2^{j+1} . Summing: $\mathbb{E}[v] = \sum_{j=1}^{\infty} j/2^j = 2$. $\square$

This folklore result appears in Lagarias and Weiss (1992) and underlies the heuristic prediction that the Syracuse map contracts by an average factor of $3 \cdot 2^{-2} = 3/4$ per step, as used by Tao (2022).

The Syracuse map embeds in the parametric family $S(n) = (xn+y)/2^{v_2(xn+y)}$ for odd parameters x, y . The net growth rate per step is $x \cdot 2^{-v}$, so the expected log-growth is $\log_2 x - \mathbb{E}[v] = \log_2 x - 2$, which changes sign at $x = 4$. For the remainder of this paper we work exclusively with $x = 3, y = 1$; the companion paper establishes 2-adic local constancy of $P_k(x, y)$ in the multiplier x .

3 The Transfer Operator on $\mathbb{Z}_2^{\text{odd}}$

The Syracuse map S extends continuously to $\mathbb{Z}_2^{\text{odd}} = 1 + 2\mathbb{Z}_2$, the odd 2-adic integers, with the sole exception $n = -1/3 \in \mathbb{Z}_2^{\text{odd}}$ where $3n+1 = 0$ and S maps outside the domain; since $-1/3$ has no preimages under S , we have $(\mathcal{L}f)(-1/3) = 0$ for all f and the operator analysis is unaffected. The transfer operator (Perron–Frobenius operator) associated to S acts on continuous functions $f: \mathbb{Z}_2^{\text{odd}} \rightarrow \mathbb{R}$:

$$(\mathcal{L}f)(n) = \sum_{S(m)=n} 2^{-v_2(3m+1)} f(m).$$

Remark (Projective limit definition). $\mathcal{L}$ is the projective limit of the finite transfer matrices P_k : the sum runs over odd integer preimages, i.e., $m \in R_k = \{1, 3, \dots, 2^k - 1\}$ satisfying $S_k(m) \equiv n \pmod{2^k}$, in the limit $k \rightarrow \infty$. Non-integer 2-adic elements are not preimage candidates; equivalently, m ranges over odd 2-adic integers that are genuine preimages of S as an endomorphism of $\mathbb{Z}_2^{\text{odd}}$ (at finite level k , these are positive odd integers in R_k). Each valid m corresponds to a branch $m = g_v(n) = (n \cdot 2^v - 1)/3$ for some $v \geq 1$, with weight 2^{-v} .

Lemma 1 (Preimage structure). *For each odd 2-adic integer n , the preimages of n under S are $m_v = (n \cdot 2^v - 1)/3$ for those $v \geq 1$ where $3 \mid (n \cdot 2^v - 1)$. Since $2 \equiv -1 \pmod{3}$:*

- (a) $n \equiv 1 \pmod{3}$: preimages at even $v = 2, 4, 6, \dots$
- (b) $n \equiv 2 \pmod{3}$: preimages at odd $v = 1, 3, 5, \dots$
- (c) $n \equiv 0 \pmod{3}$: no preimages.

Each m_v is odd, lies in $\mathbb{Z}_2^{\text{odd}}$, and satisfies $v_2(3m_v + 1) = v$.

Proof. A preimage m satisfies $3m+1 = n \cdot 2^v$, i.e., $m = (n \cdot 2^v - 1)/3$. For integrality, we need $n \cdot 2^v \equiv 1 \pmod{3}$. Since $2 \equiv -1 \pmod{3}$, we have $2^v \equiv (-1)^v \pmod{3}$. This gives $n \cdot (-1)^v \equiv 1 \pmod{3}$, yielding the three cases above.

When $m = (n \cdot 2^v - 1)/3$ exists, parity: m is odd iff $n \cdot 2^v - 1 \equiv 3 \pmod{6}$ iff $n \cdot 2^v \equiv 4 \pmod{6}$. Since $2^v \pmod{6}$ alternates as 2, 4, 2, 4, ... for $v = 1, 2, 3, 4, \dots$, the parity condition coincides exactly with the integrality condition. So whenever m is an integer, it is automatically odd.

On $\mathbb{Z}_2$, division by 3 is well-defined since 3 is a 2-adic unit ($|3|_2 = 1$). For each valid v , $m_v = (n \cdot 2^v - 1) \cdot 3^{-1}$ is a well-defined odd element of $\mathbb{Z}_2$. $\square$

Remark. *The image of S is always coprime to 3: since $3m + 1 \equiv 1 \pmod{3}$, we have $S(m) \not\equiv 0 \pmod{3}$. Consequently $(\mathcal{L}f)(n) = 0$ for all $n \equiv 0 \pmod{3}$.*

Proposition 2 (Operator norm). *The operator norm of $\mathcal{L}$ on $(C(\mathbb{Z}_2^{\text{odd}}), \|\cdot\|_\infty)$ is*

$$\|\mathcal{L}\|_{C^0 \rightarrow C^0} = 2/3.$$

More precisely, the weight function $W(n) := (\mathcal{L}\mathbf{1})(n)$ satisfies:

$$W(n) = \begin{cases} 1/3 & n \equiv 1 \pmod{3} \\ 2/3 & n \equiv 2 \pmod{3} \\ 0 & n \equiv 0 \pmod{3}. \end{cases}$$

Proof. By Lemma 1:

Case $n \equiv 1 \pmod{3}$:

$$W(n) = \sum_{v=2,4,6,\dots} 2^{-v} = \frac{1/4}{1 - 1/4} = \frac{1}{3}.$$

Case $n \equiv 2 \pmod{3}$:

$$W(n) = \sum_{v=1,3,5,\dots} 2^{-v} = \frac{1/2}{1 - 1/4} = \frac{2}{3}.$$

Case $n \equiv 0 \pmod{3}$: $W(n) = 0$ (no preimages).

The bound $|(\mathcal{L}f)(n)| \leq W(n)\|f\|_\infty \leq (2/3)\|f\|_\infty$ is achieved by $f = \mathbf{1}$ and $n \equiv 2 \pmod{3}$. $\square$

Remark. *The value $\|\mathcal{L}\| = 2/3$ corrects the claim $\|\mathcal{L}\| = 1/3$ that appeared in earlier versions of this work. The discrepancy arose from considering only preimages with even v (the $n \equiv 1 \pmod{3}$ case), which give weight sum $1/3$. The full weight sum at $n \equiv 2 \pmod{3}$ (preimages with odd v) is $2/3$.*

Proposition 3 (Spectral radius bound). $\rho(\mathcal{L}) \leq 1/2$ on $C(\mathbb{Z}_2^{\text{odd}})$.

Proof. Since $\mathcal{L}$ is a weighted composition operator and P_k represents its restriction to A_k , each eigenvalue of P_k arises from a cycle of S_k ; in the projective limit these correspond to periodic orbits of S . Every such eigenvalue has the form $\lambda = \prod_{\text{cycle}} 2^{-v_i}$. Since each $v_i \geq 1$, the mean valuation $\bar{v} \geq 1$, giving $|\lambda| = 2^{-\bar{v}} \leq 1/2$. The spectrum of $\mathcal{L}$ is the closure of the union of eigenvalues of the finite approximations P_k (Theorem~1(e) below, proved independently via Stone–Weierstrass), so $\rho(\mathcal{L}) \leq 1/2$. $\square$

Theorem 1 (Spectral properties of $\mathcal{L}$ on $C(\mathbb{Z}_2^{\text{odd}})$).

(a) $\mathcal{L}$ is bounded with $\|\mathcal{L}\| = 2/3$.

(b) $\rho(\mathcal{L}) \leq 1/2$.

- (c) $\lambda = 1/4$ is an eigenvalue with eigenfunction $\delta_1 := \mathbf{1}_{\{1\}}$ (the indicator of the clopen set $\{1\}$); simplicity is verified computationally (no other cycle with $\prod 2^{-v_i} = 1/4$ materializes through $k = 36$).
- (d) $\sigma(\mathcal{L}) \subseteq \{|\lambda| \leq 1/2\}$.
- (e) $\sigma(\mathcal{L}) = \overline{\bigcup_{k \geq 2} \sigma(P_k)}$, where P_k is the transfer matrix on odd residues mod 2^k .
- (f) For non-exceptional k (verified computationally for all non-exceptional $k \leq 36$): $\sigma(P_k) = \{0, 1/4\}$.

Proof. Parts (a)–(b) are Propositions 2–3 above.

For (c): write $\delta_1 = \mathbf{1}_{\{1\}}$ for the indicator of $\{1\}$, which is continuous on $\mathbb{Z}_2^{\text{odd}}$ since $\{1\}$ is clopen. The fixed point $S(1) = 1$ with $v_2(4) = 2$ gives $(\mathcal{L}\delta_1)(n) = (1/4)\delta_1(n)$. Simplicity is verified computationally: no other cycle with $\prod 2^{-v_i} = 1/4$ materializes at any level $k \leq 36$, so no other eigenvalue of that magnitude appears in $\bigcup_{k \leq 36} \sigma(P_k)$.

Part (d) is immediate from (b).

For (e): let $A_k \subset C(\mathbb{Z}_2^{\text{odd}})$ be the subspace of functions constant on odd residues modulo 2^k . Each A_k is finite-dimensional with $\dim A_k = 2^{k-1}$, and $\mathcal{L}$ maps A_k into $A_{k'}$ for some $k' \geq k$. The restriction of $\mathcal{L}$ to A_k , composed with projection back to A_k , is represented by P_k . Any eigenvalue of P_k is an approximate eigenvalue of $\mathcal{L}$ (the corresponding function in A_k gives $\|\mathcal{L}f - \lambda f\|$ bounded by the coarsening error, which vanishes in the projective limit). Conversely, since $P_k f \rightarrow \mathcal{L}f$ in norm for every locally constant $f \in \bigcup_k A_k$ (immediate from the construction), and $\overline{\bigcup_k A_k} = C(\mathbb{Z}_2^{\text{odd}})$ by density of locally constant functions (Stone–Weierstrass), norm-resolvent convergence on this dense domain implies that every approximate eigenvalue of $\mathcal{L}$ is approximable by eigenvalues of the P_k . Hence $\sigma(\mathcal{L}) = \overline{\bigcup_{k \geq 2} \sigma(P_k)}$. In particular, $\rho(\mathcal{L}) = \sup_k \rho_k$ (eigenvalues of P_k are L th roots of 2^{-V} with $V \geq L$, so $|\lambda| = 2^{-V/L} \leq 1/2$, giving $\rho_k \leq 1/2$ for all k).

Part (f) is verified by dense eigenvalue computation. $\square$

Remark (Sup equals limsup). *The equality $\sup_k \rho_k = \limsup_{k \rightarrow \infty} \rho_k$ follows from Theorem 5 below: case-(a) ghosts reappear at arithmetic progressions of levels, so any $\rho_k > 1/4$ is achieved at infinitely many k . Theorem 5 is proved independently of Theorem 1.*

Remark (Conditional spectral gap). *If E were finite, Theorem 1(e) would give $\rho(\mathcal{L}) = 1/4$: for non-exceptional k , $\sigma(P_k) = \{0, 1/4\}$, and finitely many exceptional levels contribute only isolated eigenvalues that do not accumulate. However, Sections 6–9 show E is infinite with density $\geq 4\%$. Case-(a) ghosts persist at arithmetic progressions of levels, and the $V = L + 1$ ghost family produces spectral radii $\rho = 2^{-(L+1)/L} \rightarrow 1/2$. Current bounds: $2^{-16/15} \leq \rho(\mathcal{L}) \leq 1/2$, where the lower bound comes from the $L = 15$, $V = 16$ ghost (the $V = L + 1$ family table in Section 8), which materializes and persists, contributing $\rho_k \geq 2^{-16/15}$ at infinitely many levels k by Theorem 1(e).*

4 The Lasota–Yorke Obstruction

A standard approach to proving spectral gap for transfer operators is the Lasota–Yorke inequality: find a Banach space $X \hookrightarrow C(\mathbb{Z}_2^{\text{odd}})$ such that $\mathcal{L}$ maps X to itself with

$$\|\mathcal{L}f\|_X \leq \alpha \|f\|_X + B \|f\|_{C^0},$$

where $\alpha < \|\mathcal{L}\|_{C^0}$. This gives quasi-compactness with essential spectral radius $\leq \alpha$, reducing the spectral analysis to finitely many isolated eigenvalues. The most natural candidate is the Lipschitz space $\text{Lip}_1(\mathbb{Z}_2^{\text{odd}})$ with respect to the 2-adic metric, since each inverse branch of S contracts 2-adically: for each valid $v \geq 1$, the inverse branch $g_v(n) = (n \cdot 2^v - 1)/3$ satisfies $|g_v(x) - g_v(y)|_2 = 2^{-v}|x - y|_2$ (because $|3|_2 = 1$).

Theorem 2 (Non-preservation of Lip_1). *The transfer operator $\mathcal{L}$ does not map $\text{Lip}_1(\mathbb{Z}_2^{\text{odd}})$ into itself. Specifically:*

- (a) *The constant function $f = \mathbf{1}$ has $|\mathbf{1}|_{\text{Lip}} = 0$.*
- (b) *$\mathcal{L}(\mathbf{1}) = W$ has $|W|_{\text{Lip}} = \infty$.*

Therefore no Lasota–Yorke inequality $|\mathcal{L}f|_{\text{Lip}} \leq \alpha|f|_{\text{Lip}} + B\|f\|_{\infty}$ can hold.

Proof. Part (a) is immediate. For (b), for each even $N \geq 2$, set $x_N = 1$ and $y_N = 1 + 2^N$. Then:

- Both are odd (since 2^N is even).
- $|x_N - y_N|_2 = 2^{-N}$.
- $x_N \equiv 1 \pmod{3}$, so $W(x_N) = 1/3$.
- $y_N = 1 + 2^N \equiv 1 + (-1)^N \equiv 2 \pmod{3}$ (since N is even), so $W(y_N) = 2/3$.

Therefore:

$$\frac{|W(x_N) - W(y_N)|}{|x_N - y_N|_2} = \frac{1/3}{2^{-N}} = \frac{2^N}{3} \rightarrow \infty.$$

This proves $|W|_{\text{Lip}} = \infty$. $\square$

Corollary 1 (Universal obstruction). *The function $W = \mathcal{L}(\mathbf{1})$ does not belong to any of the following spaces on $(\mathbb{Z}_2^{\text{odd}}, |\cdot|_2)$:*

- (a) $C^\alpha(\mathbb{Z}_2^{\text{odd}})$ for any $\alpha > 0$.
- (b) Any Banach space $X \hookrightarrow C(\mathbb{Z}_2^{\text{odd}})$ defined by a modulus-of-continuity or integral regularity condition with respect to the 2-adic metric (Hölder, Lipschitz, Zygmund, Besov, Sobolev, or BV-type spaces).

Proof. (a) The pairs from Theorem~2 give $|W(x_N) - W(y_N)|/|x_N - y_N|_2^\alpha = 2^{N\alpha}/3 \rightarrow \infty$ for any $\alpha > 0$.

- (b) Any such space requires $|f(x) - f(y)| \leq C \cdot \omega(|x - y|_2)$ for some modulus $\omega(t) \rightarrow 0$ as $t \rightarrow 0$. By (a), $|W(x_N) - W(y_N)| = 1/3$ while $|x_N - y_N|_2 = 2^{-N} \rightarrow 0$, so W violates every such condition. $\square$

Remark (Root cause: mod-3 oscillation). *The weight function $W(n)$ depends on $n \pmod{3}$. The residue $n \pmod{3}$ is a continuous function on $\mathbb{Z}_2$ (each residue class is clopen), but it is not Lipschitz: every 2-adic ball $B(x, 2^{-N})$ intersects all three residue classes mod 3 (since $\gcd(2^N, 3) = 1$). So W oscillates between 0, 1/3, and 2/3 at every scale of the 2-adic metric. This obstruction reflects the fundamental arithmetic tension in the Collatz problem: each inverse branch g_v contracts by 2^{-v} in the 2-adic metric but expands by 3 in the 3-adic metric ($|g_v(x) - g_v(y)|_3 = 3|x - y|_3$). The 3-adic expansion exactly compensates the weight sum, preventing contraction in any metric incorporating 3-adic information.*

Remark. *When $\mathcal{L}$ is restricted to pairs within the same residue class mod 3, the Lipschitz seminorm does contract: one computes $\sum_{v=1,3,5,\dots} 4^{-v} = 4/15$ and $\sum_{v=2,4,6,\dots} 4^{-v} = 1/15$, giving conditional*

contraction rates $4/15$ for pairs with $x \equiv y \equiv 2 \pmod{3}$ and $1/15$ for pairs with $x \equiv y \equiv 1 \pmod{3}$. However, the inverse branches involve division by 3, which scrambles higher powers of 3, so a global Lasota–Yorke inequality cannot be recovered from this conditional estimate.

Since standard spectral methods based on 2-adic regularity are unavailable, we now show that algebraic approaches via the Mahler basis are also obstructed.

5 The 2-Adic Unboundedness Obstruction

The Lasota–Yorke obstruction (Theorem~2) blocks spectral gap arguments based on regularity in the 2-adic metric. A natural alternative is the Mahler basis $\{\binom{x}{n}\}_{n \geq 0}$, which provides an algebraic (rather than metric) framework for functions on $\mathbb{Z}_2^{\text{odd}}$. The Amice space construction uses exponentially decaying Mahler coefficients to define nuclearity and compactness in the p -adic setting (Amice 1964, Serre 1962). We show this approach also fails, for a deeper reason: $\mathcal{L}$ is unbounded on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{Q}_2)$.

Theorem 3 (2-adic unboundedness). *The transfer operator $\mathcal{L}$ is unbounded on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{Q}_2)$ equipped with the 2-adic sup norm $\|f\|_2 = \sup_n |f(n)|_2$. Specifically, the finite-level transfer matrices satisfy $\|P_k\|_{2\text{-adic}} = 2^{k+O(1)} \rightarrow \infty$ (specifically, $\|P_k\|_{2\text{-adic}} \in \{2^k, 2^{k+1}\}$).*

Proof. Re-index the 2^{k-1} odd residues in R_k as $\{2j+1 : 0 \leq j \leq 2^{k-1}-1\}$. The matrix P_k has exactly one nonzero entry per column: $2^{-v(j)}$ in column j , where $v(j) = v_2(3(2j+1)+1)$. On $(\mathbb{Q}_2^N, \|\cdot\|_\infty)$, the operator norm of such a matrix is $\|P_k\|_2 = \max_j |2^{-v(j)}|_2 = \max_j 2^{v(j)} = 2^{\max_j v(j)}$ (each column has a single nonzero entry, and the ultrametric operator norm is $\max_j \max_i |A_{ij}|_2$).

We claim $\max_j v(j) = k + O(1)$. Write $3(2j+1)+1 = 6j+4 = 2(3j+2)$, so $v_2(6j+4) = 1 + v_2(3j+2)$. Since 3 is invertible modulo 2^{k-1} , the value $3j+2$ ranges over all residues mod 2^{k-1} as j ranges over $\{0, \dots, 2^{k-1}-1\}$. In particular, $3j+2 \equiv 0 \pmod{2^{k-1}}$ is achieved, giving $v_2(3j+2) \geq k-1$ and hence $\max_j v(j) \geq k$. For the upper bound: since $j \leq 2^{k-1}-1$ implies $3j+2 \leq 3 \cdot 2^{k-1}-1 < 2^{k+1}$, we have $v_2(3j+2) \leq k$ and hence $\max_j v(j) \leq k+1$.

Since locally constant functions of all levels are dense in $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{Q}_2)$ and the restriction of $\mathcal{L}$ to level- k functions has norm $\|P_k\|_2 \rightarrow \infty$, the operator $\mathcal{L}$ is unbounded. $\square$

Corollary 2 (Failure of the Mahler/Amice program). *The transfer operator $\mathcal{L}$ does not define a bounded operator on any 2-adic Banach space $E \hookrightarrow C(\mathbb{Z}_2^{\text{odd}}, \mathbb{Q}_2)$ containing all locally constant functions with bounded norm. In particular, $\mathcal{L}$ is not nuclear on the Amice space $\mathcal{A}(\mathbb{Z}_2, \mathbb{Q}_2)$.*

Proof. For indicator functions $\mathbf{1}_B$ of 2-adic balls of radius 2^{-k} , the preimage branches give $\|\mathcal{L}(\mathbf{1}_B)\|_2 \geq 2^{k+O(1)}$ while $\|\mathbf{1}_B\|_2 = 1$. Any space containing these indicators with bounded norm inherits the unboundedness. $\square$

Remark (Common root cause). *Both obstructions — Lasota–Yorke (Theorem 2) and 2-adic unboundedness (Theorem 3) — trace to the same arithmetic tension. Each inverse branch g_v of S carries the weight 2^{-v} , which is archimedeanly small (contraction) but 2-adically large (expansion: $|2^{-v}|_2 = 2^v$). The Lasota–Yorke obstruction manifests this as mod-3 oscillation of the weight function; the 2-adic unboundedness manifests it as divergent operator norms on 2-adic function spaces. Any spectral-theoretic approach to $\mathcal{L}$ must therefore work in the archimedean category: $\mathcal{L}$ acts on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{R})$ with operator norm $2/3$, and $\mathcal{L}$ is not compact on this space either (Proposition 6, Section 12).*

The analytical proof above establishes $\|P_k\|_{2\text{-adic}} \in \{2^k, 2^{k+1}\}$ for all k . Exact arithmetic over $\mathbb{Q}$ (Python `Fraction`) for $k = 3, \dots, 15$ further refines this to a parity pattern: $\max_j v(j) = k+1$ for odd

k and k for even k (computationally verified through $k = 15$; not required for the general bound). The Mahler matrix $M_k = C^{-1}P_kC$ (where $C[i, j] = \binom{i}{j}$) has entries with $v_2(M_k[m, j]) = -(k + \text{offset}(j))$, shifting by exactly $-\Delta k$ between levels. Since C and C^{-1} are 2-adic isometries (unipotent lower-triangular with integer entries, so $\|C\|_2 = \|C^{-1}\|_2 = 1$ in the 2-adic sup norm), the Mahler change of basis does not mitigate the unboundedness: $\|M_k\|_2 = \|P_k\|_2 = 2^{k+O(1)}$.

6 Exhaustive Cycle Enumeration

We perform exhaustive cycle enumeration for $k = 3, \dots, 36$ ($2^{35} \approx 3.4 \times 10^{10}$ residues at $k = 36$). The search uses two regimes:

- $k \leq 32$: Precomputed successor arrays (numpy uint32). Peak memory approximately 13 GB at $k = 32$.
- $k = 33$ – 36 : On-the-fly computation with Numba JIT and packed bitarrays. Peak memory approximately 4 GB at $k = 36$.

The $k = 36$ search checked $2^{35} \approx 34$ billion odd residues in approximately 14 hours.

6.1 Results

Table 1: Exceptional set E for $x = 3$, $y = 1$. Five values of k in $[3, 36]$ produce additional modular cycles beyond the fixed point $\{1\}$.

Range	$ E \cap \text{range} $	Density
$k = 3$ – 9	0	0.000
$k = 10$ – 12	3	1.000
$k = 13$ – 19	0	0.000
$k = 20$	1	—
$k = 21$ – 34	0	0.000
$k = 35$	1	—
$k = 36$	0	—
$k = 3$ – 36	5	0.147

The exceptional cycles are:

Table 2: Details of exceptional modular cycles. Even at exceptional k , the spectral radius remains well below $1/2$.

k	Extra cycles	Worst ρ	Worst cycle length	Worst mean v
10	1	0.3729	26	1.423
11	1	0.3585	25	1.480
12	2	0.4454	6	1.167
20	1	0.3886	22	1.364
35	1	0.4353	5	1.200

Ghost cycles are not transient artifacts of modular reduction. They are the modular projections of true periodic orbits of S on the 2-adic integers $\mathbb{Z}_2^{\text{odd}}$ (Figure~1).




Figure 1: Chord diagrams of the Syracuse successor map at four resolutions. Odd residues mod 2^k are arranged around a circle; each arc connects a residue to its successor. At $k = 9$ and $k = 13$ (non-exceptional), only the fixed point $\{1\}$ (gold) forms a cycle. At $k = 10$, a 26-node ghost cycle appears (cyan). At $k = 12$, two ghost cycles coexist (cyan: $L = 7$, orange: $L = 6$).

7 Ghost Cycles as 2-Adic Periodic Orbits

7.1 The Cycle Equation

Theorem 4 (Cycle equation; after Steiner 1977, Wirsching 1998²). *A modular cycle of length L at level k with valuation pattern $(v_1, \dots, v_L)$ and total valuation $V = \sum v_i$ satisfies*

$$n_1 \cdot D \equiv R \pmod{2^{k+V}},$$

where $D = 2^V - 3^L$ and $R = \sum_{i=0}^{L-1} 3^{L-1-i} \cdot 2^{\sigma_i}$, with $\sigma_0 = 0$ and $\sigma_i = v_1 + \dots + v_i$.

Proof. If the cycle visits odd residues $n_1, n_2, \dots, n_L$ with $n_{i+1} = (3n_i + 1)/2^{v_i}$, then $n_{i+1} \cdot 2^{v_i} = 3n_i + 1$ is an exact integer identity. At level k , n_{i+1} is known modulo 2^k , so this step holds modulo 2^{k+v_i} . Iterating L steps yields $n_1 \cdot 2^V = 3^L n_1 + R$, i.e., $n_1(2^V - 3^L) = R$ over $\mathbb{Z}$. With total shift $V = \sum v_i$ the precision accumulates, giving $n_1(2^V - 3^L) \equiv R \pmod{2^{k+V}}$. $\square$

Since $D = 2^V - 3^L$ is always odd (2^V is even, 3^L is odd, so their difference is odd) and nonzero (2^V and 3^L are coprime, so $2^V \neq 3^L$), the rational limit $\tilde{n}_1 = R/D$ is well-defined. For $V > L \log_2 3$, we have $D < 0$.

7.2 Case-(a) vs Case-(b) Classification

Definition 7 (Ghost type; case-(a) and case-(b)). *(Throughout, L denotes cycle length; $\mathcal{L}$ denotes the transfer operator.) A **ghost type** is a triple $(L, V, (v_1, \dots, v_L))$ where $L \geq 2$, $V = \sum v_i$ with each $v_i \geq 1$, and $D = 2^V - 3^L \neq 0$. Two ghost types are equivalent if their v -patterns are cyclic rotations of each other; henceforth, ghost types are counted up to this equivalence (each equivalence class is a canonical representative). The **denominator** of a ghost type is D ; distinct v -patterns with the same D are distinct ghost types.*

Given a ghost type with rational limit $\tilde{n}_1 = R/D$, define the **rational orbit** $\tilde{n}_1, \tilde{n}_2, \dots, \tilde{n}_L$ by $\tilde{n}_{i+1} = (3\tilde{n}_i + 1)/2^{v_i}$.

²The cycle equation in this form originates with Davison (1976), as cited by Steiner (1977). We have not been able to access the original proceedings.

- **Case (a):** $v_2(3\tilde{n}_i + 1) = v_i$ for all i . Then the rational orbit is a true periodic orbit of S on $\mathbb{Q} \cap \mathbb{Z}_2^{\text{odd}}$: since the valuation conditions determine the Syracuse map step exactly, iterating S on $\tilde{n}_1$ produces $\tilde{n}_2, \dots, \tilde{n}_L, \tilde{n}_1$.
- **Case (b):** $v_2(3\tilde{n}_i + 1) > v_i$ for some i . The rational limit has “extra” 2-adic cancellation; the orbit exists at only finitely many levels k .

The 2-adic digit structure of $\tilde{n}_1 = R \cdot D^{-1} \bmod 2^k$ stabilizes bit by bit as k increases, converging to the rational limit (Figure~2).




Figure 2: 2-adic digit stabilization for the $D = -601$ ghost ($L = 6$, $V = 7$). Each column shows the binary digits of $n_1 = R \cdot D^{-1} \bmod 2^k$. Digits stabilize from the least significant bit upward: once bit position b is determined at resolution $k = b + 1$, it never changes. The dashed line marks $k = 12$, where the ghost first materializes as a modular cycle.

Conjecture 1 (Universal case-(a)). *For all positive integers $L \geq 2$ and $L + 1 \leq V \leq 2L - 1$ (the range where $\rho = 2^{-V/L} > 1/4$), and for every composition $(v_1, \dots, v_L)$ of V into L positive parts, the rational orbit $\tilde{n}_1 = R/(2^V - 3^L)$ satisfies $v_2(3\tilde{n}_i + 1) = v_i$ for all $i = 1, \dots, L$. Equivalently, every such composition defines a true periodic orbit of the Syracuse map on $\mathbb{Z}_2^{\text{odd}}$; there are no case-(b) ghost types with $\rho > 1/4$.*

This conjecture has been verified for all $\sum_{L=2}^{15} (L - 1) = 105$ pairs (L, V) with $L = 2, \dots, 15$ and $L + 1 \leq V \leq 2L - 1$: exhaustively (all compositions) for $L \leq 12$ (up to $\sim 800,000$ compositions per pair), and by 10^6 random samples per pair for $L = 13$ – 15 . An extended survey through $L = 20$ (10^6 samples per pair, 85 million total) finds zero failures. Moreover, the case-(a) property holds even for $V \geq 2L$ (where $D > 0$), tested exhaustively through $L = 12$ at $V = 2L$ and $V = 2L + 1$ only (not a full search over all $V \geq 2L$): the sign of D does not affect the universal case-(a) property.

Remark (Towards a proof). *A proof would need to show that $v_2(3R_i + D) = v_i$ for all i and all compositions, where the R_i satisfy the orbit recurrence $R_{i+1} = (3R_i + D)/2^{v_i}$. The case-(a) condition is equivalent to each such quotient being odd; proving this uniformly across all compositions appears to require fine control on $R_i \pmod{2^{v_i+1}}$ that Baker-type bounds do not directly supply. For concentrated patterns $(1, \dots, 1, e + 1)$, this is now established unconditionally by Theorem 6 (Section 9), which derives an explicit closed form for R_i and verifies the parity conditions directly. Extending this argument to non-concentrated compositions remains open.*

7.3 Persistence of Case-(a) Ghosts

Theorem 5 (Persistence of case-(a) ghosts). *Let $(L, V, (v_1, \dots, v_L))$ be a case-(a) ghost with denominator $D = 2^V - 3^L$ and let $p = \text{ord}_2(|D|)$ be the multiplicative order of 2 modulo $|D|$. If the ghost first appears at level k_0 , then it reappears at all levels $k \equiv k_0 \pmod{p}$ with $k \geq k_0$.*

Proof. The ghost appears at level k if and only if the congruence $n_1 \cdot D \equiv R \pmod{2^{k+V}}$ has an odd solution n_1 modulo 2^k , and the resulting cycle has the prescribed valuation pattern $(v_1, \dots, v_L)$.

Since D is odd, $n_1 \equiv R \cdot D^{-1} \pmod{2^{k+V}}$, and the solution is determined by $R/D \pmod{2^k}$.

Valuation stability. The case-(a) condition states that $v_2(3\tilde{n}_i + 1) = v_i$ for all i , where $\tilde{n}_i$ are the rational orbit elements. Write $\tilde{n}_i = a_i/|D|$ with a_i an integer, $\gcd(a_i, 2) = 1$. The valuation condition $v_2(3\tilde{n}_i + 1) = v_i$ depends only on $3a_i + |D|$ modulo 2^{v_i+1} . This is independent of k .

Periodicity of the modular reduction. The residue $\tilde{n}_1 \pmod{2^k} = R \cdot D^{-1} \pmod{2^k}$ depends on $D^{-1} \pmod{2^k}$, which is periodic in k with period dividing $p = \text{ord}_2(|D|)$. Explicitly: $D^{-1} \pmod{2^{k+p}} \equiv D^{-1} \pmod{2^k} \pmod{2^k}$ because $2^p \equiv 1 \pmod{|D|}$ implies D^{-1} has a p -periodic 2-adic expansion.

Verification at level k . The cycle at level k exists if: (i) $n_1 = \tilde{n}_1 \pmod{2^k}$ is odd — this holds since $\tilde{n}_1 = R/D$ has odd numerator ($R = 3^{L-1} + \sum_{i=2}^L 3^{L-i} 2^{v_1+\dots+v_{i-1}}$: the first term is odd; all remaining terms have $v_1 + \dots + v_{i-1} \geq 1$, so are even) and odd denominator ($D = 2^V - 3^L$ is the difference of an even and an odd number); (ii) for each step i , $v_2(3n_i + 1) = v_i$ where $n_i = \tilde{n}_i \pmod{2^k}$ — this requires checking that the first v_i bits of $3n_i + 1$ match those of $3\tilde{n}_i + 1$, which holds when $k > v_i$ for all i ; and (iii) $n_{L+1} = n_1 \pmod{2^k}$ — this holds because $\tilde{n}_{L+1} = \tilde{n}_1$ (the rational orbit closes by definition), so $n_{L+1} = \tilde{n}_{L+1} \pmod{2^k} = \tilde{n}_1 \pmod{2^k} = n_1$.

Since $\tilde{n}_i \pmod{2^k}$ depends only on $k \pmod{p}$, and conditions (i)–(iii) are satisfied at k_0 , they remain satisfied at all $k \equiv k_0 \pmod{p}$ with $k \geq k_0$. $\square$

Remark (Primitivity). *The cycle at each level $k \equiv k_0 \pmod{p}$ has period exactly L , not a proper divisor. The rational orbit elements $\tilde{n}_1, \dots, \tilde{n}_L$ are distinct: materialization at level k_0 with period exactly L means the L modular residues $n_1, \dots, n_L$ are distinct modulo 2^{k_0} ; since $n_i \equiv \tilde{n}_i \pmod{2^{k_0}}$, the rational elements are pairwise distinct. Each difference $\tilde{n}_i - \tilde{n}_j$ is therefore a nonzero rational with a fixed 2-adic valuation, so $n_i \pmod{2^k} \neq n_j \pmod{2^k}$ for all $k > \max_{i \neq j} v_2(\tilde{n}_i - \tilde{n}_j)$. Since k_0 already exceeds this threshold, and since $\tilde{n}_i \pmod{2^k}$ is periodic in k with period p , the same distinctness holds at every $k \equiv k_0 \pmod{p}$.*

Remark. *For case-(b) ghosts, condition (ii) eventually fails: the “extra” valuation $v_2(3\tilde{n}_i + 1) > v_i$ means that at the rational level, the Syracuse map takes a different branch than the one prescribed by $(v_1, \dots, v_L)$. The modular cycle exists only at levels k where the extra bits are not yet visible.*

7.4 Baker–Wüstholz Bounds

Transcendence theory constrains ghost cycle denominators.

Proposition 4 (Effective lower bound on $|D|$; Baker–Wüstholz 1993, Laurent 2008, Corollary 2). *For all positive integers V, L with $V \geq 3$:*

$$|2^V - 3^L| > \max(2^V, 3^L) \cdot \exp(-25(\log V)^2).$$

In particular, $|D|$ cannot be polynomially small: for any $\epsilon > 0$, $|D| > 2^{V(1-\epsilon)}$ for all sufficiently large V .

Proof sketch. Apply Laurent (2008), Corollary 2 (real case) with $\alpha_1 = 2$, $\alpha_2 = 3$, $D = 1$, $\log A_1 = 1$, $\log A_2 = \log 3$, and $m = 30$. Table 1 of that paper gives $C_2(30) = 17.9$. With $b' = V/\log 3 + L \approx 1.54V$ (using $L \approx V/\log_2 3$), one obtains $\log b' \approx \log V + 0.43$, and for V large enough that $\log b' + 0.38 \geq 30$, the bound reduces to $\log |V \log 2 - L \log 3| \geq -17.9(\log V + 0.81)^2 \log 3 \approx -19.7(\log V)^2$. Since $|2^V - 3^L| \approx 2^V |V \log 2 - L \log 3|$ and $\max(2^V, 3^L) = 2^V$ in the ghost-cycle range, the bound with constant 25 > 19.7 follows. For V in the complementary range (below $\sim 5 \times 10^{12}$), $\exp(-25(\log V)^2)$ is smaller than $1/2^V$, so the proposition holds trivially since $|2^V - 3^L| \geq 1$. $\square$

Proposition 5 (Detection of bounded-length ghosts). *For each fixed $L_0 \geq 2$, define $K_0(L_0) = \max\{\text{ord}_2(|2^V - 3^L|) + 2L : (L, V) \text{ with } 2 \leq L \leq L_0, L+1 \leq V \leq 2L-1\}$. Then every ghost type with $L \leq L_0$ and $\rho > 1/4$ appears at some level $k \leq K_0(L_0)$: searching through $k = K_0(L_0)$ suffices to detect all such types. Explicit values: $K_0(5) \leq 279$, $K_0(10) \leq 465,259$.*

Proof. For fixed (L, V) with $\rho = 2^{-V/L} > 1/4$ (i.e., $V < 2L$), the denominator $D = 2^V - 3^L$ is fixed and has finitely many v -patterns ($\binom{V-1}{L-1}$ compositions). By Theorem~5, each case-(a) pattern reappears with period $p = \text{ord}_2(|D|)$. Since the materialization conditions depend only on $\tilde{n}_1 \bmod 2^k$, which is periodic in k with period p , if the ghost materializes at level k it also materializes at $k - p$ whenever $k - p \geq V$. Hence the first appearance satisfies $k_0 \leq p + V \leq p + 2L \leq K_0(L_0)$. Case-(b) patterns appear at finitely many levels with no periodic structure; their first appearances are bounded by direct computation of $\tilde{n}_i \bmod 2^k$ for each $(L, V, v\text{-pattern})$ within the search range. Taking the maximum over all (L, V) in the range gives $K_0(L_0)$. $\square$

Remark (Practical scope of Proposition 5). *The bounds $K_0(L_0)$ grow rapidly: $K_0(10) \leq 465,259$ already exceeds the range of exhaustive search. In practice, ghost membership beyond $k = 36$ is determined algebraically via Theorem 5 (checking $k \equiv k_0 \pmod{p}$), not by brute-force search through $k = K_0(L_0)$.*

Remark. *Propositions 4 and 5 are unconditional, but they cannot prove E is finite: that would require bounding cycle length L as a function of level k , which no known result from transcendence theory achieves.*

8 Census of Ghost Types

An extended ghost census through cycle length $L = 12$ across all 66 (L, V) pairs with $V < 2L$ (i.e., $\rho > 1/4$) identifies **157,909** canonical case-(a) ghost types. Of these, at least **88** materialize as modular cycles at some level k — a lower bound, since non-concentrated v -patterns were sampled (200 per (L, V) pair) rather than exhaustively enumerated for large parameter ranges.

The following table summarizes the materializing ghost types by denominator D . Here $p = \text{ord}_2(|D|)$ is the period of reappearance, r_{conc} is the number of residue classes mod p at which the concentrated pattern $(1, \dots, 1, e+1)$ materializes, r_{nonc} is the count for non-concentrated patterns (a lower bound for sampled (L, V) pairs), and first_k is the smallest level at which any pattern with this D appears.

Table 3: Census of materializing ghost types through $L = 12$, sorted by $|D|$. The columns r_{conc} and r_{nonc} count the number of materializing concentrated and non-concentrated v -patterns respectively; both are lower bounds for (L, V) pairs where sampling was used.

D	L	V	e	ρ	p	r_{conc}	r_{nonc}	first_k
-179	5	6	1	0.4353	178	1	0	35
-601	6	7	1	0.4454	25	1	0	12
-1675	7	9	2	0.4102	660	1	2	12
-1931	7	8	1	0.4529	1930	1	0	275
-5537	8	10	2	0.4204	84	1	0	42
-6049	8	9	1	0.4585	1441	1	0	180
-17635	9	11	2	0.4286	7052	1	2	113
-42665	10	14	4	0.3789	1716	0	6	224
-50857	10	13	3	0.4061	12714	1	11	1690
-54953	10	12	2	0.4353	9078	1	3	35
-57001	10	11	1	0.4665	5736	1	0	1147
-144379	11	15	4	0.3886	144378	0	20	434
-160763	11	14	3	0.4139	15996	0	7	515
-168955	11	13	2	0.4408	67580	1	2	2192
-400369	12	17	5	0.3746	98665	0	13	58
-498673	12	15	3	0.4204	4452	0	3	107
-515057	12	14	2	0.4454	10700	1	6	37
-523249	12	13	1	0.4719	14065	1	0	1334

Five D values — -42665 , -144379 , -160763 , -400369 , and -498673 — materialize only through non-concentrated v -patterns ($r_{\text{conc}} = 0$). This shows that restricting attention to the canonical concentrated pattern $(1, \dots, 1, e + 1)$ misses a significant fraction of materializing ghost types at higher excess values.

The full census data, including all sampled v -patterns and their materialization results, is available at <https://github.com/mysticflounder/collatz>.

8.1 The $V = L + 1$ Family

Among all ghost types with cycle length L , those with minimal valuation excess $V = L + 1$ have the largest spectral radius $\rho = 2^{-(L+1)/L}$, which approaches $1/2$ as $L \rightarrow \infty$. A systematic search for $V = L + 1$ ghosts with the canonical v -pattern $(1, 1, \dots, 1, 2)$ yields appearing ghosts at 9 of 11 fully or partially searched cycle lengths.

Table 4: The $V = L + 1$ ghost family for $L = 5, \dots, 15$. All v -patterns are algebraically case-(a). Entries $L = 5$ –13 are from complete period searches; $L = 14$ –15 are partial (r is a lower bound). The ρ column shows $2^{-(L+1)/L}$ for each L , regardless of whether the ghost materializes.

L	$ D = 3^L - 2^{L+1}$	$p = \text{ord}_2(D)$	$p/2^L$	r	First k	ρ
5	179	178	5.6	3	35	0.4353
6	601	25	0.39	1	12	0.4454
7	1,931	1,930	15	5	275	0.4529
8	6,049	1,441	5.6	10	180	0.4585
9	18,659	1,012	2.0	0	—	0.4629
10	57,001	5,736	5.6	6	1,147	0.4665
11	173,051	780	0.38	0	—	0.4695
12	523,249	14,065	3.4	4	1,334	0.4719
13	1,577,939	58,140	7.1	12	4,472	0.4740
14	4,750,201	294,712	18	≥ 4	8,087	0.4759
15	14,283,371	1,187,496	36	≥ 2	29,459	0.4774

The ghosts at $L = 9$ and $L = 11$ do not materialize at any level within their full period, despite being algebraically case-(a). A heuristic explains this: materialization requires L simultaneous binary valuation conditions on the modular reduction $R \cdot D^{-1} \bmod 2^k$, giving a target density of approximately 2^{-L} per period. The ratio $p/2^L$ thus predicts the expected number of materializations r . For $L = 11$, $p/2^L \approx 0.38$, predicting $r \approx 0$; for $L = 9$, $p/2^L \approx 2$, making $r = 0$ plausible by fluctuation. No clean algebraic criterion distinguishing appearing from non-appearing cases has been found; the phenomenon appears governed by the equidistribution of powers of 2 modulo $|D| = 3^L - 2^{L+1}$.

The nine appearing ghosts ($L = 5, 6, 7, 8, 10, 12, 13, 14, 15$) establish a proved lower bound $\limsup \rho_k \geq 2^{-16/15} \approx 0.4774$. Non-materialization at $L = 9$ and $L = 11$ appears probabilistic rather than algebraic (as the heuristic above explains), not a structural obstruction. That ghosts appear at the majority of tested cycle lengths, with non-appearance explainable by equidistribution heuristics, supports Conjecture~3 below: $\limsup \rho_k = 1/2$.

8.2 Higher Excess Families

Ghost types organize into families by the excess valuation $e = V - L$. The $V = L + 1$ family ($e = 1$) is the most extensively studied (above), but materializing ghosts also appear in the $e = 2$ and $e = 3$ families.

Notation. In the family tables below, r denotes the total number of distinct residue classes modulo p at which *any* v -pattern with the given D materializes (including both concentrated and non-concentrated patterns). This is the quantity that enters the density formula (Conjecture~2, Section 9). By contrast, the census table above lists r_{conc} and r_{nonc} , which count the number of distinct *canonical patterns* (up to cyclic rotation) that materialize. For example, $D = -1675$ has $r_{\text{conc}} = 1$, $r_{\text{nonc}} = 2$ in the census table (three canonical patterns materialize), while the family table shows $r = 9$ (nine distinct residue classes mod 660).

The $V = L + 2$ family ($e = 2$). Six denominator values produce materializing ghosts. (The “First k ” column is the first level at which the *concentrated* pattern materializes; non-concentrated patterns with the same D may appear earlier — see census table above.)

Table 5: The $V = L + 2$ ghost family for $L = 7, \dots, 12$. All v -patterns are algebraically case-(a). The ρ column shows $2^{-(L+2)/L}$ for each L , regardless of whether the ghost materializes.

L	$ D = 2^{L+2} - 3^L $	p	r	First k	ρ
7	1,675	660	9	12	0.4102
8	5,537	84	2	42	0.4204
9	17,635	7,052	5	1,567	0.4286
10	54,953	9,078	6	2,501	0.4353
11	168,955	67,580	1	8,936	0.4408
12	515,057	10,700	2	5,350	0.4454

The spectral radius $\rho = 2^{-(L+2)/L} \rightarrow 1/2$ as $L \rightarrow \infty$, though convergence is slower than for $e = 1$.

The $V = L + 3$ family ($e = 3$). One materializing type has been found with the concentrated pattern: $L = 10$, $D = -50857$, $v = (1, 1, 1, 1, 1, 1, 1, 1, 1, 4)$, with $p = 12,714$, $r = 3$, $\rho = 2^{-13/10} \approx 0.4061$.

As $L \rightarrow \infty$ with fixed excess e , the spectral radius $\rho = 2^{-1-e/L} \rightarrow 1/2$ for any fixed e . The $e = 1$ family simply converges fastest.

9 Density and Spectral Radius

9.1 Falsification of the Density-Zero Conjecture

During the development of this work, we initially conjectured that E has natural density zero and $\rho_k(3, 1) \rightarrow 1/4$, motivated by a Borel–Cantelli heuristic suggesting $|E| < \infty$. Extended computation to $k = 200$ combined with Baker–Wüstholz analysis rules out the density-zero conjecture: the exceptional set E has positive density.

The $D = -601$ ghost (case (a), $L = 6$, $V = 7$) reappears at every $k \equiv 12 \pmod{25}$, contributing $\rho \geq 2^{-7/6} \approx 0.445$ at infinitely many levels. This alone gives $\delta(E) \geq 1/25 = 4\%$.

Empirically, $|E \cap [3, 200]| = 24$ and $|E \cap [37, 200]|/164 \approx 12\%$, with six distinct case-(a) ghost types contributing. The “growing gaps” pattern observed in [3, 36] was an artifact of the short search range: beyond $k = 36$, ghost reappearances fill in the gaps.

The Borel–Cantelli heuristic $P(k \in E) \sim k^2 \cdot 2^{-k}$ was wrong because it treated ghost appearances as independent events. Case-(a) ghosts are deterministic: once identified, their reappearance pattern is exactly periodic (Figure~3).




Figure 3: Ghost cycle appearances by level k . Each row represents a ghost type (identified by denominator D). The vertical dashed line at $k = 36$ marks the boundary of exhaustive search; beyond it, ghost memberships are computed algebraically from Theorem 5. The periodic structure of case-(a) ghosts is clearly visible.

9.2 Conjecture 2 (Density of E)

Conjecture 2 (Density of E). *The exceptional set E has a well-defined natural density $\delta(E) > 0$. The density decomposes as*

$$\delta(E) = 1 - \prod_{\mathcal{G}} \left(1 - \frac{r_{\mathcal{G}}}{p_{\mathcal{G}}} \right)$$

where the product is over all case-(a) ghost types $\mathcal{G}$, $p_{\mathcal{G}} = \text{ord}_2(|D_{\mathcal{G}}|)$ is the period, and $r_{\mathcal{G}}$ is the number of residue classes within that period where $\mathcal{G}$ appears. When two ghost types have periods sharing a common factor, an inclusion-exclusion correction is needed; the product formula is a lower bound on the true density.

From the six original ghost types with $L \leq 8$, the product formula gives $\delta(E) \geq 1 - \frac{24}{25} \cdot \frac{175}{178} \cdot \frac{651}{660} \cdot \frac{1925}{1930} \cdot \frac{82}{84} \cdot \frac{1431}{1441} \approx 10.0\%$. Several period pairs share common factors ($\gcd(660, 1930) = 10$, $\gcd(660, 84) = 12$, $\gcd(660, 1441) = 11$), so the coprime assumption is not satisfied and the product formula understates the true density.

An algebraic membership scan through $k = 1000$ (using the periodicity formula of Theorem~5: ghost type $\mathcal{G}$ is present at level k iff $k \equiv k_0 \pmod{p_{\mathcal{G}}}$; this tests membership against the six ghost types with $L \leq 8$ used in the product formula above, not an exhaustive search for new cycles) gives: $k = 100$: 9.18%, $k = 200$: 10.61%, $k = 500$: 10.04%, $k = 1000$: 10.22%. The 0.2% gap between the product formula (10.0%) and the empirical density (10.2%) is consistent with the expected correction from shared period factors. The oscillation between 9% and 11% reflects finite-size effects from the various periods; convergence to a stable density requires $k \gg \text{lcm}(\text{all periods}) \sim 10^{10}$.

The $D = -601$ ghost alone gives $\delta(E) \geq 1/25 = 4\%$ unconditionally.

Remark (Towards a proof). *The main obstacle is converting the lower bound into an exact density. Two routes are available: (1) full inclusion-exclusion over all ghost types, requiring identification of every materializing family (currently bounded through $L = 15$); or (2) an equidistribution result showing that ghost memberships are asymptotically independent, which would make the product*

formula exact. The second route reduces to equidistribution of $2^k \pmod{|D|}$ for each ghost type — a classical result in multiplicative order theory that holds individually for each D , but whose uniformity across all ghost types simultaneously remains open.

9.3 Conjecture 3 (Spectral Radius)

Conjecture 3 (Spectral Radius). *The spectral radius of the transfer matrices satisfies*

$$\limsup_{k \rightarrow \infty} \rho_k = \max \left(\frac{1}{4}, \sup_{\mathcal{G}} 2^{-V_{\mathcal{G}}/L_{\mathcal{G}}} \right),$$

where the supremum is over all case-(a) ghost types. The maximum with $1/4$ accounts for the fixed point $\{1\}$. From the known ghosts, $\limsup \rho_k \geq 2^{-16/15} \approx 0.4774$ (from $D = -14283371$, $L = 15$, $V = 16$). The $V = L + 1$ ghost family gives $\rho = 2^{-(L+1)/L} \rightarrow 1/2$ as $L \rightarrow \infty$; computational evidence (ghosts appearing at $L = 5, 6, 7, 8, 10, 12, 13, 14, 15$) suggests $\limsup \rho_k = 1/2$. More generally, any family with fixed excess $e = V - L$ has $\rho = 2^{-1-e/L} \rightarrow 1/2$ as $L \rightarrow \infty$, so the spectral radius question does not depend solely on the $V = L + 1$ family.

The current proved bounds are $2^{-16/15} \leq \rho(\mathcal{L}) \leq 1/2$.

9.4 Conjecture 4 (Negative Rationality)

Conjecture 4 (Negative Rationality). *For every case-(a) ghost type with $D < 0$, all orbit elements $\tilde{n}_i = R_i/D$ are negative rationals (equivalently, $R_i > 0$ for all i).*

This conjecture has been verified computationally for all 5,996 canonical case-(a) ghost types with $D < 0$ across 66 (L, V) pairs through $L = 12$. For concentrated patterns it is now a theorem:

Theorem 6 (Negative Rationality for Concentrated Patterns). *Let $L \geq 2$, $e \geq 1$, and $D = 2^{L+e} - 3^L < 0$. For the concentrated ghost type $(L, V, (1, \dots, 1, e + 1))$ with $V = L + e$, the orbit numerators are given by the closed form*

$$R_i = 2^{L-i+1}(2^e - 1) \cdot 3^{i-1} + (3^L - 2^{L+e}), \quad i = 1, \dots, L.$$

Both terms are strictly positive, so $R_i > 0$ for all i . Since $D < 0$, all orbit elements $\tilde{n}_i = R_i/D$ are negative rationals. As a byproduct, concentrated patterns are always case-(a). (The case-(a) argument uses only the oddness of R_i and v_2 conditions, which are independent of the sign of D ; hence case-(a) holds for all concentrated patterns with $e \geq 1$, not only those with $D < 0$.)

Proof. The orbit recurrence $R_{i+1} = (3R_i + D)/2$ for $i = 1, \dots, L - 1$ is a first-order linear recurrence. Setting $Q_i = 2^{i-1}R_i$ gives $Q_{i+1} = 3Q_i + 2^{i-1}D$, which has the general solution $Q_i = C \cdot 3^{i-1} - D \cdot 2^{i-1}$. With $R_1 = \sum_{j=0}^{L-1} 3^{L-1-j}2^j = 3^L - 2^L$ (geometric series), the initial condition gives $C = 2^L(2^e - 1)$, yielding the stated formula. Orbit closure $(3R_L + D)/2^{e+1} = R_1$ is verified directly. Positivity: the first term is positive since $e \geq 1$; the second since $D < 0$ implies $3^L > 2^{L+e}$. Case-(a) requires $v_2(3R_i + D) = v_i$ exactly. For $i < L$: $3R_i + D = 2R_{i+1}$ by the recurrence, so $v_2(3R_i + D) = 1 + v_2(R_{i+1})$; we need R_{i+1} odd. For $i = L$: $3R_L + D = 2^{e+1}(3^L - 2^L)$ (verified directly), giving $v_2 = e + 1 = v_L$. Oddness of R_i for $i \leq L$: the second term $3^L - 2^{L+e}$ is always odd (odd minus even). For $i < L$: the first term $2^{L-i+1}(2^e - 1) \cdot 3^{i-1}$ has $v_2 = L - i + 1 \geq 2$, so it is even; even + odd = odd. For $i = L$: the first term is $2(2^e - 1) \cdot 3^{L-1}$, which is $2 \times \text{odd}$; so $R_L = 2 \cdot \text{odd} + \text{odd} = \text{odd}$. $\square$

The full conjecture (all $D < 0$ ghost types, not just concentrated patterns) remains open for non-concentrated compositions; a proof strategy is outlined in the Discussion (Section~12).

Remark. If a 2-adic periodic orbit $\tilde{n}_1, \dots, \tilde{n}_L$ has $\tilde{n}_i = R_i/D$ with all R_i/D positive integers, these integers form a true Collatz cycle (since the valuation conditions and the Syracuse map agree on $\mathbb{Z}_{>0} \subset \mathbb{Z}_2$). Conversely, any positive-integer Collatz cycle is a case-(a) 2-adic orbit with positive rational elements. Conjecture 4 asserts that all $D < 0$ ghost orbits are purely negative, ruling out positive or integer orbit elements from this family. A positive-integer Collatz cycle would require a case-(a) orbit with $D > 0$ (since $R_1 > 0$ is a sum of positive terms, and for $D > 0$ the recurrence $R_{i+1} = (3R_i + D)/2^{v_i}$ preserves positivity, so positivity of $\tilde{n}_i = R_i/D$ forces $D > 0$); such cycles lie outside the scope of Conjecture 4. The computational results of Steiner (1977) and Simons-de Weger (2005) exclude positive-integer cycles up to length 68 via Baker-type estimates, directly addressing the $D > 0$ case. Conjecture 4 does not address divergent trajectories.

Conjecture~1 (Universal Case-(a)) was stated in Section 7, where the case-(a)/(b) classification is introduced.

10 Eigenvalue Spectra

Dense eigenvalue computation (numpy `eig`) for $k = 3, \dots, 15$ reveals a striking pattern: for every non-exceptional k in this range, the spectrum of P_k is exactly $\{0, 1/4\}$. The only nonzero eigenvalue is $1/4$ (from the fixed point $\{1\}$); all other eigenvalues are exactly zero.

Table 6: Eigenvalue spectra of $P_k(3, 1)$ for $k = 3, \dots, 15$.

k	Matrix size N	Nonzero eigenvalues	Spectrum	Exceptional?
3	4	1	$\{0, 1/4\}$	No
4	8	1	$\{0, 1/4\}$	No
5	16	1	$\{0, 1/4\}$	No
6	32	1	$\{0, 1/4\}$	No
7	64	1	$\{0, 1/4\}$	No
8	128	1	$\{0, 1/4\}$	No
9	256	1	$\{0, 1/4\}$	No
10	512	27	27 nonzero	Yes
11	1024	26	26 nonzero	Yes
12	2048	14	14 nonzero	Yes
13	4096	1	$\{0, 1/4\}$	No
14	8192	1	$\{0, 1/4\}$	No
15	16384	1	$\{0, 1/4\}$	No

The nonzero eigenvalue count is explained by cycle lengths: the fixed point $\{1\}$ contributes 1 eigenvalue ($1/4$), and each extra cycle of length L contributes L nonzero eigenvalues (the L th roots of 2^{-V}). Thus $k = 10$ has $1 + 26 = 27$ (one extra cycle of length 26), $k = 11$ has $1 + 25 = 26$ (one extra cycle of length 25), and $k = 12$ has $1 + 7 + 6 = 14$ (two extra cycles of lengths 7 and 6).

The Fredholm determinant $F_k(z) = \det(I - z \cdot P_k)$ is a polynomial in z of degree equal to the number of nonzero eigenvalues of P_k (at most $N = 2^{k-1}$). Its zeros occur at $z = 1/\lambda_i$, the reciprocals of eigenvalues. For non-exceptional k , the only nonzero eigenvalue is $1/4$, so the only zero is at $z = 4$, giving $\det(I - zP_k) = 1 - z/4$.

11 Computational Methodology

11.1 Transfer Matrix Construction

The transfer matrix P_k is constructed as in Definition~3 (Section 2). For $x = 3$, $y = 1$: for each odd residue j , compute $\text{val}_j = 3j + 1$, extract $v_j = v_2(\text{val}_j)$, and set entry $P_k[t_j, j] = 2^{-v_j}$ where $t_j = (3j + 1)/2^{v_j} \bmod 2^k$.

11.2 Cycle Search Algorithm

Cycle detection follows the successor graph. For $k \leq 32$, the successor function is precomputed as a numpy uint32 array; for $k = 33$ – 36 , it is computed on-the-fly.

The algorithm iterates over all $N = 2^{k-1}$ odd residues in $O(N)$ time and $O(N)$ space (or $O(N/8)$ with bitpacking):

```
for start in 0..N-1:
    if visited[start]: continue
    path = []; path_set = set()
    cur = start
    while not visited[cur] and cur not in path_set:
        path.append(cur); path_set.add(cur)
        cur = successor(cur)
    if cur in path_set:
        cycle = path[path.index(cur):]
        record(cycle)
    mark all path elements visited
```

11.3 On-the-fly Computation with Numba

For $k = 33$ – 36 , storing the successor array would require more than 16 GB. Instead, we compute successors on the fly using Numba JIT compilation. The visited array is packed as a bitarray (1 bit per residue), reducing memory to approximately 4 GB at $k = 36$.

11.4 Eigenvalue Computation

Dense eigenvalue computation uses numpy’s `eig` function on the full $N \times N$ matrix. This is feasible through $k = 15$ ($N = 16384$). Sparse eigensolvers (ARPACK via `scipy.sparse.linalg.eigs`) produce artifacts for these nearly nilpotent matrices, reporting spurious nonzero eigenvalues of magnitude ~ 0.20 – 0.24 ; dense computation confirms these are numerical artifacts. All eigenvalue results in this paper use dense computation only.

11.5 Verification

All results for $k = 3, \dots, 24$ are verified against a separate implementation (113 unit tests checking cycle counts, lengths, and spectral radii). Ghost cycle persistence is verified algebraically: for each known ghost type, the rational orbit $\tilde{n}_1 = R/D$ is computed exactly and the case-(a) valuation conditions are checked. Arithmetic progression reappearance is confirmed through $k = 200$.

11.6 Reproducibility

The complete codebase, including analysis scripts, test suite, and figure generation, is available at <https://github.com/mysticflounder/collatz>. All computations use Python 3 with numpy and optional Numba acceleration.

12 Discussion

We do not address the Collatz conjecture directly. Our results concern the spectral theory of the transfer operator $\mathcal{L}$ and the structure of its exceptional set E , which is a necessary preliminary to any spectral approach to the conjecture.

12.1 The Mod-3 Restricted Approach

The Lasota–Yorke obstruction (Theorem~2) arises from the mod-3 oscillation of the weight function W . A natural salvage attempt restricts the Lipschitz seminorm to pairs within the same mod-3 class, where the conditional contraction rate is $4/15$ (Remark~4). However, the inverse branches involve division by 3, which scrambles higher powers of 3: for $x \equiv y \pmod{3}$ but $9 \nmid (x - y)$, the branches g_v do not preserve mod-3 classes, and the obstruction propagates to all levels of the 3-adic filtration.

12.2 Archimedean Compactness

The 2-adic unboundedness (Theorem~3) closes the Mahler/Amice program in the 2-adic setting. The archimedean compactness question is also now resolved:

Proposition 6 (Non-compactness of $\mathcal{L}$). *$\mathcal{L}$ is not compact on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{R})$.*

Proof. The proof is by non-equicontinuity. The mod-3 splitting of the branch set is independent of the 2-adic topology: for any $r \geq 1$ there exist $x \equiv 1 \pmod{3}$ and $y \equiv 2 \pmod{3}$ with $|x - y|_2 = 2^{-r}$. Define $f \in C(\mathbb{Z}_2^{\text{odd}})$ with $\|f\|_\infty \leq 1$ by $f = +1$ on even-valuation preimage branches of x and $f = -1$ on odd-valuation preimage branches of y (disjoint closed subsets of the compact space $\mathbb{Z}_2^{\text{odd}}$; by the Tietze extension theorem, f extends to a continuous function with $\|f\|_\infty \leq 1$; note f depends on the pair (x, y) and hence on r , though the gap magnitude is always 1). Then

$$(\mathcal{L}f)(x) = \sum_{v=2,4,\dots} 2^{-v} = \frac{1}{3}, \quad (\mathcal{L}f)(y) = - \sum_{v=1,3,\dots} 2^{-v} = -\frac{2}{3},$$

giving $|(\mathcal{L}f)(x) - (\mathcal{L}f)(y)| = 1$ for every r . Since the image $\{\mathcal{L}f : \|f\|_\infty \leq 1\}$ is not equicontinuous, $\mathcal{L}$ is not compact. $\square$

Both archimedean and 2-adic obstructions are now closed.

12.3 Projective Limit and $\sigma(\mathcal{L}) \supseteq [1/4, 1/2]$

Theorem~1(e) gives $\sigma(\mathcal{L}) = \overline{\bigcup \sigma(P_k)}$ directly, bypassing the need for quasi-compactness. Case-(a) ghosts persist across levels (Theorem~5), so exceptional eigenvalues accumulate. The $V = L + 1$ ghost family produces case-(a) ghosts with $\rho = 2^{-(L+1)/L} \rightarrow 1/2$ at the majority of tested cycle lengths $L \leq 15$, suggesting $\rho(\mathcal{L}) = 1/2$ and hence *no spectral gap* on $C(\mathbb{Z}_2^{\text{odd}})$. If the universal case-(a) property (Conjecture~1) holds for all L , and if for each (L, V) with $V < 2L$ at least one ghost type materializes (i.e., appears as a modular cycle at some level k), then $\sigma(\mathcal{L}) \supseteq [1/4, 1/2]$,

since the set $\{2^{-V/L} : L \geq 2, L+1 \leq V \leq 2L-1\}$ is dense in $[1/4, 1/2]$. Materialization is the weakest link in this chain: the $V = L+1$ family has $r = 0$ at $L = 9$ and $L = 11$, showing that case-(a) alone does not guarantee materialization. The phenomenon appears governed by equidistribution of powers of 2 modulo $|D|$, and a heuristic density argument (Section 8) predicts that most (L, V) pairs produce materializing ghosts for sufficiently large L .

12.4 Baker–Wüstholz Bounds

Effective lower bounds on $|2^V - 3^L|$ from transcendence theory (Propositions~4–5) bound ghost cycle denominators but do not prove $|E| < \infty$. Baker–Wüstholz bounds exclude case-(b) ghosts (bounded-length, finite persistence) but not case-(a) ghosts, which are true 2-adic periodic orbits that persist at arithmetic progressions of levels regardless of $|D|$.

Theorem~6 (Section~9) proves Conjecture~4 unconditionally for all concentrated patterns, with no Baker bounds required: the closed form $R_i = 2^{L-i+1}(2^e - 1) \cdot 3^{i-1} + (3^L - 2^{L+e})$ has both terms strictly positive, making the result immediate. For non-concentrated patterns, the analogous recurrence has non-constant step sizes and the resulting expression for R_i involves terms of potentially mixed sign. Extending Theorem~6 to all compositions would require new control on cancellations in these sums; Baker-type bounds in the spirit of Steiner (1977) and Simons–de Weger (2005) remain a candidate approach for that case.

12.5 Thermodynamic Formalism

Inducing schemes may provide alternative routes to spectral gap, working with the natural return map rather than the global transfer operator. The key challenge is that the inducing partition must respect both the 2-adic and mod-3 structures simultaneously.

12.6 Iwasawa Analogy

The projective system $\{P_k\}_{k \geq 2}$ of transfer matrices, with its compatible restriction maps, is structurally analogous to the tower of finite group rings in Iwasawa theory, where the Iwasawa algebra $\Lambda = \mathbb{Z}_p[[T]]$ governs the projective limit. In that classical setting, the Λ -modules of interest are the p -parts of ideal class groups ascending the $\mathbb{Z}_p$ -tower; the analogous objects here are the eigenspaces of P_k associated to ghost cycles — specifically, the exceptional eigenvalues $\lambda = 2^{-V/L}$ that persist across levels. The μ -invariant measures whether the Λ -module has p -torsion growing without bound up the tower; the analogue here is the persistence of case-(a) ghosts at arithmetic progressions of levels (Theorem~5), ensuring E is infinite with positive density rather than eventually vanishing. If E were finite, the spectral radius would collapse to $1/4$ — mirroring $\mu = 0$, where the p -part of the class group is controlled. We emphasize that this is a structural analogy, not a theorem: no Iwasawa algebra acts on the eigenspaces of $\mathcal{L}$, and the periodicity in k is multiplicative-order periodicity rather than the p -power ramification that drives Iwasawa towers.

12.7 Connection to Dhiman–Pandey (2026)

Dhiman and Pandey (2026) independently use the term “ghost cycles” for 2-adic periodic orbits of the $3x+1$ map. Our work differs in computing the density of exceptional levels, classifying ghost persistence into case-(a) and case-(b), and providing the census of materializing ghost types.

Acknowledgments

The author acknowledges extensive use of Claude (Anthropic) throughout this work, including code development, computational exploration, proof formalization, and manuscript preparation. The research directions, conjectures, and interpretation of results are the author’s own. All results are verified by an automated test suite (113 tests) and reproducible from the open-source repository. This work received no external funding.

References

- Amice, Y. (1964). Interpolation p -adique. *Bulletin de la Société Mathématique de France* 92, 117–180.
- Assani, I. (2024). Collatz Map as a Non-Singular Transformation. *Studia Mathematica* 275.
- Baker, A. and Wüstholz, G. (1993). Logarithmic forms and group varieties. *Journal für die reine und angewandte Mathematik* 442, 19–62.
- Dhiman, M. and Pandey, R. (2026). 2-Adic Obstructions to Presburger-Definable Characterizations of Collatz Cycles. arXiv:2601.12772.
- Kontorovich, A. and Lagarias, J. (2010). Stochastic Models for the $3x + 1$ and $5x + 1$ Problems. In Lagarias, J. C. (ed.), *The Ultimate Challenge: The $3x + 1$ Problem*, AMS, pp. 131–188.
- Lagarias, J. (1985). The $3x + 1$ Problem and Its Generalizations. *American Mathematical Monthly* 92, 3–23.
- Lagarias, J. (2021). The $3x + 1$ Problem: An Overview. arXiv:2111.02635.
- Lagarias, J. and Weiss, A. (1992). The $3x + 1$ Problem: Two Stochastic Models. *Annals of Applied Probability* 2(1), 229–261.
- Laurent, M. (2008). Linear forms in two logarithms and interpolation determinants II. *Acta Arithmetica* 133(4), 325–348.
- Matthews, K. and Watts, A. (1985). A Markov approach to the generalized Syracuse algorithm. *Acta Arithmetica* 45(1), 29–42.
- Mori, T. (2024). Application of Operator Theory for the Collatz Conjecture. arXiv:2411.08084.
- Neklyudov, M. (2024). Functional Analysis Approach to the Collatz Conjecture. *Results in Mathematics* 79.
- Serre, J.-P. (1962). Endomorphismes complètement continus des espaces de Banach p -adiques. *Publications Mathématiques de l’IHÉS* 12, 69–85.
- Simons, J. and de Weger, B. (2005). Theoretical and computational bounds for m -cycles of the $3n + 1$ problem. *Acta Arithmetica* 117(1), 51–70.
- Steiner, R. P. (1977). A Theorem on the Syracuse Problem. In *Proceedings of the 7th Manitoba Conference on Numerical Mathematics*, pp. 553–559.
- Tao, T. (2022). Almost All Orbits of the Collatz Map Attain Almost Bounded Values. *Forum of Mathematics, Pi* 10, e12.

Wirsching, G. (1998). *The Dynamical System Generated by the $3n + 1$ Function*. Springer LNM 1681.

Zöbelein, C. (2024). Collatz ghost cycles and multi-level sequences. Preprint.

Version history. v1 (March 2026): Initial submission. v2 (March 2026): Added Theorem~6 (unconditional proof of negative rationality for all concentrated ghost patterns; case-(a) byproduct extended to all $D \neq 0$); added proof strategy remarks for Conjectures 1, 2, and 4; qualified Theorem 1(c) simplicity as computationally verified; corrected Fredholm determinant degree statement; fixed notation and expository issues throughout. v3 (March 2026): *New results:* Added Proposition~6 (archimedean non-compactness of $\mathcal{L}$ on $C(\mathbb{Z}_2^{\text{odd}}, \mathbb{R})$, proved by non-equicontinuity via finite branch truncation); added Baker–Wüstholz proof sketch (Laurent 2008, Corollary~2); added remark clarifying $\mathcal{L}$ as projective limit of integer Syracuse maps; added primitivity remark to Theorem~5; noted $n = -1/3$ exceptional point. *Corrections:* Updated archimedean compactness discussion (resolved: not compact); corrected pair count from 91 to 105 for $L = 2, \dots, 15$; softened “closing the Mahler/Amice program entirely” to “in the 2-adic setting”; corrected Baker–Wüstholz growth claim; clarified $k = 1000$ density scan as algebraic membership testing; corrected D odd/nonzero justification; corrected Proposition~5 K_0 definition to $\max\{\text{ord}_2(|D|) + 2L : \dots\}$, closing a gap in the “in practice” step. *Proof repairs:* Repaired Proposition~3 proof to derive bound from finite matrices directly; clarified A_k projection structure in Theorem~1(e) proof; moved persistence-dependent $\sup_k = \limsup_k$ equality out of Theorem~1(e) proof into a separate remark; expanded Theorem~6 geometric series derivation; relabeled Lemma~1 cases (a)–(c); reordered conjecture list in contributions to match paper numbering; dropped imprecise total-variation bound from Corollary~1(b) and merged BV statement into revised part (b) with a rigorous one-line proof. *Attribution:* Added footnote to Theorem~4 crediting Davison (1976) as originator of the cycle equation, as cited by Steiner (1977).

Reference corrections (v3): The reference “Siegel, M. Ghost Cycles of the $3x + 1$ Map. arXiv:2601.12772” appearing in v1 and v2 was incorrect in both author and title. The correct citation is Dhiman, M. and Pandey, R. (2026), 2-Adic Obstructions to Presburger-Definable Characterizations of Collatz Cycles, arXiv:2601.12772. The paper attributed to Siegel does not exist; the error was introduced during manuscript preparation. The author apologizes to Maxwell Siegel and to Madhav Dhiman and Rohan Pandey for this misattribution. Additionally: the page range for Amice (1964) was corrected from 181–236 to 117–180 (verified from PDF); the title for Kontorovich and Lagarias (2010) was corrected to match the arXiv preprint (removed spurious “and Beyond”).