

1 Overview and Review, 8/23/16

1.1 Overview

Class field theory provides a dictionary between abelian extensions of a given number field F (i.e. Galois extensions of F with abelian Galois group) and intrinsic data about the number field, e.g. the class group of the ring of integers, $\text{Cl}(\mathcal{O}_F)$. For instance, if we let H be the union of all abelian extensions of F that are everywhere unramified, we have $\text{Gal}(H/F) \cong \text{Cl}(\mathcal{O}_F)$.

However, class field theory does not construct these abelian extensions, except for two classical constructions which were already known:

1. $F = \mathbb{Q}$, which has $H = \mathbb{Q}(\mu_\infty)$, where we have adjoined all the roots of unity. This is Kronecker-Weber theorem.
2. F a quadratic imaginary extension of \mathbb{Q} .

This is not so bad, however, since it turns out that class field theory can actually yield information about non-abelian extensions!

1.2 Topics

- Review local and global fields.
- Group and Galois cohomology.
- Local class field theory and local duality (important!).
- Global class field theory and global duality.
- Applications (Iwasawa theory).

1.3 Review

1.3.1 First Example

Let L/K be a finite Galois extension. Let \mathfrak{P} be an unramified prime of L lying over \mathfrak{p} , so that $\mathfrak{p}\mathcal{O}_L = \prod \mathfrak{P}_i^{e_i}$ with all $e_i = 1$.

Lemma 1.3.1. *There is an element $\text{Fr}_{\mathfrak{P}}$ of $\text{Gal}(L/K)$ such that*

- 1) $\text{Fr}_{\mathfrak{P}}(\mathfrak{P}) = \mathfrak{P}$, i.e. $\text{Fr}_{\mathfrak{P}}$ is in the decomposition group of \mathfrak{P} .

2) $\text{Fr}_{\mathfrak{P}}$ acts on $\mathcal{O}_L/\mathfrak{P}$ as $x \rightarrow x^{N(\mathfrak{p})}$.

Remark 1. Note that $\mathcal{O}_K/\mathfrak{p}$ is a finite field of order $N(\mathfrak{p})$, which has the Frobenius automorphism that does precisely what 2 does. We should think of $\text{Fr}_{\mathfrak{P}}$ as a lift of that map to $\mathcal{O}_L/\mathfrak{P}$.

Proof. We will construct $\text{Fr}_{\mathfrak{P}}$ explicitly. Let $\alpha \in \mathcal{O}_L$ satisfying

1. α generates $(\mathcal{O}_L/\mathfrak{P})^\times$, and
2. for all $\mathfrak{P}^o \neq \mathfrak{P}$ above \mathfrak{p} , $\alpha \in \mathfrak{P}^o$.

Set $F(X) = \prod_{\sigma \in \text{Gal}(L/K)} (X - \sigma\alpha) \in \mathcal{O}_K[X]$. Then $F(\alpha) = 0$, so $F(\alpha^{N(\mathfrak{p})}) = F(\alpha)^{N(\mathfrak{p})} = 0$.

Then for some σ , $\alpha^{N(\mathfrak{p})} \equiv \sigma\alpha \pmod{\mathfrak{P}}$. Then we claim that $\sigma\mathfrak{P} = \mathfrak{P}$. Otherwise, $\sigma^{-1}\mathfrak{P} \neq \mathfrak{P}$, so $\alpha \in \sigma^{-1}\mathfrak{P}$, so $\sigma\alpha \in \mathfrak{P}$. So $\alpha^{N(\mathfrak{p})} \equiv 0 \pmod{\mathfrak{P}}$, a contradiction.

Then for all $x \in \mathcal{O}_L/\mathfrak{P}$, we can write $x = \alpha^i + b$, for some i and $b \in \mathfrak{P}$. Then

$$\sigma(x) = \sigma(\alpha^i) + \sigma(b) = \alpha^{iN(\mathfrak{p})} + \sigma(b) \equiv x^{N(\mathfrak{p})} \pmod{\mathfrak{P}}.$$

Now define $\text{Fr}_{\mathfrak{P}} := \sigma$. Uniqueness is left to the reader. □

Remark 2. If $\mathfrak{P}^o = \tau(\mathfrak{P})$ for some $\tau \in \text{Gal}(L/K)$, $\text{Fr}_{\mathfrak{P}^o} = \tau \text{Fr}_{\mathfrak{P}} \tau^{-1}$.

Recall that L/K being Galois means $\text{Gal}(L/K)$ acts transitively on the primes \mathfrak{P} lying over \mathfrak{p} . So $\text{Fr}_{\mathfrak{P}}$ is well-defined up to conjugation. If L/K is abelian, it is well-defined.

1.3.2 $\text{Fr}_{\mathfrak{P}}$ of Cyclotomic Field

Let $L = \mathbb{Q}(\zeta_n)$ be the n -th cyclotomic field, $K = \mathbb{Q}$. Then $\text{Gal}(L/K) \cong (\mathbb{Z}/n\mathbb{Z})^\times$. Take p unramified in L/K , i.e. p not dividing n . Then Fr_p (we are in an abelian extension, so all $\text{Fr}_{\mathfrak{P}}$ are the same). By definition, Fr_p is the σ such that $\sigma(\alpha) = \alpha^p \pmod{\mathfrak{P}}$, for all \mathfrak{P} over p .

Also characterized by $\tau(\zeta_n) = \zeta_n^p$ since

$$\tau \sum a_i \zeta_n^i = \sum a_i \zeta_n^i{}^p.$$

1.3.3 $\text{Fr}_{\mathfrak{P}}$ of Quadratic Field

Here, we let $L = \mathbb{Q}(\sqrt{d})$ and K as before. Then $\text{Gal}(L/K) \cong \mathbb{Z}/2\mathbb{Z}$, and for p unramified in L , Fr_p corresponds to 1 if p splits in L , or -1 if p is inert in L . This means that

$$\text{Fr}_p = \left(\frac{d}{p} \right).$$

This connection leads us to an extremely nice proof of the quadratic reciprocity for odd primes.

Theorem 1.3.2. *Let $p \neq q$ be odd primes. Then $\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{\frac{p-1}{2} \frac{q-1}{2}}.$*

Proof. Let $L = \mathbb{Q}(\zeta_p)$, $K = \mathbb{Q}$. Then $\text{Gal}(L/K)$ is cyclic of order $p-1$, and so has a unique order two quotient which corresponds to a quadratic field F , where $F = \mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}} p}\right)$.

Since q is unramified, we can consider Fr_q which corresponds to $q \in (\mathbb{Z}/p\mathbb{Z})^\times$. Now we simply compute this quantity in two ways.

$$(i) \quad \text{Fr}_q|_F = 1 \Leftrightarrow q^{\frac{p-1}{2}} \equiv 1 \pmod{p} \Leftrightarrow \left(\frac{q}{p}\right) = 1.$$

$$(ii) \quad \text{Fr}_q|_F \text{ is also simply } \text{Fr}_q \text{ for the quadratic extension } F, \text{ hence equal to } \left(\frac{(-1)^{\frac{p-1}{2}} p}{q}\right) \text{ by the previous example.}$$

□

1.4 First Case of Fermat's Last Theorem

Theorem 1.4.1. *If p does not divide $|\text{Cl}(\mathbb{Q}(\zeta_p))|$, then $x^p + y^p = z^p$ has no integer solutions with p not dividing xyz .*

The idea is that we can factor $\prod_i (x + \zeta_p^i y) = z^p$ in $\mathbb{Z}[\zeta_p]$. It turns out that regularity gives us that the LHS factors are p -th powers, the divisibility condition giving us that the factors are coprime.