

Laboratory Assignment 3

Activities

1. (a) The Pell numbers are an infinite sequence of integers which correspond to the denominators of the closest rational approximations of $\sqrt{2}$. The Pell numbers are defined by the following recurrence relation (which looks very similar to the Fibonacci sequence):

$$P_n = \begin{cases} 0 & \text{if } n = 0 \\ 1 & \text{if } n = 1 \\ 2P_{n-1} + P_{n-2} & \text{otherwise} \end{cases}$$

Use this recurrence relation to write a recursive function, `pell`, which takes one parameter, n , and returns the n^{th} Pell number.

- (b) Write a separate function, named (`find-pell n`), which uses your Pell function to find the largest Pell number which is less than n . You can try testing it by finding the largest Pell number less than 100.
- (c) The numerator for the rational approximation of $\sqrt{2}$ corresponding to a particular Pell number is half of the corresponding number in the sequence referred to as the *companion Pell numbers* (or Pell-Lucas numbers). The companion Pell numbers are defined by the recurrence relation:

$$Q_n = \begin{cases} 2 & \text{if } n = 0 \\ 2 & \text{if } n = 1 \\ 2Q_{n-1} + Q_{n-2} & \text{otherwise} \end{cases}$$

Use this recurrence relation to write a function, named (`comp-pell n`), which returns the n^{th} companion Pell number.

- (d) Finally write a function that uses the Pell number and companion Pell number functions, as described in Part a, to write a SCHEME function, named (`sqrt-2-approx n`), to compute the n^{th} approximation for $\sqrt{2}$. You can test your new function to compute the approximation for $\sqrt{2}$ using the sixth Pell and companion Pell numbers.
2. Viète's formula is the following infinite product of nested radicals representing the mathematical constant π :

$$\frac{2}{\pi} = \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{2+\sqrt{2}}}{2} \cdot \frac{\sqrt{2+\sqrt{2+\sqrt{2}}}}{2} \dots$$

Write a SCHEME function named (`viète n`) which approximates $\frac{2}{\pi}$ using Viète's formula.

Note: You may want to use a helper function that takes an additional parameter so that you can compute the numerator of each term easily (in a `let` form perhaps) as well as pass it to the recursive call.

3. It is an interesting fact the the square-root of any number may be expressed as a *continued fraction*. For example,

$$\sqrt{x} = 1 + \frac{x-1}{2 + \frac{x-1}{2 + \frac{x-1}{\ddots}}}$$

Write a Scheme function called `new-sqrt` which takes two formal parameters x and n , where x is the number we wish to find the square root of and n is the number of continued fractions to compute recursively. Demonstrate that for large n , `new-sqrt` is very close to the builtin `sqrt` function.

4. **Nested Recursion:** The McCarthy 91 function is a recursive function, defined by the computer scientist John McCarthy as a test case for formal verification within computer science.

The McCarthy 91 function is defined as

$$m91(n) = \begin{cases} n - 10, & \text{if } n > 100 \\ m91(m91(n + 11)), & \text{if } n \leq 100 \end{cases}$$

Define a SCHEME procedure, named (m91 n), which evaluates to McCarthy's 91 function. Try a few values for n which are less than 100 and a few greater than 100.