

Getting started

The Legolas code can be found on [GitHub](#), installation instructions are on the website. Make sure to look at the prerequisites for both Legolas and Pylbo first before running the code.

1 Equilibria

Below is a list of possible setups that can be implemented in the user submodule. We've added a reference figure together with the link to the original work so you can check your implementation and compare with known results.

1.1 Internal kink modes in force-free magnetic fields

This setup is taken from [Goedbloed \(2018\)](#), which is a cylindrical equilibrium with a force-free magnetic field of constant α and profiles given by

$$\begin{aligned} \rho(r) &= \rho_0 (1 - x^2) & \rho'(r) &= -\frac{2\rho_0}{a}x \\ v_z(r) &= v_{03} (1 - x^2) & v'_z(r) &= -\frac{2v_{03}}{a}x \\ B_\theta(r) &= J_1(\alpha x) & B'_\theta(r) &= \frac{\alpha}{2a} [J_0(\alpha x) - J_2(\alpha x)] \\ B_z(r) &= J_0(\alpha x) & B'_z(r) &= -\frac{\alpha}{a} J_1(\alpha x) \\ T(r) &= \frac{p_0}{\rho(r)} & T'(r) &= \frac{2rp_0}{a^2\rho_0(1 - x^2)^2} \end{aligned}$$

where $x = r/a$ and a denotes the outer wall of the cylinder.




Figure 1: Values $\rho_0 = v_{03} = p_0 = a = m = 1$, $\alpha a = 5$, $k = 0.16\alpha$, incompressible.




Figure 2: Values $\rho_0 = p_0 = a = m = 1$, $v_{03} = 0.8$, $\alpha a = 8$, $k = 0.16\alpha$, incompressible.

References

Goedbloed, J.P.: 2018, The Spectral Web of stationary plasma equilibria. II. Internal modes. *Physics of Plasmas* **25**, 032110. [DOI](#). [ADS](#).