

TEEM-BH: An Energetic Framework for Black Hole Horizons, Entropy, and Singularity Avoidance

NS(2026 May)

Abstract

Black holes challenge the foundations of modern physics by combining extreme gravitational fields with quantum mechanical requirements such as unitarity and information preservation. The Tension–Energy Exchange Model (TEEM) offers a new energetic framework for addressing these challenges by decomposing energy into three fundamental modes: vibrational energy v , mass-condensed energy m , and spatial tension energy R . In this framework, spacetime is not geometric but an energetic configuration dominated by R , with time flow determined by the ratio v/R and gravity emerging from gradients in m and R .

We develop TEEM-BH, the black hole branch of TEEM, and show that the event horizon corresponds to an R -critical surface where spatial tension saturates and time effectively halts. Black hole entropy arises naturally from boundary tension modes on this surface, providing a microscopic explanation for the area law. Information is preserved through reconfiguration of the R -field, offering a unitary mechanism for black hole evaporation without firewalls or singularities. The interior forms a finite-tension core stabilized by R -saturation, replacing the classical singularity with a physically meaningful energetic structure.

TEEM-BH yields testable predictions, including finite horizon thickness, modified shadow profiles, non-thermal components in Hawking-like radiation, additional gravitational wave modes, and persistent near-horizon anisotropies. These signatures provide concrete avenues for distinguishing TEEM-BH from General Relativity and for validating an energetic interpretation of spacetime.

1. Introduction

Black holes represent the most extreme environments in the universe, where the known laws of physics are pushed to their limits. General Relativity (GR) provides an elegant geometric description of gravity, yet it predicts the formation of singularities—regions where curvature diverges and physical quantities lose meaning. At the same time, quantum mechanics demands unitarity and information preservation, leading to long-standing tensions such as the information paradox. Despite decades of effort, a unified and conceptually coherent framework that naturally incorporates both gravitational and quantum aspects of black holes remains elusive.

The TEEM framework (Tension–Energy Exchange Model) offers a new conceptual foundation for addressing these issues. TEEM decomposes energy into three modes—vibrational energy v ,

mass-condensed energy m , and spatial tension energy R —and treats spacetime not as a geometric background but as an energetic configuration dominated by R . Time dilation emerges from the ratio v/R , gravity arises from gradients in m and R , and cosmological evolution is interpreted as a redistribution among these modes. This perspective provides a unified language for describing gravitational, quantum, and cosmological phenomena.

Black holes are the natural first testing ground for TEEM. They represent the regime where R becomes dominant and where the interplay among the three energy modes becomes most pronounced. TEEM predicts that the event horizon corresponds to a critical value of R , that black hole entropy arises from boundary tension modes, that information is preserved through reconfiguration of R , and that singularities are avoided through saturation of spatial tension. These predictions offer a coherent alternative to the traditional geometric interpretation of GR while remaining compatible with key observational results.

This paper develops TEEM-BH, the black hole branch of the TEEM framework. We formalize the concept of an R -critical horizon, derive entropy from boundary tension degrees of freedom, propose a mechanism for information preservation, and present a non-singular interior structure. We also outline observational signatures that distinguish TEEM-BH from classical GR, including modifications to horizon structure, black hole shadows, and gravitational wave behavior.

TEEM-BH aims not to replace GR but to extend it into regimes where geometry alone becomes insufficient. By grounding black hole physics in energetic rather than geometric principles, TEEM-BH provides a unified and conceptually transparent approach that connects black holes to quantum gravity, cosmology, and the broader TEEM framework.

2. The TEEM Framework Revisited

The TEEM framework (Tension–Energy Exchange Model) proposes that all physical phenomena can be described through the interaction and redistribution of three fundamental energy modes: vibrational energy v , mass-condensed energy m , and spatial tension energy R . Unlike conventional approaches that treat spacetime as a geometric background, TEEM interprets spacetime as an energetic configuration dominated by R , with geometry emerging as a secondary description.

2.1 Three Energy Modes

TEEM decomposes total energy into a linear combination of the three modes:

$$E = \alpha v + \beta m + \gamma R.$$


Each mode has a distinct physical interpretation:

- **Vibrational energy v** Represents oscillatory or wave-like degrees of freedom. It corresponds to quantum behavior, fluctuations, and kinetic excitations.
- **Mass-condensed energy m** Represents localized, condensed, or particle-like energy. It corresponds to rest mass, matter density, and gravitational sources.
- **Spatial tension energy R** Represents the intrinsic tension or energetic state of space itself. It plays

the role traditionally attributed to spacetime curvature or vacuum energy.

The coefficients α, β, γ encode conversion efficiencies among the modes and are treated as universal constants.

TEEM Energy Modes: v, m, R



Energy modes in TEEM are interconvertible and form the basis of gravitational, quantum, and cosmological behavior.

Figure 1. TEEM Energy Modes: v, m, R .

The three fundamental energy modes of the TEEM framework—vibrational energy (v), mass-condensed energy (m), and spatial tension energy (R)—are interconvertible and form the basis of gravitational, quantum, and cosmological behavior.

Bidirectional arrows indicate dynamic exchange among the modes, governed by conversion coefficients (alpha, beta, gamma).

2.2 Time as a Ratio of Energies

A central postulate of TEEM is that the flow of time is determined by the ratio of vibrational to tension energy:

$$t \propto \frac{v}{R}.$$

This relation implies:

- Time slows down when R increases (strong tension of space).
- Time accelerates when v dominates (high vibrational activity).
- Gravitational time dilation arises naturally from spatial variations in R .

This replaces the geometric interpretation of time dilation in GR with an energetic one.

2.3 Gravity as Gradients of m and R

In TEEM, gravitational behavior emerges from spatial gradients in the energy modes:

$$\vec{g} \propto -\nabla(m + R).$$

This implies:

- Mass m contributes to gravity in the usual way.
- Spatial tension R also contributes, acting as a distributed gravitational source.
- Dark matter-like effects can arise from variations in R without requiring new particles.

Thus, gravity is not curvature but energetic imbalance.

2.4 Energy Exchange and the Partial Reset Principle

TEEM assumes that the three energy modes can convert into one another:

$$v \leftrightarrow m \leftrightarrow R.$$

This exchange underlies cosmological evolution and extreme gravitational phenomena.

The partial reset principle states that during high-energy events—such as the early universe or black hole interiors—energy modes can be redistributed without complete erasure of prior information. This provides a natural mechanism for:

- avoiding singularities,
- preserving information,
- generating initial fluctuations in cosmology.

2.5 Why TEEM Is Suited for Black Hole Physics

Black holes represent the regime where:

- R becomes dominant,
- v/R approaches zero (time freezing),
- energy mode conversion becomes significant,
- information preservation becomes critical.

3. Horizon as an R -Critical Surface

In General Relativity, the event horizon is defined geometrically as a null surface beyond which causal signals cannot escape. TEEM replaces this geometric definition with an energetic one: the horizon is the surface where the spatial tension energy R reaches a critical value. At this surface, the flow of time effectively halts, and the conversion dynamics among the energy modes v, m , and R undergo a qualitative change.

3.1 Time Freezing Condition: $v/R \rightarrow 0$

TEEM defines the local rate of time flow as

$$t \propto \frac{v}{R}.$$

As an object approaches a black hole, the spatial tension energy R increases due to the gravitational compression of space. When R reaches a critical value R_{crit} , the ratio v/R approaches zero:

$$\frac{v}{R_{\text{crit}}} \rightarrow 0.$$

This condition corresponds to complete time dilation, matching the GR prediction that time stops at the horizon for an external observer. However, TEEM interprets this not as a geometric effect but as an energetic saturation phenomenon.

3.2 Definition of the R -Critical Horizon

We define the TEEM horizon as:

$$R(r_h) = R_{\text{crit}}.$$

Here, r_h is the radial coordinate of the horizon. The value R_{crit} is universal and corresponds to the maximum sustainable spatial tension before a phase transition occurs in the energy configuration of space.

This definition implies:

- The horizon is a phase boundary in the distribution of R .
- Inside the horizon, R remains near saturation.
- Outside the horizon, R decreases smoothly with radius.

Thus, the horizon is not a geometric singularity but a critical surface in the energy landscape.




Figure 2. R -Critical Horizon in TEEM-BH.

Cross-sectional representation of a non-rotating black hole showing the spatial tension energy R approaching its critical value R_{crit} at the horizon radius r_h . The color gradient illustrates the increase of R toward the horizon, where $v/R \rightarrow 0$ and time effectively halts. Inside the horizon, R remains saturated, forming a finite-tension core that replaces the classical singularity. The inset graph shows the sharp rise of $R(r)$ at r_h .

3.3 Relation to the Schwarzschild Radius

For a non-rotating black hole, TEEM predicts that the R -critical surface coincides with the classical Schwarzschild radius:

$$R_{\text{crit}} \propto \frac{1}{r_s}.$$

This proportionality ensures compatibility with GR at large scales while providing a deeper energetic interpretation of the horizon.

Unlike GR, TEEM does not require the metric to diverge at the horizon; instead, the divergence is replaced by R-saturation, which naturally halts time flow.

3.4 Horizon as a Phase Transition in R

The behavior of R near the horizon resembles a second-order phase transition:

- The gradient ∇R becomes steep near r_h .
- The system becomes dominated by tension energy.
- Conversion between v and R becomes suppressed.
- Local excitations freeze out as $v/R \rightarrow 0$.

This provides a physical explanation for:

- the freezing of infalling matter,
- the universality of horizon behavior,
- the emergence of surface degrees of freedom (entropy),
- the robustness of the horizon under perturbations.

3.5 Causal Structure from Energetic Constraints

In TEEM, causal disconnection arises because:

$$\frac{v}{R_{\text{crit}}} = 0 \Rightarrow \text{no propagation of vibrational modes.}$$

Since information is carried by vibrational energy v , the vanishing of v/R implies:

- no signal propagation outward,
- no local time evolution at the horizon,
- effective causal isolation.

Thus, the TEEM horizon reproduces the causal structure of GR without requiring null surfaces or metric singularities.

3.6 Summary

The TEEM horizon is defined not by geometry but by energetic saturation:

$$R = R_{\text{crit}}.$$

This definition:

- reproduces GR's time dilation,
- explains causal isolation,
- provides a natural mechanism for entropy and information storage,
- avoids geometric singularities,

- and sets the stage for the TEEM interpretation of black hole interiors.

4. Entropy as Boundary Degrees of Freedom of R

Black hole entropy is one of the most profound clues about the microscopic structure of spacetime. In General Relativity, the Bekenstein–Hawking entropy is proportional to the area of the event horizon, yet GR itself provides no explanation for the origin of these degrees of freedom. TEEM offers a natural energetic interpretation: black hole entropy arises from the boundary tension modes of the spatial tension energy R at the R -critical surface.

4.1 Spatial Tension Energy as the Source of Microstates

In TEEM, the spatial tension energy R represents the intrinsic energetic configuration of space. Near the horizon, R approaches its critical value R_{crit} , and the system becomes dominated by tension energy. This saturation leads to the emergence of boundary tension modes, which behave as microscopic degrees of freedom localized on the horizon.

These modes arise because:

- R cannot increase beyond R_{crit} ,
- but small fluctuations in R along the horizon are still allowed,
- creating a large number of energetically equivalent configurations.

Thus, the horizon becomes a high-degeneracy surface, naturally giving rise to entropy.

4.2 Area Law from Boundary Mode Counting

Let the horizon surface support N independent tension modes. Since each mode corresponds to a local fluctuation of R , the number of microstates grows exponentially with the number of surface patches.

If each patch of area ΔA supports one independent mode, then:

$$N \propto \frac{A}{\Delta A}.$$

The entropy is then:

$$S = k_B \ln \Omega \propto A,$$

where Ω is the number of microstates.

Thus, the area law emerges directly from the structure of R at the horizon, without requiring string theory, loop quantum gravity, or holography as assumptions.

4.3 Relation to Bekenstein–Hawking Entropy

TEEM predicts:

$$S_{\text{BH}} = \eta A,$$

where η is a constant determined by the density of tension modes. Matching this with the Bekenstein–Hawking formula,

$$S_{\text{BH}} = \frac{k_B c^3}{4G\hbar} A,$$

implies that η corresponds to the inverse of the fundamental area scale ΔA . In TEEM, this scale is interpreted as the minimum resolvable patch of R-tension, not as a geometric Planck area.

This provides a physical interpretation of the entropy coefficient: it reflects the quantization of spatial tension, not the quantization of geometry.

4.4 Horizon as a Tension-Dominated Membrane

The R-critical surface behaves like a membrane with:

- high tension,
- suppressed vibrational activity ($v/R \rightarrow 0$),
- localized boundary excitations,
- and a large number of microstates.

This membrane-like behavior is reminiscent of the membrane paradigm in GR, but TEEM provides a microscopic energetic origin rather than a phenomenological one.

4.5 Entropy as a Measure of R-Configuration Degeneracy

In TEEM, entropy counts the number of distinct configurations of R on the horizon that:

- satisfy $R = R_{\text{crit}}$,
- preserve the macroscopic mass and radius,
- differ only in microscopic tension patterns.

Thus, entropy is not a property of matter inside the black hole but a property of the horizon's tension field. This interpretation naturally aligns with holographic principles: information is stored on the boundary because R-tension modes live on the boundary.

4.6 Summary

TEEM explains black hole entropy as:

- arising from boundary tension modes of R ,
- obeying an area law due to surface mode counting,
- matching the Bekenstein–Hawking formula,
- providing a microscopic origin for horizon thermodynamics,
- and offering a natural foundation for holography.

This sets the stage for the next section, where we show how information is preserved through reconfiguration of R .

Boundary Tension Modes as the Origin of Black Hole Entropy




Figure 3. Boundary Tension Modes as the Origin of Black Hole Entropy.

The horizon surface is composed of discrete patches, each representing one R-tension mode. The large number of energetically equivalent configurations of these boundary modes gives rise to entropy. The area law emerges naturally from counting the number of surface patches, consistent with the proportionality $S \propto A$.

5. Information Preservation via R-Reconfiguration

The black hole information paradox arises from the apparent conflict between the unitarity of quantum mechanics and the classical prediction that information falling into a black hole is irretrievably lost. TEEM resolves this tension by proposing that information is not destroyed but reconfigured into the spatial tension field R at and within the horizon. Because R encodes the energetic state of space itself, its configuration retains the essential information of infalling matter even when vibrational and mass-condensed modes are suppressed.

5.1 Energy Mode Conversion as the Basis for Information Flow

In TEEM, the three energy modes are interconvertible:

$$v \leftrightarrow m \leftrightarrow R.$$

As matter approaches the horizon:

- vibrational energy v is suppressed due to $v/R \rightarrow 0$,
- mass-condensed energy m becomes unstable under extreme tension,
- both modes are gradually converted into spatial tension energy R .

This conversion does not erase information; instead, it transfers information from matter and radiation into the configuration of the tension field.

Thus, information flow is preserved through:

$$\mathcal{I}_{\text{matter}} \rightarrow \mathcal{I}_R.$$

5.2 *R*-Configuration as an Information-Bearing Field

The spatial tension energy R is not a scalar quantity but a field with internal structure. Its configuration includes:

- local tension gradients,
- boundary mode patterns,
- anisotropies induced by infalling matter,
- residual vibrational imprints suppressed but not erased.

These features allow R to encode a vast amount of information.

In TEEM, the horizon acts as a storage surface where the incoming information is reorganized into boundary tension modes. This provides a natural mechanism for holographic behavior.

5.3 *TEEM Interpretation of Hawking Radiation*

Hawking radiation arises in TEEM not from pair creation at the horizon but from slow relaxation of the R -configuration.

As the black hole evaporates:

- the tension field R gradually decreases,
- stored information is released as vibrational energy v ,
- the outgoing radiation carries subtle correlations reflecting the original R -pattern.

Thus, Hawking radiation is information-bearing, consistent with unitarity.

This avoids the need for firewalls, nonlocal wormholes, or exotic quantum gravity mechanisms.

5.4 *No-Loss Principle: Information Is Never Destroyed*

TEEM asserts a No-Loss Principle:

$$\frac{d\mathcal{I}}{dt} = 0,$$

where $\mathcal{I}$ is the total information encoded across all energy modes.

Information may change form:

$$\mathcal{I}_v + \mathcal{I}_m + \mathcal{I}_R = \text{constant},$$

but it is never annihilated.

This principle is consistent with:

- quantum unitarity,
- holographic entropy bounds,
- the partial reset mechanism in cosmology.

5.5 Interior Information Storage and the Role of R Saturation

Inside the horizon, R remains near its critical value. This saturated region acts as a high-capacity information reservoir.

Key properties:

- R saturation prevents singularity formation,
- the interior retains a stable tension configuration,
- information is stored in the internal structure of R ,
- the horizon acts as the interface for information exchange.

Thus, the black hole interior is not a singular point but a finite-tension core with structured information content.

5.6 Resolution of the Information Paradox

TEEM resolves the information paradox through three mechanisms:

1. **Information transfer** Infalling matter transfers its information to the R -field.
2. **Information storage** The R -configuration at and inside the horizon stores this information.
3. **Information release** Hawking radiation gradually releases information as the R -field relaxes.

No information is lost at any stage.

5.7 Summary

In TEEM-BH:

- information is encoded in the spatial tension field R ,
- the horizon acts as a boundary where information is reorganized,
- Hawking radiation carries information through R -relaxation,
- the interior stores information in a finite-tension core,
- and the total information across $v/m/R$ modes is conserved.

This provides a coherent, energetic, and non-geometric solution to the black hole information paradox.

Information Preservation via R -Reconfiguration




Figure 4. Information Preservation via R -Reconfiguration.

Conceptual diagram illustrating how information is conserved in TEEM-BH through the dynamics of the spatial tension field R . Infalling matter transfers its information to R at the horizon boundary (R -conversion zone), forming R -configuration patterns that store information within the finite-tension core. As the R -field gradually relaxes, stored information is released as correlated Hawking-like radiation, ensuring unitary evolution and resolving the information paradox.

6. Singularity Avoidance through R Saturation

General Relativity predicts that the gravitational collapse of matter inevitably leads to a singularity—a region where curvature diverges and physical quantities become ill-defined. Such singularities signal a breakdown of the theory rather than a physical feature of nature. TEEM provides a natural mechanism for avoiding singularities through the saturation of spatial tension energy R . As matter collapses, the conversion of vibrational and mass-condensed energy into spatial tension prevents R from diverging, stabilizing the interior of the black hole into a finite-tension core.

6.1 Collapse Dynamics in TEEM

During gravitational collapse:

- mass-condensed energy m increases in density,
- vibrational energy v becomes suppressed as $v/R \rightarrow 0$,
- spatial tension energy R increases due to compression of space.

However, unlike curvature in GR, R cannot grow without bound. The conversion dynamics among the three energy modes enforce a natural upper limit:

$$R \leq R_{\text{crit}}.$$

Once R approaches this critical value, further collapse does not increase tension but instead triggers energy redistribution.

6.2 Saturation Mechanism: Redistribution Among $v/m/R$

When R reaches R_{crit} , the system enters a saturation regime characterized by:

- suppression of further $m \rightarrow R$ conversion,
- stabilization of R at its critical value,
- partial conversion of excess m into v ,
- redistribution of energy into internal tension modes.

This prevents the runaway growth of density and curvature that would otherwise lead to a singularity.

The saturation condition can be expressed as:

$$\frac{dR}{dt} \rightarrow 0 \text{ as } R \rightarrow R_{\text{crit}}.$$

6.3 Formation of a Finite-Tension Core

Instead of collapsing to a point, the interior of a TEEM black hole forms a finite-sized core where:

- R remains near R_{crit} ,
- m is partially converted into internal tension modes,
- v is suppressed but nonzero,
- the energy distribution reaches a stable equilibrium.

This core is not a geometric singularity but a high-tension, high-information region with finite density and finite curvature.

The radius of the core depends on the mass of the black hole and the conversion efficiencies α, β, γ .

6.4 Comparison with GR Singularities

In GR:

- curvature diverges as $r \rightarrow 0$,
- density becomes infinite,
- time evolution breaks down.

In TEEM:

- R saturates at R_{crit} ,
- density remains finite due to energy redistribution,
- time evolution continues internally (though highly suppressed),
- no divergence occurs in any physical quantity.

Thus, TEEM replaces the singularity with a physically meaningful energetic structure.

6.5 Implications for Black Hole Interiors

The existence of a finite-tension core has several consequences:

- **No singularity:** the interior remains regular.
- **Information storage:** the core acts as a reservoir for R -encoded information.

- **Stability:** the core resists further collapse due to tension saturation.
- **Continuity:** the interior evolution remains well-defined.

This structure also provides a natural endpoint for collapse and a foundation for understanding black hole evaporation.

6.6 Connection to Cosmology: Partial Reset Analogy

The saturation mechanism parallels the partial reset process proposed in TEEM cosmology:

- both involve redistribution among $v/m/R$,
- both avoid singularities through R -saturation,
- both preserve information in the tension field.

This suggests a deep connection between black hole interiors and the early universe, hinting at a unified energetic origin for both phenomena.

6.7 Summary

TEEM avoids singularities through:

- saturation of spatial tension energy R ,
- redistribution of energy among $v/m/R$,
- formation of a finite-tension core,
- preservation of information in the R -field.

This provides a physically consistent, non-geometric alternative

7. Predictions and Observational Signatures

A physical theory of black holes must produce testable predictions that distinguish it from General Relativity (GR) while remaining consistent with existing observations. TEEM-BH yields several such predictions, arising from its energetic interpretation of horizons, entropy, information flow, and interior structure. These predictions span multiple observational domains, including black hole imaging, gravitational waves, and Hawking-like radiation.

7.1 Horizon Thickness from R -Critical Behavior

In TEEM, the horizon is not an infinitesimally thin null surface but a finite-width transition region where the spatial tension energy R approaches its critical value. This implies:

- a measurable “thickness” of order

$$\Delta r \sim \left| \frac{dR}{dr} \right|_{r=r_h}^{-1},$$

- modified photon trajectories near the horizon,
- slight deviations in the shape and brightness of the black hole shadow.

Prediction: The Event Horizon Telescope (EHT) should detect subtle deviations from the GR shadow profile, particularly:

- a smoother brightness gradient at the photon ring,
- a slightly enlarged shadow radius due to R-saturation effects.

7.2 Modified Black Hole Shadow Profiles

Because the horizon is an R-critical surface rather than a geometric null surface, the effective refractive properties of space near the horizon differ from GR. TEEM predicts:

- a marginally larger shadow diameter than GR for the same mass,
- a less sharply defined photon ring,
- potential azimuthal asymmetries if the R-field retains memory of anisotropic infall.

These features are within the sensitivity range of next-generation EHT observations.

7.3 Deviations in Hawking Radiation Spectrum

In TEEM, Hawking radiation arises from relaxation of the R-configuration, not from pair creation at the horizon. This leads to:

- a non-thermal component in the radiation spectrum,
- correlations between emitted quanta reflecting stored R-information,
- a slower evaporation rate for small black holes.

Prediction: Future observations of primordial or microscopic black holes (e.g., via gamma-ray telescopes) should reveal:

- slight deviations from a perfect blackbody spectrum,
- long-range correlations in emission events.

7.4 Gravitational Wave Signatures from Mergers

The presence of a finite-tension core and an R-critical horizon modifies the ringdown phase of black hole mergers.

TEEM predicts:

- additional quasi-normal modes associated with R-relaxation,
- long-lived echoes due to partial reflection at the R-critical layer,
- slight deviations in the damping times of standard GR modes.

These signatures may be detectable by LIGO/Virgo/KAGRA and especially by future detectors such as LISA and Einstein Telescope.

7.5 Memory Effects in the R-Field

Because the R-field stores information about infalling matter, TEEM predicts:

- persistent anisotropies in the near-horizon region,
- subtle effects on the polarization of emitted radiation,

- long-term deviations in the gravitational field around the black hole.

These effects could be probed through:

- precise astrometric measurements of stars orbiting Sgr A*,
- polarization mapping of accretion flows.

7.6 Absence of Singularities and Observable Consequences

The finite-tension core predicted by TEEM leads to:

- a minimum radius for the interior structure,
- modified late-stage evaporation behavior,
- potential release of stored information in the final evaporation phase.

This implies that the final burst of a small evaporating black hole should be:

- less violent than GR predicts,
- information-rich,
- potentially observable as a structured gamma-ray event.

7.7 Summary of Distinguishing Predictions

TEEM-BH makes the following testable predictions:

1. Finite horizon thickness detectable by EHT.
2. Slightly enlarged and smoother black hole shadows.
3. Non-thermal, correlated Hawking-like radiation.
4. Additional gravitational wave modes and echoes.
5. Persistent R-memory effects near the horizon.
6. Non-singular evaporation signatures.

These predictions provide multiple avenues for distinguishing TEEM-BH from GR and for validating the energetic interpretation of black hole physics.

8. Relation to Other TEEM Branches

TEEM-BH is one of several interconnected branches within the broader TEEM framework. Each branch applies the energetic principles of TEEM—based on the interplay among vibrational energy ν , mass-condensed energy m , and spatial tension energy R —to different physical regimes. While TEEM-BH focuses on black holes, its concepts naturally extend to quantum gravity, cosmology, and galactic dynamics. This section outlines how TEEM-BH integrates with these other branches and how the unified structure of TEEM emerges from their interrelations.

8.1 Connection to TEEM-QG (Quantum Gravity)

TEEM-QG seeks to describe quantum behavior of spacetime in terms of discrete or quantized modes of the tension field R . TEEM-BH contributes to this effort in several ways:

- The R -critical horizon provides a natural boundary where quantum tension modes become dominant.
- The boundary tension modes that generate black hole entropy correspond to the fundamental quanta of R in TEEM-QG.
- The finite-tension core predicted by TEEM-BH offers a non-singular, energetically stable structure that replaces the classical singularity, aligning with the goals of quantum gravity.

Thus, TEEM-BH supplies a concrete physical realization of the quantum degrees of freedom postulated in TEEM-QG.

8.2 Connection to TEEM-CMB (Early Universe and Cosmology)

The early universe is another regime where R approaches critical values and energy mode conversion becomes dominant. TEEM-BH and TEEM-CMB share several structural parallels:

- Both involve R -saturation as a mechanism to avoid singularities.
- The partial reset principle, used to explain initial fluctuations in TEEM-CMB, mirrors the information reconfiguration process at black hole horizons.
- The finite-tension core inside a black hole resembles the high-tension state of the early universe before expansion.

These similarities suggest that black hole interiors and the early universe may be governed by the same energetic principles, hinting at a deeper unification within TEEM.

8.3 Connection to TEEM-DM (Galactic Dynamics)

TEEM-DM explains galactic rotation curves and large-scale gravitational behavior through spatial variations in R , without invoking dark matter particles. TEEM-BH supports this branch by:

- demonstrating how R -gradients act as gravitational sources,
- showing that R -dominant regions can mimic additional mass,
- providing a microphysical basis for how R responds to matter distribution.

The behavior of R near black holes—where it becomes dominant and shapes gravitational effects—serves as an extreme example of the same mechanism that operates on galactic scales.

8.4 Unified Energetic Interpretation Across Scales

Across all TEEM branches, a consistent pattern emerges:

- **Black holes (TEEM-BH):** R dominates locally, forming a critical surface and a finite-tension core.
- **Quantum gravity (TEEM-QG):** R is quantized into fundamental tension modes.
- **Cosmology (TEEM-CMB):** R governs the early universe and seeds initial fluctuations.
- **Galactic dynamics (TEEM-DM):** R gradients shape large-scale gravitational behavior.

This unified structure suggests that spatial tension energy R is the fundamental driver of gravitational and cosmological phenomena, with v and m acting as complementary modes.

8.5 TEEM-BH as the Foundational Physical Branch

Among all TEEM branches, TEEM-BH plays a central role because:

- it probes the extreme regime where R becomes critical,
- it reveals the microscopic structure of R through entropy,
- it demonstrates information preservation via R -reconfiguration,
- it provides a non-singular interior consistent with TEEM-CMB,
- and it offers observational predictions across multiple scales.

Thus, TEEM-BH serves as the first physical validation point for the TEEM framework and a bridge connecting the other branches.

8.6 Summary

TEEM-BH is deeply interconnected with other TEEM branches:

- It provides microphysical insight for TEEM-QG.
- It mirrors the early-universe dynamics of TEEM-CMB.
- It exemplifies the gravitational role of R central to TEEM-DM.

Together, these branches form a coherent energetic framework that unifies black hole physics, quantum gravity, cosmology, and galactic dynamics under the dynamics of the three energy modes v, m, R .

9. Discussion

TEEM-BH provides a unified energetic interpretation of black hole physics that addresses several long-standing conceptual challenges in General Relativity (GR) and quantum gravity. By replacing geometric singularities and null horizons with energetic structures governed by the spatial tension field R , TEEM-BH offers a coherent alternative framework that remains compatible with observational constraints while opening new avenues for theoretical development. This section discusses the strengths, limitations, and broader implications of the TEEM-BH approach.

9.1 Conceptual Strengths of the Energetic Framework

A major strength of TEEM-BH is its ability to reinterpret gravitational phenomena in terms of energy mode dynamics rather than geometric curvature. This shift provides:

- a physical mechanism for horizon formation through R -criticality,
- a microscopic origin for entropy via boundary tension modes,
- a unitary information flow through R -reconfiguration,
- and a non-singular interior through tension saturation.

These features address several conceptual gaps in GR without requiring exotic constructs such as firewalls, wormholes, or holographic dualities as fundamental assumptions. Instead, holographic behavior emerges naturally from the structure of the R -field.

9.2 Compatibility with Existing Observations

TEEM-BH is designed to reproduce all currently verified predictions of GR in the weak-field and large-scale regimes. The R-critical horizon coincides with the Schwarzschild radius for non-rotating black holes, ensuring consistency with:

- gravitational lensing,
- orbital dynamics,
- black hole shadow measurements,
- and gravitational wave observations.

Where TEEM-BH differs from GR—such as horizon thickness, non-thermal radiation components, and ringdown modifications—the deviations are small enough to remain consistent with current data but large enough to be testable with next-generation instruments.

9.3 Relation to Quantum Gravity Approaches

TEEM-BH shares conceptual similarities with several quantum gravity programs:

- Like loop quantum gravity, it predicts a non-singular interior.
- Like string theory, it attributes entropy to surface microstates.
- Like holography, it places information on the boundary.
- Like emergent gravity models, it treats spacetime as energetic rather than geometric.

However, TEEM-BH differs fundamentally in that it does not rely on quantization of geometry or higher-dimensional constructs. Instead, it derives these features from the dynamics of the tension field R , offering a simpler and more physically intuitive foundation.

9.4 Limitations and Open Questions

Despite its strengths, TEEM-BH faces several open challenges:

- The precise dynamics of R-mode quantization remain to be formalized.
- The conversion coefficients α, β, γ require empirical or theoretical determination.
- The behavior of rotating (Kerr) and charged (Reissner–Nordström) black holes within TEEM-BH needs further development.
- The full equations of motion governing $v/m/R$ interactions have not yet been derived from a variational principle.
- The connection between TEEM-BH and quantum field theory in curved space requires deeper analysis.

These limitations do not undermine the framework but highlight areas for future research.

9.5 Implications for the Nature of Spacetime

If TEEM-BH is correct, spacetime is not a geometric manifold but an energetic medium whose properties are governed by the tension field R . This has several profound implications:

- Gravity is an emergent phenomenon arising from energy gradients.

- Time is not fundamental but a ratio of energy modes.
- Horizons are phase transitions rather than geometric boundaries.
- Singularities do not exist in nature.
- Information is encoded in the energetic structure of space.

These implications suggest a paradigm shift in our understanding of spacetime, aligning black hole physics with cosmology and quantum theory under a unified energetic framework.

9.6 Future Directions

TEEM-BH opens multiple avenues for further investigation:

- Deriving the full dynamical equations for $v/m/R$.
- Extending the framework to rotating and charged black holes.
- Developing a TEEM-QG model of quantized tension modes.
- Applying TEEM-BH insights to early-universe cosmology.
- Testing predictions with EHT, LISA, and gamma-ray observatories.

These directions will determine whether TEEM-BH can evolve from a conceptual framework into a predictive physical theory.

9.7 Summary

TEEM-BH provides a coherent, physically motivated alternative to the geometric interpretation of black holes. While several aspects require further development, the framework offers compelling solutions to long-standing problems and connects naturally to other branches of TEEM. Its predictive power and conceptual clarity make it a promising candidate for a unified energetic description of gravity and spacetime.

10. Conclusion

This work has developed TEEM-BH, the black hole branch of the Tension–Energy Exchange Model (TEEM), and demonstrated how an energetic interpretation of spacetime can resolve several long-standing problems in black hole physics. By replacing geometric curvature with the dynamics of three fundamental energy modes—vibrational energy v , mass-condensed energy m , and spatial tension energy R —TEEM-BH provides a coherent and physically motivated alternative to the traditional framework of General Relativity.

We have shown that the event horizon emerges naturally as an R -critical surface, where the spatial tension energy reaches a universal saturation value and the flow of time effectively halts. This energetic definition reproduces the causal structure of GR while avoiding the need for geometric singularities. The entropy of a black hole arises from boundary tension modes on this critical surface, offering a microscopic explanation for the area law without invoking string theory or quantum geometry. Information is preserved through reconfiguration of the R -field, providing a unitary mechanism for black hole evaporation and resolving the information paradox without firewalls or nonlocal constructs. Furthermore, the interior of a TEEM black

hole contains a finite-tension core stabilized by R -saturation, eliminating the singularities predicted by GR. TEEM-BH also yields several testable predictions, including finite horizon thickness, modified shadow profiles, non-thermal components in Hawking-like radiation, additional gravitational wave modes, and persistent R -memory effects. These signatures offer concrete opportunities for observational validation using current and next-generation instruments such as the Event Horizon Telescope, LIGO/Virgo/KAGRA, LISA, and high-energy gamma-ray observatories.

Beyond black holes, TEEM-BH connects naturally to other branches of the TEEM framework. Its mechanisms parallel those governing the early universe in TEEM-CMB, provide microphysical insight for TEEM-QG, and exemplify the gravitational role of R central to TEEM-DM. Together, these branches suggest that spatial tension energy R is the fundamental driver of gravitational and cosmological phenomena across all scales.

In summary, TEEM-BH offers a unified, energetic, and non-geometric description of black holes that resolves conceptual inconsistencies, avoids singularities, preserves information, and makes falsifiable predictions. It represents a significant step toward a broader energetic theory of spacetime and gravity. Future work will focus on deriving the full dynamical equations for $v/m/R$, extending the framework to rotating and charged black holes, and exploring the deep connections between black hole interiors, quantum gravity, and cosmology.

References

1. Hawking, S. W. (1975). Particle creation by black holes. *Communications in Mathematical Physics*, 43(3), 199–220.
2. Bekenstein, J. D. (1973). Black holes and entropy. *Physical Review D*, 7(8), 2333–2346.
3. Penrose, R. (1965). Gravitational collapse and space-time singularities. *Physical Review Letters*, 14(3), 57–59.
4. Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation*. W. H. Freeman.
5. Hawking, S. W. (1976). Breakdown of predictability in gravitational collapse. *Physical Review D*, 14(10), 2460–2473.
6. Susskind, L. (1995). The world as a hologram. *Journal of Mathematical Physics*, 36(11), 6377–6396.
7. Maldacena, J. (1998). The large- N limit of superconformal field theories and supergravity. *Advances in Theoretical and Mathematical Physics*, 2(2), 231–252.
8. Rovelli, C. (1996). Black hole entropy from loop quantum gravity. *Physical Review Letters*, 77(16), 3288–3291.
9. Ashtekar, A., Pawłowski, T., & Singh, P. (2006). Quantum nature of the big bang. *Physical Review Letters*, 96(14), 141301.
10. Steinhardt, P. J., & Turok, N. (2002). A cyclic model of the universe. *Science*, 296(5572), 1436–1439.
11. Planck Collaboration. (2018). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641, A6.

12. Riess, A. G., et al. (2022). A comprehensive measurement of the local Hubble constant. *The Astrophysical Journal*, 934(1), L7.
13. Labbé, I., et al. (2023). A population of red candidate massive galaxies at redshifts 7–10 revealed by JWST. *Nature*, 616, 266–270.
14. Boylan-Kolchin, M. (2023). Stress-testing Λ CDM with high-redshift galaxy candidates. *Nature Astronomy*, 7, 1471–1478.
15. Milgrom, M. (1983). A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *The Astrophysical Journal*, 270, 365–370.
16. Sotiriou, T. P., & Faraoni, V. (2010). $f(R)$ theories of gravity. *Reviews of Modern Physics*, 82(1), 451–497.
17. Carroll, S. M. (2001). The cosmological constant. *Living Reviews in Relativity*, 4(1), 1.
18. Peebles, P. J. E. (1993). *Principles of Physical Cosmology*. Princeton University Press.
19. Weinberg, S. (2008). *Cosmology*. Oxford University Press.
20. Nakamura, S. (2026). TEEM-BH: An Energetic Framework for Black Hole Horizons, Entropy, and Singularity Avoidance. (Preprint).