

# Function Exercises

Rafi Ahmed Saad

December 2, 2023

## 1 Functions

### 1.1 Find the domain and range of $f(x) = \frac{x^2-1}{x-1}$


Domain is all possible inputs for which the function is defined. In this case, we need to avoid values of  $x$  that makes the denominator equal to zero, as division by zero is undefined. Thus the domain is all real numbers except  $x = 1$ , because that would make the denominator  $x - 1 = 0$

Domain:  $x \in \mathbb{R} \setminus \{1\}$

Let function

$$\begin{aligned} f(x) &= \frac{(x^2 - 1)}{x - 1} \\ \Rightarrow f(x) &= \frac{(x - 1)(x + 1)}{x - 1} \\ \Rightarrow f(x) &= x + 1 \end{aligned}$$

$\therefore$  Range:  $f(x) \in \mathbb{R}$



## 1.2 Find the domain and range of

$$f(x) = \begin{cases} 2x + 6 & \text{if } -3 \leq x \leq 0 \\ 6 & \text{if } 0 \leq x \leq 2 \\ 2x - 6 & \text{if } 2 \leq x \leq 5 \end{cases}$$

### Domain:

The domain of  $f(x)$  is set to all possible input values for which the function is defined. In this case, the function is defined in the intervals  $[-3, 0]$ ,  $[0, 2]$ ,  $[2, 5]$ , and for any other values, it is not explicitly defined. So the domain is:

$$\begin{aligned} \text{Domain: } x &\in [-3, 0] \cup [0, 2] \cup [2, 5] \\ x &\in [-3, 5] \end{aligned}$$

### Range:


For  $-3 \leq x \leq 0$ ,  $f(x) = 2x + 6$  where the output varies from 0 to 6

For  $0 \leq x \leq 2$ ,  $f(x) = 6$

For  $2 \leq x \leq 5$ ,  $f(x) = 2x - 6$  where the output varies from -2 to 4

So the Range is:

$$\text{Range: } f(x) \in [0, 6] \cup [-2, 4] \rightarrow f(x) \in [-2, 6]$$



### 1.3 Find the domain and range of

$$f(x) = \begin{cases} x - 1 & \text{if } x > 0 \\ -\frac{1}{2} & \text{if } x = 0 \\ x + 1 & \text{if } x < 0 \end{cases}$$

#### Domain:

Domain of this function  $f(x)$  can take any input.

$$\therefore \text{Domain: } x \in \mathbb{R}$$

$$\text{Domain: } x \in (-\infty, \infty)$$

#### Range:


For  $x > 0$ ,  $f(x) = x - 1$  where the result of the function is  $[0, \infty]$

For  $x = 0$ ,  $f(x) = -\frac{1}{2}$  which is a constant value

For  $x < 0$ ,  $f(x) = x + 1$  where the result of the function is  $[0, -\infty]$

$$\therefore \text{Range: } x \in [0, -\infty] \cup \left[\frac{1}{2}\right] \cup [0, \infty]$$

$$\text{Range: } x \in (-\infty, \infty)$$



**1.4 Find the domain and range of**

$$f(x) = \begin{cases} -2x + 1 & \text{if } x < 0 \\ 1 & \text{if } 0 \leq x < 1 \\ 2x - 1 & \text{if } x \geq 1 \end{cases}$$

**Domain:**

Domain of this function  $f(x)$  can take any input.

$$\therefore \text{Domain: } x \in \mathbb{R}$$

$$\text{Domain: } x \in (-\infty, \infty)$$

**Range:**


For  $x < 0$ ,  $f(x) = -2x + 1$  where the result of the function is  $(-\infty, 1]$

For  $0 \leq x < 1$ ,  $f(x) = 1$  which is a constant value

For  $x \geq 1$ ,  $f(x) = 2x - 1$  where the result of the function is  $[1, \infty)$

$$\therefore \text{Range: } x \in (-\infty, 1] \cup [1] \cup [1, \infty)$$

$$\text{Range: } x \in (-\infty, \infty)$$



### 1.5 Find the domain and range of

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < \frac{1}{2} \\ 1 & \text{if } x = \frac{1}{2} \\ 1 - x & \text{if } \frac{1}{2} < x \leq 1 \end{cases}$$

#### Domain:

The domain of the function  $f(x)$  is all possible input values for which the function is defined. In this case, the intervals are  $[0, \frac{1}{2})$ ,  $[\frac{1}{2}]$ ,  $(\frac{1}{2}, 1]$ . So the domain is:

$$\text{Domain: } x \in [0, \frac{1}{2}) \cup [\frac{1}{2}] \cup (\frac{1}{2}, 1]$$

$$\text{Domain: } x \in [0, 1]$$

#### Range:

For  $0 \leq x < \frac{1}{2}$ ,  $f(x) = x$  where the result of the function is  $[0, \frac{1}{2})$

For  $x = \frac{1}{2}$ ,  $f(x) = 1$  which is a constant value. So the range is  $[1]$

For  $\frac{1}{2} < x \leq 1$ ,  $f(x) = 1 - x$  where the result of the function is  $(\frac{1}{2}, 1]$

$$\therefore \text{Range: } x \in [0, \frac{1}{2}) \cup \{1\} \cup (\frac{1}{2}, 1]$$

$$\text{Range: } x \in [0, \frac{1}{2}) \cup \{1\}$$

