

Probability Theory Lecture 01

Padmini Mukkamala

Budapest University of Technology and Economics

Overview

- 1 What is probability?
- 2 Definition of Probability
- 3 Some tools for finding Probabilities
 - Inclusion-Exclusion
 - Calculus and Combinatorics

Birthday Paradox

Will you bet?

Consider the following bet: if two people in this class have their birthday on the same day, then every student has to give me 1 dollar, while if no two students have a birthday on the same day, then I will give the class 1000 dollars.

Birthday Paradox

Will you bet?

Consider the following bet: if two people in this class have their birthday on the same day, then every student has to give me 1 dollar, while if no two students have a birthday on the same day, then I will give the class 1000 dollars.

There are 365 days in a year and 53 students in this class. Do you think its a fair bet? Who do you think will win? How high is the probability that I will win?

Three probability problems

- Suppose we roll a fair dice once. What is the probability of rolling 1?

Three probability problems

- Suppose we roll a fair dice once. What is the probability of rolling 1?
- At a grocery store, we observe the weight of potatoes bought by customers. It is some number between, say, 0 and 2 kilograms. What is the probability that the next customer bought 1 kg? less than 1 kg?

Three probability problems

- Suppose we roll a fair dice once. What is the probability of rolling 1?
- At a grocery store, we observe the weight of potatoes bought by customers. It is some number between, say, 0 and 2 kilograms. What is the probability that the next customer bought 1 kg? less than 1 kg?
- Suppose we pick a point at random from the square $[-1, 1] \times [-1, 1]$. What is the probability it will lie in a unit circle centered at the origin?

Basic definitions

Experiment

Eg. Rolling of a dice, tossing a coin, buying potatoes.

Outcome

A possible result of the experiment.

Sample Space Ω (Omega)

The set of all possible outcomes.

Event

A subset of the sample space $A \subset \Omega$.

Examples

Experiment

Example 1: single dice roll, Example 2: weight of potatoes,
Example 3: randomly picking a point.

Outcome

Six possible outcomes: 1, 2, 3, 4, 5, 6, Infinitely many outcomes:
any real number in the interval $[0, 2]$, Infinitely many outcomes:
any point in the square.

Sample space Ω

$$\{1, 2, 3, 4, 5, 6\}, [0, 2], [-1, 1] \times [-1, 1]$$

Examples

Events

Even dice roll $A = \{2, 4, 6\}$

Odd dice roll $B = \{1, 3, 5\}$

Prime dice roll $C = \{2, 3, 5\}$

Dice roll of 1 $D = \{1\}$

Dice roll of a number bigger than 10 $E = \phi$ (Empty event).

Weight of potatoes is exactly 1 kg $F = \{1\}$

Weight of potatoes is less than 1 kg $G = [0, 1]$

Set operations, Mutually exclusive events

The sample space, in mathematical terms, is a set, and events are subsets of the sample space. So naturally, set operations Union, Intersection, Complement etc can be done to events.

Set operations, Mutually exclusive events

The sample space, in mathematical terms, is a set, and events are subsets of the sample space. So naturally, set operations Union, Intersection, Complement etc can be done to events.

Mutually exclusive events

Two events are said to be mutually exclusive if their intersection is ϕ , the empty event. Which of A, B, C, D are mutually exclusive?

Set operations, Mutually exclusive events

The sample space, in mathematical terms, is a set, and events are subsets of the sample space. So naturally, set operations Union, Intersection, Complement etc can be done to events.

Mutually exclusive events

Two events are said to be mutually exclusive if their intersection is ϕ , the empty event. Which of A, B, C, D are mutually exclusive?

De Morgan's laws

For any two events A and B , $\overline{A \cup B} = \overline{A} \cap \overline{B}$. etc.

Intuitive definition of probability

It is the fraction of favorable outcomes out of all possible outcomes of an experiment. (Also called classical probability)

Three probability problems

- Suppose we roll a fair dice once. What is the probability of rolling 1?

Three probability problems

- Suppose we roll a fair dice once. What is the probability of rolling 1? $\frac{1}{6}$
- If I roll the dice 6 times, will the outcome be 1 exactly once?

Relative Frequency

Relative Frequency

The relative frequency is the fraction of times we get a favorable result out of the total number of times an experiment is performed.

In the case of the dice roll outcome being 1, its relative frequency is:

$$\frac{\text{number of 1 outcomes}}{\text{number of dice rolls}}$$

Relative Frequency

Relative Frequency

The relative frequency is the fraction of times we get a favorable result out of the total number of times an experiment is performed.

In the case of the dice roll outcome being 1, its relative frequency is:

$$\frac{\text{number of 1 outcomes}}{\text{number of dice rolls}}$$

This number converges to our intuitive definition of probability for a large number of experiments. (Central Limit Theorem, Lecture 10)

How long have human beings been gambling?

How long have human beings been gambling?

Sheep bones and other materials have been used as dice for as far back as 4000 BC. Probably even earlier.

When did the above intuitive definition of Probability get formulated?

How long have human beings been gambling?

Sheep bones and other materials have been used as dice for as far back as 4000 BC. Probably even earlier.

When did the above intuitive definition of Probability get formulated?

Intuitive and easy as it may seem, this definition along with the first systematic treatment of probability was published (posthumously) by Cardano in 1663.

Compare this with Pythagorus theorem whose oldest axiomatic proof dates from Euclid's elements from 300 BC!

Two probability problems: revisited

The intuitive definition of Probability is not useful when the Sample space Ω is infinite.

- At a grocery store, we observe the weight of potatoes bought by customers. It is some number between, say, 0 and 2 kilograms. Assume that we can measure any weight between 0 and 2 (any real number) and that based on previous data, all weights are equally likely.
- What is the probability that the next customer bought less than 1 kg?

Two probability problems: revisited

The intuitive definition of Probability is not useful when the Sample space Ω is infinite.

- At a grocery store, we observe the weight of potatoes bought by customers. It is some number between, say, 0 and 2 kilograms. Assume that we can measure any weight between 0 and 2 (any real number) and that based on previous data, all weights are equally likely.
- What is the probability that the next customer bought less than 1 kg? $\frac{1}{2}$
- Suppose we pick a point at random from the square $[-1, 1] \times [-1, 1]$. What is the probability it will lie in a unit circle centered at the origin?

Two probability problems: revisited

The intuitive definition of Probability is not useful when the Sample space Ω is infinite.

- At a grocery store, we observe the weight of potatoes bought by customers. It is some number between, say, 0 and 2 kilograms. Assume that we can measure any weight between 0 and 2 (any real number) and that based on previous data, all weights are equally likely.
- What is the probability that the next customer bought less than 1 kg? $\frac{1}{2}$
- Suppose we pick a point at random from the square $[-1, 1] \times [-1, 1]$. What is the probability it will lie in a unit circle centered at the origin? $\frac{\pi}{4}$

Definition of Probability

Probability is a measure (a function), $P : \mathcal{F} \mapsto [0, 1]$, on a 'proper' collection of subsets (events), \mathcal{F} , over the sample space Ω . The following are the Axioms of Probability.

- $P(\Omega) = 1$
- (Sigma additivity) For a collection $A_1, A_2, A_3, \dots, A_i, \dots$ of mutually exclusive events, that is, $A_i \cap A_j = \phi, \forall i \neq j$,

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$$

Definition of Probability

- What is $P(\phi) = ?$

Definition of Probability

- What is $P(\phi) = ?$
- What is $P(A) + P(\bar{A}) = ?$

Definition of Probability

- What is $P(\phi) = ?$
- What is $P(A) + P(\bar{A}) = ?$
- If $A \subseteq B$, what can you say about $P(A)$ and $P(B)$?

Definition of Probability

- What is $P(\phi) = ?$
- What is $P(A) + P(\bar{A}) = ?$
- If $A \subseteq B$, what can you say about $P(A)$ and $P(B)$?

Definition of Probability

- What is $P(\phi) = ?$
- What is $P(A) + P(\bar{A}) = ?$
- If $A \subseteq B$, what can you say about $P(A)$ and $P(B)$?
 $P(A) \leq P(B)$. Proof?
- What is $P(A \cap B) + P(A \cap \bar{B})$?

Definition of Probability

- What is $P(\phi) = ?$
- What is $P(A) + P(\bar{A}) = ?$
- If $A \subseteq B$, what can you say about $P(A)$ and $P(B)$?
 $P(A) \leq P(B)$. Proof?
- What is $P(A \cap B) + P(A \cap \bar{B})$?

Definition of Probability

- What is $P(\phi) = ?$
- What is $P(A) + P(\bar{A}) = ?$
- If $A \subseteq B$, what can you say about $P(A)$ and $P(B)$?
 $P(A) \leq P(B)$. Proof?
- What is $P(A \cap B) + P(A \cap \bar{B}) = P(A)$. Proof?

Examples of different measures

Fair coin - Uniform measure

$P(H) = P(T) = \frac{1}{2}$, here our previous intuition of probability, that it is number of favorable outcomes divided by total number of outcomes, holds.

Examples of different measures

Fair coin - Uniform measure

$P(H) = P(T) = \frac{1}{2}$, here our previous intuition of probability, that it is number of favorable outcomes divided by total number of outcomes, holds.

Unfair coin - Non-uniform measure

$P(H) = \frac{1}{3}, P(T) = \frac{2}{3}$. Note, here also $P(\Omega) = 1$ and sigma additivity holds. But now, probability of an outcome is not number of favorable outcomes divided by total number of outcomes.

Principle of Inclusion-Exclusion (Poincare's formula)

Case of two events

Given two events A_1, A_2 and a probability measure P , the following always holds: $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2)$.

Proof: Exercise, use Venn diagram and sigma additivity.

Principle of Inclusion-Exclusion (Poincare's formula)

Case of two events

Given two events A_1, A_2 and a probability measure P , the following always holds: $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2)$.

Proof: Exercise, use Venn diagram and sigma additivity.

Case of three events

Given three events A_1, A_2, A_3 and a probability measure P , the following always holds: $P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3)$.

Proof: Exercise, use Venn diagram and sigma additivity.

Example of Inclusion-Exclusion

Let $P(A) = \frac{1}{2}$, $P(B) = \frac{1}{2}$, $P(A \cap B) = \frac{1}{3}$. What is $P(A \cup B)$?

Example of Inclusion-Exclusion

Let $P(A) = \frac{1}{2}$, $P(B) = \frac{1}{2}$, $P(A \cap B) = \frac{1}{3}$. What is $P(A \cup B)$?
So, $P(A \cup B) = \frac{1}{2} + \frac{1}{2} - \frac{1}{3} = \frac{2}{3}$

Union bound or Boole's inequality

Union bound

Given n events A_1, \dots, A_n and a probability measure P , the following holds:

$$P(\cup_{i=1}^n A_i) \leq \sum_{i=1}^n P(A_i)$$

Union bound or Boole's inequality

Union bound

Given n events A_1, \dots, A_n and a probability measure P , the following holds:

$$P(\cup_{i=1}^n A_i) \leq \sum_{i=1}^n P(A_i)$$

Proof: Use new events B_1, \dots, B_n , where $B_1 = A_1$ and $B_i = A_i \setminus (\cup_{j=1}^{i-1} A_j)$, $\forall i > 1$.

Limit Properties

Property 1

Given events $A_1 \subseteq A_2 \subseteq \dots \subseteq A_n \subseteq \dots$, and a probability measure P , the following holds:

$$P(\cup_i A_i) = \lim_{n \rightarrow \infty} P(A_n)$$

Proof

Describe new events $B_1 = A_1$ and $B_i = A_i \setminus A_{i-1}$. Then use sigma additivity for B_i 's that $P(\cup_i B_i) = \sum_i P(B_i)$.

Limit Properties

Property 2

Given events $A_1 \supseteq A_2 \supseteq \dots \supseteq A_n \supseteq \dots$, and a probability measure P , the following holds:

$$P(\cap_i A_i) = \lim_{n \rightarrow \infty} P(A_i)$$

Geometric probability

Example 1

Given a point in the plane with both coordinates in the interval $[0, 1]$, what is the probability that the point is below the curve $y = x^2$?

Example 2

Given a point in a square of side 1, what is the probability that its distance to the center of the square is less than $\frac{1}{2}$?

Combinatorics

Picking k out of n items.

Experiment	With replacement	Without replacement
Ordered	n^k	$n(n-1)\dots(n-k+1)$
Unordered	$\binom{n+k-1}{k}$	$\binom{n}{k}$

Table: Some basic ways of counting

Combinatorics

Examples

What is the probability of drawing two Aces if two cards are drawn from a standard deck with replacement?

Combinatorics

Examples

What is the probability of drawing two Aces if two cards are drawn from a standard deck with replacement?

Without replacement?

Combinatorics

Examples

What is the probability of drawing two Aces if two cards are drawn from a standard deck with replacement?

Without replacement?

What is the probability that the top three scores on the final exams are of three girls, if 25 percent of the students in a class of 100 are girls?

Combinatorics

Examples

What is the probability of drawing two Aces if two cards are drawn from a standard deck with replacement?

Without replacement?

What is the probability that the top three scores on the final exams are of three girls, if 25 percent of the students in a class of 100 are girls?

You want to randomly distribute 6 identical candies among 3 children. What is the probability the first child will get all 6 candies?

Summary

- Classical probability: fraction of favorable outcomes out of all possible outcomes of an experiment.
- (Sigma additivity) For mutually exclusive A_i 's,

$$P(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i).$$
- Inclusion-Exclusion:

$$P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3).$$
- Union bound: For any A_i 's, $P(\bigcup_{i=1}^n A_i) \leq \sum_{i=1}^n P(A_i)$
- Limit Property: For increasing events A_i ,

$$P(\bigcup_i A_i) = \lim_{n \rightarrow \infty} P(A_n)$$

The End