

On the Cut, a Radial Weight Beats a Flat Kernel; the CHM Parabola Is Not Selected

Robert W. McGwier
CTO, Cohere Technology Group
GitHub: [n4hy](#)

15 August 2026 — Version 1

Abstract

Paper 27 found that on the horizon, K_{CHM} beats a flat hop kernel. Paper 25 found that for the modular kernel itself, a linear $R - r$ weight can beat the CHM parabola on a small ball. This note asks the first-law form of that question, on a new grid, discarding any hop that flips n_{occ} .

$N = 12$, $R = 4$, 294 surface hops, zero occupancy flips. $R_{\text{CHM}} = 0.808 < R_{\text{flat}} = 0.994$ (C3 PASS). $R_{\text{CHM}} = 0.8082$ versus $R_{\text{lin}} = 0.8099$ (C4 is a tie). Tracking floors fail ($|\rho| = 0.589 < 0.60$).

An auditor on $N = 14$, $R = 3$ kept only 8 hops after 142 occupancy flips and is not a C4 confirmation.

The cut prefers a radially decreasing weight to a shapeless local kernel. It does not prefer $R^2 - r^2$ to $R - r$. Not $\eta = 1/4G$. Not Einstein. Not the Bloch guess.

1 Numbers

	n	R_{CHM}	R_{lin}	R_{flat}
$N = 12, R = 4$	294	0.808	0.810	0.994
audit $N = 14, R = 3$	8	0.959	0.959	0.982

C3 is the Paper 27 fact, now at a second radius with a stable Fermi sea. C4 is the new question and it does not select the quadratic.

Admission. *I asked whether the parabola was doing Clausius work beyond any decreasing radial weight. On this grid it is not. I did not call a 0.002 residual gap a theorem, and I did not write Einstein.*