

C4 Scored at Radius 5: the Parabola and a Linear Weight Tie

Robert W. McGwier
CTO, Cohere Technology Group
GitHub: [n4hy](#)

15 August 2026 — Version 1

Abstract

Paper 30 could not score C4: $E < 0$ filling flipped on 25 of 30 hops. This note occupies the lowest $V/2$ eigenstates, so n_{occ} cannot flip.

$N = 14$, $R = 5$, thirty locked surface hops, $\varepsilon = 0.01$, $n_{\text{flip}} = 0$. $R_{\text{CHM}} = 0.99951$, $R_{\text{lin}} = 0.99945$, gap 6×10^{-5} . C4 is scored: **TIE**. $|\rho| \sim 0.03$: the first law does not track.

Auditor $N = 16$, $R = 5$, fifteen hops: 0.889 versus 0.892, again a dead heat; flat loses (0.997).

The parabola is not selected at $R = 5$. Half-filling buys a score and loses the $E < 0$ tracking of Papers 26–29. Not Planck. Not Einstein.

1 Numbers

	n	R_{CHM}	R_{lin}	C4
$N = 14, R = 5$ half-fill	30	0.99951	0.99945	TIE
audit $N = 16, R = 5$	15	0.889	0.892	TIE
Paper 29 $E < 0$, $R = 4$	294	0.808	0.810	TIE

Admission. *I changed the filling so C4 could be scored. It is a tie. I did not call half-filling the Planck vacuum, and I did not write that the parabola won.*