

Add the Masses, Then Take One Gradient. Do Not Add the Fields.

Robert W. McGwier
CTO, Cohere Technology Group
GitHub: [n4hy](#)

16 August 2026 — Version 1

Abstract

The pair field of Papers 69–70 is $\mathbf{g} = -(M/A)\hat{n}$ with $\hat{n} = \nabla_c M / |\nabla_c M|$. That operation is not linear:

$$\mathbf{g}_{AB} \propto M_{AB} \frac{\nabla M_A + \nabla M_B}{|\nabla M_A + \nabla M_B|} \neq \mathbf{g}_A + \mathbf{g}_B$$

unless $\nabla M_A \parallel \nabla M_B$.

Exact open-hop basis, `mpmath` $dps = 80$, no eigensolve. Mass maps add to 2×10^{-7} . Median angle $\angle(\mathbf{g}_{AB}, \mathbf{g}_A + \mathbf{g}_B) = 5.43^\circ$. Relative vector residual 0.255: C_rel fails. Far from both packets the angle is 0.008° . Auditor $N = 11$: map and angle confirmed; residual 0.45 refuted.

To use two packets: $M = M_A + M_B$, then one $\hat{n} = \nabla_c M$. Not Einstein.

Admission. *I hoped the derived $\mathbf{g}$ would add. The residual gate failed. I did not move it.*