

Inherited Newton: Attractive $1/r^2$ Passes. Isolated $\Phi = -GM/r$ Does Not.

Robert W. McGwier
CTO, Cohere Technology Group
GitHub: `n4hy`

15 August 2026 — Version 1

Abstract

The NSM metric sector is Einstein–Cartan. For a spinless source that *is* Einstein–Hilbert. This note runs the locked M9.2 gates on a 3-d discrete Poisson problem, Dirichlet $\Phi = 0$ on a cube of half-width L .

$n = 97$. Probes at $0.30L$, $0.35L$, $0.40L$. $\mathbf{a} = -\nabla\Phi$ is attractive GM/r^2 to 1.3%, 1.0%, 1.1%. Log-log slope -2.008 . C1 PASS. Vacuum, 2-d mutation, and $G \rightarrow -G$ pass.

Φ is not $-GM/r$ (residuals 26–35%). A Dirichlet wall at L is not isolated space. C2 FAIL.

Auditor, Gaussian source: C1 CONFIRMED, C2 REFUTED.

Inherited, not discovered. Not FGHMV. Not de Sitter. Not a `MODELS.md` column.

Admission. *I ran the Newton lock. The force gate passed. I did not call a Dirichlet-box potential isolated Newton, and I did not write that entanglement made G .*